

# Generation of chiral solitons in antiferromagnetic chains by a quantum quench

Barbara Bravo,<sup>1</sup> Ariel Dobry,<sup>1</sup> Diego Mastrogiuseppe,<sup>2,3</sup> and Claudio Gazza<sup>1</sup>

<sup>1</sup>*Facultad de Ciencias Exactas Ingeniería and Agrimensura, Universidad Nacional de Rosario and Instituto de Física Rosario, Bv. 27 de Febrero 210 bis, 2000 Rosario, Argentina.*

<sup>2</sup>*Department of Physics and Astronomy, and Nanoscale and Quantum Phenomena Institute, Ohio University, Athens, Ohio 45701–2979*

<sup>3</sup>*Dahlem Center for Complex Quantum Systems and Fachbereich Physik, Freie Universität Berlin, 14195 Berlin, Germany*

(Dated: November 15, 2013)

We analyze the time evolution of a magnetic excitation in a spin- $\frac{1}{2}$  antiferromagnetic Heisenberg chain after a quantum quench. By a proper modulation of the magnetic exchange coupling, we prepare a static soliton of total spin  $\frac{1}{2}$  as an initial spin state. Using bosonization and a numerical time dependent density matrix renormalization group algorithm, we show that the initial excitation evolves to a state composed of two counter-propagating chiral states, which interfere to yield  $\langle S^z \rangle = \frac{1}{4}$  for each mode. We find that these dynamically generated states remain considerably stable as time evolution is carried out. We propose spin-Peierls materials and ultracold-atom systems as suitable experimental scenarios in which to conduct and observe this mechanism.

PACS numbers: 75.10.Jm, 75.10.Pq, 03.75.Lm

## I. INTRODUCTION

Thinking about classical nonlinear physics, solitons are peculiar solutions which can be characterized by constant velocity and shape. Recently, Wöllert and Honecker<sup>1</sup> pursuing the understanding of the extension of the soliton concept to the quantum regime, chose the easy-axis ferromagnetic  $XXZ$  model as the scenario in which to analyze how a localized quantum wave packet evolves in time. They have shown that besides the quantum mechanical delocalization due to the uncertainty principle, they are in qualitative agreement with its classical counterpart. Following this objective of deciphering the quantum soliton term, we tackle an alternative problem in which we study the time evolution of a one-dimensional topological quantum soliton after a quench.

The study of nonequilibrium phenomena in one-dimensional systems has become a very active area of research in recent years due to new advances in the experiments with ultracold atoms in optical lattices,<sup>2</sup> and the latest studies on thermalization after quantum quenches.<sup>3,4</sup> With these ideas in mind, we propose a frustrated  $J_1$ - $J_2$  spin- $\frac{1}{2}$  antiferromagnetic Heisenberg chain as a suitable framework in which to conduct the analysis. It is known that this model undergoes a phase transition from a quasi-long-range ordered ground state to a product of localized singlets clusters as a function of the next-nearest neighbor parameter. Moreover, this Hamiltonian has an exact ground state at the Majumdar-Ghosh (MG) point  $J_1 = 2J_2$ ,<sup>5</sup> and a variational approach describes the elementary excitations adequately.<sup>6</sup> More specifically, in this work we propose a fine-tuned magnetic soliton as the initial pattern which will be evolved in time after a quantum quench of the model parameters. In order to create this initial spin- $\frac{1}{2}$  excitation, we choose a one-dimensional chain with spin-phonon coupling as a witness case, which is realized in quasi-one dimensional spin-Peierls systems such as  $\text{CuGeO}_3$  (Ref. 7) and  $\text{TiOX}$  ( $X = \text{Cl, Br}$ ).<sup>8</sup> An alternative approach would be to prepare the initial excitation and conduct the time evolution in a setup of ultracold atoms in an optical lattice.

Once the initial soliton is prepared, we follow the dynamics of this excitation on the uniform zig-zag Heisenberg chain, driven by a quench of the spin-phonon coupling. We observe that, in the gapless phase of the model, the initial soliton evolves into two counterpropagating modes, indicating a quantum superposition of left- and right- moving components of the original soliton. As time evolves, the excitation remains quite stable despite the quantum-mechanical spreading. On the other hand, for a highly localized soliton generated in the MG point, the excitation shows quick spreading with time evolution.

We select the density matrix renormalization group (DMRG)<sup>9</sup> and bosonization<sup>10</sup> as the numerical and analytical techniques to conduct our study. There are plenty of examples in the literature showing that both methods are convenient for giving a reliable description of spin chain systems, particularly, when the coupling to the lattice is also considered.<sup>11</sup> Among different DMRG options,<sup>12</sup> we use the algorithm introduced by Manmana *et al.*,<sup>4</sup> that allows us to perform non equilibrium simulations for systems with interactions beyond the nearest neighbors. It is worth mentioning that related studies were done on spin transport, even at finite temperature.<sup>13</sup>

## II. THE MODEL AND STRATEGY TO GENERATE EXCITED STATES

In order to generate a soliton-like topological excitation, we introduce a one dimensional antiferromagnetic Heisenberg Hamiltonian with first and second neighbor interactions and spin-lattice coupling, which reads

$$H = \sum_i [1 + \delta_i] \mathbf{S}_i \cdot \mathbf{S}_{i+1} + \beta \mathbf{S}_{i-1} \cdot \mathbf{S}_{i+1}. \quad (1)$$

$\mathbf{S}_i$  is a spin-1/2 operator for the  $i$ th lattice site, and we have set  $J_1 = 1$  as the energy scale, such that  $\beta = J_2/J_1$  is the second neighbor exchange coupling;  $\delta_i = \lambda(u_{i+1} - u_i)/J_1$  is the dimensionless bond length variation, where  $u_i$  are the displacements of the magnetic ions from their equilibrium




FIG. 1. (Color online) Collective excitation obtained by DMRG in a finite lattice of  $N_s = 99$  sites, in the absence of frustration ( $\beta = 0$ ). The soliton, represented by  $\langle S_i^z \rangle$  (circles), was tuned by selecting  $\delta_i$  in a given configuration (squares). The parameters are  $\xi = 1$ ,  $\delta_0 = 0.3$ . We also show the cumulative magnetization up to a given site (diamonds). The bosonization result for this magnitude is represented by the solid line.

positions, and  $\lambda$  is the spin-lattice coupling parameter. Although frustration is not necessary to present our ideas, we introduce it in order to compare it with the well-known MG limit<sup>5</sup> to achieve a better understanding of the process. So, with the intention of promoting a sort of quantum topological soliton, we break the lattice symmetry to manipulate the nearest neighbor magnetic interaction. From the studies of models with spin-phonon coupling it is known that, when phonons are treated adiabatically, the stable pattern for  $\delta_i$  in the  $S^z = 1/2$  sector is the one called the static lattice soliton, given by  $\delta_i = (-1)^i \delta_0 \tanh(\frac{i-i_0}{\xi})$ . The domain wall is centered at site  $i_0$  and its width is given by  $\xi$ , which produces an interpolation between two dimerized patterns. The  $S^z = \frac{1}{2}$  magnetic soliton generated by this lattice arrangement is also centered at site  $i_0$ , and the number of sites involved within the wall is controlled by the parameters  $\xi$  and  $\delta_0$ . In Fig. 1 we show the lattice configuration generated with the set of parameters  $\{\beta = 0, \delta_0 = 0.3, \xi = 1\}$ , together with the associated spin pattern given by  $\langle S_i^z \rangle$ , which was obtained from Eq. (1) using DMRG in a lattice of size  $N_s = 99$ , where the odd number of sites is to set the total magnetization to  $\frac{1}{2}$ . We use open boundary conditions keeping  $m = 300$  states, enough to assure the accuracy, with a truncation error of order  $O(10^{-9})$  in the worst case. We see that most of the spins arrange in localized singlet clusters, producing zero local magnetization. However, in the center of the chain where the bonds interpolate between the two possible dimerized states,  $\langle S_i^z \rangle \neq 0$ . We also show the cumulative magnetization up to a given site  $I$ , which is defined as  $M_I = \sum_{i=1}^I \langle S_i^z \rangle$ . A clear solitonic profile can be observed in this quantity. Let us now resort to the bosonization technique to analyze the  $M_I$  parameter. In this representation, the  $z$ -component of the spin is connected to a bosonic field  $\phi(x)$  by<sup>10</sup>

$$S_i^z = \frac{1}{2\pi} \partial_x \phi(x = ia) + \frac{(-1)^i}{\pi\alpha} \cos[2\phi(x = ia)], \quad (2)$$

where  $a$  is the lattice constant and  $\alpha$  is a short range cutoff in the bosonization procedure. The quantity  $M_I$  is obtained by integration of the previous equation up to a point  $X$ . Being oscillatory at the lattice level, the last term vanishes. In

the  $S^z = \frac{1}{2}$  subspace the field goes from  $\phi(-\infty) = -\frac{\pi}{2}$  to  $\phi(\infty) = \frac{\pi}{2}$ . Therefore, the expression for  $M_I$  is

$$M_I = \frac{1}{2\pi} \langle \phi(X = Ia) \rangle + \frac{1}{4}. \quad (3)$$

As the bosonized version of the Hamiltonian (1) is not exactly solvable for general displacements  $\delta_i$ , we resort to a semiclassical solution in order to calculate the mean value of the field in the corresponding subspace and compare with the DMRG results. This solution can be obtained by adding an elastic energy term  $\frac{K}{2} \sum_i \delta_i^2$  to the Hamiltonian and treating  $\delta_i$  in the adiabatic approximation.<sup>14,15</sup> The classical solutions for the continuous fields are  $\delta(x) = \delta_0 \tanh(x/\xi)$  and  $\phi(x) = \arcsin[\tanh(x/\xi)]$ . With the inclusion of an elastic energy term,  $\delta_0$  and  $\xi$  depend on the values of the microscopic parameters  $K$  and  $J_{1,2}$ . In the previous DMRG calculation we chose  $\delta_i$  with arbitrary  $\delta_0$  and  $\xi$ , i.e. not subject to fulfilling an adiabatic equation. In the spirit of a semi-classical quantization<sup>16</sup> we assume the following ansatz for the mean value of the bosonic field:  $\langle \phi \rangle \equiv \phi_S(x) = \arcsin[\tanh(x/\xi')]$ , where the subindex  $S$  refers to a solitonic pattern. In Fig. 1 we show a fitting to the numerical results using the analytical expression, for which we obtain a good agreement with  $\xi' = 1.773$ .

### III. QUENCH AND TIME EVOLUTION OF THE SOLITON

We now turn off the spin-lattice coupling and study the dynamics of this magnetic excitation, which will be conducted by a homogeneous Heisenberg Hamiltonian  $H' \equiv H[\delta_i = 0]$ . Once the quantum soliton is constructed, the time-dependent DMRG (t-DMRG) algorithm enables us to evolve in time under this new homogeneous Hamiltonian.<sup>17</sup> As a gapless phase is stable for  $0 \leq \beta \leq \beta_c = 0.245$ , we analyze different tuned solitons in this zone where the bosonization analysis is valid. In Fig. 2, we show the dynamics of two witness cases which are defined by  $\{\beta = 0, \delta_0 = 0.3, \xi = 1\}$  and  $\{\beta = 0.24, \delta_0 = 0.3, \xi = 3\}$ . One can observe that the excitations behave in a subtle way. In both cases, the time evolution shows how the original soliton evolves in such a way that, at a given time, two spin clouds are observed, each of them carrying  $\langle S^z \rangle = \frac{1}{4}$ . Remarkably, these right and left modes are quite stable: as observed in Fig. 2, these modes barely disperse as time evolves. We will return to this interpretation later.

These left and right excitations travel at a velocity which agrees very well with the spin wave velocity of the low energy excitations of the homogeneous chain. This can be seen in Fig. 2 by comparing the evolution of the maximum of  $\langle S_i^z \rangle$  with the spin wave velocity  $v_s = \frac{\pi}{2}$  predicted by the Bethe ansatz for the one dimensional homogeneous Heisenberg chain,<sup>10</sup> and the renormalized  $v_s = \frac{\pi}{2}(1 - 1.12\beta)$  calculated in Ref. 18 using exact diagonalization in small chains.

Let us analyze the origin of this behavior by studying the time evolution in the bosonization language. The Hamiltonian  $H'$  in the bosonic representation corresponds to a (1+1) free bosonic field theory, up to marginally irrelevant operators.<sup>10</sup> These marginal operators vanish at  $\beta = \beta_c$ .




FIG. 2. (Color online) Time evolution of  $\langle S_i^z \rangle$  in the collective excitation obtained by DMRG with and without frustration for the two witness cases, (a)  $\beta = 0$ ,  $\delta_0 = 0.3$ ,  $\xi = 1$  and  $\beta = 0.24$ ,  $\delta_0 = 0.3$ ,  $\xi = 3$ . In (a), the black line corresponds to the slope in the time-space diagram of the spin velocity  $v_s = \frac{\pi}{2}$  obtained by the Bethe ansatz for a homogeneous Heisenberg chain. In (b), the black line shows the spin velocity renormalized by the frustration. In both cases one observes the splitting of the soliton into two chiral modes.

The time evolution of the mean value of  $\phi$  is given by  $\langle \phi(x, t) \rangle = \langle e^{-iH't} \phi_R(x) e^{iH't} \rangle + \langle e^{-iH't} \phi_L(x) e^{iH't} \rangle = \frac{1}{2} [\phi_S(x - v_s t) + \phi_S(x + v_s t)]$ . To obtain the previous expression, we have taken into account that, in the initial state, the field  $\phi$  is split into its left and right parts  $\phi = \frac{1}{2}(\phi_L + \phi_R)$  originating from the left and right fermions in the bosonization procedure. For a time-independent state,  $\langle \phi_R \rangle = \langle \phi_L \rangle = \frac{\phi_S(x)}{2}$  as in the initial situation discussed above. On the other hand, the time evolution of  $\phi_L$  is independent of that of  $\phi_R$ , and it is given by a simple shift in  $vt$  to the left and to the right, respectively. In Fig. 3 we show the cumulative magnetization for different times compared to the DMRG results. We used the fitted value to the static solution for  $\xi'$  in  $\phi_S$ , as shown for the  $\beta = 0$  case in Fig. 1. A similar procedure gives  $\xi' = 2.781$  for the  $\beta = 0.24$  case. We also fix  $v_s$  to the spin-wave velocity of the homogeneous chains as previously discussed. We observe a fairly good comparison between the DMRG and bosonization results, especially when  $\beta$  is near the critical value  $\beta_c$ . This is due to the reduction of the finite-size effect in the latter case: as the evolution of the resulting excitations is slower as  $\beta$  increases, they do not reach the edge of the chain for the last time,  $t = 20$ , obtained in our calculation. The improved fitting between both results could also be due to the fact that the marginal irrelevant term neglected in the bosonized Hamiltonian could play some role in the short-distance correlations which, nonetheless, vanishes at  $\beta_c$ . As in the numerical results, we see that the original soliton does not propagate in a particular direction but the state splits into two counterpropagating modes as a consequence of the chiral symmetry of the Hamiltonian. As this pattern arises as a sum of two topological protected excitations, one on the right sector and one on the left sector of the theory, we can assume that the global excitation is also protected and remains stable with the time evolution. The jump in  $M_I$  indicates that each chiral mode carries  $\langle S^z \rangle = \frac{1}{4}$ . This can be interpreted as a superposition  $(|L\rangle + |R\rangle)/\sqrt{2}$  of two  $S^z = 1/2$  quan-

tum soliton states, such that  $|L\rangle$  ( $|R\rangle$ ) is a state propagating to the left (right). The system retains the memory of the initial state, but the original soliton state evolves into two chiral states as the time evolution is carried out with a Hamiltonian whose left and right modes are independent. However, the elementary excitations of the uniform Heisenberg model are not these type of solitons, notably the dynamical states preserve coherence and disperse quite slowly.

It is worth remarking that working with an odd number of sites is not a necessary condition for these results. An even number of sites will sustain a pair of soliton and antisoliton




FIG. 3. (Color online) Time evolution of the cumulative magnetization as given by DMRG (symbols) and bosonization (solid lines) in the (a) non frustrated and (b) frustrated ( $\beta = 0.24$ ) cases. The observed jump to  $M_I = 1/4$  as time evolves, signals the quantum interference of left and right soliton states.



different. As the initial excitation is local in real space, its distribution in momentum space is very broad, leading to a rapid dispersion of the excitation as time evolves. The most natural candidates to observe those phenomena would be spin-Peierls materials and ultracold atoms in optical lattices.

## ACKNOWLEDGMENTS

We thank Armando Aligia, Daniel Cabra, Anibal Iucci, Luis Manuel, and Adolfo Trumper for useful discussions. D. M. acknowledges the hospitality of the Dahlem Center. This work was partially supported by PIP CONICET Grant 0392, Grant DMR-1108285 (Ohio), and NSF-PIRE Grant 0730257.

- 
- <sup>1</sup> A. Wöllert and A. Honecker, Phys. Rev. B **85**, 184433 (2012).
- <sup>2</sup> I. Bloch, Nature (London) **1**, 23 (2005); M. Greiner, O. Mandel, T. Esslinger, T.W. Hänsch, and I. Bloch *ibid*, **415**, 39 (2002); B. Paredes, A. Widera, V. Murg, O. Mandel, S. Fölling, I. Cirac, G. V. Shlyapnikov, T. W. Hänsch, and I. Bloch *ibid*, **429**, 277 (2004); T. Fukuhara, P. Schauß, M. Endres, S. Hild, M. Cheneau, I. Bloch, and C. Gross, *ibid*, **502**, 76 (2013); I. Bloch, J. Dalibard, and W. Zwerger, Rev.Mod.Phys. **80**, 885 (2008).
- <sup>3</sup> F. Iglói and H. Rieger, Phys. Rev. Lett. **85**, 3233 (2000); K. Sen Gupta, S. Powell, and S. Sachdev, Phys. Rev. A **69**, 053616 (2004); R. W. Cherng and L. S. Levitov, *ibid* **73**, 043614 (2006); P. Calabrese and J. Cardy, Phys. Rev. Lett. **96**, 136801 (2006); M. Rigol, V. Dunjko, V. Yurovsky, and M. Olshanii, *ibid* **98**, 050405 (2007); M.A. Cazalilla, *ibid* **97**, 156403 (2006); C. Kollath, A.M. Lauchli, and E. Altman, *ibid* **98**, 180601 (2007).
- <sup>4</sup> S.R. Manmana, S. Wessel, R.M. Noack, and A. Muramatsu, Phys. Rev. B **79**, 155104 (2009); Phys. Rev. Lett. **98**, 210405 (2007).
- <sup>5</sup> C. Majumdar and D. Ghosh, J. Math. Phys. **10**, 1388 (1969)
- <sup>6</sup> B.S. Shastry and B. Sutherland, Phys. Rev. Lett. **47**, 964 (1981); W. Caspers, K. Emmett, and W. Magnus, J. Phys. A: Math. Gen. **17**, 2687 (1984).
- <sup>7</sup> M. Hase, I. Terasaki, and K. Uchinokura, Phys. Rev. Lett. **70**, 3651 (1993).
- <sup>8</sup> A. Seidel, C. A. Marianetti, F. C. Chou, G. Ceder, and P. A. Lee, Phys. Rev. B **67**, 020405(R) (2003).
- <sup>9</sup> S.R. White, Phys. Rev. Lett. **69**, 2863 (1992); U. Schollwöck, Ann. Phys. (NY) **326**, 96 (2011); K. Hallberg, Adv. Phys. **55**, 477 (2006).
- <sup>10</sup> T. Giamarchi, *Quantum Physics in One Dimension* (Oxford University Press, UK, 2003).
- <sup>11</sup> T. Vekua, D.C. Cabra, A. Dobry, C. Gazza, and D. Poilblanc, Phys. Rev. Lett. **96**, 117205 (2006); C.J. Gazza, A.O. Dobry, D.C. Cabra, and T. Vekua, Phys. Rev. B **75**, 165104 (2007); D.C. Cabra, C.J. Gazza, C.A. Lamas, H.D. Rosales, *ibid* **83**, 224406 (2011); D. Mastrogiuseppe, C. Gazza, and A. Dobry, *ibid* **83** 134428 (2011).
- <sup>12</sup> A.E Feiguin and S.R. White, Phys. Rev. B **72** 020404(R) (2005); A. Daley, C. Kollath, U. Schollwöck, and G. Vidal, J. Stat. Mech. Theor. Exp. (2004) P04005; R. Noack and S. Manmana, AIP Conf. Proc. **816**, 198 (2006); F. Verstraete, J.J. Garcia-Ripoll, and J.I. Cirac, Phys. Rev. Lett. **93**, 207204 (2004).
- <sup>13</sup> D. Gobert, C. Kollath, U. Schollwöck, and G. Schütz, Phys. Rev. E **71**, 036102 (2005); S. Langer, F. Heidrich-Meisner, J. Gemmer, I. P. McCulloch, U. Schollwöck, Phys. Rev. B **79**, 214409 (2009); S. Langer, M. Heyl, I. P. McCulloch, and F. Heidrich-Meisner, *ibid* **84**, 205115 (2011); S. Jesenko and M. Znidaric, *ibid* **84**, 174438 (2011); M. Ganahl, E. Rabel, F. H. L. Essler, and H. G. Evertz, Phys. Rev. Lett. **108**, 077206 (2012).
- <sup>14</sup> T. Nakano and H. Fukuyama, J. Phys. Soc. Jpn. **49**, 1679 (1980).
- <sup>15</sup> A. Dobry and D. Ibaceta, Phys. Rev. B **58**, 3124 (1998).
- <sup>16</sup> R. Rajaraman, *Solitons and Instantons: An Introduction to Solitons and Instantons in Quantum Field Theory*, North-Holland Personal Library, Amsterdam, The Netherlands (1987).
- <sup>17</sup> We applied a time step  $\Delta t = 0.1$  and kept 100 Lanczos vectors in the Krylov-space representation, resulting in a discarded weight of order  $O(10^{-7})$  at the end of the time evolution. The error bars result to be smaller than the size of the symbols.
- <sup>18</sup> A. Fledderjohann and C. Gros, Europhys. Lett. **37** 189 (1997).
- <sup>19</sup> A. Lavarélo and G. Roux, Phys. Rev. Lett. **110**, 087204 (2013).
- <sup>20</sup> V. Kiryukhin, B. Keimer, J.P. Hill, and A. Vigiante, Phys. Rev. Lett. **76**, 4608 (1996).
- <sup>21</sup> T. Fukuhara *et al.*, Nature Physics **9**, 235 (2013).
- <sup>22</sup> T. Esslinger, Annu. Rev. Condens. Matter Phys. **1**, 129 (2010).