

Published in final edited form as:

*Biochim Biophys Acta*. 2014 December ; 1842(0): 2610–2620. doi:10.1016/j.bbadis.2014.07.013.

## Central role of soluble adenylyl cyclase and cAMP in sperm physiology

Mariano G. Buffone<sup>1,\*</sup>, Eva V. Wertheimer<sup>2</sup>, Pablo E. Visconti<sup>3,\*,#</sup>, and Dario Krapf<sup>4,\*</sup>

<sup>1</sup>Instituto de Biología y Medicina Experimental, National Research Council of Argentina (CONICET), Buenos Aires, Argentina

<sup>2</sup>Centro de Estudios Farmacológicos y Botánicos (CEFYO), Facultad de Medicina, Universidad de Buenos Aires, Argentina

<sup>3</sup>Department of Veterinary and Animal Sciences, ISB, University of Massachusetts, Amherst MA 01003, USA

<sup>4</sup>Instituto de Biología Molecular y Celular de Rosario (CONICET) and Facultad de Ciencias Biológicas y Farmacéuticas, UNR, Rosario, Argentina

### Abstract

Cyclic adenosine 3',5'-monophosphate (cAMP), the first second messenger to be described, plays a central role in cell signaling in a wide variety of cell types. Over the last decades, a wide body of literature addressed the different roles of cAMP in cell physiology, mainly in response to neurotransmitters and hormones. cAMP is synthesized by a wide variety of adenylyl cyclases that can generally be grouped in two types: transmembrane adenylyl cyclase and soluble adenylyl cyclases. In particular, several aspects of sperm physiology are regulated by cAMP produced by a single atypical adenylyl cyclase (Adcy10, aka sAC, SACY). The signature that identifies sAC among other ACs, is their direct stimulation by bicarbonate. The essential nature of cAMP in sperm function has been demonstrated using gain of function as well as loss of function approaches. This review unifies state of the art knowledge of the role of cAMP and those enzymes involved in cAMP signaling pathways required for the acquisition of fertilizing capacity of mammalian sperm.

### Keywords

cAMP; sperm capacitation; soluble adenylyl cyclase; transmembrane adenylyl cyclase

---

© 2014 Elsevier B.V. All rights reserved.

<sup>#</sup>to whom correspondence should be addressed: pvisconti@vasci.umass.edu.

<sup>\*</sup>These authors equally contributed to this work

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## 1. Introduction

More than 50 years of investigation have passed since the discovery of cyclic adenosine 3', 5'-monophosphate (cAMP). This molecule stands as the first second messenger to be described<sup>1</sup>. Over the last decades, a wide body of literature addressed the different roles of cAMP in cell physiology, mainly in response to neurotransmitters and hormones. Although protein kinase A (PKA) was initially believed to be the sole effector of cAMP, new cAMP targets were later described. Examples of these are the RAP guanine-nucleotide-exchange factor (GEF) EPAC<sup>2,3</sup> and the cyclic-nucleotide-gated ion channels<sup>4</sup> which are also directly responsive to cAMP. Added to the complexity of the pathway, cAMP synthesis is conducted by products of ten distinct genes encoding adenylyl cyclase. Nine of them (Adcy1–9) encode the family of transmembrane adenylyl cyclase (tmAC) isoforms with differential tissue distribution and regulatory responsiveness. All isoforms possess a common predicted topology with high primary structure homology of their catalytic sites. Moreover, they are regulated by G proteins and activated by the diterpene forskolin<sup>5</sup>.

In contrast to the tmAC family, a single atypical adenylyl cyclase (Adcy10, aka sAC, SACY)<sup>6</sup> generates multiple isoforms by alternative splicing<sup>7,8</sup>. The signature that identifies sAC among other ACs, is their direct stimulation by bicarbonate<sup>9,10,11,12</sup> and as such serves as a CO<sub>2</sub> sensor in many cellular systems<sup>6</sup>. Unlike tmACs, sAC is insensitive to G protein regulation and its enzymatic activity is over ten times more effective in the presence of Mn<sup>2+</sup>/ATP than in the presence of Mg<sup>2+</sup>/ATP<sup>12</sup>. sAC was first described as a soluble adenylyl cyclase present in testicular extracts<sup>13</sup>. This property was used to purify sAC from rat testes and to identify it by microsequencing<sup>6</sup>. Not surprisingly, sAC is essential for fertilization; it is abundant in mature sperm and as such is the main source of cAMP in these cells. This review will be centered on the role of cAMP and those enzymes involved in cAMP signaling pathways in mammalian sperm. For more information on other aspects of the molecular basis of sperm function, other recent reviews can be found<sup>14,15,16,17,18,19,20,21,22</sup>

## 2. Role of cAMP in sperm function

Cyclic AMP orchestrates different aspects of sperm function required for the acquisition of fertilizing capacity. For a detailed list of principal findings and references, see Table I. After leaving the testis, mammalian sperm are morphologically mature; however, they are not able to fertilize. Fertilization capacity is acquired after completion of two post-testicular maturational processes. The first one, known as epididymal maturation, is poorly understood at the molecular level. It involves the activation of signaling pathways and the incorporation of proteins secreted by the epididymal epithelium in the male reproductive tract. During this process, the sperm acquire the ability to move progressively. However, after epididymal maturation, mammalian sperm are still incapable of fertilizing an egg. In the '50s, Chang<sup>23</sup> and Austin<sup>24</sup> independently reported that mammalian sperm should reside in the female tract for an obligatory period of time to acquire 'fertilizing capacity' in a process known as capacitation. This discovery was essential for the development of mammalian *in vitro* fertilization which was achieved by Chang eight years later<sup>25</sup>. Following Chang<sup>23</sup> and

Austin<sup>24</sup> pioneering work, capacitation was defined as the physiological processes that renders the sperm fertile<sup>26</sup>.

Cyclic AMP has been reported to be essential for events occurring during capacitation, including activation of motility, changes in the motility pattern known as hyperactivation and for development of the ability to undergo the acrosome reaction. There is clear evidence that many of the cAMP effects on sperm are mediated by activation of PKA. However, other cAMP-binding proteins have been proposed to participate in some of the processes<sup>27</sup>. In addition, cAMP and its targets regulate a variety of capacitation-induced signaling events including: 1) membrane lipid remodeling<sup>28,29</sup>; 2) hyperpolarization of the sperm plasma membrane<sup>30</sup>; 3) increase in intracellular pH<sup>30,31,32,33</sup>; 4) increase in intracellular Ca<sup>2+</sup><sup>34</sup>; and 5) increase in protein tyrosine phosphorylation<sup>35</sup>. Although one important aspect of the sperm capacitation process is the preparation of the sperm to undergo a physiologically relevant exocytotic reaction known as acrosome reaction, the acrosome reaction *per se* is usually considered outside of the capacitation process<sup>36</sup>. Therefore, the cAMP action on the acrosome reaction will be mentioned independently of capacitation<sup>27,37,38,39,40,41</sup>.

Capacitation can be mimicked in simple culture media containing Ca<sup>2+</sup>, HCO<sub>3</sub><sup>-</sup>, energy sources and a cholesterol acceptor that is usually bovine serum albumin. One of the first signaling events observed upon exposing sperm to capacitating conditions is a fast increase in cAMP content<sup>42</sup>. Interestingly, these initial high cAMP levels are rapidly reduced<sup>43</sup> for human sperm,<sup>44</sup> for boar sperm. Decrease of cAMP levels could be attributed either to hydrolysis by phosphodiesterases (see below) or to cAMP extrusion by specific transporters<sup>45</sup>. As expected, the increase in cAMP immediately activates PKA activity<sup>44</sup>. Although the first attempts to show PKA activation relied on direct PKA measurements<sup>46</sup>, more recently, development of antibodies against phosphorylated consensus sequences allowed for more precise kinetic studies. In particular, antibodies against phosphorylated PKA substrates recognizing either RXXpS/pT or RRXpS/pT were instrumental to demonstrate a fast PKA activation<sup>44,47,48,49,50,51</sup>. In all mammalian species studied, PKA is observed to be at maximum levels in less than a minute of exposure to capacitating conditions. This rapid up-regulation of cAMP signaling pathways is induced by HCO<sub>3</sub><sup>-</sup> in the capacitation medium, suggesting that it is mediated by sAC<sup>52</sup>.

Direct cAMP measurements and Western blots using anti phospho-PKA substrates antibodies have been used to reveal how the cAMP pathway is regulated. However, these experiments were silent regarding the role of cAMP in sperm function. To study the consequences of cAMP increase and PKA activation in sperm, loss and gain of function strategies were used. Gain of function experiments compared the sperm response in the absence or presence of HCO<sub>3</sub><sup>-</sup>, and also employed a combination of permeable cAMP agonists and phosphodiesterase inhibitors to mimic the cAMP response in the absence of an upstream stimulus. On the other hand, loss of function was achieved using pharmacological<sup>35</sup> and genetic knock-out approaches targeting sAC<sup>52,53,54</sup>, the PKA catalytic subunit<sup>49</sup> or the anchoring of PKA to A-kinase anchoring proteins (AKAPs)<sup>55</sup>. Altogether these approaches revealed an essential role of cAMP-signaling pathways, in particular of PKA, in the activation of sperm motility and in the induction of a vigorous asymmetrical movement (*i.e.* hyperactivated motility) necessary for sperm to fertilize.

Although the role of cAMP in the acrosome reaction is less conclusive, recent work presented evidence that the GDP/GTP exchanger EPAC, another target of cAMP is present in sperm and is involved in the regulation of the acrosome reaction<sup>37,56</sup>. It is still not clear whether cAMP synthesis in the sperm anterior head is due to the activity of sAC<sup>37</sup> or tmACs<sup>41</sup>. In the following sections, we will discuss individually the role of the enzymes involved in cAMP signaling.

### 3. G proteins

A positive correlation between cAMP and mammalian sperm motility has been first observed by the group of Henry Lardy in 1971 in bovine samples<sup>57</sup>. These first investigations were followed by active research on the enzymes involved in cAMP synthesis. See Table II for a summarized detail of molecules regulating cAMP levels in sperm and corresponding references. Even though cAMP was known to be important for sperm physiology, conflicting reports were published regarding the presence and activity of tmACs in spermatozoa. These enzymes are both positively and negatively regulated by heterotrimeric G proteins, composed of a  $G\alpha$ -subunit and a  $G\beta\gamma$ -dimer. The  $G\alpha$ -subunit can either stimulate ( $\alpha_s$  and  $\alpha_{olf}$ ), or inhibit ( $\alpha_{i1}$ ,  $\alpha_{i2}$  and  $\alpha_{i3}$ ) tmACs<sup>58</sup>. Activation of the  $\alpha$  subunit promotes the exchange of bound GDP for GTP. Then, GTP binding alters the topology within the  $G\alpha$ -subunit, releasing both an active  $\alpha$ -subunit and  $\beta\gamma$ -dimer from the formerly inactive trimeric G protein. Separately, both the  $\alpha$ -subunit and  $\beta\gamma$ -dimer act on different downstream effectors, such as phosphodiesterases, phospholipases, ion channels and tmACs. Both stimulatory and inhibitory  $\alpha$  subunits can be post-translationally modified by bacterial toxins. While all  $G\alpha_i$  subunits are ADP ribosylated by pertussis toxin, both  $G\alpha_s$  and  $G\alpha_{olf}$  can be modified by cholera toxin<sup>59,60,61</sup>. In both cases, toxin modifications inhibit the respective G protein function and can be followed *in vitro* using the substrate NAD(<sup>32</sup>P). The differential ADP ribosylation properties of G protein  $\alpha$ -subunits have been used over the years to modulate  $G\alpha_s$  and  $G\alpha_i$  activities and also as a molecular signature for their identification in many cell types<sup>62</sup> including sperm<sup>63,64</sup>.

Regarding the presence of G proteins in sperm, few direct experiments have been published. A first report by Hildebrandt *et al.* failed to detect either  $G\alpha_s$  or  $G\alpha_i$  by ADP ribosylation studies in total dog sperm extracts using cholera or pertussis toxins respectively<sup>65</sup>. Accordingly, Western blots using antibodies raised against  $G\alpha_s$  in bovine and human sperm also failed to find positive evidence<sup>66,67</sup>. However, opposite results can be found in the literature; Kopf *et al.*<sup>68</sup>, using a modification of the ADP-ribosylation methodology, showed that Lubrol PX-extracted 48,000  $\times g$  supernatant fractions of sperm from sea urchin, abalone, mouse and human contained substrates for ADP-ribosylation by pertussis toxin. Moreover, the pertussis toxin substrates were of same molecular weight as that reported for  $G\alpha_i$  (41 KDa) and could be immunoprecipitated with anti  $G\alpha_i$  antibodies<sup>68</sup>. It was argued that the discrepancy with Hildebrandt *et al.*<sup>65</sup> was as a result of a detergent extraction previous to the ADP-ribosylation, indicating that in the native form, G proteins might be masked.

Soon after, Kopf's results were corroborated in other vertebrate and invertebrate species<sup>69</sup>. On the other hand, none of the above presented groups were able to detect  $G\alpha_s$  in sperm. It

was many years later that Fraser and Adeoya-Osiguwa<sup>70</sup> reported the presence of  $G\alpha_s$  by Western blot analysis. A later report<sup>71</sup> showed that the addition of cholera toxin to cell lysates prepared from permeabilized sperm increased cAMP production by 10% suggesting that ADP-ribosylation of  $G\alpha_s$  by cholera toxin lead to enhanced stimulation of tmAC activity. The same group detected positive ADP-ribosylation of  $G\alpha_s$  by cholera toxin in permeabilized human and mouse sperm, also consistent with the presence of  $G\alpha_s$ <sup>71</sup>. Supporting this conclusion, it was recently shown that a cholera toxin ADP-ribosylation substrate protein can be immunoprecipitated from mouse sperm extracts, showing the expected molecular weight for  $G\alpha_s$ <sup>41</sup>. Immunofluorescence experiments localized  $G\alpha_s$  only to the anterior head of mouse sperm. Moreover,  $G\alpha_s$  immunoreactivity was lost in acrosome-reacted sperm suggesting that  $G\alpha_s$  resides in the acrosomal region<sup>41</sup>. Unfortunately, in the light of high sequence similarities, the above mentioned techniques did not allow to discriminate between  $G\alpha_s$  and  $G\alpha_{olf}$ -subunits mainly because: 1) both of them can be ADP-ribosylated in the presence of cholera toxin; 2) they are likely to be recognized by the same antibodies; and 3) once activated, both can stimulate tmACs. Noteworthy, positive data regarding the presence of  $G\alpha_s$  in mouse sperm also comes from mass spectrometry analyses of sperm protein extracts<sup>72</sup>.

Besides  $G\alpha_s$ , studies were also conducted to address the presence of  $G_q$ , which acts stimulating phospholipase  $C_\beta$ . Immunodetection analysis showed the localization of  $G_q$  to the anterior acrosomal region of mouse<sup>73</sup> and equatorial segment of human sperm<sup>74</sup> suggesting that it might have a role in acrosomal exocytosis. Recently, another  $G\alpha$  member,  $G\alpha_{13}$  (akaGNA13) has been identified in bovine sperm by Western blot<sup>75</sup>. Their roles in sperm physiology are yet to be determined.

## 4. cAMP synthesis: soluble and transmembrane adenylyl cyclases

### 4.1. sAC

Multiple studies have shown that sperm capacitation is a  $HCO_3^-$  and  $Ca^{2+}$ -dependent process<sup>29,76,77,78,79,80</sup>. The first connection between  $HCO_3^-$ ,  $Ca^{2+}$  and cAMP metabolism was demonstrated by the Garbers group<sup>81</sup>. In this manuscript it was shown that in the presence of  $HCO_3^-$ , extracellular  $Ca^{2+}$  increases intracellular cAMP within 1 min in guinea pig spermatozoa. When either  $HCO_3^-$  or  $Ca^{2+}$  was removed from the medium, only a slightly increase of cAMP was observed. Later on, Okamura and collaborators<sup>82</sup> reported that a low molecular weight factor present in boar semen was able to induce pig sperm motility. The low molecular weight molecule was later identified as  $HCO_3^-$  and was shown to directly activate adenylyl cyclase activity and consequently increase cAMP levels in sperm<sup>83</sup>. Similar results were later reported in bovine and hamster spermatozoa<sup>84,85</sup>. Independently of these reports, other groups have described the presence of a soluble adenylyl cyclase in testicular extracts<sup>13,86</sup>. This enzyme activity presented certain properties similar to the particulate sperm enzyme. Analogous to the sperm adenylyl cyclase, the soluble testicular enzyme did not appear to be responsive to G proteins and was more active when assayed in the presence of  $Mn^{2+}$  than in the presence of  $Mg^{2+}$ <sup>86</sup>. Using a biochemical approach, the group of Drs. Lonny Levin and Jochen Buck<sup>6</sup> purified a protein responsible for the soluble cyclase activity from 950 rat testes and microsequenced it by mass

spectrometry. The derived peptide sequence was used to build degenerate primers to clone the enzyme. In a follow-up manuscript, they demonstrated that the testicular sAC is present in sperm and that it is responsive to  $\text{HCO}_3^-$ <sup>10</sup>. The elucidated sequence revealed conservation throughout evolution. Most interestingly, the sAC catalytic domain presents homology to cyanobacterial adenylyl cyclase which is also  $\text{HCO}_3^-$ -dependent<sup>10,32</sup>. Bicarbonate stimulates the enzyme's  $V_{\max}$ <sup>12</sup> by fostering a conformational rearrangement of the active site, allowing a  $\text{Mg}^{2+}$ -ATP to properly interact with it<sup>87</sup>. The sensitivity to  $\text{HCO}_3^-$ , turns sAC into a  $\text{CO}_2$  sensor widely distributed in nature. Interestingly, sAC can also function as a pH sensor, due to the effect of pH on the  $\text{CO}_2/\text{HCO}_3^-$  equilibrium<sup>88</sup>. Thus, alkalization of the intracellular milieu during capacitation might also be expected to impact on sAC activity, through modulation of  $\text{HCO}_3^-$  levels. As mentioned, cAMP levels peak at 1 min after sperm exposure to  $\text{HCO}_3^-$ <sup>43,89</sup> and then return to basal levels over the incubation period. A self-regulatory loop has been proposed in which PKA exerts a negative feedback on sAC activity<sup>90,91,92</sup>. A self-regulatory loop has been proposed in which PKA phosphorylates sAC or an intermediary protein, exerting a negative feedback on its activity. In addition to responding to  $\text{HCO}_3^-$ , sAC is also a  $\text{Ca}^{2+}$  sensor. In the presence of  $\text{Ca}^{2+}$ , the  $K_m$  for  $\text{Mg}^{2+}$ -ATP is reduced to levels that are close to the ATP concentration found in cells, turning sAC into a signal transducer highly sensitive to physiologically changes of  $\text{HCO}_3^-$ <sup>93</sup>. Finally, it is important to consider that although sAC was originally found in the testicular soluble fraction, the  $\text{HCO}_3^-$  dependent adenylyl cyclase activity in mature sperm is associated with the plasma membrane<sup>85</sup>. More research is needed to understand the nature of this association and how sAC is transferred from the soluble to the particulate fraction during spermiogenesis or epididymal maturation. Alternatively, sAC could be represented by two populations: a soluble and a particulate one.

Although the aforementioned works strongly suggested a role of sAC in sperm motility, loss of function experiments were needed to conclusively demonstrate its role in sperm capacitation. To obliterate sAC function genetic and pharmacological approaches were used. First, elimination of the sAC gene by homologous recombination resulted in a sterile phenotype<sup>52,53</sup>. Sperm from the sAC knock-out were immotile and were unable to undergo the cAMP/PKA-dependent increase in tyrosine phosphorylation associated with the capacitation process<sup>52</sup>. In addition, basal levels of cAMP in sAC<sup>-/-</sup> sperm were under detection limits when these cells were incubated in capacitation-supporting media containing  $\text{HCO}_3^-$ <sup>41</sup>. Second, sAC was used as bait in high throughput assays designed to find small molecules which specifically inhibit sAC. The first compound found, KH7, was a competitive inhibitor with  $\text{Mg}^{2+}$ -ATP and blocked recombinant sAC activity with an  $\text{IC}_{50}$  of 10  $\mu\text{M}$ <sup>94</sup>. KH7 was then used in sperm and was shown to block cAMP production, the increase in tyrosine phosphorylation and sperm motility<sup>52</sup>. Altogether, these results are consistent with the hypothesis that sAC mediates the  $\text{HCO}_3^-$ -induced increase in cAMP necessary to initiate capacitation. Moreover, in the absence of sAC, sperm have reduced ATP levels, do not undergo hyperactivation and failed to fertilize *in vitro*<sup>52,54</sup>.

#### 4.2. tmACs

The presence of tmAC isoforms in sperm has been investigated using enzymatic assays<sup>95</sup>, immunological approaches<sup>95</sup>, mass spectrometry<sup>72</sup> and genetic knock-outs<sup>96</sup>. Despite

these efforts, contrary to sAC, the function as well as the presence of tmACs in mammalian sperm remains controversial. Although some studies presented evidence of their presence in mammalian gametes<sup>41,96,97,98,99,100</sup>, others reported negative results<sup>65,89,101,102,103,104,105</sup> (see Brenker et al<sup>89</sup> for a thorough discussion of these arguments). In many of these studies, the authors used the diterpene, forskolin, which known to increase cAMP levels in most cell types as a surrogate to identify the presence of tmAC activity. Initially described by Seamon and Daly<sup>106,107</sup>, forskolin activates all tmACs except Adcy9. Most importantly, forskolin does not stimulate sAC activity. In mammalian sperm incubated with forskolin, several groups were unable to observe an increase in cAMP<sup>11,52,89,104,105,108</sup>. However, other groups reported a forskolin-dependent elevation of intracellular cAMP concentrations<sup>41,71,95,96</sup>. These last results were in agreement with data from Leclerc and Kopf<sup>99,109</sup> which showed a forskolin effect in adenylyl cyclase activity assayed in membrane fractions of capacitated mouse sperm. Interestingly, this work also showed an effect of both GTP analogues and mastoparan, a wasp venom toxin known to activate G proteins. Considering the abundance of sAC, one possibility is that the effect of forskolin in mammalian sperm is masked by higher levels of endogenous forskolin-independent cAMP synthesis. Considering this possibility, our group recently showed that forskolin increased cAMP levels in sperm from sAC knock-out mice<sup>41</sup>.

Despite these data, little is known regarding the function of specific tmACs in sperm. Considering that tmACs are regulated by G proteins, the finding of  $G\alpha_s$  only in the sperm acrosome<sup>41</sup> suggests that tmACs have a role in the acrosome reaction. Consistently, incubation with forskolin stimulated the acrosome reaction in capacitated mouse sperm and, in addition, induced an increase in intracellular  $Ca^{2+}$  concentration<sup>41</sup>. Because tmACs may have overlapping functions in many cell types, it is difficult to pinpoint the specific isoform involved in a given process. Due to the low abundance of tmACs in sperm and the lack of good specific antibodies, results on specific isoforms in sperm have been inconclusive<sup>95</sup>. Regarding loss of function experiments, in addition to sAC knock-out mice<sup>52</sup>, only *Adcy3*-null mice are infertile<sup>96</sup>. Knockout mice for *Adcy1*, 5, 6 and 8 are fertile; *Adcy7* knock-out shows complete postnatal lethality<sup>110</sup> and null mice for *Adcy2* and *Adcy9* have not yet been produced. In the case of *Adcy3* null mice, the ability of their sperm to penetrate cumulus-enclosed oocytes *in vitro* is compromised<sup>96</sup>.

## 5. cAMP Degradation: Phosphodiesterases in sperm physiology

Levels of intracellular cAMP are highly dynamic; its availability depends on the concerted action of both synthesis and degradation. In sperm, addition of permeable cAMP analogues trigger capacitation-associated events<sup>111</sup>. However, in many cases, depending on the species, no matter how high the concentration of permeable cAMP analogues are used, unless phosphodiesterases (PDE) inhibitors are added, no effect is observed. In mammals, 20 genes grouped in 11 PDE families have been described. Each PDE family shows different substrate specificity and pharmacological sensitivity. PDE4, PDE7, and PDE8 family members are more efficient hydrolyzing cAMP, whereas PDE5, PDE6, and PDE9 are cGMP-specific. On the other hand, PDE1, PDE2, PDE3, PDE10, and PDE11 are active toward both cAMP and cGMP<sup>112</sup>. In sperm, PDE1 was first reported by Wasco and Orr<sup>113</sup> and shown to be stimulated by calmodulin and to associate with the head and tailpieces of

demembrated rat caudal epididymal sperm. Using immunological techniques, six additional PDE types have been shown to be present in mammalian sperm (1, 4, 6, 8, 10 and 11) <sup>114</sup>. Inhibition of either PDE1 (by MMPX) or PDE4 (by RS-25344, Rolipram ) augmented intracellular levels of cAMP <sup>114</sup>. In human sperm, inhibition of PDE4 by RS-25344 and Rolipram enhanced sperm motility without affecting the acrosome reaction <sup>115</sup>. On the other hand, PDE1 inhibitors selectively stimulated the acrosome reaction <sup>115</sup>. Although more research is needed, these results are consistent with the hypothesis that, in sperm, the molecules involved in cAMP signaling which include adenylyl cyclases, phosphodiesterases and cAMP targets are readily compartmentalized and regulate different sperm processes, some in the head (*e.g.* acrosome reaction); and some in the tail (*e.g.* regulation of motility). This compartmentalization would be possible through local cAMP generation within independently regulated microdomains. It can be hypothesized that a variety of cAMP modules are modulated by AKAPs (see below), which tether not only PKA but many enzymes and substrates to specific cell locations <sup>116</sup>. Among these tethered enzymes, PDEs can function to prevent cAMP diffusion, reinforcing barriers between microdomains. In the light of the restricted cAMP microdomains hypotheses, conclusions from experiments in which sperm are flooded with permeable cAMP agonists should be carefully analyzed.

## 6. cAMP-dependent protein kinase (PKA)

The best characterized cAMP target is the holoenzyme PKA, which is essential in both, sperm and egg biology <sup>90</sup>. Please see Table III for a complete list of known cAMP targets in sperm. PKA is a heterotetramer composed of two catalytic subunits (C) containing the enzyme's active site, and two regulatory subunits (R), which block PKA activity when bound to the C subunit. The active catalytic subunit is dissociated as an active kinase upon cAMP binding to R subunits. In mammals, there are four R (*RI $\alpha$* , *RI $\beta$* , *RII $\alpha$*  and *RII $\beta$* ) and five C genes (*PRKACA*, *PRKACB*, *PRKACG*, *PRKX*, and *PRKY*). Expression of C genes gives rise to isoforms C $\alpha$ , C $\beta$ , C $\gamma$ , PRKX and PRKY <sup>117</sup>. However, the functions of PRKX, PRKY and C $\gamma$  are not known. Interestingly, C $\gamma$  is an intronless gene exclusively expressed in the testis <sup>117,118</sup>. C $\alpha$  and C $\beta$  can combine with the different R subunits to form a variety of PKA holoenzymes. Added to this complexity is the finding that both C $\alpha$  and C $\beta$  can be found as alternative spliced variants. While C $\beta$  has at least ten different variants, C $\alpha$  has only two (C $\alpha$ 1 and C $\alpha$ 2).

Because mature sperm are transcriptionally and translationally inactive, gene expression studies are usually conducted in testicular germ cells <sup>90</sup>. Throughout spermatogenesis, only *RI $\alpha$* , *RII $\alpha$* , C $\alpha$ 1, C $\alpha$ 2 and C $\gamma$  mRNAs are found. Interestingly, while *RI $\alpha$*  is expressed in all germ cell stages, *RII $\alpha$*  expression patterns are restricted to later stages suggesting a role in sperm function. Regarding the C subunits, C $\alpha$ 1 is expressed in spermatogonias which correspond to the mitotic stages of spermatogenesis; however, the C $\alpha$ 1 expression declines in meiotic spermatocytes and is completely replaced by splice variant C $\alpha$ 2 in pachytene spermatocytes and in spermatids. C $\alpha$ 2 contains 7 unique amino acids in the N-terminal domain and lack the myristylation site present in C $\alpha$ 1. It has been proposed that these changes may confer special properties to the PKA holoenzyme in the sperm. Finally,



although  $C\gamma$  mRNA is found in testicular germ cells, protein expression of this mRNA has not been demonstrated yet.

As mentioned above, as soon as the sperm comes in contact with  $HCO_3^-$  in the seminal plasma there is a rapid increase in cAMP synthesis mediated by sAC. Design of antibodies against PKA substrate consensus phosphorylation sites has permitted the kinetics of PKA activation in sperm to be monitored. Using these antibodies, it was shown in sperm from different species that PKA activity reaches maximum activity in less than 1 min after exposure to  $HCO_3^-$ <sup>43,44,48</sup>. The role of  $HCO_3^-$  in this activation strongly suggests that sAC is the adenylyl cyclase involved in this effect. This conclusion is also supported by the lack of effect of forskolin on PKA activation and by the finding that, while  $G\alpha_s$  localized to the sperm anterior head, the PKA catalytic subunit is only found in the flagellum<sup>41</sup>. The role of cAMP in the regulation of sperm capacitation has been supported using a battery of inhibitors such as H89, rpScAMP and a permeable stearated version of the peptide protein kinase inhibitor PKI<sup>35,119</sup>. In particular, it is well established that PKA activation is upstream the regulation of other signaling events involved in capacitation (see below). However, although PKA activation is a fast event, it should be noted that some of these PKA-dependent signaling pathways known to be downstream of PKA activation such as the increase in protein tyrosine phosphorylation and the induction of hyperactivated motility are slow processes<sup>120</sup>. The delayed effects of cAMP-dependent pathways suggest that PKA action on this pathway is not direct.

The pharmacological approaches have been later validated using genetic knock-out models. Although elimination of the whole  $C\alpha$  gene by homologous recombination resulted in infertile sperm<sup>121</sup>, these mice have many other defects that preclude an accurate assessment of the reproductive phenotype. Considering that  $C\alpha_2$  is the only C subunit present in mature mammalian sperm<sup>91,122</sup>, it was possible to obliterate PKA activity specifically from sperm by eliminating the unique N-terminus  $C\alpha_2$  exon by homologous recombination. These mice, containing a functional somatic  $C\alpha_1$ , have only a reproductive phenotype. They are sterile and the infertile phenotype is related to defects in sperm motility and capacitation<sup>91, 121</sup>. On the other hand, target disruption of the type II $\alpha$  regulatory subunit (*RII $\alpha$* ) resulted in a compensatory increase of RI $\alpha$  protein and a dramatic relocalization of PKA such that the majority of the holoenzyme now appears in the soluble fraction and colocalizes to the cytoplasmic droplet. Unexpectedly the *RII $\alpha$*  null mice are fertile and have no significant effect on sperm motility<sup>123</sup>. Further confirmation of the role of  $C\alpha_2$  in sperm motility was achieved by a pharmacological and genetics combined approach. First described by Bishop *et al.*<sup>124</sup>, this approach consists of a point mutation of the ATP binding domain of a given protein kinase, changing a specific amino acid of the ATP binding site for a smaller one such as alanine or glycine. This mutation makes the respective protein kinase sensitive to bulky ATP competitive inhibitors which otherwise cannot block the wild type kinase activity. Morgan *et al.*<sup>49</sup> used a knock-in approach to swap the endogenous  $C\alpha$  gene for one in which the “gate-keeper” methionine had been mutated to alanine (called *CaM120A*). Sperm from the mutant  $C\alpha$  but not those from wild type mice were sensitive to the bulky inhibitor NMPP1<sup>49</sup>. As expected, NMPP1 blocked PKA activation in the mutant sperm; in addition,

this compound inhibited the  $\text{HCO}_3^-$ -dependent increase in flagellar beat frequency and the downstream increase in tyrosine phosphorylation.<sup>49</sup>

Similar to other protein kinases and phosphoprotein phosphatases, PKA has multiple substrates. This property highlights an important problem in signal transduction: How is the phosphorylated state of a given substrate regulated without affecting the phosphorylation state of other possible substrates? One answer to this question was given by the discovery of proteins that can anchor PKA. These proteins are known as AKAPs (for A Kinase Anchoring Proteins) for review see 116 and references therein AKAPs are a family of proteins that anchor the regulatory subunit of PKA, confining its activity to discrete locations within the cell, crucial for successful localized signaling activity of PKA and other enzymes<sup>116,125</sup>. Although AKAPs can bind a series of enzymes allowing scaffolding of many signaling pathways, their discovery was facilitated by the AKAP property of binding RII PKA subunits *in vitro*. Using RII binding assays, two sperm-specific AKAPs were described, AKAP3<sup>126,127,128</sup> and AKAP4<sup>129</sup>. These proteins are distributed throughout the sperm flagellum, and have been found to become phosphorylated during capacitation<sup>129,130,131,132</sup>. The role of these phosphorylations has not yet been elucidated but it has been proposed that these post translational modifications may regulate the activity or localization of different signaling complexes needed for sperm capacitation.

AKAPs role in sperm was analyzed using modified permeable peptides such as St-Ht31. This inhibitor peptide is a steared form of the peptide Ht-31 derived from the human thyroid AKAP and inhibits the interaction between the RII subunit of PKA and AKAP in cell extracts. This peptide has been first used to study PKA signaling in sperm<sup>119,133</sup>, and later expanded to other cell types. The steared peptide HT31 impairs sperm motility and the onset of signaling pathways associated with sperm capacitation<sup>47,119,134</sup>. Interestingly, this blocking peptide produced a stronger inhibition of motility that obtained solely with the use of PKA inhibitors such as H89. The blocking peptide St-Ht31 was designed to mimic the RII sequence that binds to AKAPs. Homologous RII sequences can be found in other proteins and this AKAP-interacting domain has been named R2D2<sup>135</sup>. Therefore, the stronger phenotype from St-Ht31 has been attributed to the combined disruption of other R2D2-containing proteins present in sperm cells<sup>126,136</sup>. The role of AKAPs in sperm physiology has been also validated using knock out models. Sperm lacking AKAP4 are infertile and present morphological flagellar defects an reduced progressive motility due to the role of this protein in tail assembly<sup>55</sup>. Sperm numbers were not reduced in male mice lacking AKAP4 but proteins usually associated with the fibrous sheath were absent or substantially reduced in amount. Other cytoskeletal components of the tail appeared fully developed.

## 7. Other targets of cAMP in spermatozoa

In addition to PKA, cAMP is able to bind and regulate other proteins. These proteins are characterized by a cyclic nucleotide binding domain (CNBD) characterized by depicting a conserved structure which binds cyclic nucleotides and differs in their specificity towards cAMP and cGMP. CNBD-containing proteins are involved in signaling. Some of the best characterized belong to the families of phosphodiesterases, ion channels and G protein

exchange factors. CNBD-containing proteins have been identified in mammalian sperm and shown to play a role in the regulation of different aspects of sperm function.

Cyclic nucleotide-gated channels (CNGc) are ion channels whose activity responds to the binding of cyclic nucleotides. These channels result from the combination of two  $\alpha$  subunits (CNG $\alpha$ 1 to 4) and two  $\beta$  subunits (CNG $\beta$ 1 to 3), which are differentially expressed in different tissues<sup>137</sup>. These channels have been long known for participating in sensory transduction processes, but their function in other tissues is not well established yet. From a structural point of view, each subunit possesses six transmembrane segments and a single cyclic nucleotide-binding site near the C-terminal region of the protein<sup>138,139</sup>. Binding of cAMP or cGMP to nucleotide binding domain promotes the opening of the CNG channels. Depending on subunit combinations, CNGc displays differential selectivity towards cAMP or cGMP<sup>140</sup> and often carry inward Na<sup>+</sup> and Ca<sup>2+</sup> currents<sup>141</sup>.

The role of these channels in mammalian sperm capacitation is not clear. However, their role in sea urchin sperm chemotaxis is well known<sup>142,143</sup>. Interestingly, the first sperm ion channel cloned from mouse testis was a CNG channel<sup>144</sup>. After this discovery, others have been cloned<sup>145</sup>. The  $\alpha$ 3 and  $\beta$ 1 subunits were localized to the flagellum of mouse spermatozoa<sup>145</sup>. Addition of permeable cGMP analogues to murine sperm results in an increase of intracellular Ca<sup>2+</sup><sup>146</sup>. The same effect was observed with permeable cAMP analogues, but with less effectiveness than with cGMP analogues<sup>147</sup>. Even though CNGc were originally proposed to carry Ca<sup>2+</sup> inward currents upon direct stimulation with cAMP/cGMP<sup>146</sup>, it was later shown that sperm lacking the sperm specific Ca<sup>2+</sup> channel CatSper do not show this response, even though CNGc are still present<sup>148</sup>. Currently the role of CNG channels in sperm physiology is not clear because mice bearing targeted deletion of  $\alpha$ 3 showed normal fertility rates<sup>149</sup>.

Another cAMP effector is the exchange protein directly activated by cAMP (Epac). Epac is a Rap-specific guanine-nucleotide exchange factor that is activated by the interaction of cAMP with the cyclic nucleotide monophosphate-binding domain<sup>2,3</sup>. Epac1 and Epac2, the two isoforms so far described in mammalian cells, possess a regulatory and a catalytic region in the N- and C-terminal segments respectively. The regulatory domain contains the cAMP binding site, which in the absence of cAMP, self inhibits its catalytic activity<sup>150</sup>. In sperm, EPAC has been shown to be present using immunological as well as pharmacological approaches.

Epac1 and Epac2 have been detected by immunoblotting in sperm from different species including human<sup>27,56,151,152</sup>. In addition, immunofluorescence experiments suggested that these proteins localize to the sperm head. In addition, the Epac G protein target Rap1A was also found to be present in sperm<sup>56,151</sup>. Functionally, the Epac-selective 8-(p-chlorophenylthio)-2'-O-methyladenosine-3',5'-cyclic monophosphate (8-pCPT-2-O-Me-cAMP) analogue is routinely used to stimulate Epac without inducing PKA activation. This cAMP analogue induced the acrosome reaction in permeabilized human sperm in a Ca<sup>2+</sup>-independent manner<sup>27</sup>. These data suggest that extracellular Ca<sup>2+</sup> entry might be upstream of Epac activation. Moreover, using a streptolysin-permeabilized sperm model, it was shown that addition of Ca<sup>2+</sup> failed to trigger acrosomal exocytosis when intracellular cAMP was

depleted by phosphodiesterase activity or when Epac was sequestered by specific blocking antibodies<sup>27</sup>. Interestingly, the PKA catalytic subunit was not found in the sperm head<sup>41</sup> indicating that the effect of cAMP in the sperm head is PKA-independent and suggesting a role for alternative targets such as Epac in the regulation of the acrosome reaction. A third CNBD protein known to be present in mouse sperm is a specific member of the mammalian NHE superfamily of Na<sup>+</sup>/H<sup>+</sup> exchangers, sNHE<sup>31</sup>. This exchanger localized to the principal piece of the sperm flagellum, and is predicted to contain 14 putative transmembrane segments, although its actual topology remains to be determined. One key feature of this exchanger is that close to its intracellular C-terminus, there is a nucleotide-binding domain indicating that modulation by cyclic nucleotides might also be functional<sup>31</sup>. Targeted deletion of *sNHE* gene results in infertility<sup>31</sup>. Homozygous-null males mated with females, produced vaginal plugs with the same frequency as wild-type males, but did not result in pups. Interestingly, sperm from sNHE null mice lack motility but either permeable cAMP analogs or NH<sub>4</sub>Cl rescues this phenotype. Later reports from the same group showed that sNHE is necessary for the full length expression of SACY suggesting that these two proteins may form a protein complex involved in sperm intracellular pH (pH<sub>i</sub>) control<sup>32</sup>. Regulation of sperm pH<sub>i</sub> is fundamental for motility, capacitation, and acrosome reaction in different mammalian species<sup>30,33,153</sup>. It has been proposed that cAMP affects the sperm pH<sub>i</sub> by controlling the sperm-specific Na<sup>+</sup>/H<sup>+</sup> exchanger through its cAMP-binding domain<sup>20</sup>.

Finally, a novel CNBD containing protein named CRIS (for cyclic nucleotide receptor involved in sperm function) has been recently identified<sup>154</sup>. CRIS is exclusively expressed in testicular germ cells while it is not present in mature sperm. Interestingly, mouse genetic models lacking CRIS are either infertile or subfertile. While infertility is related to spermatogenic arrest, some of the mice completed this process and contain sperm in their epididymis. These sperm, however, have defects in motility due to problems in Ca<sup>2+</sup> transport and flagellar bending. At present, the role of cAMP in CRIS regulation has not been established.

## 8. cAMP-dependent signalling pathways

In the previous sections, we have reviewed the current knowledge on those enzymes that directly participate in cAMP metabolism in sperm. We have also summarized the role of these enzymes and of cAMP in the physiology of the sperm, in particular in sperm capacitation and the acrosome reaction. As mentioned in several sections, sAC is stimulated immediately after the sperm is exposed to HCO<sub>3</sub><sup>-</sup> upon ejaculation. In addition, the relevance of sAC, intracellular cAMP concentrations and the consequent PKA activation in the regulation of the sperm fertilizing capacity is well established. However, less is known about those pathways downstream of cAMP and PKA. In the following sections we will address some of these signaling events.

### 8.1. Lipid remodeling

Similar than what occurs in somatic cells, sperm phospholipids are distributed asymmetrically within the plasma membrane<sup>155</sup> by the concerted activity of several phospholipid transferases. An aminophospholipid transferase that moves phosphatidylserine and phosphatidylethanolamine from the outer to the inner leaflet<sup>156</sup>, a non-specific

transferase (floppase) that transfers phospholipids from the inner to the outer leaflet and a scramblase that moves all phospholipids in both directions. The increase of scramblase isoform 1 activity was observed during sperm capacitation associated with phospholipids scrambling<sup>157</sup>. This change in asymmetry is stimulated by  $\text{HCO}_3^-$  through sAC and the cAMP/PKA pathway, and promotes a fast exposure of phosphatidyl ethanolamine and phosphatidyl serine not related to apoptosis<sup>157</sup>. Regarding the functional implications of this event, it has been proposed that phospholipid scrambling facilitates cholesterol efflux, an event required for capacitation<sup>158,159</sup>.

## 8.2. Increase in tyrosine phosphorylation

Capacitation can be mimicked in vitro in a simple medium containing a combination of ions such as  $\text{Na}^+$ ,  $\text{Cl}^-$ ,  $\text{Ca}^{2+}$  and  $\text{HCO}_3^-$ , energy substrates such as glucose and a cholesterol acceptor (usually bovine serum albumin). Absence of any of these molecules prevents the sperm from acquiring fertilizing capacity. Omission of any of these molecules also blocks an increase in tyrosine phosphorylation<sup>80,92,160,161</sup>. The capacitation-associated increase in mouse sperm tyrosine phosphorylation is downstream of a cAMP/PKA-dependent pathway<sup>35</sup> and is also the case for capacitation in sperm from other mammalian species including human<sup>130,162,163,164</sup>. Because forskolin is unable to induce this pathway, it is clear that the only AC involved in the regulation of tyrosine phosphorylation is sAC<sup>41</sup>. Although the role of PKA upstream this pathway the mechanism by which PKA activates tyrosine phosphorylation is not known. Considering that PKA is a ser/thr protein kinase, it is clear that a tyrosine kinase should mediate the role of PKA. However, the identity of this tyrosine kinase has not been conclusively elucidated yet. One of the candidates proposed is cSrc<sup>165</sup>. However, sperm from cSrc null mice undergo tyrosine phosphorylation to similar levels than the ones observed in wild type sperm<sup>48,166</sup>. In addition, the cSrc family kinases (SFK) inhibitors SKI606 and SU6656 also blocked phosphorylation of PKA substrates. Interestingly, both PKA and tyrosine phosphorylation pathways were recovered in the presence of 0.1 nM of okadaic acid. This reversal of SFK inhibition by addition of okadaic acid strongly suggests that the role of SFK in capacitation is mediated by down-regulation of a ser/thr phosphatase. Consistently, it has been shown in other cell types that SFKs can inactivate PP2A by tyrosine phosphorylation of its C-terminus domain (TPDYFL)<sup>167,168,169</sup>.

More recently, a second family of tyrosine kinases has been proposed to mediate the role of PKA in sperm tyrosine phosphorylation<sup>170</sup>. The inhibitor PF-431396 has high specificity for members of the focal adhesion kinases, FAK1 and FAK2 (aka PYK2). It was shown that this inhibitor blocks the increase in tyrosine phosphorylation in horse sperm<sup>170</sup>. Moreover, anti-phospho antibodies recognizing activated FAK1 and PYK2 forms detected proteins in capacitated stallion sperm. More research is needed to understand the mechanism of action of these kinases and whether their action is downstream or upstream cAMP-dependent pathways.

## 8.3. Crosstalk between cAMP and $\text{Ca}^{2+}$ pathways

There is a complex interaction between the regulation of cAMP and  $\text{Ca}^{2+}$ -dependent pathways. In mouse sperm, the cAMP-dependent increase in tyrosine phosphorylation does not occur in  $\text{Ca}^{2+}$ -free media. However, in human sperm, the tyrosine phosphorylation is up-

regulated under these conditions<sup>171</sup>. These different effects of  $\text{Ca}^{2+}$  in the regulation of cAMP-dependent pathways are not surprising. On one hand,  $\text{Ca}^{2+}$  is known to activate sAC and consequently positively modulate cAMP synthesis; on the other hand,  $\text{Ca}^{2+}$  also activates calmodulin-dependent enzymes including PDE1; therefore  $\text{Ca}^{2+}$  can also stimulate cAMP degradation. Added to the  $\text{Ca}^{2+}$  modulation of cAMP synthesis and degradation, it has also been shown that addition of cGMP or cAMP induces a fast increase in intracellular  $\text{Ca}^{2+}$ <sup>147,148,172,173,174,175</sup>. Although this increase was initially believed to be mediated by cyclic nucleotide-dependent channels, it has been recently shown that cAMP and cGMP analogues can directly and promiscuously activate Catsper from outside the cell<sup>89</sup>. The same authors demonstrated that intracellular elevation of cAMP levels using either  $\text{HCO}_3^-$ , IBMX or caged cAMP compounds is unable to induce intracellular  $\text{Ca}^{2+}$  elevations<sup>105</sup>. Finally, it is well established using pharmacological and genetic knock-out approaches that  $\text{Ca}^{2+}$  and cAMP are required for the sperm to undergo hyperactivated motility necessary for spermatozoa to fertilize the egg. Therefore, a crosstalk between these pathways has been proposed. Consistent with a role of cAMP upstream of  $\text{Ca}^{2+}$  signaling, when intracellular  $\text{Ca}^{2+}$  is transiently increased by exposing sperm for 10 min to  $\text{Ca}^{2+}$  ionophore A23187, sperm acquired fertilizing capacity when incubated in either the absence of  $\text{HCO}_3^-$  or in the presence of H89<sup>34</sup>. These experiments indicate that  $\text{Ca}^{2+}$  can overcome the need for PKA activation during capacitation. Altogether, results from Brenker *et al*<sup>89</sup> indicates that there is no direct role of cAMP in the regulation of intracellular  $\text{Ca}^{2+}$ . However, similarly to the cAMP involvement in tyrosine phosphorylation,  $\text{Ca}^{2+}$  pathways required for the sperm to achieve fertilizing capacity could be indirectly regulated downstream of cAMP production in a slow time-scale (see above).

#### 8.4. Autocrine and paracrine cAMP signaling

As described throughout this review, many events that occur during capacitation rely on an increase in intracellular cAMP. Interestingly, the expression of members of the multidrug resistance-associated proteins (MRPs) family has been reported in mammalian sperm<sup>45</sup>. MRPs, also members of the ATP binding cassette (ABC) transporter subfamily C, are known to actively transport cAMP and cGMP and other small non-structurally-related compounds to the extracellular environment. One member of this family, MRP4 is localized to the sperm flagellum and to the post acrosomal region of bovine sperm<sup>45</sup>. In addition, this work showed that cAMP is extruded to the extracellular milieu and that MRP inhibition by probenecid abolished this efflux resulting in the accumulation of intracellular cAMP. Interestingly, these authors presented evidence indicating that extracellular cAMP mediates some of the events associated with sperm capacitation through the activation of adenosine receptors. The action of extracellular cAMP as a modulator of sperm capacitation may suggest a possible paracrine or autocrine modulation of the capacitation process. Further experimentation is necessary to understand this mechanism and its relevance in the physiological context of *in vivo* capacitation.

## 9. Conclusions

Initiated by the discovery of sperm capacitation more than 50 years ago, investigations in sperm physiology have been fundamental for the advances in reproductive medicine.

Despite these advances, many translational challenges remain in the clinic, in male contraception and in the development of reliable *in vitro* fertilization protocols for many mammalian species. The central role of sAC in the fast up-regulation of cAMP and consequently in sperm capacitation is well established. Figure 1 illustrates the main pathways of sperm capacitation where cAMP is directly involved. As mentioned throughout the review, the essential nature of cAMP in this process has been demonstrated using gain of function as well as loss of function approaches. However, many questions remain unanswered, among them: which is the molecular basis for sAC inactivation by PKA? What is the reason for the significant temporal lag between PKA activation and the onset of other downstream pathways? What are the identities of PKA substrates? How are the cAMP microdomains established and compartmentalized in the sperm? New research using novel technologies and molecular strategies that will help to answer these questions and understand the molecular mechanisms involved in sperm capacitation.

## Acknowledgements

This work was supported by NIH R01-TW008662 and Agencia Nacional de Promoción Científica y Tecnológica de Argentina PICT 2012-1175 (to MGB) and PICT 2011-0540 (to DK), NIH R01 HD44044 and HD038082 (to PEV), National Research Council of Argentina PIP 112-201101-00740 (to MGB and DK).

## References

1. Sutherland EW, Rall TW. Fractionation and Characterization of a Cyclic Adenine Ribonucleotide Formed by Tissue Particles. *J. Biol. Chem.* 1958; 232:1077–1091. [PubMed: 13549488]
2. Kawasaki H, Springett GM, Mochizuki N, Toki S, Nakaya M, Matsuda M, Housman DE, Graybiel AM. A Family of cAMP-Binding Proteins That Directly Activate Rap1. *Science.* 1998; 282:2275–2279. [PubMed: 9856955]
3. de Rooij J, Zwartkruis FJ, Verheijen MH, Cool RH, Nijman SM, Wittinghofer A, Bos JL. Epac Is a Rap1 Guanine-Nucleotide-Exchange Factor Directly Activated by Cyclic AMP. *Nature.* 1998; 396:474–477. [PubMed: 9853756]
4. Ludwig J, Margalit T, Eismann E, Lancet D, Kaupp UB. Primary Structure of cAMP-Gated Channel From Bovine Olfactory Epithelium. *FEBS Lett.* 1990; 270:24–29. [PubMed: 1699793]
5. Dessauer CW, Scully TT, Gilman AG. Interactions of Forskolin and ATP With the Cytosolic Domains of Mammalian Adenylyl Cyclase. *J. Biol. Chem.* 1997; 272:22272–22277. [PubMed: 9268376]
6. Buck J, Sinclair ML, Schapal L, Cann MJ, Levin LR. Cytosolic Adenylyl Cyclase Defines a Unique Signaling Molecule in Mammals. *Proc. Natl. Acad. Sci. U. S. A.* 1999; 96:79–84. [PubMed: 9874775]
7. Geng W, Wang Z, Zhang J, Reed BY, Pak CY, Moe OW. Cloning and Characterization of the Human Soluble Adenylyl Cyclase. *Am. J. Physiol Cell Physiol.* 2005; 288:C1305–C1316. [PubMed: 15659711]
8. Jaiswal BS, Conti M. Identification and Functional Analysis of Splice Variants of the Germ Cell Soluble Adenylyl Cyclase. *J. Biol. Chem.* 2001; 276:31698–31708. [PubMed: 11423534]
9. Chaloupka JA, Bullock SA, Iourgenko V, Levin LR, Buck J. Autoinhibitory Regulation of Soluble Adenylyl Cyclase. *Mol. Reprod. Dev.* 2006; 73:361–368. [PubMed: 16250004]
10. Chen Y, Cann MJ, Litvin TN, Iourgenko V, Sinclair ML, Levin LR, Buck J. Soluble Adenylyl Cyclase As an Evolutionarily Conserved Bicarbonate Sensor. *Science.* 2000; 289:625–628. [PubMed: 10915626]
11. Jaiswal BS, Conti M. Calcium Regulation of the Soluble Adenylyl Cyclase Expressed in Mammalian Spermatozoa. *Proc. Natl. Acad. Sci. U. S. A.* 2003; 100:10676–10681. [PubMed: 12958208]

12. Litvin TN, Kamenetsky M, Zarifyan A, Buck J, Levin LR. Kinetic Properties of "Soluble" Adenylyl Cyclase. Synergism Between Calcium and Bicarbonate. *J. Biol. Chem.* 2003; 278:15922–15926. [PubMed: 12609998]
13. Braun T, Dods RF. Development of a Mn<sup>2+</sup>-Sensitive, "Soluble" Adenylate Cyclase in Rat Testis. *Proc. Natl. Acad. Sci. U. S. A.* 1975; 72:1097–1101. [PubMed: 1055368]
14. Abou-haila A, Tulsiani DR. Signal Transduction Pathways That Regulate Sperm Capacitation and the Acrosome Reaction. *Arch. Biochem. Biophys.* 2009; 485:72–81. [PubMed: 19217882]
15. Aitken RJ, Baker MA. The Role of Proteomics in Understanding Sperm Cell Biology. *Int. J. Androl.* 2008; 31:295–302. [PubMed: 18179557]
16. Brewis IA, Gadella BM. Sperm Surface Proteomics: From Protein Lists to Biological Function. *Mol. Hum. Reprod.* 2010; 16:68–79. [PubMed: 19717474]
17. Darszon A, Labarca P, Nishigaki T, Espinosa F. Ion Channels in Sperm Physiology. *Physiol Rev.* 1999; 79:481–510. [PubMed: 10221988]
18. Fraser LR. The "Switching on" of Mammalian Spermatozoa: Molecular Events Involved in Promotion and Regulation of Capacitation. *Mol. Reprod. Dev.* 2010; 77:197–208. [PubMed: 19908247]
19. Buffone MG, Ijiri TW, Cao W, Merdiushev T, Aghajanian HK, Gerton GL. Heads or Tails? Structural Events and Molecular Mechanisms That Promote Mammalian Sperm Acrosomal Exocytosis and Motility. *Mol. Reprod. Dev.* 2012; 79:4–18. [PubMed: 22031228]
20. Lishko PV, Kirichok Y, Ren D, Navarro B, Chung JJ, Clapham DE. The Control of Male Fertility by Spermatozoan Ion Channels. *Annu. Rev. Physiol.* 2012; 74:453–475. [PubMed: 22017176]
21. Visconti PE, Krapf D, De LaVega-Beltran JL, Acevedo JJ, Darszon A. Ion Channels, Phosphorylation and Mammalian Sperm Capacitation. *Asian J. Androl.* 2011; 13:395–405. [PubMed: 21540868]
22. Bailey JL. Factors Regulating Sperm Capacitation. *Syst. Biol. Reprod. Med.* 2010; 56:334–348. [PubMed: 20849222]
23. Chang MC. Fertilizing Capacity of Spermatozoa Deposited into the Fallopian Tubes. *Nature.* 1951; 168:697–698. [PubMed: 14882325]
24. Austin CR. The Capacitation of the Mammalian Sperm. *Nature.* 1952; 170:326. [PubMed: 12993150]
25. Chang MC. Fertilization of Rabbit Ova in Vitro. *Nature.* 1959; 184(Suppl 7):466–467. [PubMed: 13809155]
26. Yanagimachi, R. *Physiology of Reproduction.* In: Knobil, E.; Neill, J., editors. *Mammalian Fertilization.* New York: Raven Press; 1994.
27. Branham MT, Mayorga LS, Tomes CN. Calcium-Induced Acrosomal Exocytosis Requires CAMP Acting Through a Protein Kinase A-Independent, Epac-Mediated Pathway. *J. Biol. Chem.* 2006; 281:8656–8666. [PubMed: 16407249]
28. Flesch FM, Brouwers JF, Nievelstein PF, Verkleij AJ, van Golde LM, Colenbrander B, Gadella BM. Bicarbonate Stimulated Phospholipid Scrambling Induces Cholesterol Redistribution and Enables Cholesterol Depletion in the Sperm Plasma Membrane. *J. Cell Sci.* 2001; 114:3543–3555. [PubMed: 11682613]
29. Gadella BM, Harrison RA. The Capacitating Agent Bicarbonate Induces Protein Kinase A-Dependent Changes in Phospholipid Transbilayer Behavior in the Sperm Plasma Membrane. *Development.* 2000; 127:2407–2420. [PubMed: 10804182]
30. Demarco IA, Espinosa F, Edwards J, Sosnik J, De La Vega-Beltran JL, Hockensmith JW, Kopf GS, Darszon A, Visconti PE. Involvement of a Na<sup>+</sup>/HCO<sub>3</sub><sup>-</sup> Cotransporter in Mouse Sperm Capacitation. *J. Biol. Chem.* 2003; 278:7001–7009. [PubMed: 12496293]
31. Wang D, King SM, Quill TA, Doolittle LK, Garbers DL. A New Sperm-Specific Na<sup>+</sup>/H<sup>+</sup> Exchanger Required for Sperm Motility and Fertility. *Nat. Cell Biol.* 2003; 5:1117–1122. [PubMed: 14634667]
32. Wang D, Hu J, Bobulescu IA, Quill TA, McLeroy P, Moe OW, Garbers DL. A Sperm-Specific Na<sup>+</sup>/H<sup>+</sup> Exchanger (SNHE) Is Critical for Expression and in Vivo Bicarbonate Regulation of the Soluble Adenylyl Cyclase (SAC). *Proc. Natl. Acad. Sci. U. S. A.* 2007; 104:9325–9330. [PubMed: 17517652]



33. Zeng Y, Oberdorf JA, Florman HM. PH Regulation in Mouse Sperm: Identification of Na(+)-, Cl(-), and HCO<sub>3</sub>(-)-Dependent and Arylamino benzoate-Dependent Regulatory Mechanisms and Characterization of Their Roles in Sperm Capacitation. *Dev. Biol.* 1996; 173:510–520. [PubMed: 8606009]
34. Tateno H, Krapf D, Hino T, Sanchez-Cardenas C, Darszon A, Yanagimachi R, Visconti PE. Ca<sup>2+</sup> Ionophore A23187 Can Make Mouse Spermatozoa Capable of Fertilizing in Vitro Without Activation of CAMP-Dependent Phosphorylation Pathways. *Proc. Natl. Acad. Sci. U. S. A.* 2013; 110:18543–18548. [PubMed: 24128762]
35. Visconti PE, Moore GD, Bailey JL, Leclerc P, Connors SA, Pan D, Olds-Clarke P, Kopf GS. Capacitation of Mouse Spermatozoa. II. Protein Tyrosine Phosphorylation and Capacitation Are Regulated by a CAMP-Dependent Pathway. *Development.* 1995; 121:1139–1150. [PubMed: 7538069]
36. Florman, HM.; Babcock, DF. Progress Toward Understanding the Molecular Basis of Capacitation. Chapter 5. In: Wassarman, PM., editor. *Elements of Mammalian Fertilization*. Vol. 1. Boca Raton, FL: CRC Press; 1991.
37. Branham MT, Bustos MA, De Blas GA, Rehmann H, Zarelli VE, Trevino CL, Darszon A, Mayorga LS, Tomes CN. Epac Activates the Small G Proteins Rap1 and Rab3A to Achieve Exocytosis. *J. Biol. Chem.* 2009; 284:24825–24839. [PubMed: 19546222]
38. Breitbart H, Spungin B. The Biochemistry of the Acrosome Reaction. *Mol. Hum. Reprod.* 1997; 3:195–202. [PubMed: 9237245]
39. de Lamirande E, Leclerc P, Gagnon C. Capacitation As a Regulatory Event That Primes Spermatozoa for the Acrosome Reaction and Fertilization. *Mol. Hum. Reprod.* 1997; 3:175–194. [PubMed: 9237244]
40. Lefievre L, Jha KN, de Lamirande E, Visconti PE, Gagnon C. Activation of Protein Kinase A During Human Sperm Capacitation and Acrosome Reaction.
41. Wertheimer E, Krapf D, De La Vega-Beltran JL, Sanchez-Cardenas C, Navarrete F, Haddad D, Escoffier J, Salicioni AM, Levin LR, Buck J, Mager J, Darszon A, Visconti PE. Compartmentalization of Distinct CAMP Signaling Pathways in Mammalian Sperm. *J. Biol. Chem.* 2013; 288:35307–35320. [PubMed: 24129574]
42. Harrison RA, Miller NG. CAMP-Dependent Protein Kinase Control of Plasma Membrane Lipid Architecture in Boar Sperm. *Mol. Reprod. Dev.* 2000; 55:220–228. [PubMed: 10618662]
43. Battistone MA, Da Ros VG, Salicioni AM, Navarrete FA, Krapf D, Visconti PE, Cuasnicu PS. Functional Human Sperm Capacitation Requires Both Bicarbonate-Dependent PKA Activation and Down-Regulation of Ser/Thr Phosphatases by Src Family Kinases. *Mol. Hum. Reprod.* 2013; 19:570–580. [PubMed: 23630234]
44. Harrison RA. Rapid PKA-Catalysed Phosphorylation of Boar Sperm Proteins Induced by the Capacitating Agent Bicarbonate. *Mol. Reprod. Dev.* 2004; 67:337–352. [PubMed: 14735495]
45. Osycka-Salut C, Diez F, Burdet J, Gervasi MG, Franchi A, Bianciotti LG, Davio C, Perez-Martinez S. Cyclic AMP Efflux, Via MRPs and A1 Adenosine Receptors, Is Critical for Bovine Sperm Capacitation. *Mol. Hum. Reprod.* 2014; 20:89–99. [PubMed: 23907162]
46. Visconti PE, Johnson LR, Oyaski M, Fornes M, Moss SB, Gerton GL, Kopf GS. Regulation, Localization, and Anchoring of Protein Kinase A Subunits During Mouse Sperm Capacitation. *Dev. Biol.* 1997; 192:351–363. [PubMed: 9441673]
47. Kaneto M, Krisfalusi M, Eddy EM, O'Brien DA, Miki K. Bicarbonate-Induced Phosphorylation of P270 Protein in Mouse Sperm by CAMP-Dependent Protein Kinase. *Mol. Reprod. Dev.* 2008; 75:1045–1053. [PubMed: 18357561]
48. Krapf D, Arcelay E, Wertheimer EV, Sanjay A, Pilder SH, Salicioni AM, Visconti PE. Inhibition of Ser/Thr Phosphatases Induces Capacitation-Associated Signaling in the Presence of Src Kinase Inhibitors. *J. Biol. Chem.* 2010; 285:7977–7985. [PubMed: 20068039]
49. Morgan DJ, Weisenhaus M, Shum S, Su T, Zheng R, Zhang C, Shokat KM, Hille B, Babcock DF, McKnight GS. Tissue-Specific PKA Inhibition Using a Chemical Genetic Approach and Its Application to Studies on Sperm Capacitation. *Proc. Natl. Acad. Sci. U. S. A.* 2008; 105:20740–20745. [PubMed: 19074277]

50. O'Flaherty C, de Lamirande E, Gagnon C. Phosphorylation of the Arginine-X-X-(Serine/Threonine) Motif in Human Sperm Proteins During Capacitation: Modulation and Protein Kinase A Dependency. *Mol. Hum. Reprod.* 2004; 10:355–363. [PubMed: 14997001]
51. O'Brien ED, Krapf D, Cabada MO, Visconti PE, Arranz SE. Transmembrane Adenylyl Cyclase Regulates Amphibian Sperm Motility Through Protein Kinase A Activation. *Dev. Biol.* 2011; 350:80–88. [PubMed: 21126515]
52. Hess KC, Jones BH, Marquez B, Chen Y, Ord TS, Kamenetsky M, Miyamoto C, Zippin JH, Kopf GS, Suarez SS, Levin LR, Williams CJ, Buck J, Moss SB. The "Soluble" Adenylyl Cyclase in Sperm Mediates Multiple Signaling Events Required for Fertilization. *Dev. Cell.* 2005; 9:249–259. [PubMed: 16054031]
53. Esposito G, Jaiswal BS, Xie F, Krajnc-Franken MA, Robben TJ, Strik AM, Kuil C, Philipsen RL, van Duin M, Conti M, Gossen JA. Mice Deficient for Soluble Adenylyl Cyclase Are Infertile Because of a Severe Sperm-Motility Defect. *Proc. Natl. Acad. Sci. U. S. A.* 2004; 101:2993–2998. [PubMed: 14976244]
54. Xie F, Garcia MA, Carlson AE, Schuh SM, Babcock DF, Jaiswal BS, Gossen JA, Esposito G, van Duin M, Conti M. Soluble Adenylyl Cyclase (SAC) Is Indispensable for Sperm Function and Fertilization. *Dev. Biol.* 2006; 296:353–362. [PubMed: 16842770]
55. Miki K, Willis WD, Brown PR, Goulding EH, Fulcher KD, Eddy EM. Targeted Disruption of the Akap4 Gene Causes Defects in Sperm Flagellum and Motility. *Dev. Biol.* 2002; 248:331–342. [PubMed: 12167408]
56. Miro-Moran A, Jardin I, Ortega-Ferrusola C, Salido GM, Pena FJ, Tapia JA, Aparicio IM. Identification and Function of Exchange Proteins Activated Directly by Cyclic AMP (Epac) in Mammalian Spermatozoa. *PLoS. One.* 2012; 7:e37713. [PubMed: 22662198]
57. Garbers DL, Lust WD, First NL, Lardy HA. Effects of phosphodiesterase inhibitors and cyclic nucleotides on sperm respiration and motility. *Biochemistry.* 1971; 10(10):1825–1831.
58. Sunahara RK, Dessauer CW, Whisnant RE, Kleuss C, Gilman AG. Interaction of G $\alpha$  With the Cytosolic Domains of Mammalian Adenylyl Cyclase. *J. Biol. Chem.* 1997; 272:22265–22271. [PubMed: 9268375]
59. Hepler JR, Gilman AG. G Proteins. *Trends Biochem. Sci.* 1992; 17:383–387. [PubMed: 1455506]
60. Kaslow HR, Burns DL. Pertussis Toxin and Target Eukaryotic Cells: Binding, Entry, and Activation. *FASEB J.* 1992; 6:2684–2690. [PubMed: 1612292]
61. Alfano M, Rizzi C, Corti D, Adduce L, Poli G. Bacterial Toxins: Potential Weapons Against HIV Infection. *Curr. Pharm. Des.* 2005; 11:2909–2926. [PubMed: 16101445]
62. Lahiri SS. Bacterial Toxins--an Overview. *J. Nat. Toxins.* 2000; 9:381–408. [PubMed: 11126516]
63. Glassner M, Jones J, Kligman I, Woolkalis MJ, Gerton GL, Kopf GS. Immunocytochemical and Biochemical Characterization of Guanine Nucleotide-Binding Regulatory Proteins in Mammalian Spermatozoa. *Dev. Biol.* 1991; 146:438–450. [PubMed: 1907582]
64. Ward CR, Storey BT, Kopf GS. Selective Activation of Gi1 and Gi2 in Mouse Sperm by the Zona Pellucida, the Egg's Extracellular Matrix. *J. Biol. Chem.* 1994; 269:13254–13258. [PubMed: 8175755]
65. Hildebrandt JD, Codina J, Tash JS, Kirchick HJ, Lipschultz L, Sekura RD, Birnbaumer L. The Membrane-Bound Spermatozoal Adenylyl Cyclase System Does Not Share Coupling Characteristics With Somatic Cell Adenylyl Cyclases. *Endocrinology.* 1985; 116:1357–1366. [PubMed: 3918851]
66. Hinsch KD, Habermann B, Just I, Hinsch E, Pfisterer S, Schill WB, Aktories K. ADP-Ribosylation of Rho Proteins Inhibits Sperm Motility. *FEBS Lett.* 1993; 334:32–36. [PubMed: 8224222]
67. Hinsch KD, Schwerdel C, Habermann B, Schill WB, Muller-Schlosser F, Hinsch E. Identification of Heterotrimeric G Proteins in Human Sperm Tail Membranes. *Mol. Reprod. Dev.* 1995; 40:345–354. [PubMed: 7772345]
68. Kopf GS, Woolkalis MJ, Gerton GL. Evidence for a Guanine Nucleotide-Binding Regulatory Protein in Invertebrate and Mammalian Sperm. Identification by Islet-Activating Protein-Catalyzed ADP-Ribosylation and Immunocytochemical Methods. *J. Biol. Chem.* 1986; 261:7327–7331. [PubMed: 3086310]

69. Bentley JK, Garbers DL, Domino SE, Noland TD, Van Dop C. Spermatozoa Contain a Guanine Nucleotide-Binding Protein ADP-Ribosylated by Pertussis Toxin. *Biochem. Biophys. Res. Commun.* 1986; 138:728–734. [PubMed: 3091021]
70. Fraser LR, Adeoya-Osiguwa S. Modulation of Adenylyl Cyclase by FPP and Adenosine Involves Stimulatory and Inhibitory Adenosine Receptors and g Proteins. *Mol. Reprod. Dev.* 1999; 53:459–471. [PubMed: 10398422]
71. Baxendale RW, Fraser LR. Immunolocalization of Multiple Galpha Subunits in Mammalian Spermatozoa and Additional Evidence for Galphas. *Mol. Reprod. Dev.* 2003; 65:104–113. [PubMed: 12658639]
72. Spehr M, Schwane K, Riffell JA, Barbour J, Zimmer RK, Neuhaus EM, Hatt H. Particulate Adenylate Cyclase Plays a Key Role in Human Sperm Olfactory Receptor-Mediated Chemotaxis. *J. Biol. Chem.* 2004; 279:40194–40203. [PubMed: 15271985]
73. Walensky LD, Snyder SH. Inositol 1,4,5-Trisphosphate Receptors Selectively Localized to the Acrosomes of Mammalian Sperm. *J. Cell Biol.* 1995; 130:857–869. [PubMed: 7642703]
74. Merlet F, Weinstein LS, Goldsmith PK, Rarick T, Hall JL, Bisson JP, de Mazancourt P. Identification and Localization of G Protein Subunits in Human Spermatozoa. *Mol. Hum. Reprod.* 1999; 5:38–45. [PubMed: 10050660]
75. Fiedler SE, Bajpai M, Carr DW. Identification and Characterization of RHOA-Interacting Proteins in Bovine Spermatozoa. *Biol. Reprod.* 2008; 78:184–192. [PubMed: 17928627]
76. Boatman DE, Robbins RS. Bicarbonate: Carbon-Dioxide Regulation of Sperm Capacitation, Hyperactivated Motility, and Acrosome Reactions. *Biol. Reprod.* 1991; 44:806–813. [PubMed: 1907858]
77. Lee MA, Storey BT. Bicarbonate Is Essential for Fertilization of Mouse Eggs: Mouse Sperm Require It to Undergo the Acrosome Reaction. *Biol. Reprod.* 1986; 34:349–356. [PubMed: 3082381]
78. Neill JM, Olds-Clarke P. A Computer-Assisted Assay for Mouse Sperm Hyperactivation Demonstrates That Bicarbonate but Not Bovine Serum Albumin Is Required. *Gamete Res.* 1987; 18:121–140. [PubMed: 3507366]
79. Shi QX, Roldan ER. Bicarbonate/CO<sub>2</sub> Is Not Required for Zona Pellucida- or Progesterone-Induced Acrosomal Exocytosis of Mouse Spermatozoa but Is Essential for Capacitation. *Biol. Reprod.* 1995; 52:540–546. [PubMed: 7756449]
80. Visconti PE, Bailey JL, Moore GD, Pan D, Olds-Clarke P, Kopf GS. Capacitation of Mouse Spermatozoa I. Correlation Between the Capacitation State and Protein Tyrosine Phosphorylation. *Development.* 1995; 121:1129–1137. [PubMed: 7743926]
81. Garbers DL, Tubb DJ, Hyne RV. A Requirement of Bicarbonate for Ca<sup>2+</sup>-Induced Elevations of Cyclic AMP in Guinea Pig Spermatozoa. *J. Biol. Chem.* 1982; 257:8980–8984. [PubMed: 6284752]
82. Okamura N, Sugita Y. Activation of Spermatozoan Adenylate Cyclase by a Low Molecular Weight Factor in Porcine Seminal Plasma. *J. Biol. Chem.* 1983; 258:13056–13062. [PubMed: 6630219]
83. Okamura N, Tajima Y, Soejima A, Masuda H, Sugita Y. Sodium Bicarbonate in Seminal Plasma Stimulates the Motility of Mammalian Spermatozoa Through Direct Activation of Adenylate Cyclase. *J. Biol. Chem.* 1985; 260:9699–9705. [PubMed: 2991260]
84. Garty NB, Galiani D, Aharonheim A, Ho YK, Phillips DM, Dekel N, Salomon Y. G-Proteins in Mammalian Gametes: an Immunocytochemical Study. *J. Cell Sci.* 1988; 91(Pt 1):21–31. [PubMed: 3150972]
85. Visconti PE, Muschietti JP, Flawia MM, Tezon JG. Bicarbonate Dependence of CAMP Accumulation Induced by Phorbol Esters in Hamster Spermatozoa. *Biochim. Biophys. Acta.* 1990; 1054:231–236. [PubMed: 2169311]
86. Kornblihtt AR, Flawia MM, Torres HN. Manganese Ion Dependent Adenylate Cyclase Activity in Rat Testes: Purification and Properties. *Biochemistry.* 1981; 20:1262–1267. [PubMed: 7225326]
87. Steegborn C, Litvin TN, Levin LR, Buck J, Wu H. Bicarbonate Activation of Adenylyl Cyclase Via Promotion of Catalytic Active Site Closure and Metal Recruitment. *Nat. Struct. Mol. Biol.* 2005; 12:32–37. [PubMed: 15619637]

88. Tresguerres M, Parks SK, Salazar E, Levin LR, Goss GG, Buck J. Bicarbonate-Sensing Soluble Adenylyl Cyclase Is an Essential Sensor for Acid/Base Homeostasis. *Proc. Natl. Acad. Sci. U. S. A.* 2010; 107:442–447. [PubMed: 20018667]
89. Brenker C, Goodwin N, Weyand I, Kashikar ND, Naruse M, Krahling M, Muller A, Kaupp UB, Strunker T. The CatSper Channel: a Polymodal Chemosensor in Human Sperm. *EMBO J.* 2012; 31:1654–1665. [PubMed: 22354039]
90. Burton KA, McKnight GS. PKA, Germ Cells, and Fertility. *Physiology.* (Bethesda. ). 2007; 22:40–46. [PubMed: 17289929]
91. Nolan MA, Babcock DF, Wennemuth G, Brown W, Burton KA, McKnight GS. Sperm-Specific Protein Kinase A Catalytic Subunit Calpha2 Orchestrates CAMP Signaling for Male Fertility. *Proc. Natl. Acad. Sci. U. S. A.* 2004; 101:13483–13488. [PubMed: 15340140]
92. Wertheimer EV, Salicioni AM, Liu W, Trevino CL, Chavez J, Hernandez-Gonzalez EO, Darszon A, Visconti PE. Chloride Is Essential for Capacitation and for the Capacitation-Associated Increase in Tyrosine Phosphorylation. *J. Biol. Chem.* 2008; 283:35539–35550. [PubMed: 18957426]
93. Zippin JH, Chen Y, Straub SG, Hess KC, Diaz A, Lee D, Tso P, Holz GG, Sharp GW, Levin LR, Buck J. CO<sub>2</sub>/HCO<sub>3</sub><sup>(-)</sup>- and Calcium-Regulated Soluble Adenylyl Cyclase As a Physiological ATP Sensor. *J. Biol. Chem.* 2013; 288:33283–33291. [PubMed: 24100033]
94. Han H, Stessin A, Roberts J, Hess K, Gautam N, Kamenetsky M, Lou O, Hyde E, Nathan N, Muller WA, Buck J, Levin LR, Nathan C. Calcium-Sensing Soluble Adenylyl Cyclase Mediates TNF Signal Transduction in Human Neutrophils. *J. Exp. Med.* 2005; 202:353–361. [PubMed: 16043520]
95. Baxendale RW, Fraser LR. Evidence for Multiple Distinctly Localized Adenylyl Cyclase Isoforms in Mammalian Spermatozoa. *Mol. Reprod. Dev.* 2003; 66:181–189. [PubMed: 12950106]
96. Livera G, Xie F, Garcia MA, Jaiswal B, Chen J, Law E, Storm DR, Conti M. Inactivation of the Mouse Adenylyl Cyclase 3 Gene Disrupts Male Fertility and Spermatozoon Function. *Mol. Endocrinol.* 2005; 19:1277–1290. [PubMed: 15705663]
97. Fraser LR, Duncan AE. Adenosine Analogues With Specificity for A<sub>2</sub> Receptors Bind to Mouse Spermatozoa and Stimulate Adenylate Cyclase Activity in Uncapacitated Suspensions. *J. Reprod. Fertil.* 1993; 98:187–194. [PubMed: 8393927]
98. Fraser LR, Dudley K. New Insights into the T-Complex and Control of Sperm Function. *Bioessays.* 1999; 21:304–312. [PubMed: 10377892]
99. Leclerc P, Kopf GS. Mouse Sperm Adenylyl Cyclase: General Properties and Regulation by the Zona Pellucida. *Biol. Reprod.* 1995; 52:1227–1233. [PubMed: 7632830]
100. Monks NJ, Stein DM, Fraser LR. Adenylate Cyclase Activity of Mouse Sperm During Capacitation in Vitro: Effect of Calcium and a GTP Analogue. *Int. J. Androl.* 1986; 9:67–76. [PubMed: 3744587]
101. Cheng CY, Boettcher B. Effects of Cholera Toxin and 5'-Guanylylimidodiphosphate on Human Spermatozoal Adenylate Cyclase Activity. *Biochem. Biophys. Res. Commun.* 1979; 91:1–9. [PubMed: 518608]
102. Forte LR, Bylund DB, Zahler WL. Forskolin Does Not Activate Sperm Adenylate Cyclase. *Mol. Pharmacol.* 1983; 24:42–47. [PubMed: 6683355]
103. Hanski E, Garty NB. Activation of Adenylate Cyclase by Sperm Membranes. The Role of Guanine Nucleotide Binding Proteins. *FEBS Lett.* 1983; 162:447–452. [PubMed: 6313428]
104. Rojas FJ, Bruzzone ME. Regulation of Cyclic Adenosine Monophosphate Synthesis in Human Ejaculated Spermatozoa. I. Experimental Conditions to Quantitate Membrane-Bound Adenylyl Cyclase Activity. *Hum. Reprod.* 1992; 7:1126–1130. [PubMed: 1328276]
105. Strunker T, Goodwin N, Brenker C, Kashikar ND, Weyand I, Seifert R, Kaupp UB. The CatSper Channel Mediates Progesterone-Induced Ca<sup>2+</sup> Influx in Human Sperm. *Nature.* 2011; 471:382–386. [PubMed: 21412338]
106. Seamon KB, Padgett W, Daly JW. Forskolin: Unique Diterpene Activator of Adenylate Cyclase in Membranes and in Intact Cells. *Proc. Natl. Acad. Sci. U. S. A.* 1981; 78:3363–3367. [PubMed: 6267587]

107. Seamon KB, Daly JW. Forskolin: a Unique Diterpene Activator of Cyclic AMP-Generating Systems. *J. Cyclic. Nucleotide. Res.* 1981; 7:201–224. [PubMed: 6278005]
108. Aitken RJ, Mattei A, Irvine S. Paradoxical Stimulation of Human Sperm Motility by 2-Deoxyadenosine. *J. Reprod. Fertil.* 1986; 78:515–527. [PubMed: 2433439]
109. Leclerc P, Kopf GS. Evidence for the Role of Heterotrimeric Guanine Nucleotide-Binding Regulatory Proteins in the Regulation of the Mouse Sperm Adenylyl Cyclase by the Egg's Zona Pellucida. *J. Androl.* 1999; 20:126–134. [PubMed: 10100483]
110. Duan B, Davis R, Sadat EL, Collins J, Sternweis PC, Yuan D, Jiang LI. Distinct Roles of Adenylyl Cyclase VII in Regulating the Immune Responses in Mice. *J. Immunol.* 2010; 185:335–344. [PubMed: 20505140]
111. Harayama H. Roles of Intracellular Cyclic AMP Signal Transduction in the Capacitation and Subsequent Hyperactivation of Mouse and Boar Spermatozoa. *J. Reprod. Dev.* 2013; 59:421–430. [PubMed: 24162806]
112. Mehats C, Andersen CB, Filopanti M, Jin SL, Conti M. Cyclic Nucleotide Phosphodiesterases and Their Role in Endocrine Cell Signaling. *Trends Endocrinol. Metab.* 2002; 13:29–35. [PubMed: 11750860]
113. Wasco WM, Orr GA. Function of Calmodulin in Mammalian Sperm: Presence of a Calmodulin-Dependent Cyclic Nucleotide Phosphodiesterase Associated With Demembrated Rat Caudal Epididymal Sperm. *Biochem. Biophys. Res. Commun.* 1984; 118:636–642. [PubMed: 6322756]
114. Baxendale RW, Fraser LR. Mammalian Sperm Phosphodiesterases and Their Involvement in Receptor-Mediated Cell Signaling Important for Capacitation. *Mol. Reprod. Dev.* 2005; 71:495–508. [PubMed: 15856425]
115. Fisch JD, Behr B, Conti M. Enhancement of Motility and Acrosome Reaction in Human Spermatozoa: Differential Activation by Type-Specific Phosphodiesterase Inhibitors. *Hum. Reprod.* 1998; 13:1248–1254. [PubMed: 9647555]
116. Carnegie GK, Means CK, Scott JD. A-Kinase Anchoring Proteins: From Protein Complexes to Physiology and Disease. *IUBMB. Life.* 2009; 61:394–406. [PubMed: 19319965]
117. Soberg K, Jahnsen T, Rognes T, Skalhegg BS, Laerdahl JK. Evolutionary Paths of the CAMP-Dependent Protein Kinase (PKA) Catalytic Subunits. *PLoS. One.* 2013; 8:e60935. [PubMed: 23593352]
118. Morris RC, Morris GZ, Zhang W, Gellerman M, Beebe SJ. Differential Transcriptional Regulation by the Alpha- and Gamma-Catalytic Subunit Isoforms of CAMP-Dependent Protein Kinase. *Arch. Biochem. Biophys.* 2002; 403:219–228. [PubMed: 12139971]
119. Vijayaraghavan S, Goueli SA, Davey MP, Carr DW. Protein Kinase A-Anchoring Inhibitor Peptides Arrest Mammalian Sperm Motility. *J. Biol. Chem.* 1997; 272:4747–4752. [PubMed: 9030527]
120. Visconti PE. Understanding the Molecular Basis of Sperm Capacitation Through Kinase Design. *Proc. Natl. Acad. Sci. U. S. A.* 2009; 106:667–668. [PubMed: 19144927]
121. Skalhegg BS, Huang Y, Su T, Idzerda RL, McKnight GS, Burton KA. Mutation of the Calpha Subunit of PKA Leads to Growth Retardation and Sperm Dysfunction. *Mol. Endocrinol.* 2002; 16:630–639. [PubMed: 11875122]
122. Desseyn JL, Burton KA, McKnight GS. Expression of a Nonmyristylated Variant of the Catalytic Subunit of Protein Kinase A During Male Germ-Cell Development. *Proc. Natl. Acad. Sci. U. S. A.* 2000; 97:6433–6438. [PubMed: 10841548]
123. Burton KA, Treash-Osio B, Muller CH, Dunphy EL, McKnight GS. Deletion of Type IIalpha Regulatory Subunit Delocalizes Protein Kinase A in Mouse Sperm Without Affecting Motility or Fertilization. *J. Biol. Chem.* 1999; 274:24131–24136. [PubMed: 10446185]
124. Bishop AC, Ubersax JA, Petsch DT, Matheos DP, Gray NS, Blethrow J, Shimizu E, Tsien JZ, Schultz PG, Rose MD, Wood JL, Morgan DO, Shokat KM. A Chemical Switch for Inhibitor-Sensitive Alleles of Any Protein Kinase. *Nature.* 2000; 407:395–401. [PubMed: 11014197]
125. Scott JD, Pawson T. Cell Signaling in Space and Time: Where Proteins Come Together and When They'Re Apart. *Science.* 2009; 326:1220–1224. [PubMed: 19965465]
126. Carr DW, Fujita A, Stentz CL, Liberty GA, Olson GE, Narumiya S. Identification of Sperm-Specific Proteins That Interact With A-Kinase Anchoring Proteins in a Manner Similar to the

- Type II Regulatory Subunit of PKA. *J. Biol. Chem.* 2001; 276:17332–17338. [PubMed: 11278869]
127. Vijayaraghavan S, Liberty GA, Mohan J, Winfrey VP, Olson GE, Carr DW. Isolation and Molecular Characterization of AKAP110, a Novel, Sperm-Specific Protein Kinase A-Anchoring Protein. *Mol. Endocrinol.* 1999; 13:705–717. [PubMed: 10319321]
  128. Mandal A, Naaby-Hansen S, Wolkowicz MJ, Klotz K, Shetty J, Retief JD, Coonrod SA, Kinter M, Sherman N, Cesar F, Flickinger CJ, Herr JC. FSP95, a Testis-Specific 95-Kilodalton Fibrous Sheath Antigen That Undergoes Tyrosine Phosphorylation in Capacitated Human Spermatozoa. *Biol. Reprod.* 1999; 61:1184–1197. [PubMed: 10529264]
  129. Carrera A, Gerton GL, Moss SB. The Major Fibrous Sheath Polypeptide of Mouse Sperm: Structural and Functional Similarities to the A-Kinase Anchoring Proteins. *Dev. Biol.* 1994; 165:272–284. [PubMed: 8088444]
  130. Carrera A, Moos J, Ning XP, Gerton GL, Tesarik J, Kopf GS, Moss SB. Regulation of Protein Tyrosine Phosphorylation in Human Sperm by a Calcium/Calmodulin-Dependent Mechanism: Identification of A Kinase Anchor Proteins As Major Substrates for Tyrosine Phosphorylation. *Dev. Biol.* 1996; 180:284–296. [PubMed: 8948591]
  131. Ficarro S, Chertihin O, Westbrook VA, White F, Jayes F, Kalab P, Marto JA, Shabanowitz J, Herr JC, Hunt DF, Visconti PE. Phosphoproteome Analysis of Capacitated Human Sperm. Evidence of Tyrosine Phosphorylation of a Kinase-Anchoring Protein 3 and Valosin-Containing Protein/P97 During Capacitation. *J. Biol. Chem.* 2003; 278:11579–11589. [PubMed: 12509440]
  132. Platt MD, Salicioni AM, Hunt DF, Visconti PE. Use of Differential Isotopic Labeling and Mass Spectrometry to Analyze Capacitation-Associated Changes in the Phosphorylation Status of Mouse Sperm Proteins. *J. Proteome. Res.* 2009; 8:1431–1440. [PubMed: 19186949]
  133. Harrison DA, Carr DW, Meizel S. Involvement of Protein Kinase A and A Kinase Anchoring Protein in the Progesterone-Initiated Human Sperm Acrosome Reaction. *Biol. Reprod.* 2000; 62:811–820. [PubMed: 10684828]
  134. Hillman P, Ickowicz D, Vizek R, Breitbart H. Dissociation Between AKAP3 and PKARII Promotes AKAP3 Degradation in Sperm Capacitation. *PLoS. One.* 2013; 8:e68873. [PubMed: 23894359]
  135. Newell AE, Fiedler SE, Ruan JM, Pan J, Wang PJ, Deininger J, Corless CL, Carr DW. Protein Kinase A RII-Like (R2D2) Proteins Exhibit Differential Localization and AKAP Interaction. *Cell Motil. Cytoskeleton.* 2008; 65:539–552. [PubMed: 18421703]
  136. Fiedler SE, Dudiki T, Vijayaraghavan S, Carr DW. Loss of R2D2 Proteins ROPN1 and ROPN1L Causes Defects in Murine Sperm Motility, Phosphorylation, and Fibrous Sheath Integrity. *Biol. Reprod.* 2013; 88:41. [PubMed: 23303679]
  137. Cukkemane A, Seifert R, Kaupp UB. Cooperative and Uncooperative Cyclic-Nucleotide-Gated Ion Channels. *Trends Biochem. Sci.* 2011; 36:55–64. [PubMed: 20729090]
  138. Brown RL, Strassmaier T, Brady JD, Karpen JW. The Pharmacology of Cyclic Nucleotide-Gated Channels: Emerging From the Darkness. *Curr. Pharm. Des.* 2006; 12:3597–3613. [PubMed: 17073662]
  139. Weitz D, Ficek N, Kremmer E, Bauer PJ, Kaupp UB. Subunit Stoichiometry of the CNG Channel of Rod Photoreceptors. *Neuron.* 2002; 36:881–889. [PubMed: 12467591]
  140. Pifferi S, Boccaccio A, Menini A. Cyclic Nucleotide-Gated Ion Channels in Sensory Transduction. *FEBS Lett.* 2006; 580:2853–2859. [PubMed: 16631748]
  141. Mazzolini M, Marchesi A, Giorgetti A, Torre V. Gating in CNGA1 Channels. *Pflugers Arch.* 2010; 459:547–555. [PubMed: 19898862]
  142. Darszon A, Acevedo JJ, Galindo BE, Hernandez-Gonzalez EO, Nishigaki T, Trevino CL, Wood C, Beltran C. Sperm Channel Diversity and Functional Multiplicity. *Reproduction.* 2006; 131:977–988. [PubMed: 16735537]
  143. Kaupp UB, Kashikar ND, Weyand I. Mechanisms of Sperm Chemotaxis. *Annu. Rev. Physiol.* 2008; 70:93–117. [PubMed: 17988206]
  144. Weyand I, Godde M, Frings S, Weiner J, Muller F, Altenhofen W, Hatt H, Kaupp UB. Cloning and Functional Expression of a Cyclic-Nucleotide-Gated Channel From Mammalian Sperm. *Nature.* 1994; 368:859–863. [PubMed: 7512693]


145. Kaupp UB, Seifert R. Cyclic Nucleotide-Gated Ion Channels. *Physiol Rev.* 2002; 82:769–824. [PubMed: 12087135]
146. Wiesner B, Weiner J, Middendorff R, Hagen V, Kaupp UB, Weyand I. Cyclic Nucleotide-Gated Channels on the Flagellum Control Ca<sup>2+</sup> Entry into Sperm. *J. Cell Biol.* 1998; 142:473–484. [PubMed: 9679145]
147. Kobori H, Miyazaki S, Kuwabara Y. Characterization of Intracellular Ca(2+) Increase in Response to Progesterone and Cyclic Nucleotides in Mouse Spermatozoa. *Biol. Reprod.* 2000; 63:113–120. [PubMed: 10859249]
148. Ren D, Navarro B, Perez G, Jackson AC, Hsu S, Shi Q, Tilly JL, Clapham DE. A Sperm Ion Channel Required for Sperm Motility and Male Fertility. *Nature.* 2001; 413:603–609. [PubMed: 11595941]
149. Biel M, Seeliger M, Pfeifer A, Kohler K, Gerstner A, Ludwig A, Jaissle G, Fauser S, Zrenner E, Hofmann F. Selective Loss of Cone Function in Mice Lacking the Cyclic Nucleotide-Gated Channel CNG3. *Proc. Natl. Acad. Sci. U. S. A.* 1999; 96:7553–7557. [PubMed: 10377453]
150. de Rooij J, Rehmann H, van Triest M, Cool RH, Wittinghofer A, Bos JL. Mechanism of Regulation of the Epac Family of CAMP-Dependent RapGEFs. *J. Biol. Chem.* 2000; 275:20829–20836. [PubMed: 10777494]
151. Amano R, Lee J, Goto N, Harayama H. Evidence for Existence of CAMP-Epac Signaling in the Heads of Mouse Epididymal Spermatozoa. *J. Reprod. Dev.* 2007; 53:127–133. [PubMed: 17077580]
152. McPartlin LA, Visconti PE, Bedford-Guaus SJ. Guanine-Nucleotide Exchange Factors (RAPGEF3/RAPGEF4) Induce Sperm Membrane Depolarization and Acrosomal Exocytosis in Capacitated Stallion Sperm. *Biol. Reprod.* 2011; 85:179–188. [PubMed: 21471298]
153. Parrish JJ, Susko-Parrish JL, First NL. Capacitation of Bovine Sperm by Heparin: Inhibitory Effect of Glucose and Role of Intracellular PH. *Biol. Reprod.* 1989; 41:683–699. [PubMed: 2620077]
154. Krahling AM, Alvarez L, Debowski K, Van Q, Gunkel M, Irsen S, Al Amoudi A, Strunker T, Kremmer E, Krause E, Voigt I, Wortge S, Waisman A, Weyand I, Seifert R, Kaupp UB, Wachten D. CRIS-a Novel CAMP-Binding Protein Controlling Spermiogenesis and the Development of Flagellar Bending. *PLoS. Genet.* 2013; 9:e1003960. [PubMed: 24339785]
155. Daleke DL. Regulation of Transbilayer Plasma Membrane Phospholipid Asymmetry. *J. Lipid Res.* 2003; 44:233–242. [PubMed: 12576505]
156. Tanaka K, Fujimura-Kamada K, Yamamoto T. Functions of Phospholipid Flippases. *J. Biochem.* 2011; 149:131–143. [PubMed: 21134888]
157. de Vries KJ, Wiedmer T, Sims PJ, Gadella BM. Caspase-Independent Exposure of Aminophospholipids and Tyrosine Phosphorylation in Bicarbonate Responsive Human Sperm Cells. *Biol. Reprod.* 2003; 68:2122–2134. [PubMed: 12606386]
158. Visconti PE, Ning X, Fornes MW, Alvarez JG, Stein P, Connors SA, Kopf GS. Cholesterol Efflux-Mediated Signal Transduction in Mammalian Sperm: Cholesterol Release Signals an Increase in Protein Tyrosine Phosphorylation During Mouse Sperm Capacitation. *Dev. Biol.* 1999; 214:429–443. [PubMed: 10525345]
159. Visconti PE, Galantino-Homer H, Ning X, Moore GD, Valenzuela JP, Jorgez CJ, Alvarez JG, Kopf GS. Cholesterol Efflux-Mediated Signal Transduction in Mammalian Sperm. Beta-Cyclodextrins Initiate Transmembrane Signaling Leading to an Increase in Protein Tyrosine Phosphorylation and Capacitation. *J. Biol. Chem.* 1999; 274:3235–3242. [PubMed: 9915865]
160. Hernandez-Gonzalez EO, Sosnik J, Edwards J, Acevedo JJ, Mendoza-Lujambio I, Lopez-Gonzalez I, Demarco I, Wertheimer E, Darszon A, Visconti PE. Sodium and Epithelial Sodium Channels Participate in the Regulation of the Capacitation-Associated Hyperpolarization in Mouse Sperm. *J. Biol. Chem.* 2006; 281:5623–5633. [PubMed: 16407190]
161. Travis AJ, Tutuncu L, Jorgez CJ, Ord TS, Jones BH, Kopf GS, Williams CJ. Requirements for Glucose Beyond Sperm Capacitation During *In Vitro* Fertilization in the Mouse. *Biol. Reprod.* 2004; 71:139–145. [PubMed: 14985248]

162. Leclerc P, de LE, Gagnon C. Cyclic Adenosine 3',5'Monophosphate-Dependent Regulation of Protein Tyrosine Phosphorylation in Relation to Human Sperm Capacitation and Motility. *Biol. Reprod.* 1996; 55:684–692. [PubMed: 8862788]
163. Mahony MC, Gwathmey T. Protein Tyrosine Phosphorylation During Hyperactivated Motility of Cynomolgus Monkey (*Macaca Fascicularis*) Spermatozoa. *Biol. Reprod.* 1999; 60:1239–1243. [PubMed: 10208990]
164. Si Y, Okuno M. Role of Tyrosine Phosphorylation of Flagellar Proteins in Hamster Sperm Hyperactivation. *Biol. Reprod.* 1999; 61:240–246. [PubMed: 10377055]
165. Baker MA, Hetherington L, Aitken RJ. Identification of SRC As a Key PKA-Stimulated Tyrosine Kinase Involved in the Capacitation-Associated Hyperactivation of Murine Spermatozoa. *J. Cell Sci.* 2006; 119:3182–3192. [PubMed: 16835269]
166. Krapf D, Ruan YC, Wertheimer EV, Battistone MA, Pawlak JB, Sanjay A, Pilder SH, Cuasnicu P, Breton S, Visconti PE. CSrc Is Necessary for Epididymal Development and Is Incorporated into Sperm During Epididymal Transit. *Dev. Biol.* 2012; 369:43–53. [PubMed: 22750823]
167. Chen J, Martin BL, Brautigam DL. Regulation of Protein Serine-Threonine Phosphatase Type-2A by Tyrosine Phosphorylation. *Science.* 1992; 257:1261–1264. [PubMed: 1325671]
168. Chen J, Parsons S, Brautigam DL. Tyrosine Phosphorylation of Protein Phosphatase 2A in Response to Growth Stimulation and V-Src Transformation of Fibroblasts. *J. Biol. Chem.* 1994; 269:7957–7962. [PubMed: 7510677]
169. Hu X, Wu X, Xu J, Zhou J, Han X, Guo J. Src Kinase Up-Regulates the ERK Cascade Through Inactivation of Protein Phosphatase 2A Following Cerebral Ischemia. *BMC. Neurosci.* 2009; 10:74. [PubMed: 19602257]
170. Gonzalez-Fernandez L, Macias-Garcia B, Loux SC, Varner DD, Hinrichs K. Focal Adhesion Kinases and Calcium/Calmodulin-Dependent Protein Kinases Regulate Protein Tyrosine Phosphorylation in Stallion Sperm. *Biol. Reprod.* 2013; 88:138. [PubMed: 23595906]
171. Luconi M, Krausz C, Forti G, Baldi E. Extracellular Calcium Negatively Modulates Tyrosine Phosphorylation and Tyrosine Kinase Activity During Capacitation of Human Spermatozoa. *Biol. Reprod.* 1996; 55:207–216. [PubMed: 8793077]
172. Liu J, Xia J, Cho KH, Clapham DE, Ren D. CatSperbeta, a Novel Transmembrane Protein in the CatSper Channel Complex. *J. Biol. Chem.* 2007; 282:18945–18952. [PubMed: 17478420]
173. Machado-Oliveira G, Lefievre L, Ford C, Herrero MB, Barratt C, Connolly TJ, Nash K, Morales-Garcia A, Kirkman-Brown J, Publicover S. Mobilisation of Ca<sup>2+</sup> Stores and Flagellar Regulation in Human Sperm by S-Nitrosylation: a Role for NO Synthesised in the Female Reproductive Tract. *Development.* 2008; 135:3677–3686. [PubMed: 18842814]
174. Xia J, Reigada D, Mitchell CH, Ren D. CATSPER Channel-Mediated Ca<sup>2+</sup> Entry into Mouse Sperm Triggers a Tail-to-Head Propagation. *Biol. Reprod.* 2007; 77:551–559. [PubMed: 17554080]
175. Xia J, Ren D. Egg Coat Proteins Activate Calcium Entry into Mouse Sperm Via CATSPER Channels. *Biol. Reprod.* 2009; 80:1092–1098. [PubMed: 19211808]



### Highlights

1. Cyclic AMP is essential in sperm physiology
2. Soluble adenylyl cyclase (sAC) is the main source of cAMP in mammalian sperm cells
3. Soluble AC knock-out mice are infertile
4. Signaling events associated with cAMP are highly compartmentalized within the sperm



**Figure 1. Model of main pathways of sperm capacitation where cAMP is directly involved**  
 Model of cAMP pathways acting during mammalian sperm capacitation. Two different sources of cAMP orchestrate signaling pathways in the sperm head (A) and in the principal piece (B). In the head, a tmAC increases cAMP levels which act primarily on EPAC, leading to the preparation to undergo the acrosome reaction. In the sperm flagellum, cAMP is increased through the synergic effect of Ca<sup>2+</sup> and HCO<sub>3</sub><sup>-</sup> stimulation of sAC. The increased pool of cAMP acts on PKA, as well as on CNG channels and NHE channels.

These molecules impact directly on the tyrosine phosphorylation pathway that correlates to sperm capacitation. See text for details.

**Table I**

## cAMP in sperm physiology

<b>role</b>	<b>Refs</b>
cAMP regulation of capacitation	28,29,30,31,32,33,34
cAMP regulation of acrosome reaction	27,37,38,39,40,41,80
Membrane lipid remodeling	28,29,157
Hyperpolarization of sperm plasma membrane	30
Increase in intracellular pH	30,31,32,33
Increase in intracellular Ca <sup>2+</sup>	34,147,148,172,173,174, 175
Increase in protein tyrosine phosphorylation	35
cAMP role in ATP synthesis	54

Table II

molecules involved in cAMP regulation

Molecule	Conclusion	Experimental Approach	Refs
<b>G proteins</b>	present	WB	70
		CTX and cAMP levels measurements	71
		CTX ADP-ribosylation	95
		Proteomics	72
		ADP-ribosylation, IP, WB, IF, acrosome reaction and $[Ca^{2+}]_i$ levels measurements	41
	absent	CTX ADP-ribosylation	65
		WB	66,67
		Lubrol PX extraction and PTX ADP ribosylation, WB, IP	68,69
		PTX ADP-ribosylation	65
		WB	68,69
<b>Gi</b>	present	Immunodetection	73,74
	absent	WB	75
	present	WB	6,13,86
	present	Purification from testis, microsequencing and mass spectrometry and cloning	
<b>Gα13 (akaGNA13)</b>	present	IP, cyclase activity	10
	present	Loss of function sAC KO and specific inhibitors, gain of function experiments with cAMP permeable analogs	52,53
<b>Soluble Adenylyl Cyclase</b>	present	Enzymatic and immunological assays	95
		Mass spectrometry	72
		Loss of function AC3 KO	96
		FSK stimulation and cAMP levels determination	41,71,95,96
		FSK stimulation and AC activity determination, use of GTP analogs	99,109
<b>Transmembrane adenylyl cyclases</b>	absent	FSK stimulation and cAMP levels determination	11,52,89,104,105,108
	present	Immunodetection, PDE inhibitors, CTC assay, acrosome reaction measurements	111,121,113
	present	WB, immunofluorescence, loss of function by MRP inhibitors and extracellular cAMP levels determination	45
<b>Phosphodiesterases</b>	PDEs 1, 4, 6, 8, 10 and 11		
<b>MRP</b>			

Abbreviations: WB: Western Blot; CTX: cholera toxin; IP: immunoprecipitation; IF: immunofluorescence; IP: immunoprecipitation; FSK: forskolin; CTC: clortetracycline; PTX: Pertussin toxin

Table III

cAMP targets

Molecule	assay	experimental approach	Refs
<b>PKA</b>	Presence in sperm flagella	WB, IF, PKA activity and cAMP measurements	41
	Activation during capacitation	WB	44,47,48,49,50,51
	Dependence on sAC	Loss of function sAC KO	52
	Role in tyrosine phosphorylation	Loss and gain of function experiments with PKA Inhibitors and cAMP permeable analogs	35
<b>CNGs</b>	Role in capacitation and sperm motility	Loss of function by pharmacological and genetic knock-out approaches targeting sAC, the PKA catalytic subunit or the anchoring of PKA to AKAPs	35,49,91, 119
	Presence in sperm	Cloning, electrophysiological characterization, Ca <sup>2+</sup> imaging and patch clamp recording	144,145
	Role in chemotaxis	Sperm motility recording, fluorescent dyes, Caged compounds and flash photolysis techniques	142,143
	Increase [Ca <sup>2+</sup> ] <sub>i</sub> in sperm	Ca <sup>2+</sup> imaging	146,147
<b>EPAC</b>	Presence in sperm and involvement in acrosome reaction	Gain of function by EPAC-selective cAMP analogs, acrosome reaction measurements, WB, Immunofluorescence	27,56,151,152
	Presence in sperm and role in internal alkalization and hyperpolarization	Loss of function by NHE-targeted deletion	31
<b>NHE</b>	Role in sAC expression during spermiogenesis	Experiments with pH <sub>i</sub> -sensitive fluorescence probes	21
		Loss of function NHE KO	32

Abbreviations: WB: Western Blot; IF: immunofluorescence