



CrossMark  
click for updates

## Research

**Cite this article:** Martínez RN, Apaldetti C, Colombi CE, Praderio A, Fernandez E, Malnis PS, Correa GA, Abelin D, Alcober O. 2013 A new sphenodontian (Lepidosauria: Rhynchocephalia) from the Late Triassic of Argentina and the early origin of the herbivore opisthodontians. *Proc R Soc B* 280: 20132057. <http://dx.doi.org/10.1098/rspb.2013.2057>

Received: 7 August 2013

Accepted: 19 September 2013

### Subject Areas:

palaeontology

### Keywords:

Lepidosauria, Sphenodontia, Opisthodontia, Upper Triassic, Argentina

### Author for correspondence:

Ricardo N. Martínez

e-mail: [martinez@unsj.edu.ar](mailto:martinez@unsj.edu.ar)

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rspb.2013.2057> or via <http://rspb.royalsocietypublishing.org>.

# A new sphenodontian (Lepidosauria: Rhynchocephalia) from the Late Triassic of Argentina and the early origin of the herbivore opisthodontians

Ricardo N. Martínez<sup>1</sup>, Cecilia Apaldetti<sup>1,2</sup>, Carina E. Colombi<sup>1,2</sup>, Angel Praderio<sup>1</sup>, Eliana Fernandez<sup>1,2</sup>, Paula Santi Malnis<sup>1,2</sup>, Gustavo A. Correa<sup>1,2</sup>, Diego Abelin<sup>1</sup> and Oscar Alcober<sup>1</sup>

<sup>1</sup>Instituto y Museo de Ciencias Naturales, Universidad Nacional de San Juan, Avenida España 400 Norte, 5400 San Juan, Argentina

<sup>2</sup>Consejo Nacional de Investigaciones Científicas y Técnicas, CONICET, Buenos Aires, Argentina

Sphenodontians were a successful group of rhynchocephalian reptiles that dominated the fossil record of Lepidosauria during the Triassic and Jurassic. Although evidence of extinction is seen at the end of the Laurasian Early Cretaceous, they appeared to remain numerically abundant in South America until the end of the period. Most of the known Late Cretaceous record in South America is composed of opisthodontians, the herbivorous branch of Sphenodontia, whose oldest members were until recently reported to be from the Kimmeridgian–Tithonian (Late Jurassic). Here, we report a new sphenodontian, *Sphenotitan leyesi* gen. et sp. nov., collected from the Upper Triassic Quebrada del Barro Formation of northwestern Argentina. Phylogenetic analysis identifies *Sphenotitan* as a basal member of Opisthodontia, extending the known record of opisthodontians and the origin of herbivory in this group by 50 Myr.

## 1. Introduction

Lepidosaurian reptiles, comprising Squamata and Rhynchocephalia, constitute a large proportion of extant terrestrial vertebrate fauna. Squamates include more than 7000 extant species (lizards, snakes and amphisbaenas), whereas the only living rhynchocephalian is *Sphenodon punctatus* from New Zealand [1]. Contrary to the extant record, the fossil record demonstrates that the sphenodontians—the most successful group of rhynchocephalian reptiles—were the dominant lepidosaurians during the Triassic and Jurassic [1–7]. During the early Cretaceous, sphenodontians nearly became extinct in Laurasia, replaced by the diversifying squamates [8]. By contrast, South American sphenodontians remained abundant until at least the end of the Campanian, and it was only during the Palaeogene that squamates began to dominate small reptilian fauna [9]. Most Argentinean Upper Cretaceous sphenodontians are eilenodontine opisthodontians [9–11], a herbivorous branch of Sphenodontia characterized by lateromedially expanded dentary teeth and advanced propalinal jaw movements. The opisthodontians have been observed across several regions and periods, from the Upper Jurassic of North America [12–14], Europe [15] and South Africa [16], to the Upper Cretaceous of South America [9]. Until now, the oldest known opisthodontian was *Opisthias*, a widely distributed genus from the Kimmeridgian–Tithonian of North America, and Kimmeridgian and Berriasian of Europe [12–16]. Here, we report partial skeletons of a new opisthodontian from the Upper Triassic of South America, which extends the oldest record of this group by 50 Myr.

The specimens described here were collected from the Upper Triassic Quebrada del Barro Formation in northwestern Argentina (see electronic supplementary material, figure S1). This unit was originally dated as Norian [17], although that age was later questioned [18]. A recent stratigraphic

survey and the discovery of a new vertebrate association confirm the Norian age of the Quebrada del Barro Formation [19] (see electronic supplementary material).

## 2. Systematic palaeontology

Lepidosauria Haeckel, 1866 [20]

Rhynchocephalia Günther, 1867 [21]

Sphenodontia Williston, 1925 [22]

Opisthodontia Apesteguía and Novas, 2003 [9]

*Sphenotitan leyesi* gen. et sp. nov.

### (a) Etymology

*Spheno* refers to Sphenodontia. *Titan*, meaning giant (Greek), refers to the large size of the new taxon (largest specimen of *Sphenotitan* reached more than 100 mm in skull length). The species name is dedicated to the Leyes family from the small town of Balde de Leyes, which is located close to the type locality.

### (b) Holotype

PVSJ 886 (Museo de Ciencias Naturales, Universidad Nacional de San Juan, San Juan, Argentina) consists of a partially articulated posterior portion of the skull of an adult individual. The specimen preserves partial pterygoids, ectopterygoids, palatines, maxillae, right premaxilla, parietals, squamosals, quadrates, left jugal and postorbital, and the rear half of the lower jaws, from the level of posterior-most dentary teeth.

### (c) Referred material


PVSJ 887, left dentary; PVSJ 888, both incomplete maxillae, anterior part of both lower jaws, both premaxillae; PVSJ 889, left maxilla; PVSJ 890, right quadrate; PVSJ 891, incomplete lower jaws; PVSJ 892, right palatine, pterygoid and ectopterygoid; PVSJ 893, right premaxilla and postorbital, left incomplete palatine and pterygoid; PVSJ 895, articulated sequence of three distal dorsal vertebrae, sacral vertebrae 1 and 2, and first caudal vertebra.

### (d) Locality and horizon

Upper layers of the Quebrada del Barro Formation (Norian) of the Marayes–El Carrizal Basin, ‘Balde de Leyes’ fossil locality, San Juan Province, Argentina (see electronic supplementary material, figure S1). The specimens were found in fluvial sandstones, in association with basal sauropodomorph dinosaurs, tritheledontid cynodonts and ‘sphenosuchid’ pseudosuchians.

### (e) Diagnosis

*Sphenotitan* differs from other sphenodontians in the following autapomorphies: large supratemporal fenestra—one-third of the skull length; posterodorsal process of the premaxilla, which ends posteriorly as a projection wedged between the anterolateral process of the nasal and the premaxillary process of the maxilla; palatine bearing a small cluster of 15 conical teeth lying medial to the main tooth row and posterior to the internal naris; four prominent rows of dozens of small teeth on the anterior palatal shelf of the pterygoids; large quadrate–quadratojugal foramen, consisting of more than one-third of the maximum length of the quadrate; straight orbital border of the maxilla; long posterior maxillary teeth, reaching 90% of the total dorsoventral height of the maxilla;




**Figure 1.** *Sphenotitan leyesi* gen. et sp. nov. (a) Right maxilla and premaxilla (PVSJ 886) in lateral view. (b) Left maxilla (PVSJ 889) in occlusal view. (c) Anterior palatal fragment (PVSJ 892) in palatal view. (d) Posterior palatal fragment (PVSJ 886) in palatal view. Scale bars equal 10 mm. White dashed lines indicate sutures. ctpt, cluster of pterygoid teeth; ec, ectopterygoid; mx, maxilla; pal, palatine; palt, palatine teeth; pdp, posterodorsal process of the premaxilla; pmx, premaxilla; pt, pterygoid; ptt, pterygoid teeth; wf, wear facet.

pronounced, stout, square-section and dorsally curved retro-articular process of the lower jaw; additional wide teeth with a transverse width : anteroposterior length ratio of five.

## 3. Description

The skull of *Sphenotitan* is lightly built, roughly triangular in dorsal view and characterized by the presence of large supratemporal fenestrae—making up over one-third of the skull length—separated by very narrow and fused parietals, and a short preorbital region (figures 1–4). Adult specimens of *Sphenotitan* reached more than 100 mm in skull length, larger than any previously known sphenodontian besides the eilenodontine *Kaikaifilusauros* [9].

The premaxillae of both small and large specimens of *Sphenotitan* form a beak comprising a single chisel-like cutting edge (figures 1a and 4c), similar to that of *Kaikaifilusauros* and very large specimens of *Clevosaurus* [23]. The posterodorsal process of the premaxilla is long, excluding the maxilla from the margin of the external naris (figures 1a and 4c) as seen




**Figure 2.** *Sphenotitan leyesi* gen. et sp. nov. (a) Fused parietals and right squamosal (PVSJ 886) in dorsal view. (b) Left quadrate (PVSJ 890) in posterior view. Scale bars equal 10 mm. p, parietal; pc, parietal crest; pf, parietal foramen; qf, quadrate foramen; sq, squamosal.

in *Clevosaurus*, *Godavarisaurus* [24], *Pamizinsaurus* [25] and *Kaikaifilusaurus*. However, this differs from all other sphenodontians in that the posterodorsal process of the premaxilla is curved and hook-like, and wedged between the anterolateral process of the nasal and the premaxillary process of the maxilla (figures 1a and 4c).

The maxilla is stout, with a short premaxillary process (figure 1a). Over half of the maxillary length forms the straight lower border of the orbit. The maxilla of a large individual of *Sphenotitan* has 17–18 long, closely packed teeth, anteriorly decreasing in size, with long anterolateral flanges (figure 1a,b). The maxillary teeth vary from 8.7 mm long and 2.2 mm wide for the posterior-most to 2.6 mm long and 1.6 mm wide for the anterior-most, which are the most worn (figure 1a,b). The ratio of the length of the longest tooth to the corresponding dorsoventral depth of the maxilla is 0.9, greater than in any other sphenodontian (which are generally less than 0.3).

The palatine widens anteriorly and bears a row of at least 13 large teeth on its lateral margin, located parallel to those of the maxilla (figures 1c,d and 4b). The palatal teeth are blade-like, lateromedially wide, posteriorly pointed, and with extended lateral and medial flanges. The size of teeth increases posteriorly in a manner similar to that of *Kaikaifilusaurus* (although in the latter, the row of teeth is straight while in the former it is markedly curved with lateral convexity). Anteromedially, the palatine also bears a small cluster of 15 conical teeth lying medial to the main tooth row and posterior to the internal naris (figures 1c and 4b)—a similar location to that of the isolated tooth present in *Clevosaurus* [23]. These small palatine teeth are closely packed in a subcircular area 2 mm wide (figures 1c and 4b). The pterygoids have broad anterior contact with each other and are excluded from the margin of the suborbital fenestra (figures 1d and 4b), as in most other derived sphenodontians (e.g. *Kaikaifilusaurus*, *Clevosaurus* and *Sphenodon*). The palatal shelf bears four prominent rows of dozens of small teeth (figures 1d and 4b). Each pterygoid has two anteroposteriorly oriented double rows of teeth that are roughly parallel to each other (figures 1d and 4b). The lateral double-row, which ends more posteriorly than the medial double-row, caudally ends in a small subcircular cluster of 12 very small teeth (figures 1d and 4b). The arrangement of two double rows of pterygoid teeth and the aforementioned cluster of palatine teeth are unique features among Rhynchocephalia.




**Figure 3.** Lower jaw of *S. leyesi* gen. et sp. nov. Left dentary (PVSJ 887) in (a) lateral, (b) medial and (c) occlusal view. (d) Posterior portion of the right lower jaw (PVSJ 886) in occlusal view. Scale bar equals 10 mm. adfos, adductor fossa; adt, additional teeth; ar, articular; cp, coronoid process; d, dentary; dpp, dentary posterior process; mdfo, mandibular foramen; mkc, meckelian channel; rp, retroarticular process; sa, surangular; sas, surface for articulation with surangular; sy, symphysis.

The quadrate–quadratojugal foramen, located between the quadrate and the quadratojugal (figure 2b) as in *Sphenodon* and *Zapatadon* [26], is proportionally large—over one-third of the maximum length of the quadrate (figure 2b). The size of the foramen is larger than that of other rhynchocephalians (except for the dorsoventrally reduced quadrate of *Sphenodon* [27]).

The lower jaw is stout and deep (figures 3 and 4c,d), as in all opisthodontians (e.g. *Opisthias* [12], *Kaikaifilusaurus* [9,10] and *Toxolophosaurus* [13,28]). *Sphenotitan* also shares the anterior edentulous area of the dentary and the absence of dental regionalization with opisthodontians, lacking caniniform and hatchling teeth (figure 3a–c). Tooth implantation is acrodont without evidence of replacement. As in all eilenodontines (e.g. *Eilenodon*, *Toxolophosaurus* and *Kaikaifilusaurus*), in *Pelecymala* [3] and *Clevosaurus latidens* [29] the teeth of the dentary are transversally wide, uniform in shape, closely packed and increase in size progressively towards the posterior additional series (figures 3 and 4d). However, unlike eilenodontines, the teeth of *Sphenotitan* are prism-like, with straight labiolingual cutting edges parallel to each other. The teeth are extremely wide (labiolingual width : mesiodistal length ratio = 5), closely packed (5 teeth  $\text{cm}^{-1}$ ) and lack lateral flanges (figures 3 and 4d). The lateral surfaces of the dentary teeth have vertical wear facets as a result of the shearing action of the maxillary dentition (figures 3a,c and 5), similar to that reported in *Pelecymala* and *C. latidens*. The continuous rasp-like surface formed by the row of teeth faces dorso-medially at an angle of  $45^\circ$  with respect to the horizontal at the posterior-most region, and decreases progressively toward the anterior end. The coronoid process is moderately high, triangular and anterodorsally pointed (figures 3a,b and 4c),






**Figure 4.** Reconstruction of the skull and lower jaw of a mid-range size individual of *S. leyesi* gen. et sp. nov. in (a) lateral, (b) ventral and (c) dorsal view. (d) Lower jaw in occlusal view. Scale bar equals 20 mm. Dark grey tone indicates cavities. Light grey tone indicates missing fragments. a, angular; ar, articular; c, coronoid; ctpt, cluster of pterygoid teeth; d, dentary; ec, ectopterygoid; f, frontal; j, jugal; mx, maxilla; n, nasal; p, parietal; pa, prearticular; pal, palatine; palt, palatine teeth; pf, prefrontal; pfo, parietal foramen; pmx, premaxilla; po, postorbital; pof, postfrontal; pt, pterygoid; ptt, pterygoid teeth; q, quadrate; qj, quadratojugal; rp, retroarticular process; sa, surangular; sq, squamosal.

similar to that of *Kallimodon* and *Sapheosaurus* [2], but different from the low coronoid process of all opisthodontians. The glenoid cavity is markedly symmetrical with a central anteroposteriorly ridge (figures 3d and 4d) as in the eilenodontines *Eilenodon* and *Kaikaiifilusaurus* (although in the latter the glenoid is anteroposteriorly longer). The retroarticular process is pronounced as in basal sphenodontians, but in *Sphenotitan* it is dorsally curved and very stout, with uniformly roughly squared cross-sections along its entire length (figures 3d and 4c,d).

#### 4. Phylogenetic analysis


The phylogenetic relationship of *S. leyesi* was analysed by adding it and the Triassic taxa *C. latidens* and *Pelecymala* to the data matrix of Apesteguía *et al.* [7] (see the electronic supplementary materials).

The strict consensus tree (see electronic supplementary material, figure S3) depicted *Sphenotitan* as the basal-most



**Figure 5.** Schematic posterior view cross-section through the snout level of *S. leyesi* gen. et sp. nov. Scale bar equals 10 mm. d, dentary; dt, dentary teeth; mx, maxilla; mxt, maxillary teeth; pal, palatine; palt, palatine teeth; pt, pterygoid; ptt, pterygoid teeth.

opisthodontian sphenodontian related to the Upper Jurassic *Opisthias*, *Toxolophosaurus* and *Eilenodon*, and to the Upper Cretaceous *Kaikaiifilusaurus*. In this context, opisthodontians share five synapomorphies: a rounded and well-developed mandibular symphysis (Character 34.2), a well-developed



**Figure 6.** Calibrated reduced strict consensus tree summarizing the results obtained in the constrained searches. Calibrated phylogeny of sphenodontians, indicating different tooth morphotypes and dietary interpretations.

anterodorsally projecting mandibular spur (Character 37.1), dentary regionalization absent in juveniles and adults (Character 45.3), posteromedial flanges on posterior maxillary teeth present as extensive flanges (Character 51.2) and the presence of anteromedial flanges on mandibular teeth (Character 58.1). With the exception of *Sphenotitan*, all opisthodontians (*Opisthias* and more derived Opisthodontia) share an unambiguous synapomorphy: the presence of dental ridges on additional mandibular teeth (Character 59.1). The presence of the plesiomorphic state in that character—the absence of dental ridges in additional mandibular teeth (Character 59.0)—supports the basal position of *Sphenotitan* within Opisthodontia.

This study classifies Opisthodontia as part of a polytomy (see electronic supplementary material, figure S3) that includes several derived crown-sphenodontian taxa (i.e. *Pelecymala*, *C. latidens*, *Ankylosphenodon*, *Kawasphenodon*, *Sphenodon*, *Cynosphenodon*, *Zapatodon*, *Homoeosaurus*, *Sphenovipera*, *Theretairus*, *Sapeosaurus* and *Kallimodon*). In order to evaluate the cause of this polytomy, pruned trees were compared to obtain a reduced consensus tree [30] (see electronic supplementary

material). The result shows three unstable taxa: *Pelecymala*, *Kawasphenodon* and *Cynosphenodon*. The instability of the three taxa is owing to missing data (unknown character states) and character conflict (some characters support alternative positions in different trees), as determined by iterative PCR [31]. Excluding the three unstable taxa from the consensus shows a completely resolved reduced strict consensus tree (figure 6). Owing to the close relation of *Ankylosphenodon* and *C. latidens* with basal opisthodontians in most parsimonious trees (see electronic supplementary material), the strict consensus shows these two taxa in a polytomy with *Sphenotitan* and more derived opisthodontians (figure 6). Therefore, according to the stem-based definition of Opisthodontia [9], the topology of the reduced strict consensus suggests *Ankylosphenodon* and *C. latidens* may be regarded as basal opisthodontians.

Until now, the oldest known opisthodontians were from the Late Jurassic (e.g. *Eilenodon* and *Opisthias* from the Kimmeridgian–Tithonian Morrison Formation of Wyoming [12–14]; an opisthodontian mandible from the Kimmeridgian Alcobaça Formation of Portugal [16]). *Sphenotitan* is the earliest

record of this group, extending its origin back by at least 50 Myr to the Norian. This suggestion is also strengthened by the position of the Upper Triassic *C. latidens* in a polytomy with the basal opisthodontians (figure 6). Moreover, the relationships of opisthodontians and sphenodontines (the branch that includes *Sphenodon*), coupled with the Norian record of *Sphenotitan*, suggest a Triassic origin of the lineage leading to sphenodontines, as was previously suggested [27,28].

## 5. Discussion

Sphenodontians are characterized by substantial variation in body size, body shape, habit, skull structure and size, and size and shape of teeth [32]. From these different traits, the diversity of tooth shape and arrangement suggests that Sphenodontia pursued a wide range of feeding strategies [33]. Based on tooth dimensions (mesiodistal length versus labiolingual width), three main morphotypes have been defined, each of them suited for a different mechanism of food processing: (i) small columnar teeth for piercing, indicative of an insectivorous diet; (ii) large elongated teeth for cutting and slicing, indicative of insectivory and/or a carnivorous diet; (iii) large wide teeth for shearing and shredding, indicative of an herbivorous diet [32]. The dentary of *Sphenotitan* has a large number of wide and closely packed teeth, increasing the surface area in contact with food (figures 1*a,b* and 3). This arrangement is present in all opisthodontians, including *C. latidens* and *Pelecymala*, although in *Sphenotitan* the teeth are wider and more closely packed than in any other sphenodontian. *Clevosaurus latidens* (which has not previously been proposed as Opisthodontia) and *Pelecymala* are the only known wide-toothed sphenodontians located out of Opisthodontia, which suggests that this feature has evolved twice within Sphenodontia [32]. Although *Pelecymala* was found to be unstable in this analysis owing to a lack of information on several character states, in several MPTs of the reduced strict consensus it is located close to the opisthodontians (see electronic supplementary material, figure S4). On the other hand, *C. latidens* is depicted in a polytomy with Opisthodontia and *Ankylosphenodon* (figure 6), which suggests an affinity towards Opisthodontia according to its stem-based

definition [7]. Although the known material of *C. latidens* and *Pelecymala* is limited and more material is needed to definitively solve this problem, the result of the present analysis suggests that wide-toothed herbivorous sphenodontians evolved just once in the evolutionary story of Sphenodontia.

The shared landmarks of *C. latidens* with Opisthodontia are not surprising, because both are wide-toothed, but the distance from *Clevosaurus hudsoni* is notable (figure 6), especially considering the fact that both have been related to the same genus. These results suggest that critical review of these materials is needed in order to determine the validity of the assignment of *C. latidens* to the genus *Clevosaurus*, which exceeds the scope of this work.

The anteroposteriorly enlarged and symmetrical glenoid articular cavity (figures 3*d* and 4*d*), the continuous shear facets found on the ventromedial side of the maxillary tooth row (figures 1*b* and 5), and the lateral surface of the dentary tooth row (figure 3*a*) indicate that a propalinal component of jaw movement played a major role in intraoral food processing in *Sphenotitan*. This seems to have been produced by two principal modes: slicing occurring between the ventromedial side of the maxillary teeth and the labial face of the dentary tooth row, and shredding being done by the ventrolateral side of the palatine tooth row and the dorsolingual face of the dentary tooth row, coupled with conspicuous rows of pterygoid teeth (figure 5). This masticatory mechanism is similar to that proposed for the most derived opisthodontians, the eilenodontines (e.g. *Toxolophosaurus* [28] and *Eilenodon* [34]). *Sphenotitan* shows that the most relevant advanced herbivorous traits present in the younger eilenodontines (e.g. large adult size, dentary teeth much more wide than long and eupropalinity) were acquired at least 50 Myr earlier than previously documented, at the end of the Late Triassic.

**Acknowledgements.** We thank S. Apesteguía, M. Colbert, D. Pol and J. Carballido for their early comments on the manuscript. We are indebted to M. Hassell (Editor), Zhe-Xi Luo (Associate Editor) and anonymous referees for their suggestions, which considerably improved the quality of this work.

**Funding statement.** We thank FONCYT and Instituto y Museo de Ciencias Naturales of the Universidad Nacional de San Juan for research support.

## References

- Evans SE, Jones MEH. 2010 The origin, early history and diversification of lepidosauromorph reptiles. In *New aspects of Mesozoic biodiversity: lecture notes in Earth Sciences 132* (ed. S Bandyopadhyay), pp. 27–44. Berlin, Germany: Springer.
- Cocude-Michel M. 1963 Les rhynchocephales et les sauriens des calcaires lithographiques (Jurassique supérieur) d'Europe occidentale. *Nouv. Arch. Mus. d'Hist. Nat. Lyon* **7**, 1–187.
- Fraser NC. 1986 New Triassic sphenodontids from south west England and a review of their classification. *Palaeontology* **29**, 165–186.
- Whiteside D. 1986 The head skeleton of the Rhaetian sphenodontid *Diphyodontosaurus avonensis* gen. et sp. nov. and the modernizing of a living fossil. *Phil. Trans. R. Soc. Lond. B* **312**, 379–430. (doi:10.1098/rstb.1986.0014)
- Fraser NC. 1989 The Triassic reptiles Brachyrhinodon and Polysphenodon and the relationships of the sphenodontids. *Zool. J. Linn. Soc.* **96**, 413–445. (doi:10.1111/j.1096-3642.1989.tb02521.x)
- Wu X-C. 1994 Late Triassic–Early Jurassic sphenodontians (*Clevosaurus*) from China and the phylogeny of the Sphenodontia. In *In the shadow of the dinosaurs: Early Mesozoic tetrapods* (eds NC Fraser, H-D Sues), pp. 38–69. Cambridge, UK: Cambridge University Press.
- Apesteguía S, Gómez RO, Rougier GW. 2012 A basal sphenodontian (Lepidosauria) from the Jurassic of Patagonia: new insights on the phylogeny and biogeography of Gondwanan rhynchocephalians. *Zool. J. Linn. Soc.* **166**, 342–360. (doi:10.1111/j.1096-3642.2012.00837.x)
- Carroll RL. 1988 *Vertebrate paleontology and evolution*. New York, NY: WH Freeman and Company.
- Apesteguía S, Novas FE. 2003 Large Cretaceous sphenodontian from Patagonia provides insight into lepidosaur evolution in Gondwana. *Nature* **425**, 609–612. (doi:10.1038/nature01995)
- Simón ME, Kellner AWA. 2003 New sphenodontid (Lepidosauria, Rhynchocephalia, Eilenodontinae) from the Candeleros Formation, Cenomanian of Patagonia, Argentina. *Bol. Mus. Nac. Geol.* **68**, 1–12.
- Apesteguía S, Carballido JL. In press. *Kaikafilusaurus minimus*, a dwarf Eilenodontine

- (Lepidosauria, Sphenodontidae) from the Cretaceous of Central Patagonia. *J. Vert. Paleontol.*
12. Gilmore CW. 1909 A new rhynchocephalian reptile from the Jurassic of Wyoming, with notes on the fauna of 'Quarry 9'. *Proc. US Nat. Mus.* **37**, 35–42. (doi:10.5479/si.00963801.37-1698.35)
  13. Rasmussen TE, Callison G. 1981 A new herbivorous sphenodontid (Rhynchocephalia: Reptilia) from the Jurassic of Colorado. *J. Paleontol.* **55**, 1109–1116.
  14. Kowallis BJ, Christiansen EH, Deino AL, Peterson F, Turner CE, Kunk MJ, Obradovich JD. 1998 The age of the Morrison Formation. *Mod. Geol.* **22**, 235–260.
  15. Evans SE, Fraser NC. 1992 A sphenodontid jaw (Reptilia: Lepidosauria) from the Upper Jurassic of Dorset. *Proc. Dorset Nat. Hist. Soc.* 199–200.
  16. Ortega F, Malafaia E, Escaso F, Perez García A, Dantas P. 2000 Faunas de répteis do Jurássico Superior de Portugal. *Paleolusitana* **1**, 43–56.
  17. Bossi GE, Bonaparte JF. 1978 Sobre la presencia de un dinosaurio prosauropodo en la Formación Quebrada del Barro, en el Borde Austral de la Cuenca Marayes-El Carrizal (Triásico Superior de San Juan). *Acta Geol. Lilloana XV* **1**, 41–47.
  18. Apaldetti C, Martínez RN, Alcober OA, Pol D. 2011 A new basal sauropodomorph (Dinosauria: Saurischia) from Quebrada del Barro Formation (Marayes-El Carrizal Basin), northwestern Argentina. *PLoS ONE* **6**, e26964. (doi:10.1371/journal.pone.0026964)
  19. Martínez RN, Apaldetti C, Praderio A, Fernández E, Colombi C, Alcober OA, Santi Malnis P, Correa G, Abelin D. 2013 *New vertebrate findings shed light on the controversial age of the Quebrada del Barro Formation (Marayes–El Carrizal Basin)*. Chubut, Argentina: V Simp. Arg. Jur. Trelew.
  20. Haeckel E. 1866 *Generelle Morphologie der Organismen. Zweiter Band: Allgemeine Entwicklungsgeschichte der Organismen*. Berlin, Germany: Reimer.
  21. Günther A. 1867 Contribution to the anatomy of *Hatteria* (Rhynchocephalus, Owen). *Phil. Trans. R. Soc.* **157**, 595–629. (doi:10.1098/rstl.1867.0019)
  22. Williston SW. 1925 *The osteology of the reptiles*. Cambridge, MA: Harvard University Press.
  23. Fraser NC. 1988 The osteology and relationships of *Clevosaurus* (Reptilia: Sphenodontida). *Phil. Trans. R. Soc. Lond. B* **321**, 125–178. (doi:10.1098/rstb.1988.0092)
  24. Evans SE, Prasad GVR, Manhas BK. 2001 Rhynchocephalians (Diapsida: Lepidosauria) from the Jurassic Kota Formation of India. *J. Linn. Soc.* **133**, 309–334. (doi:10.1111/j.1096-3642.2001.tb00629.x)
  25. Reynoso VH. 1997 A 'beaded' sphenodontian (Diapsida: Lepidosauria) from the Early Cretaceous of Central Mexico. *J. Vert. Paleontol.* **17**, 52–59. (doi:10.1080/02724634.1997.10010953)
  26. Reynoso V-H, Clark JM. 1998 A dwarf sphenodontian from the Jurassic La Boca formation of Tamaulipas, México. *J. Vert. Paleontol.* **18**, 333–339. (doi:10.1080/02724634.1998.10011061)
  27. Apesteguía S. 2008 Esfenodontes (Reptilia, Lepidosauria) del Cretácico Superior de Patagonia: Anatomía y Filogenia, p. 535. Thesis, Universidad Nacional de Buenos Aires, Buenos Aires, Argentina.
  28. Throckmorton GS, Hopson JA, Parks P. 1981 A redescription of *Toxolophosaurus claudi* Olson, a Lower Cretaceous herbivorous sphenodontid reptile. *J. Paleontol.* **55**, 586–597.
  29. Fraser NC. 1993 A new Sphenodontian from the early Mesozoic of England and North America: implications for correlating early Mesozoic continental deposits. In *The nonmarine Triassic* (eds SG Lucas, M Morales), pp. 135–139. Albuquerque, NM: New Mexico Museum of Natural History and Science.
  30. Wilkinson M. 1994 Common cladistic information and its consensus representation: reduced Adams and reduced cladistic consensus trees and profiles. *Syst. Biol.* **43**, 343–368.
  31. Pol D, Escapa IH. 2009 Untestable taxa in cladistic analysis: identification and the assessment of relevant characters. *Cladistics* **25**, 515–527. (doi:10.1111/j.1096-0031.2009.00258.x)
  32. Jones MEH. 2009 Dentary tooth shape in *Sphenodon* and its fossil relatives (Diapsida: Rhynchocephalia). *Front. Oral Biol.* **13**, 9–15. (doi:10.1159/000242382)
  33. Jones MEH. 2008 Skull shape and feeding strategy in *Sphenodon* and other Rhynchocephalia (Diapsida: Lepidosauria). *J. Morphol.* **269**, 945–966. (doi:10.1002/jmor.10634)
  34. Foster JR. 2003 New specimens of Eilenodon (Reptilia, Sphenodontia) from the Morrison Formation (Upper Jurassic) of Colorado and Utah. *BYU Geol. Stud.* **47**, 17–22.