

Phosphate Mediated Adsorption and Electron Transfer of Cytochrome *c*. A Time-Resolved SERR Spectroelectrochemical Study

Daiana Capdevila, Waldemar Marmisollé, Federico J. Williams and Daniel H. Murgida*

Supplementary Information




Figure S11. C1s XPS spectra of $(\text{NH}_2)_{0.5}(\text{OH})_{0.5}\text{C}_6$ SAM modified gold electrode incubated overnight in 0.5M ATP (top) and 1mM Cyt in 0.5M ATP solution (bottom).




Figure S12. C1s XPS spectra of $(\text{NH}_2)_{0.5}(\text{OH})_{0.5}\text{C}_6$ SAM modified gold electrode incubated overnight in 0.5M ATP (top) and 1mM Cyt in 0.5M ATP solution (bottom).




Figure SI3. Component analysis of SERR spectra of Cyt on $\text{NH}_2\text{-C6}$ SAM coated silver electrode at three different applied potentials. The components were obtained from RR spectra of Cyt^{2+} and Cyt^{3+} .




Figure SI4. Nernst plot for Cyt on NH_2 -C6 SAM coated silver electrode. Relative concentrations of reduced and oxidized forms were obtained by component analysis as indicated in Figure SI3.




Figure SI5. Time dependence of relative concentration of Cyt^{2+} after a potential jump from -100 mV to the redox potential for Cyt on NH_2 -C6 SAM coated silver electrode. Data of two independent experiments are included and fitted to a monoexponential decay function.