

Classification of Cubic Symmetric Tricirculants

István Kovács^{a,1} Klavdija Kutnar^{a,1,2} Dragan Marušič^{a,b,1,*} Steve Wilson^c

^aUniversity of Primorska, FAMNIT, Glagoljaška 8, 6000 Koper, Slovenia

^bUniversity of Ljubljana, PEF, Kardeljeva pl. 16, 1000 Ljubljana, Slovenia

^cNorthern Arizona University, Flagstaff, AZ 86011, USA

Submitted: Dec 16, 2011; Accepted: May 21, 2012; Published: May 31, 2012

Abstract

A *tricirculant* is a graph admitting a non-identity automorphism having three cycles of equal length in its cycle decomposition. A graph is said to be *symmetric* if its automorphism group acts transitively on the set of its arcs. In this paper it is shown that the complete bipartite graph $K_{3,3}$, the Pappus graph, Tutte's 8-cage and the unique cubic symmetric graph of order 54 are the only connected cubic symmetric tricirculants.

Keywords: symmetric graph, semiregular, tricirculant.

1 Introductory remarks

A graph is said to be *arc-transitive*, or *symmetric*, the term that will be used in this paper, if its automorphism group acts transitively on the set of arcs of the graph. A Cayley graph on a cyclic group, that is, a graph admitting an automorphism with a single cycle in its cycle decomposition, is said to be a *circulant*. A graph admitting a non-identity automorphism with two (respectively, three) cycles of equal length in its cycle decomposition is said to be a *bicirculant* (respectively, *tricirculant*).

It is known that the complete graph K_4 , the complete bipartite graph $K_{3,3}$ and the cube Q_3 are the only connected cubic symmetric graphs with girth (the length of shortest cycle) less than 5 (see [8, Proposition 3.4.]). Since any cubic circulant, being a Cayley

¹Supported in part by “Agencija za raziskovalno dejavnost Republike Slovenije”, research program P1-0285.

²The author gratefully acknowledges support by the US Department of State and the Fulbright Scholar Program, and thanks the Ohio State University for hospitality during her visit in Autumn 2010.

*Corresponding author e-mail: dragan.marusic@upr.si.

graph on an abelian group, has girth less than or equal to 4, one can easily see that K_4 and $K_{3,3}$ are the only examples of connected cubic symmetric circulants.

We may think of the classical result of Frucht, Graver and Watkins [7] in which they have classified all symmetric generalized Petersen graphs as the main step in the classification of all cubic connected symmetric bicirculants. The remaining cases were then completed in [19] and [21]. In particular, a connected cubic symmetric graph is a bicirculant if and only if it is isomorphic to one of the following graphs: the complete graph K_4 , the complete bipartite graph $K_{3,3}$, the seven symmetric generalized Petersen graphs $GP(4, 1)$, $GP(5, 2)$, $GP(8, 3)$, $GP(10, 2)$, $GP(10, 3)$, $GP(12, 5)$, and $GP(24, 5)$ (see [7, 20]), the Heawood graph F014A, and a Cayley graph $\text{Cay}(D_{2n}, \{b, ba, ba^{r+1}\})$ on a dihedral group $D_{2n} = \langle a, b \mid a^n = b^2 = baba = 1 \rangle$ of order $2n$ with respect to the generating set $\{b, ba, ba^{r+1}\}$, where $n \geq 11$ is odd and $r \in \mathbb{Z}_n^*$ such that $r^2 + r + 1 \equiv 0 \pmod{n}$. (Hereafter the notation FnA, FnB, etc. will refer to the corresponding graphs in the Foster census [3, 4].)

The aim of this paper is to move from bicirculants to tricirculants. A complete classification of connected cubic symmetric tricirculants is given. In particular, it is shown that the complete bipartite graph $K_{3,3}$ (which is also a circulant and a bicirculant), the Pappus graph F018A, Tutte's 8-cage F030A (sometimes also called the Tutte-Coxeter graph), and the graph F054A are the only connected cubic symmetric tricirculants, see Theorem 1.1. (For brevity in this paper, we will refer to these graphs as $K_{3,3}$, F018A, F030A and F054A, respectively.)

Theorem 1.1 *A connected cubic symmetric graph X is a tricirculant if and only if it is isomorphic to one of the following four graphs: $K_{3,3}$, F018A, F030A, and F054A.*

The classification is obtained by first considering the so-called *core-free tricirculants*, that is, tricirculants admitting a non-identity automorphism ρ , having three cycles of equal length in its cycle decomposition, such that the subgroup generated by ρ is core-free in the full automorphism group of the graph. A remarkable group-theoretic result of Herzog and Kaplan [11], which says that 'sufficiently large' cyclic subgroups are never corefree (see Lemma 3.2), combined together with the well-known fact that the automorphism group of a connected cubic symmetric graph of order n is of order $3 \cdot 2^{s-1}n$, where $s \leq 5$, enable us to prove that $K_{3,3}$, F018A, and F030A are the only connected cubic symmetric core-free tricirculants (see Theorem 3.3). Whereas, for non-core-free cubic symmetric tricirculants Lorimer's result about cubic symmetric graphs admitting a normal subgroup in its automorphism group implies that any such graph is a regular cyclic cover of a cubic symmetric core-free tricirculant (see Lemma 3.4). This then enables us to use graph covering techniques, which we recall in Subsection 2.2, to prove that the graph F054A is the only connected cubic symmetric non-core-free tricirculant, and the classification follows.

2 Preliminaries

Throughout this paper graphs are simple, finite, undirected and connected. Given a

graph X we let $V(X)$, $E(X)$, $A(X)$ and $\text{Aut}X$ be the vertex set, the edge set, the arc set and the automorphism group of X , respectively. A sequence of $k + 1$ vertices in X , not necessarily all distinct, such that any two consecutive vertices are adjacent and any three consecutive vertices are distinct is called a k -arc.

A subgroup $G \leq \text{Aut}X$ is said to be *vertex-transitive*, *edge-transitive* and *arc-transitive* provided it acts transitively on the sets of vertices, edges and arcs of X , respectively. The graph X is said to be *vertex-transitive*, *edge-transitive*, and *arc-transitive* if its automorphism group is vertex-transitive, edge-transitive and arc-transitive, respectively. An arc-transitive graph is also called *symmetric*. A subgroup $G \leq \text{Aut}X$ is said to be *s-arc-transitive* if it acts transitively on the set of s -arcs, and it is said to be *s-regular* if it is s -arc-transitive and the stabilizer of an s -arc in G is trivial. A graph is said to be *s-regular* if its automorphism group is *s-regular*. By Tutte's result [22] every connected cubic symmetric graph is s -regular for some $s \leq 5$.

For a partition \mathcal{W} of $V(X)$, we let $X_{\mathcal{W}}$ be the associated *quotient graph* of X relative to \mathcal{W} , that is, the graph with vertex set \mathcal{W} and edge set induced naturally by the edge set $E(X)$. In the case when \mathcal{W} corresponds to the set of orbits of a subgroup N of $\text{Aut}X$, the symbol $X_{\mathcal{W}}$ will be replaced by X_N .

2.1 Semiregular automorphisms

A non-identity automorphism of a graph is *semiregular*, in particular, (k, n) -*semiregular* if it has k cycles of equal length n in its cycle decomposition. An n -*trirculant* (*trirculant* for short) is a graph with a $(3, n)$ -semiregular automorphism.

Let X be a connected graph admitting a (k, n) -semiregular automorphism

$$\rho = (u_0^0 u_0^1 \cdots u_0^{n-1})(u_1^0 u_1^1 \cdots u_1^{n-1}) \cdots (u_{k-1}^0 u_{k-1}^1 \cdots u_{k-1}^{n-1}),$$

and let $\mathcal{W} = \{W_i \mid i \in \mathbb{Z}_k\}$ be the set of orbits $W_i = \{u_i^s \mid s \in \mathbb{Z}_n\}$ of ρ . Using Frucht's notation [6] X may be represented in the following way. Each orbit of ρ is represented by a circle. Inside a circle corresponding to the orbit W_i the symbol n/T , where $T = T^{-1} \subseteq \mathbb{Z}_n \setminus \{0\}$, indicates that for each $s \in \mathbb{Z}_n$, the vertex u_i^s is adjacent to all the vertices u_i^{s+t} where $t \in T$. When $|T| \leq 2$ we use a simplified notation n/t , $n/(n/2)$ and n , when, respectively, $T = \{t, -t\}$, $T = \{n/2\}$ and $T = \emptyset$. Finally, an arrow pointing from the circle representing the orbit W_i to the circle representing the orbit W_j , $j \neq i$, labeled by $y \in \mathbb{Z}_n$ means that for each $s \in \mathbb{Z}_n$, the vertex $u_i^s \in W_i$ is adjacent to the vertex u_j^{s+y} . When the label is 0, the arrow on the line may be omitted. Examples illustrating this notation are given in Figure 1.

2.2 Graph Covers

A *covering projection* of a graph \tilde{X} is a surjective mapping $p: \tilde{X} \rightarrow X$ such that for each $\tilde{u} \in V(\tilde{X})$ the set of arcs emanating from \tilde{u} is mapped bijectively onto the set of arcs emanating from $u = p(\tilde{u})$. The graph \tilde{X} is called a *covering graph* of the *base graph* X . The set $\text{fib}_u = p^{-1}(u)$ is a *fibre* of a vertex $u \in V(X)$. The subgroup K of




Figure 1: The graphs $K_{3,3}$, F018A, F030A and F054A shown in Frucht's notation with respect to a $(3, 2)$ -, $(3, 6)$ -, $(3, 10)$ - and $(3, 18)$ -semiregular automorphism, respectively.

all those automorphisms of \tilde{X} which fix each of the fibres setwise is called the *group of covering transformations*. The graph \tilde{X} is also called a *K-cover* of X . It is a simple observation that the group of covering transformations of a connected covering graph acts semiregularly on each of the fibres. In particular, if the group of covering transformations is regular on the fibres of \tilde{X} , we say that \tilde{X} is a *regular K-cover*. We say that $\alpha \in \text{Aut}X$ *lifts* to an automorphism of \tilde{X} if there exists an automorphism $\tilde{\alpha} \in \text{Aut}\tilde{X}$, called a *lift* of α , such that $\tilde{\alpha}p = p\alpha$. If the covering graph \tilde{X} is connected then K is the lift of the trivial subgroup of $\text{Aut}X$. Note that a subgroup $G \leq \text{Aut}\tilde{X}$ projects if and only if the partition of $V(\tilde{X})$ into the orbits of K is G -invariant.

A combinatorial description of a K -cover was introduced through a voltage graph by Gross and Tucker [10] as follows. Let X be a graph and K be a finite group. A *voltage assignment* of X is a mapping $\zeta: A(X) \rightarrow K$ with the property that $\zeta(u, v) = \zeta(v, u)^{-1}$ for any arc $(u, v) \in A(X)$ (here, and in the rest of the paper, $\zeta(u, v)$ is written instead of $\zeta((u, v))$ for the sake of brevity). The voltage assignment ζ extends to walks in X in a natural way. In particular, for any walk $D = u_1u_2 \cdots u_t$ of X we let $\zeta(D)$ denote the product voltage $\zeta(u_1, u_2)\zeta(u_2, u_3) \cdots \zeta(u_{t-1}, u_t)$ of D , that is, the ζ -voltage of D .

The values of ζ are called *voltages*, and K is the *voltage group*. The *voltage graph* $X \times_{\zeta} K$ derived from a voltage assignment $\zeta: A(X) \rightarrow K$ has vertex set $V(X) \times K$, and edges of the form $\{(u, g), (v, \zeta(x)g)\}$, where $x = (u, v) \in A(X)$. Clearly, $X \times_{\zeta} K$ is a covering of X with the first coordinate projection. By letting K act on $V(X \times_{\zeta} K)$ as $(u, g)^{g'} = (u, gg')$, $(u, g) \in V(X \times_{\zeta} K)$, $g' \in K$, one obtains a semiregular group of automorphisms of $X \times_{\zeta} K$, showing that $X \times_{\zeta} K$ can in fact be viewed as a K -cover of X .

Given a spanning tree T of X , the voltage assignment $\zeta: A(X) \rightarrow K$ is said to be *T-reduced* if the voltages on the tree arcs equal the identity element in K . In [9] it is shown that every regular covering graph \tilde{X} of a graph X can be derived from a T -reduced voltage assignment ζ with respect to an arbitrary fixed spanning tree T of X .

The problem of whether an automorphism α of X lifts or not is expressed in terms of voltages as follows (see Proposition 2.1). Given $\alpha \in \text{Aut}X$ and the set of fundamental closed walks \mathcal{C} based at a fixed vertex $v \in V(X)$, we define $\bar{\alpha} = \{(\zeta(C), \zeta(C^\alpha)) \mid C \in \mathcal{C}\} \subseteq K \times K$. Note that if K is abelian, $\bar{\alpha}$ does not depend on the choice of the base vertex, and the fundamental closed walks at v can be substituted by the fundamental cycles generated by the cotree arcs of X . Also, from the definition, it is clear that for a T -reduced voltage assignment ζ the derived graph $X \times_{\zeta} K$ is connected if and only if the voltages of the cotree arcs generate the voltage group K .

We wrap up this section with four propositions dealing with lifting of automorphisms in graph covers. The first one may be deduced from [17, Theorem 4.2], the second one from [12] whereas the third one is taken from [5, Proposition 2.2], but it may also be deduced from [18, Corollaries 9.4, 9.7, 9.8].

Proposition 2.1 [17] *Let K be a finite group, and let $X \times_{\zeta} K$ be a connected regular cover of a graph X derived from a voltage assignment ζ with the voltage group K . Then an automorphism α of X lifts if and only if $\bar{\alpha}$ is a function which extends to an automorphism α^* of K .*

For a connected regular cover $X \times_{\zeta} K$ of a graph X derived from a T -reduced voltage assignment ζ with an abelian voltage group K and an automorphism $\alpha \in \text{Aut}X$ that lifts, $\bar{\alpha}$ will always denote the mapping from the set of voltages of the fundamental cycles on X to the voltage group K and α^* will denote the automorphism of K arising from $\bar{\alpha}$.

Two coverings $p_i: \tilde{X}_i \rightarrow X$, $i \in \{1, 2\}$, are said to be *isomorphic* if there exists a graph isomorphism $\phi: \tilde{X}_1 \rightarrow \tilde{X}_2$ such that $\phi p_2 = p_1$.

Proposition 2.2 [12] *Let K be a finite group. Two connected regular covers $X \times_{\zeta} K$ and $X \times_{\varphi} K$, where ζ and φ are T -reduced, are isomorphic if and only if there exists an automorphism $\sigma \in \text{Aut}K$ such that $\zeta(u, v)^{\sigma} = \varphi(u, v)$ for any cotree arc (u, v) of X .*

Proposition 2.3 [5] *Let K be a finite group, and let $X \times_{\zeta} K$ be a connected regular cover of a graph X derived from a voltage assignment ζ with the voltage group K , and let the lifts of $\alpha \in \text{Aut}X$ centralize K , considered as the group of covering transformations. Then for any closed walk W in X , there exists $k \in K$ such that $\zeta(W^\alpha) = k\zeta(W)k^{-1}$. In particular, if K is abelian, $\zeta(W^\alpha) = \zeta(W)$ for any closed walk W of X .*

Given a voltage assignment ζ of X and $\beta \in \text{Aut } X$, we set ζ^β for the voltage assignment of X such that $\zeta^\beta(u, v) = \zeta(u^{\beta^{-1}}, v^{\beta^{-1}})$, $(u, v) \in A(X)$; and set $\tilde{\beta}$ for the permutation of $V(X) \times K$ acting as $(u, k)^{\tilde{\beta}} = (u^\beta, k)$. Our last proposition is straightforward.

Proposition 2.4 *Let K be a finite group, and let $\tilde{X} = X \times_\zeta K$ be a connected regular cover of a graph X derived from a voltage assignment ζ with the voltage group K , and let $\beta \in \text{Aut } X$. Then the following hold.*

- (i) $\tilde{\beta}$ is an isomorphism from \tilde{X} to $X \times_{\zeta^\beta} K$.
- (ii) If $\tilde{\alpha}$ is in $\text{Aut } \tilde{X}$ which projects to α , then $\tilde{\beta}^{-1}\tilde{\alpha}\tilde{\beta}$ is in $\text{Aut}(X \times_{\zeta^\beta} K)$, and it projects to $\beta^{-1}\alpha\beta$.
- (iii) If $\tilde{\alpha} \in \text{Aut } \tilde{X}$ centralizes the group K of covering transformations, then also $\tilde{\beta}^{-1}\tilde{\alpha}\tilde{\beta}$ centralizes K .

3 Cubic symmetric tricirculants

Let \mathcal{TC} be the family of connected cubic symmetric tricirculants. We are going to show that $K_{3,3}$, F018A, F030A and F054A, the four graphs shown in Figure 1, are the only connected cubic symmetric tricirculants. The graph F018A is the unique connected cubic symmetric graph of order 18. It is 3-regular, has girth 6 and is the Levi graph of the Pappus configuration. The graph F030A is the unique connected cubic symmetric graph of order 30. It is 5-regular, has girth 8 and diameter 4. As the unique smallest cubic graph of girth 8 it is a cage and a Moore graph (see also [13]). It is bipartite, and can be constructed as the Levi graph of the generalized quadrangle $GQ(2, 2)$. The graph F054A is the unique cubic symmetric graph of order 54. It is 2-regular and has girth 6. For more information on these graphs we refer the reader to [3, 4, 14, 16].

Using the table of cubic symmetric graphs of order up to 768 in [3, 4] and a program package MAGMA [2] one can see that the following lemma holds.

Lemma 3.1 *There is no cubic 1-regular tricirculant of order less than 27. The graphs F024A, F048A, F090A, F096B, F102A, F204A, and F234B are not tricirculants.*

Recall that the *core* of a subgroup K in a group G (denoted by $\text{core}_G(K)$) is the largest normal subgroup of G contained in K . A graph $X \in \mathcal{TC}$ with a $(3, n)$ -semiregular automorphism is said to be *core-free* if there exists a $(3, n)$ -semiregular automorphism $\rho \in \text{Aut } X$ such that the cyclic subgroup $\langle \rho \rangle$ has trivial core in $\text{Aut } X$. To obtain the classification of cubic symmetric core-free tricirculants (see Theorem 3.3) the following group-theoretical result will be used.

Lemma 3.2 [11, Theorem B] *If H is a cyclic subgroup of a finite group G with $|H| \geq \sqrt{|G|}$, then H contains a non-trivial normal subgroup of G .*

Theorem 3.3 *A graph $X \in \mathcal{TC}$ is core-free if and only if it is isomorphic to one of the following three graphs: $K_{3,3}$, F018A, and F030A.*

PROOF. Let X be cubic s -arc-transitive trirculant of order $o = 3n$, and let $G = \text{Aut } X$. Then, by [1, Proposition 18.1]), G is regular on the set of s -arcs of X . By Tutte's theorem (see [1, Theorem 18.6]) we have that $s \leq 5$, and therefore $|G| = 9 \cdot 2^{s-1} \cdot n \leq 144n$. Since X is core-free, Lemma 3.2 implies that $n^2 < |G|$, and consequently $n < 144$.

In particular, if $s = 1$ then $n^2 < |G| = 9n$, implying that $n < 9$ and $o = 3n < 27$, which in view of Lemma 3.1 is impossible.

If $s = 2$ then $n^2 < |G| = 18n$, implying that $n < 18$ and $o = 3n < 54$. By the table of cubic symmetric graphs [4] the only cubic 2-regular graphs of order less than 54 (and divisible by 3) are F024A and F048A. However, by Lemma 3.1, F024A and F048A are not trirculants.

If $s = 3$ then $n^2 < |G| = 36n$, and so $n < 36$ and $o = 3n < 108$. The only cubic 3-regular graphs satisfying this condition are the graphs $K_{3,3}$, F018A and F096B. The first two graphs are clearly trirculants whereas the latter graph is not a trirculant by Lemma 3.1.

If $s = 4$ then $n^2 < |G| = 72n$, and so $n < 72$ and $o = 3n < 216$. The only cubic 4-regular graphs satisfying this condition are the graphs F102A and F204A, which, by Lemma 3.1, are not trirculants.

If $s = 5$ then $n^2 < |G| = 144n$, and so $n < 144$ and $o = 3n < 432$. The only cubic 5-regular graphs satisfying this condition are the graphs F030A, F090A and F234B. However, the last two graphs, by Lemma 3.1, are not trirculants.

That $K_{3,3}$, F018A, and F030A are indeed core-free can be easily checked with the use of MAGMA [2]. ■

The following lemma follows from [15, Theorem 9].

Lemma 3.4 *Let $X \in \mathcal{TC}$ with a $(3, n)$ -semiregular automorphism $\rho \in \text{Aut } X$, and let N be the core of $\langle \rho \rangle$ in $\text{Aut } X$. Then N is the kernel of $\text{Aut } X$ acting on the set of orbits of N , $\text{Aut } X/N$ acts arc-transitively on X_N , $X_N \in \mathcal{TC}$ with a $(3, n/|N|)$ -semiregular automorphism and X_N is core-free.*

We are now ready to prove the main theorem of this paper.

PROOF OF THEOREM 1.1. Let X be a cubic symmetric trirculant with a $(3, n)$ -semiregular automorphism $\rho \in \text{Aut } X$. If X is core-free then, by Theorem 3.3, X is isomorphic to $K_{3,3}$, or F018A, or F030A.

Suppose now that X is not core-free. Then there exists a nontrivial subgroup N of $\langle \rho \rangle$ which is normal in $\text{Aut } X$. By Lemma 3.4, the quotient graph X_N is a connected cubic symmetric core-free trirculant, and hence, by Theorem 3.3, it is isomorphic to $K_{3,3}$, F018A or F030A. In fact, X is isomorphic to a regular \mathbb{Z}_r -cover of one of these three graphs, where $|N| = r$. Note also that ρ projects to a $(3, n/r)$ -semiregular automorphism of X_N . (Below, all arithmetic operations are to be taken modulo r if at least one argument is from \mathbb{Z}_r and the symbol mod r is always omitted.)




Figure 2: The voltage assignment ζ on $K_{3,3}$. The spanning tree consists of undirected bold edges, all carrying trivial voltage.

CASE 1. $X_N \cong K_{3,3}$.

The graph $K_{3,3}$ is illustrated in Figure 2. It is known that $K_{3,3}$ is the unique connected cubic symmetric graph of order 6 and that this graph is in fact 3-regular (see [4]). Let us choose the following automorphisms of $K_{3,3}$

$$\alpha = (1)(2, 4, 3)(5)(6) \text{ and } \beta = (1, 2)(3, 5)(4, 6),$$

and let $H = \langle \alpha, \beta \rangle$. It can be checked directly, using Magma [2], that every $(3, 2)$ -semiregular automorphism of $K_{3,3}$ is conjugate to β , and that every arc-transitive subgroup of $\text{Aut } K_{3,3}$, which contains β , must contain also the subgroup H . Because of Proposition 2.4 we may assume without loss of generality that ρ projects to β (therefore, the lifts of β centralize the group N of covering transformations) and that H lifts to a subgroup of $\text{Aut } X$.

Any such cover X can be derived from $K_{3,3}$ through a suitable voltage assignment $\zeta : A(K_{3,3}) \rightarrow \mathbb{Z}_r$. To find this voltage assignment ζ fix the spanning tree T of $K_{3,3}$ as the one consisting of the edges

$$\{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 5\}, \{2, 6\}$$

(see also Figure 2). There are four fundamental cycles in $K_{3,3}$, which are generated, respectively, by four cotree arcs $(4, 6)$, $(4, 5)$, $(3, 5)$, and $(3, 6)$. By Proposition 2.3, β^* is the identity automorphism of \mathbb{Z}_r , and thus we get from Table 1, where all these cycles and their voltages are listed, that $2x_1 = 2x_3 = 0$ and $x_2 = -x_4$. Moreover, since α lifts, Proposition 2.1 implies that $\bar{\alpha}$ is a function which extends to an automorphism α^* of \mathbb{Z}_r . Therefore, since $\zeta(C_3^\alpha) = -x_2$ and $\zeta(C_4^\alpha) = -x_1$, it follows that all non-trivial voltages are of order at most 2. Since X is assumed to be connected at least one non-trivial voltage exists and the set of all non-trivial voltages generates the voltage group. Since the voltage group is cyclic, it follows that $r = 2$. But, however, there is no connected cubic symmetric graph of order 12, and so this case is impossible.

	C	$\zeta(C)$	C^α	$\zeta(C^\alpha)$	C^β	$\zeta(C^\beta)$
C_1	1, 4, 6, 2, 1	x_1	1, 3, 6, 4, 1	$x_4 - x_1$	2, 6, 4, 1, 2	$-x_1$
C_2	1, 4, 5, 2, 1	x_2	1, 3, 5, 4, 1	$x_3 - x_2$	2, 6, 3, 1, 2	$-x_4$
C_3	1, 3, 5, 2, 1	x_3	1, 2, 5, 4, 1	$-x_2$	2, 5, 3, 1, 2	$-x_3$
C_4	1, 3, 6, 2, 1	x_4	1, 2, 6, 4, 1	$-x_1$	2, 5, 4, 1, 2	$-x_2$

Table 1: Fundamental cycles and their images with corresponding voltages in $K_{3,3}$.




Figure 3: The voltage assignment ζ on F018A. The spanning tree consists of undirected bold edges, all carrying trivial voltage.

CASE 2. $X_N \cong \text{F018A}$.

The graph F018A is illustrated in Figure 3. It is known that F018A is the unique connected cubic symmetric graph of order 18 and that this graph is in fact 3-regular (see [4]). Let us choose the following automorphisms of F018A

$$\begin{aligned}
\alpha &= (1, 2)(3, 6)(4, 5)(7, 8)(9, 12)(10, 11)(13, 14)(15, 18)(16, 17), \\
\beta &= (1, 7, 14, 9, 3, 2)(4, 8, 6, 18, 17, 16)(5, 15, 12, 11, 10, 13), \\
\gamma_1 &= (3, 8)(4, 15)(5, 18)(6, 7)(9, 13)(12, 14), \\
\gamma_2 &= (2, 6)(3, 5)(8, 12)(9, 11)(14, 18)(15, 17), \\
\gamma_3 &= (1, 2)(3, 6, 8, 7)(4, 12, 15, 14)(5, 13, 18, 9)(10, 17)(11, 16), \\
\delta &= (4, 9)(5, 14)(6, 7)(10, 16)(11, 17)(12, 18)(13, 15).
\end{aligned}$$

Then $\text{AutF018A} = \langle \alpha, \beta, \gamma_3, \delta \rangle = \langle \gamma_1, \gamma_2, \gamma_3, \delta \rangle$. Each subgroup of AutF018A generated by a $(3, 6)$ -semiregular automorphism is conjugate to $\langle \beta \rangle$, and each proper arc-transitive subgroup of AutF018A is conjugate in AutF018A to one of the three subgroups $H_1 = \langle \alpha, \beta \rangle$, $H_2 = \langle \alpha, \beta, \gamma_1 \rangle$ and $H_3 = \langle \gamma_1, \gamma_2, \gamma_3 \rangle$. In addition, H_1 is 1-regular, H_2 and H_3 are 2-regular,

$\langle \beta \rangle \leq H_1 \leq H_2$, and H_3 does not contain a $(3, 6)$ -semiregular automorphism. These can be checked directly using Magma [2]. Thus it suffices to find those arc-transitive regular \mathbb{Z}_r -covers of F018A for which the subgroup H_1 lifts and the lifts of the automorphism β centralizing the group of covering transformations and having precisely 3 orbits of size n .

Row		C	$\zeta(C)$	C^α	$\zeta(C^\alpha)$
A.1	C_0	1, 2, 3, 4, 5, 6, 1	x	2, 1, 6, 5, 4, 3, 2	$-x$
A.2	C_1	1, 7, 14, 9, 3, 4, 5, 6, 1	y_1	2, 8, 13, 12, 6, 5, 4, 3, 2	$-y_6$
A.3	C_2	2, 8, 15, 10, 4, 3, 2	y_2	1, 7, 18, 11, 5, 6, 1	$-y_5$
A.4	C_3	3, 9, 16, 11, 5, 4, 3	y_3	6, 12, 17, 10, 4, 5, 6	$-y_4$
A.5	C_4	4, 10, 17, 12, 6, 5, 4	y_4	5, 11, 16, 9, 3, 4, 5	$-y_3$
A.6	C_5	5, 11, 18, 7, 1, 6, 5	y_5	4, 10, 15, 8, 2, 3, 4	$-y_2$
A.7	C_6	6, 12, 13, 8, 2, 3, 4, 5, 6	y_6	3, 9, 14, 7, 1, 6, 5, 4, 3	$-y_1$
A.8	C_7	1, 7, 18, 15, 10, 4, 5, 6, 1	z_1	2, 8, 15, 18, 11, 5, 4, 3, 2	$y_2 - z_1 - y_5$
A.9	C_8	2, 8, 13, 16, 11, 5, 4, 3, 2	z_2	1, 7, 14, 17, 10, 4, 5, 6, 1	$y_1 + z_3 - y_4$
A.10	C_9	3, 9, 14, 17, 12, 6, 5, 4, 3	z_3	6, 12, 13, 16, 9, 3, 4, 5, 6	$y_6 + z_2 - y_3$
		C	$\zeta(C)$	C^β	$\zeta(C^\beta)$
B.1	C_0	1, 2, 3, 4, 5, 6, 1	x	7, 1, 2, 8, 15, 18, 7	$x + y_2 - z_1$
B.2	C_1	1, 7, 14, 9, 3, 4, 5, 6, 1	y_1	7, 14, 9, 3, 2, 8, 15, 18, 7	$y_1 + y_2 - z_1$
B.3	C_2	2, 8, 15, 10, 4, 3, 2	y_2	1, 6, 12, 13, 8, 2, 1	$y_6 - x$
B.4	C_3	3, 9, 16, 11, 5, 4, 3	y_3	2, 3, 4, 10, 15, 8, 2	$-y_2$
B.5	C_4	4, 10, 17, 12, 6, 5, 4	y_4	8, 13, 16, 11, 18, 15, 8	$z_2 + y_5 + z_1 - y_2$
B.6	C_5	5, 11, 18, 7, 1, 6, 5	y_5	15, 10, 17, 14, 7, 18, 15	$y_4 - z_3 - y_1 + z_1$
B.7	C_6	6, 12, 13, 8, 2, 3, 4, 5, 6	y_6	18, 11, 5, 6, 1, 2, 8, 15, 18	$-y_5 + x + y_2 - z_1$
B.8	C_7	1, 7, 18, 15, 10, 4, 5, 6, 1	z_1	7, 14, 17, 12, 13, 8, 15, 18, 7	$y_1 + z_3 + y_6 + y_2 - z_1$
B.9	C_8	2, 8, 13, 16, 11, 5, 4, 3, 2	z_2	1, 6, 5, 4, 10, 15, 8, 2, 1	$-y_2 - x$
B.10	C_9	3, 9, 14, 17, 12, 6, 5, 4, 3	z_3	2, 3, 9, 16, 11, 18, 15, 8, 2	$y_3 + y_5 + z_1 - y_2$

Table 2: Fundamental cycles and their images with corresponding voltages in F018A.

The graph X can be derived from F018A through a suitable voltage assignment $\zeta: A(\text{F018A}) \rightarrow \mathbb{Z}_r$. To find this voltage assignment ζ fix the spanning tree T of F018A as the one consisting of the edges

$$\{1, 7\}, \{2, 8\}, \{3, 9\}, \{4, 10\}, \{5, 11\}, \{6, 12\}, \{7, 18\}, \{8, 13\}, \\ \{9, 14\}, \{10, 15\}, \{11, 16\}, \{12, 17\}, \{2, 3\}, \{3, 4\}, \{4, 5\}, \{5, 6\}, \{6, 1\}$$

(see also Figure 3). There are ten fundamental cycles in F018A, which are generated, respectively, by ten cotree arcs $(1, 2)$, $(7, 14)$, $(8, 15)$, $(9, 16)$, $(10, 17)$, $(11, 18)$, $(12, 13)$, $(18, 15)$, $(13, 16)$ and $(14, 17)$ (see Table 2 where all these cycles and their voltages are listed).

Now let us consider the mappings $\bar{\alpha}$ and $\bar{\beta}$ from the set $S = \{x, y_i, z_j \mid i \in \{1, 2, \dots, 6\}, j \in \{1, 2, 3\}\}$ of voltages of the ten fundamental cycles of F018A to the voltage group \mathbb{Z}_r . Since X is connected we have $\mathbb{Z}_r = \langle S \rangle$. Proposition 2.1 implies that the mappings $\bar{\alpha}$ and $\bar{\beta}$ are extended to automorphisms α^* and β^* of \mathbb{Z}_r , respectively. Also, since the lifts of β centralize the group of covering transformations, Proposition 2.3 implies that β^* is the identity automorphism of \mathbb{Z}_r . Therefore, it follows from Rows B.1 and B.4 of Table 2 that $y_2 = z_1 = -y_3$. By Rows A.3 and A.4 of Table 2, we get that $y_2^{\alpha^*} = -y_5$ and $y_3^{\alpha^*} = -y_4$, and so $y_5 = -y_4$. In other words, y_2, y_3, y_4, y_5 and z_1 are of the same

order and so $y_2, y_3, y_4, z_1 \in \langle y_5 \rangle$. Further, by Rows B.5 and B.7 of Table 2 we get that $z_2 = 2y_4 = -2y_5 \in \langle y_5 \rangle$ and $x = y_5 + y_6$, respectively, implying that in fact $\mathbb{Z}_r = \langle y_i, z_3 \mid i \in \{1, 5, 6\} \rangle$. In addition, by Row B.10 of Table 2, we have $z_3 = y_3 + y_5 \in \langle y_5 \rangle$, and Rows B.3 and B.9 of Table 2 combined together imply that $z_2 = -y_2 - x = -y_6 \in \langle y_5 \rangle$. Thus, since, by Row A.2 of Table 2, we have $y_1^{\alpha^*} = -y_6$, we can conclude that $\mathbb{Z}_r = \langle y_5 \rangle$.

By Proposition 2.2 we can, without loss of generality, assume that $y_5 = 1$. Since, by Row B.9 of Table 2, $z_2 = -y_2 - x$, the automorphism α^* gives that $y_1 + z_3 - y_4 = y_5 + x$ (see Rows A.1, A.3 and A.9 of Table 2) and so, since $y_5 = -y_4$, we have that $x = y_1 + z_3 = y_5 + y_6$. Since $-x = x^{\alpha^*} = (y_5 + y_6)^{\alpha^*} = -y_1 - y_2$, it follows that $y_1 + z_3 = x = y_1 + y_2$, and thus $z_3 = y_2 = z_1$. Row B.10 of Table 2 now implies that $0 = y_3 + y_5 - y_2$. Since $y_3 = -y_2$ it follows that $2y_2 = y_5 = -y_4$. Applying Rows A.3 and A.5 of Table 2 to this equality gives that $-2y_5 = y_3 = -y_2$. Therefore $y_2 = 2$, and consequently, by Row A.6 of Table 2, it follows that $1^{\alpha^*} = -2$. Now we get from Row A.1 of Table 2 that $-2x = -x$, and so $x = 0$. This implies that $y_6 = -y_5 = -1$ (since $x = y_5 + y_6$). However, by Row B.3 of Table 2, $y_6 = y_2 + x = y_2 = 2$ and so $2 = -1$. This shows that $r = 3$, and therefore X is isomorphic to F054A, the unique cubic symmetric graph of order 54.




Figure 4: The voltage assignment ζ on F030A. The spanning tree consists of undirected bold edges, all carrying trivial voltage.

CASE 3. $X_N \cong F030A$.

The graph F030A is illustrated in Figure 4. It is known that F030A is the unique connected cubic symmetric graph of order 30 and that this graph is in fact 5-regular (see [4]). Each subgroup of $\text{Aut}F030A$ generated by a (3, 10)-semiregular automorphism is conjugate to $\langle \beta \rangle$ where

$$\begin{aligned} \beta = & (1, 3, 7, 15, 28, 13, 30, 20, 9, 4)(2, 8, 16, 24, 22, 6, 17, 23, 19, 10) \\ & (5, 18, 27, 12, 25, 14, 26, 11, 29, 21), \end{aligned}$$

and $\text{Aut}F030A$ has two proper arc-transitive subgroups, both of order 720 acting 4-regularly. In addition, one of these two proper arc-transitive subgroups does not contain a (3, 10)-semiregular automorphism whereas the other is generated by β and

$$\begin{aligned} \alpha = & (2)(5)(11)(12)(23)(25)(1, 6)(3, 13)(4, 14)(7, 30)(8, 28) \\ & (9, 27)(10, 29)(15, 17)(16, 20)(18, 22)(19, 21)(24, 26). \end{aligned}$$

These can be checked directly using Magma [2].

Let $H = \langle \alpha, \beta \rangle$. Then, in order to show that this case is impossible, that is, that X is not a \mathbb{Z}_r -cover of F030A, it suffices to show that there is no connected \mathbb{Z}_r -cover X of F030A such that ρ projects to β (therefore, the lifts of β centralizes the group N of covering transformations) and that H lifts to a subgroup of $\text{Aut} X$.

For this purpose observe that any such cover can be derived from F030A through a suitable voltage assignment $\zeta: A(F030A) \rightarrow \mathbb{Z}_r$. To find this voltage assignment ζ fix the spanning tree T of F030A as the one consisting of the edges

$$\begin{aligned} & \{1, 2\}, \{2, 5\}, \{3, 8\}, \{8, 18\}, \{7, 16\}, \{16, 27\}, \{15, 24\}, \{12, 24\}, \{22, 28\}, \{22, 25\}, \\ & \{6, 13\}, \{6, 14\}, \{17, 30\}, \{17, 26\}, \{20, 23\}, \{11, 23\}, \{9, 19\}, \{19, 29\}, \{4, 10\}, \{10, 21\}, \\ & \{3, 7\}, \{7, 15\}, \{15, 28\}, \{13, 28\}, \{13, 30\}, \{20, 30\}, \{9, 20\}, \{4, 9\}, \{1, 4\}. \end{aligned}$$

(see also Figure 4). There are sixteen fundamental cycles in F030A, which are generated, respectively, by sixteen cotree arcs (1, 3), (2, 6), (8, 17), (16, 23), (24, 19), (22, 10), (5, 12), (12, 26), (26, 21), (21, 27), (27, 14), (14, 29), (29, 18), (18, 25), (25, 11) and (11, 5) (see Table 3 where all these cycles and their voltages are listed).

The set $S = \{x, y_i, z_j \mid i \in \{1, 2, \dots, 5\}, j \in \{1, 2, \dots, 10\}\}$ of voltages of the sixteen fundamental cycles of F030A generates the voltage group \mathbb{Z}_r . By Proposition 2.3, the mapping $\bar{\beta}$ extends to the identity automorphism of \mathbb{Z}_r . Thus, Rows B.2 – B.6 of Table 3 imply that $y_2 = y_3 = y_4 = y_5 = -y_1$ and $x = 2y_1$, whereas Rows B.7 – B.16 of Table 3 imply that $z_1 = z_4 = z_7$ and $z_2 = z_3 = z_5 = z_6 = z_8 = z_9 = z_{10} = z_1 - x$. It follows that $\mathbb{Z}_r = \langle y_1, z_1 \rangle$. Moreover, applying the automorphism α^* to $z_2 = z_5$ we get that $-z_2 + y_2 = x + y_2$ and so $x = -z_2$ (see Rows A.8 and A.11 of Table 3). This shows that $z_1 = 0$ and therefore $\mathbb{Z}_r = \langle y_1 \rangle$. But, however, Row A.4 of Table 3 implies that $y_3^{\alpha^*} = 0$, and so $y_1 = -y_3 = 0$, a contradiction. ■

Row		C	$\zeta(C)$	C^α	$\zeta(C^\alpha)$
A.1	C_0	1, 3, 7, 15, 28, 13, 30, 20, 9, 4, 1	x	6, 13, 30, 17, 8, 3, 7, 16, 27, 14, 6	$-y_2 + z_5$
A.2	C_1	1, 2, 6, 13, 30, 20, 9, 4, 1	y_1	6, 2, 1, 3, 7, 16, 27, 14, 6	$-y_1 + x + z_5$
A.3	C_2	3, 8, 17, 30, 13, 28, 15, 7, 3	y_2	13, 28, 15, 7, 3, 8, 17, 30, 13	y_2
A.4	C_3	7, 16, 23, 20, 30, 13, 28, 15, 7	y_3	30, 20, 23, 16, 7, 3, 8, 17, 30	$-y_3 + y_2$
A.5	C_4	15, 24, 19, 9, 20, 30, 13, 28, 15	y_4	17, 26, 21, 27, 16, 7, 3, 8, 17	$z_3 + z_4 + y_2$
A.6	C_5	28, 22, 10, 4, 9, 20, 30, 13, 28	y_5	8, 18, 29, 14, 27, 16, 7, 3, 8	$-z_7 - z_6 - z_5$
A.7	C_6	1, 2, 5, 12, 24, 15, 28, 13, 30, 20, 9, 4, 1	z_1	6, 2, 5, 12, 26, 17, 8, 3, 7, 16, 27, 14, 6	$-y_1 + z_1 + z_2 - y_2 + z_5$
A.8	C_7	15, 24, 12, 26, 17, 30, 13, 28, 15	z_2	17, 26, 12, 24, 15, 7, 3, 8, 17	$-z_2 + y_2$
A.9	C_8	30, 17, 26, 21, 10, 4, 9, 20, 30	z_3	7, 15, 24, 19, 29, 14, 27, 16, 7	$y_4 - z_6 - z_5$
A.10	C_9	4, 10, 21, 27, 16, 7, 15, 28, 13, 30, 20, 9, 4	z_4	14, 29, 19, 9, 20, 30, 17, 8, 3, 7, 16, 27, 14	$z_6 - y_2 + z_5$
A.11	C_{10}	7, 16, 27, 14, 6, 13, 28, 15, 7	z_5	30, 20, 9, 4, 1, 3, 8, 17, 30	$x + y_2$
A.12	C_{11}	13, 6, 14, 29, 19, 9, 20, 30, 13	z_6	3, 1, 4, 10, 21, 27, 16, 7, 3	$-x + z_4$
A.13	C_{12}	9, 19, 29, 18, 8, 3, 7, 15, 28, 13, 30, 20, 9	z_7	27, 21, 10, 22, 28, 13, 30, 17, 8, 3, 7, 16, 27	$-z_4 - y_5 - y_2$
A.14	C_{13}	3, 8, 18, 25, 22, 28, 15, 7, 3	z_8	13, 28, 22, 25, 18, 8, 17, 30, 13	$-z_8 + y_2$
A.15	C_{14}	28, 22, 25, 11, 23, 20, 30, 13, 28	z_9	8, 18, 25, 11, 23, 16, 7, 3, 8	$z_8 + z_9 - y_3$
A.16	C_{15}	20, 23, 11, 5, 2, 1, 4, 9, 20	z_{10}	16, 23, 11, 5, 2, 6, 14, 27, 16	$y_3 + z_{10} + y_1 - z_5$
		C	$\zeta(C)$	C^β	$\zeta(C^\beta)$
B.1	C_0	1, 3, 7, 15, 28, 13, 30, 20, 9, 4, 1	x	3, 7, 15, 28, 13, 30, 20, 9, 4, 1, 3	x
B.2	C_1	1, 2, 6, 13, 30, 20, 9, 4, 1	y_1	3, 8, 17, 30, 20, 9, 4, 1, 3	$y_2 + x$
B.3	C_2	3, 8, 17, 30, 13, 28, 15, 7, 3	y_2	7, 16, 23, 20, 30, 13, 28, 15, 7	y_3
B.4	C_3	7, 16, 23, 20, 30, 13, 28, 15, 7	y_3	15, 24, 19, 9, 20, 30, 13, 28, 15	y_4
B.5	C_4	15, 24, 19, 9, 20, 30, 13, 28, 15	y_4	28, 22, 10, 4, 9, 20, 30, 13, 28	y_5
B.6	C_5	28, 22, 10, 4, 9, 20, 30, 13, 28	y_5	13, 6, 2, 1, 4, 9, 20, 30, 13	$-y_1$
B.7	C_6	1, 2, 5, 12, 24, 15, 28, 13, 30, 20, 9, 4, 1	z_1	3, 8, 18, 25, 22, 28, 13, 30, 20, 9, 4, 1, 3	$z_8 + x$
B.8	C_7	15, 24, 12, 26, 17, 30, 13, 28, 15	z_2	28, 22, 25, 11, 23, 20, 30, 13, 28	z_9
B.9	C_8	30, 17, 26, 21, 10, 4, 9, 20, 30	z_3	20, 23, 11, 5, 2, 1, 4, 9, 20	z_{10}
B.10	C_9	4, 10, 21, 27, 16, 7, 15, 28, 13, 30, 20, 9, 4	z_4	1, 2, 5, 12, 24, 15, 28, 13, 30, 20, 9, 4, 1	z_1
B.11	C_{10}	7, 16, 27, 14, 6, 13, 28, 15, 7	z_5	15, 24, 12, 26, 17, 30, 13, 28, 15	z_2
B.12	C_{11}	13, 6, 14, 29, 19, 9, 20, 30, 13	z_6	30, 17, 26, 21, 10, 4, 9, 20, 30	z_3
B.13	C_{12}	9, 19, 29, 18, 8, 3, 7, 15, 28, 13, 30, 20, 9	z_7	4, 10, 21, 27, 16, 7, 15, 28, 13, 30, 20, 9, 4	z_4
B.14	C_{13}	3, 8, 18, 25, 22, 28, 15, 7, 3	z_8	7, 16, 27, 14, 6, 13, 28, 15, 7	z_5
B.15	C_{14}	28, 22, 25, 11, 23, 20, 30, 13, 28	z_9	13, 6, 14, 29, 19, 9, 20, 30, 13	z_6
B.16	C_{15}	20, 23, 11, 5, 2, 1, 4, 9, 20	z_{10}	9, 19, 29, 18, 8, 3, 1, 4, 9	$z_7 - x$

Table 3: Fundamental cycles and their images with corresponding voltages in F030A.

References

- [1] N. Biggs, “Algebraic graph theory”, 2nd ed., Cambridge University Press, Cambridge, 1993.
- [2] W. Bosma, J. Cannon, and C. Playoust, The MAGMA Algebra System I: The User Language, *J. Symbolic Comput.* **24** (1997) 235–265.
- [3] I. Z. Bouwer (ed.): The Foster Census, Winnipeg, 1988.
- [4] M. D. E. Conder and P. Dobcsányi, Trivalent symmetric graphs on up to 768 vertices, *J. Combin. Math. & Combin. Comp.* **40** (2002), 41–63.
- [5] S. F. Du, J. H. Kwak and M. Y. Xu, On 2-arc-transitive covers of complete graphs with covering transformation group \mathbb{Z}_p^3 , *J. Combin. Theory Ser. B* **93** (2005), 73–93.
- [6] R. Frucht, How to describe a graph, *Ann. N. Y. Acad. Sci.* **175** (1970), 159–167.
- [7] R. Frucht, J. E. Graver and M. E. Watkins, The groups of the generalized Petersen graphs, *Proc. Camb. Phil. Soc.* **70** (1971), 211–218.
- [8] H. H. Glover and D. Marušič, Hamiltonicity of cubic Cayley graph, *J. Eur. Math. Soc.* **9** (2007), 775–787.
- [9] J. L. Gross and T. W. Tucker, Generating all graph coverings by permutation voltage assignment, *Discrete Math.* **18** (1977), 273–283.
- [10] J. L. Gross and T. W. Tucker, “Topological Graph Theory”, Wiley - Interscience, New York, 1987.
- [11] M. Herzog and G. Kaplan, Large cyclic subgroups contain non-trivial normal subgroups, *J. Group Theory* **4** (2001), 247–253.
- [12] S. Hong, J. H. Kwak and J. Lee, Regular graph coverings whose covering transformation groups have the isomorphism extension property, *Discrete Math* **168** (1996), 85–105.
- [13] R. Jajcay and J. Širáň, Small vertex-transitive graphs of given degree and girth, *Ars Math. Contemp.* **4** (2011), 375–384.
- [14] I. N. Kagno, Desargues’ and Pappus’ Graphs and Their Groups, *Amer. J. Math.* **69** (1947), 859–862.
- [15] P. Lorimer, Vertex-transitive graphs: symmetric graphs of prime valency, *J. Graph Theory* **8** (1984), 55–68.
- [16] P. Lorimer, The construction of Tutte’s 8-cage and the Conder graph, *J. Graph Theory* **13** (1989), 553–557.
- [17] A. Malnič, Group actions, coverings and lifts of automorphisms, *Discrete Math.* **182** (1998), 203–218.
- [18] A. Malnič, R. Nedela and M. Škoviera, Lifting graph automorphisms by voltage assignments, *Europ. J. Combin.* **21** (2000), 927–947.
- [19] D. Marušič and T. Pisanski, Symmetries of hexagonal molecular graphs on the torus, *Croat. Chem. Acta* **73** (2000), 969–981.
- [20] D. Marušič and T. Pisanski, The remarkable generalized Petersen graph $G(8, 3)$, *Mathematica Slovaca* **50** (2000), 117–121.
- [21] T. Pisanski, A classification of cubic bicirculants, *Discrete Math.* **307** (2007), 567–578.
- [22] W. T. Tutte, A family of cubical graphs, *Proc. Cambridge Philos. Soc.* **43** (1947), 459–474.