

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.



ISSN: 0015-5659

e-ISSN: 1644-3284

Two cases of variations in inferior thyroid arterial pattern and their clinical implications

Authors: S. S. Novakov, S. D. Delchev

DOI: 10.5603/FM.a2022.0032

Article type: Case report

Submitted: 2022-02-08

Accepted: 2022-03-14

Published online: 2022-03-29

This article has been peer reviewed and published immediately upon acceptance. It is an open access article, which means that it can be downloaded, printed, and distributed freely, provided the work is properly cited.

Articles in "Folia Morphologica" are listed in PubMed.

Two cases of variations in inferior thyroid arterial pattern and their clinical implications

S.S. Novakov, S.D. Delchev, Two cases of rare thyroid arterial pattern

S.S. Novakov, S..D. Delchev

Department of Human Anatomy, Histology and Embryology, Medical University, Plovdiv, Bulgaria

Address for correspondence: S.S. Novakov, Chief Assistant Professor, Department of Anatomy, Histology and Embryology, Medical University, 15A V. Aprilov bul., 4002 Plovdiv, Bulgaria, e-mail: stoyan.novakov@mu-plovdiv.bg

ABSTRACT

Vascular variations are the most common ones in humans. Inferior thyroid artery arises from the thyrocervical trunk in 90.5%, from subclavian in 7.5%, and very rarely from the common carotid, aortic arch, brachiocephalic, internal thoracic, pericardiacophrenic, or vertebral. Thyroid ima artery is more common variety found in up to 12.2% of the population. Two cadavers dissected in the anatomy department are presented with variations in the blood supply of the thyroid gland. The first case was a 61-year-old man with middle thyrothyroid artery arising from the common carotid on the right side and inferior thyroid as a branch of the common carotid on the left. The second case was an 85-year-old female without inferior thyroid arteries bilaterally, replaced by thyroid ima arising from brachiocephalic artery. The awareness of such arterial pattern is crucial for the specialists in imaging and preoperative diagnosing and escaping eventual iatrogenic complications of thyroid gland.

Key words: middle thyrothyroid artery, thyroid ima artery, human anatomy variations

INTRODUCTION

The human anatomy variations are part of the human anatomy science from its dawn till nowadays and are still an essential part of the gross anatomy research. The main significance of that type of evidence is for the invasive medical procedures and especially surgery [5]. The thyroid gland is an endocrine cervical gland which is well vascularized. Arterial supply comprises 2 superior thyroid arteries (STAs) originating from the external carotid arteries and 2 inferior thyroid arteries (ITAs) originating from the thyrocervical trunk, branch of the subclavian arteries [3, 6, 9]. Occasionally there is an accessory artery termed thyroid ima (TIA), which when present (0.4 to 12.2%), originates from the brachiocephalic trunk (43.3–86.7%), from the common carotid artery (CCA) (2–50%) or aortic arch (0–66.7%) [11, 13]. Though rare in humans, middle thyroid artery is found in some animals, such as guinea pig and rabbit [10] and avian species like budgerigar [8]. The awareness of the anatomic variations in the blood supply of thyroid gland are of crucial significance when planning a surgical approach in the infrahyoid region. Knowledge of variability in thyroid vascularization will facilitate the surgical procedure itself, make it safer and minimize the risk of hemorrhage [5].

CASE 1 PRESENTATION

Formalin/ethanol-embalmed cadavers were used for dissection during the anatomy lessons. The length and diameter of the structures were measured by calipers.

A 61-year-old male cadaver was dissected in the department of anatomy and multiple variations in the bilateral arterial blood supply of the thyroid gland were registered. An accessory artery supplying both thyroid and thymic glands was found on the right side. We agreed with previous authors and named it middle thymothyroid artery [1]. That was a spatial variation also belonging to the category of consistency [12]. It originated from the CCA 48 mm proximal to its bifurcation. The diameter of the CCA was 8 mm and the length of the unusual branch was 17 mm till its division into 3 thyroid branches (upper, middle and lower). The diameter of the middle thymothyroid artery was 2.5 mm and its length with its longest branch as it enters thyroid gland was 63 mm. The

continuation of the artery to the thymus (thymic branch) was 1.5 mm in diameter. The STA and ITA had their standard origin and position. On the left side of the same cadaver the ITA had rare origin from the CCA. The former started at 50 mm from the bifurcation of CCA, which was 6 mm wide. The length of the variable ITA to the release of its branches was 8 mm and its superior branch, which directly continued to the gland was 10 mm long and 1.5 mm wide, while the inferior one was 16 mm long and 0.8 mm wide. The diameter of the ITA was 2.5 mm. The rest of the structures on this side were with standard topography (Fig.1).

CASE 2 PRESENTATION

During the same semester another variation was found in 85-year-old female cadaver. The routine dissection of the neck exposed bilateral absence of the ITAs. The arterial blood supply was compensated by the presence of relatively larger STAs, especially the one on the right side, and prominent TIA arising from the innominate artery. The TIA emerged in the midline 28 mm from the origin of the brachiocephalic trunk measured along its left border. It had a spiral course and diameter of 4 mm as well as a length of 21 mm to its division into: 1. Right branch – 3 mm long and 2 mm wide common trunk, which divides into thymic artery and right ITA. 2. Left branch – it was a continuation of the main artery as left ITA (2 mm in width), which climbed to the left lobe of the thyroid gland and before reaching it released a small branch for the isthmus of the gland. The rest of the topographical anatomy of the neck was standard (Fig. 2).

DISCUSSION

The origin of the ITA from CCA is very rare. Therefore, finding it in the gross anatomy dissecting lab is worth presenting it. The Case 1 of this report describes bilateral variants of the blood supply of the thyroid gland, which are both rare aberrant origins of ITA from CCA (about 1%) [5]. There is an embryological basis of such variations as they are associated with time and stages of the development of the neck arteries by the aortic arches and the appearance and migration of the thyroid gland. That period between the third and seventh week of development is when the formation of the aortic arch branches

and the synchronous descend of the thyroid gland occur and it is not unreasonable to presume that the vascularization of the gland would occasionally come from other vessels such as the subclavian, the vertebral or the CCA [4,5]. The thyroid ima found in Case 2 is more common (up to 12.2%) but not usual which makes it an obstacle in thyroid surgery. Phylogenetically, the origin of this variable artery is supposed to appear from the original vascular network which connects brachiocephalic, aortic arch and carotids. During the later development the primary vessels disappearing by fusion may either supplement or substitute the regular blood supply of the thyroid gland, namely the superior and inferior thyroid arteries as the thyroid ima [7].

Though very rare the ITA arising from CCA exposes two major risks for surgical intervention in the area: hemorrhage by injuring this artery when on unusual position and increased possibility of injuring recurrent laryngeal nerve [2, 6].

Carotid endarterectomy as an established procedure for stroke prevention is also related to the above-mentioned arterial pattern of the CCA, which uniqueness could be crucial for successful removal of the atherosclerotic plaque and the avoidance of postoperative complications [5].

Interventional radiology procedures like endovascular aneurysm reconstruction, chemoembolization techniques like tumor embolization, and arterial embolization for the treatment of thyroid conditions requiring ablation might be affected in case of such variations [2, 5].

CONCLUSIONS

The role of anatomy science on variations and especially those of the vascular system and in the neck region is essential. Surgeons and interventional radiologists might come upon such rare variations during procedures in the neck area and any preceding awareness will be valuable to limit the incidence of possible iatrogenic complications.

Acknowledgements

The authors would like to thank the people who donated their bodies for teaching students and scientific studies, and the technical staff of the Anatomy, Histology and Embryology Department for their support.

Conflict of interest: None declared

REFERENCES

1. Banneheka S, Chiba S, Fukazawa M et al. Middle thymothyroid artery arising from the common carotid artery: case report of a rare variation. *Anat Sci Int.* 2010; 85(4):241-4. doi: 10.1007/s12565-009-0055-9.
2. Branca JJV, Lascialfari Bruschi A, Pilia AM, Carrino D, Guarnieri G, Gulisano M, Pacini A, Paternostro F. The Thyroid Gland: A Revision Study on Its Vascularization and Surgical Implications. *Medicina (Kaunas).* 2022 Jan 17;58(1):137. doi: 10.3390/medicina58010137.
3. Burlakoti A, Massy-Westropp N. Bilateral variant thyroid arteries *Int J Anat Var.* 2015; 8:43–46.
4. Cigali BS, Ulucam E, Bozer C. Accessory inferior thyroid artery and internal thoracic artery originating from the thyrocervical trunk. *Anat Sci Int.* 2008 Dec;83(4):283-5. doi: 10.1111/j.1447-073X.2007.00206.x.
5. Mariolis-Sapsakos T, Kalles V, Papapanagiotou I et al., Bilateral aberrant origin of the inferior thyroid artery from the common carotid artery. *Surg Radiol Anat.* 2014; 36(3):295-7. doi: 10.1007/s00276-013-1153-x.
6. Ngo Nyeki AR, Pelsoni G, Karenovics W et al. Aberrant origin of the inferior thyroid artery from the common carotid artery: a rare anatomical variation. *Gland Surg.* 2016; 5(6):644-646. doi: 10.21037/g.s.2016.12.03.
7. Raj S, Mohiyuddin A, Merchant S et al. Thyroidea ima Artery: A Report of Two Cases. *Int J Head Neck Surg.* 2014; 5(2):89-90.
8. Radek T, Piasecki T. The topographical anatomy and arterial supply of the thyroid and parathyroid glands in the budgerigar (*Melopsittacus undulatus*). *Folia Morphol.* 2004; 63(2):163-71.
9. Standring S. *Gray's Anatomy: The Anatomical Basis of Clinical Practice.* 40th Ed., Spain, Churchill Livingstone. 2008: 462.

10. Won HS, Han SH, Oh CS, Chung IH. Superior and middle thyroid arteries arising from the common carotid artery. *Surg Radiol Anat.* 2011 Sep;33(7):645-7. doi: 10.1007/s00276-011-0781-2.
11. Yalcin B. Thyroid gland. In: Tubbs RS, Shoja MM, Loukas M (eds.) *Bergman's comprehensive encyclopedia of human anatomic variation*, 1st edn. Wiley-Blackwell, New Jersey 2016: 1199–1200.
12. Yammine K. Evidence-based anatomy. *Clin Anat.* 2014; 27(6):847-852. doi:10.1002/ca.22397. doi: 10.5005/jp-journals-10001-1188.
13. Yohannan DG, Rajan R, Chandran AB et al. An unusual origin and course of the thyroidea ima artery, with absence of inferior thyroid artery bilaterally. *Surg Radiol Anat.* 2019;41(2):235-237. doi: 10.1007/s00276-018-2122-1.

Figure 1. Middle thymothyroid artery on the right side and atypical origin of inferior thyroid artery on the left side; RCTM/LCTM — right/left cricothyroid muscle; CC — cricoid cartilage; Thr — thyroid gland; Thm — thymus; RCCA/LCCA — right/left common carotid artery; RBV/LBV — right/left brachiocephalic vein; RSV/LSV — right/left subclavian vein; LIJV — left internal jugular vein; 1 — middle thymothyroid artery; 2 — upper thyroid branch; 3 — middle thyroid branch; 4 — lower thyroid branch; 5 — thymic branch; 6 — inferior thyroid artery; 7 — superior branch of ITA; 8 — inferior branch of ITA; 9 — inferior thyroid veins.

Figure 2. Thyroid ima artery replacing missing inferior thyroid arteries; RSTM — right sternothyroid muscle; Thr — thyroid gland; Thm – thymus; RCCA/LCCA — right/left common carotid artery; RSA — right subclavian artery; BCT — brachiocephalic trunk; RIJV/LIJV — right/left internal jugular vein; RSV/LSV — right/left subclavian vein; RBV/LBV — right/left brachiocephalic vein; 1 — thyroid ima artery; 2 — right branch; 3 — thymic artery; 4 — right inferior thyroid branch; 5 — left inferior thyroid branch; 6 — thyrocervical trunk; 7 — suprascapular artery; 8 — transverse cervical artery; 9 — ascending cervical artery; 10 — right superior thyroid artery; 11 — left superior thyroid artery.



