



KARDIOLOGIA POLSKA

Polish Heart Journal
The Official Peer-reviewed Journal
of the Polish Cardiac Society
since 1957

Online first

This is a provisional PDF only. Copyedited and fully
formatted version will be made available soon

ISSN 0022-9032

e-ISSN 1897-4279

Arrhythmia coming from the transplanted heart: What problems does it generate?

Authors: Anna Drohomirecka, Michał M Farkowski, Aleksander Maciąg, Mariusz Pytkowski,
Tomasz Zieliński

Article type: Letter to the Editor

Received: June 3, 2022

Accepted: June 6, 2022

Early publication date: June 6, 2022

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

Arrhythmia coming from the transplanted heart: What problems does it generate?

Short title: Arrhythmia coming from the transplanted heart

Anna Drohomirecka¹, Michał M Farkowski², Aleksander Maciąg², Mariusz Pytkowski², Tomasz Zieliński¹

¹Department of Heart Failure and Transplantation, National Institute of Cardiology, Warszawa, Poland

²²nd Department of Heart Arrhythmia, National Institute of Cardiology, Warszawa, Poland

Correspondence to:

Anna Drohomirecka, PhD, MD,
Department of Heart Failure and Transplantation,
National Institute of Cardiology,
Alpejska 42, 04–628 Warszawa, Poland,
phone: +48 22 343 44 62,
e-mail: adrohomirecka@tlen.pl

Conflict of interest: None declared.

We have recently read with a great interest the article by Myrda et al. [1]. The authors described a case of a 63-year-old patient who was diagnosed in the second year after heart transplantation with asymptomatic intermittent preexcitation syndrome and subsequently underwent radiofrequency catheter ablation of an left-sided accessory pathway (AP). Therefore we would like to place some more emphasis on few aspects of the issue.

Although the problem of cardiac arrhythmias due to pre-existing AP or dual atrioventricular nodal physiology in donor hearts is not new and occurs in less than 0.5% of patients [2], it still raises some controversies. Usually, the history of supraventricular arrhythmia in the donor is absent or not known and the first episode of arrhythmia in the recipient may be provoked by changes in autonomic tone (due to cardiac innervation and catecholamine excess) which affect the

substrate [3, 4]. Symptomatic patients should be diagnosed and treated in the same manner as non-transplanted patients. In asymptomatic recipients, even with a negative result of an invasive electrophysiological study, there remains the possibility that the clinical significance of arrhythmia is underestimated, especially if the heart was harvested after donor death caused by accidental head injury. In those cases loss of consciousness due to arrhythmia episode often can not be excluded.

Classic indications for ablation of accessory pathways include supraventricular tachyarrhythmia or a short refractory period [5]. Moreover, in patient after heart transplantation, psychological aspects of recipient's attitude should be taken into account. An elevated level of patient anxiety could be expected as soon as the patient is informed that "the new heart is not completely healthy". The decision about ablation or its abandonment should be made after careful consideration of risks related to arrhythmia, intervention and patient preferences. At this point, it is worth noting that an invasive strategy has an advantage over pharmacotherapy in patients receiving immunosuppressive treatment: drugs such as verapamil, diltiazem, propafenone interact with calcineurin inhibitors, increasing the risk of complications of immunosuppressive agents.

As mentioned above, some patients remained asymptomatic and, on the other hand, first signs or symptoms of arrhythmia coming from the transplanted heart can occur months or years after transplantation. When the episodes are short and not frequent, the time to diagnosis may be delayed. Thus, we would like to ask the Authors if they have any protocol in standard care settings for monitoring of arrhythmias occurrence in the early and late posttransplantation period.

Based on our center experience we have observed that in the last few years on average one patient annually is diagnosed with an AP or dual atrioventricular nodal physiology, which was practically not observed more than 10 years ago. Do the Authors have any knowledge about the prevalence of this type arrhythmia in their center?

And last but not least, we would like to encourage the discussion about harvesting a heart from a donor with known AP. In our opinion it is completely acceptable solution as long as the patients give their informed consent and monitoring is provided along with the possibility of invasive treatment. In the era of the shortage of organ donation, when in more and more countries programs of heart donation after cardiac death are implemented, an easy treatable electrophysiological abnormality should not be a contraindication for harvesting a heart.

REFERENCES

1. Myrda K, Nowowiejska-Wiewióra A, Zembala M. Radiofrequency catheter ablation of an asymptomatic intermittent accessory pathway after heart transplantation. *Kardiol Pol.* 2022 [epub ahead of print], doi: [10.33963/KP.a2022.0112](https://doi.org/10.33963/KP.a2022.0112), indexed in Pubmed: [35475463](https://pubmed.ncbi.nlm.nih.gov/35475463/).
2. Vaseghi M, Boyle NG, Kedia R, et al. Supraventricular tachycardia after orthotopic cardiac transplantation. *J Am Coll Cardiol.* 2008; 51(23): 2241–2249, doi: [10.1016/j.jacc.2008.02.065](https://doi.org/10.1016/j.jacc.2008.02.065), indexed in Pubmed: [18534271](https://pubmed.ncbi.nlm.nih.gov/18534271/).
3. Joglar JA, Wan EY, Chung MK, et al. Management of arrhythmias after heart transplant: current state and considerations for future research. *Circ Arrhythm Electrophysiol.* 2021; 14(3): e007954, doi: [10.1161/CIRCEP.120.007954](https://doi.org/10.1161/CIRCEP.120.007954), indexed in Pubmed: [33685207](https://pubmed.ncbi.nlm.nih.gov/33685207/).
4. Hamon D, Taleski J, Vaseghi M, et al. Arrhythmias in the heart transplant patient. *Arrhythm Electrophysiol Rev.* 2014; 3(3): 149–155, doi: [10.15420/aer.2014.3.3.149](https://doi.org/10.15420/aer.2014.3.3.149), indexed in Pubmed: [26835083](https://pubmed.ncbi.nlm.nih.gov/26835083/).
5. Brugada J, Katritsis DG, Arbelo E, et al. 2019 ESC Guidelines for the management of patients with supraventricular tachycardiaThe Task Force for the management of patients with supraventricular tachycardia of the European Society of Cardiology (ESC). *Eur Heart J.* 2020; 41(5): 655–720, doi: [10.1093/eurheartj/ehz467](https://doi.org/10.1093/eurheartj/ehz467), indexed in Pubmed: [31504425](https://pubmed.ncbi.nlm.nih.gov/31504425/).

