

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.

# REPORTS OF PRACTICAL ONCOLOGY AND RADIOTHERAPY

ISSN: 1507-1367

e-ISSN: 2083-4640

## **The use of the normal tissue non-complication probability (NTCP0) methodology as a new alternative of assessing side-effects in brachytherapy treatments**

**Authors:** Terman Frometa-Castillo, Anil Pyakuryal, Ganesh Narayanasamy, Asghar Mesbahi, Amadeo Wals-Zurita

**DOI:** 10.5603/RPOR.a2022.0063

**Article type:** Research paper

**Published online:** 2022-06-01

This article has been peer reviewed and published immediately upon acceptance. It is an open access article, which means that it can be downloaded, printed, and distributed freely, provided the work is properly cited.

# The use of the normal tissue non-complication probability (NTCP0) methodology as a new alternative of assessing side-effects in brachytherapy treatments

**Running title:** NTCP0 in the BT

10.5603/RPOR.a2022.0063

Terman Frometa-Castillo<sup>1</sup>, Anil Pyakuryal<sup>2</sup>, Ganesh Narayanasamy<sup>3</sup>, Asghar Mesbahi<sup>4</sup>, Amadeo Wals-Zurita<sup>5</sup>

<sup>1</sup>*Owner at Statistical models project, LLC, Chicago, IL, United States*

<sup>2</sup>*University of District of Columbia, Division of Science and Mathematics, Washington, DC, United States*

<sup>3</sup>*University of Arkansas for Medical Sciences, Little Rock, United States*

<sup>4</sup>*Tabriz University of Medical Sciences, Tabriz, Iran*

<sup>5</sup>*Hospital Universitario Virgen Macarena, Sevilla, Spain*

**Address for correspondence:** Terman Frometa-Castillo, Statistical models project, LLC, Chicago, IL, United States, tel: 312-687-6422; e-mail: [terman.frometa@gmail.com](mailto:terman.frometa@gmail.com)

## Abstract

**Background:** The NTCP methodology evaluating side-effects (S-Es) was initially used in radiotherapy (RT), and later was extended to brachytherapy (BT). The NTCP0 methodology has been recently introduced in RT. Given the advantages, this methodology could replace NTCP.

**Materials and methods:** Revisions of studies related to use of NTCP in the evaluations of S-Es in BT. Development of the first versions of two Matlab applications of the NTCP0 methodology. These applications have three options. Two of them employ the well-known aspects of a phenomenological model, or the probabilistic relationship between NTCP0 and total NTCP (TNTCP) that is the  $sum(NTCP(x_i))$   $i$ :  $i^{th}$  complication  $i:1..nc$ : Number of complications; where  $NTCP0 = 100\% - TNTCP$ ; and the third option assumes a  $NTCP(x_i)$  discrete probabilistic distribution generated by the binomial distribution, where one of its parameters is automatically obtained from a database of the Disease locations Vs. Late complications.

**Results:** The NTCP0cal and NTCP0calDr Matlab applications have been developed, and respectively used for fractional continuous low dose-rate BT.

**Conclusions:** NTCP0 is defined as the ratio of the number of patients without acute/late complications and total of them, and also can be obtained using our Matlab applications. NTCP0 works do not disregard the last 10–15 years of NTCP research; but NTCP0 was not considered during these years. A generic example was used for showing the variations of the late complications and NTCP0 for a BT treatment of a constant number of fractions and six different dose per fraction values.

**Key words:** NTCP; binomial distribution; LKB NTCP model; side-effect; brachytherapy

## Introduction

The safety of treatments with drugs is an aspect that must be evaluated in the pre-clinical phases of development of a drug before using it in humans; and must be reported during the clinical treatments. As a widely used drug treatment, BT has probabilistic levels of cure and side-effects (S-Es). The normal tissue complication probability (NTCP) is a way of evaluating S-E in radiation treatments. Regardless of the level of toxicity of any treatment, there is a probabilistic level of safety, which is a complement of the global toxicity; i.e., total NTCP (TNTCP) that is the  $\sum(NTCP(x_i))$   $i: i^{th}$  complication  $i:1..nc$ : Number of complications.

Whatever specific BT treatment given to a homogenous population with specific patients having a specific tumor has its own  $NTCP(x_i)$  discrete probabilistic distribution (DPD), where  $NTCP0 = NTCP(0)$ .

Individual  $NTCP(x_i)$  has been modeled with complex analytical models, like Lyman-Kutcher-Burman (LKB) NTCP model, as shown in [1–2]; function of an independent variable (IV), then it was necessary to formulate analytical expressions for these IVs in order to determine an equivalent uniform dose (EUD) or an effective dose (Deff).

As a result of a radiation treatment, the volume of an organ at risk (OAR) generally receives a heterogenous distribution of dose. Based on this distribution, some NTCP models have been developed, such as the LKB and Relative seriality of [3].

NTCP0 is a metric associated to safety, which is the ratio between the number of patients without acute/late complications and the total number of them given a radiation treatment, well-characterized by its variables and factors. This is not associated with OARs, but non-complications. The NTCP0

phenomenological model of [4], the SMp NTCP0(D), is a function of the prescribed dose ( $D_{pres}$  or  $D=n*d$ ). This model should be used for a constant number of fractions (n) and a range of dose per fraction (d), or vice versa.

NTCP0 value can be determined from experimental/observational data; or from assuming a determined NTCP( $x_i$ ) DPD. There are developed methodologies that mathematically generate DPDs, as described in [5] and [6]. Introducing NTCP0 and its phenomenological SMp models in the BT will be advantageous compared to the current NTCP methodologies.

The SMp NTCP0(D) and SMp NTCP0(R0) phenomenological models, where R0 is the initial dose-rate, are simple and not dose-volume histogram (DVH)-based; i.e., the DVHs of the OARs are irrelevant for these models. In other words, the new NTCP0 methodologies of evaluating S-E will not require the current DVH calculations for the OARs. NTCP0 is a new alternative of evaluating S-Es, instead of the habitual NTCP methodologies.

Given inherent probabilistic aspects of a specific stochastic process (SP) with more than one outcome, like normal complications in a BT treatment given to a specific population under specific circumstances; then:

- whatever specific BT treatment is associated with NTCP( $x_i$ ) DPD;
- $NTCP0 = NTCP(0)$ , and  $NTCP0 = 100\% - TNTCP$ ;
- as a SP, the normal complications have their deterministic and stochastic regions. The SMp NTCP0 parameters ( $TD_{min}$ ,  $TD_{max}$ ,  $R0_{min}$  and  $R0_{max}$ ) are respectively the lower and upper limits of the stochastic region.


NTCP0cal and NTCP0calDr applications calculate NTCP0 using three options. The first of them is related to phenomenological models, in particular SMp NTCP0(D) and SMp NTCP0(R0) that are probabilistic-decreasing functions, and appropriate for describing the mean radiobiological behavior of NTCP0 in the function of D and R0, respectively. The second option is based on the probabilistic relationship between NTCP0 and TNTCP like  $NTCP0 = 100\% - TNTCP$ .

Contrary to TCP calculations that can be done with computational simulations, for NTCP0 it is very difficult or impossible due to numerous parameters and variables involved; for this reason, the second and third options use an assumed or known NTCP( $x_i$ ) DPDs. In the third, we employ the binomial distribution (BD). As described in [5], the BD is an excellent-mathematical generator of these kind distributions

## Results

### *The NTCP0cal application*

This application provides three options, two of them employ the well-known aspects of a phenomenological model, or the relationship with TNTCP; and the third option determines NTCP0 from an assumed  $NTCP(x_i)$  DPD generated from the BD, where one of its parameters is automatically defined from a databased of the Disease locations Vs. Late complications. Figure 1 is the flow chart for determining NTCP0 in a fractionated BT treatment with  $D_{pres}$ .



**Figure 1.** Diagram of procedures for determining NTCP0 in a fractional BT treatment

The steps for executing the NTCP0cal are:

— select one of the three panels pressing the “Use” button of the desired panel;

If the selection is Panel 1 “Using the SMp NTCP0 parameters”; introduce  $d$  of the  $D_{pres}$ , and the SMp NTCP0 parameters ( $TD_{min}$ ,  $TD_{max}$  and  $pN0$ ).

If the selection is Panel 2 “Using an assuming NTCP(x) DPD”; select the disease location, and introduce the BD parameter  $p$ .

If the selection is Panel 3 “Using a known/assumed NTCPi DPD”; introduce the values of probabilities (VPs) for each complication  $C_i$  ( $i = 1..7$ ), and introduce the VP for Other complications  $OCs$ ;

- if the selection is “Using an assuming NTCP(x) DPD”, one can define the legend of the numerical and graphical information. Each disease location has its number of possible cases ( $X_{max}$ ).  $X_{max}$  is equal to BD parameter  $n$ ;
- by pressing the “Finish” button of the selected panel you return to the main screen.

### ***The NTCP0calDr application***

The essential difference between this application and NTCP0cal is given in Panel 1, where  $SM_p$  NTCP0 is in the function of  $R_0$ , instead of  $D$ , and expressed as

$$SM_p NTCP_0(R_0) = \left[ \frac{TR_{0max} - R_0}{TR_{0max} - TR_{0min}} \right]^{pN_0} \quad (1)$$

$TR_{0min}$  — maximum value of  $R_0$  for  $NTCP_0 = 100\%$ . ( $TR_{0min} \geq 0$ );  $TR_{0max}$  — minimum value of  $R_0$  for  $NTCP_0 = 0\%$ ;  $pN_0$  — Power in this model.  $pN_0 > 0$ .

In  $R_0 < TR_{0min}$  and  $R_0 > TR_{0max}$ ,  $SM_p NTCP_0(R_0)$  is respectively equal to 100% and 0%.

The flow chart for determining NTCP0 in a CDLR treatment is similar to a fractionated one with  $D_{pres}$ ; and they differ in their respective  $SM_p$  NTCP0 models.

The steps for executing the NTCP0 calculation are:

- select one of the three panels pressing the “Use” button of the desired panel.  
If the selection is Panel 1 “Using the  $SM_p$  NTCP0 parameters”; select the radionuclide used, introduce the initial dose-rate  $R_0$  in Gy/h, and introduce the  $SM_p$  NTCP0 parameters ( $TR_{0min}$ ,  $TR_{0max}$  and  $pN_0$ ).

If the selection is Panel 2 “Using an assuming NTCP(x) DPD (Discrete probabilistic distribution)”; select the disease location, and introduce the BD parameter  $p$ .

If the selection is Panel 3 “Using a known NTCPi DPD”; introduce the values of probabilities for each complication  $C_i$  ( $i = 1..7$ ), and introduce the value of probability for other complications  $OCs$ ;

- press the “For calculating NTCP0” button for obtaining the result of NTCP0;
- if the selection is “Using an assuming NTCP(x) DPD”, one can define the legend of the numerical and graphical information. Each disease location has its number of possible cases ( $X_{max}$ ).  $X_{max}$  is equal to BD parameter  $n$ ;
- by pressing the “Finish” button of the selected panel you return to the main screen.

## Discussion

### *The SMp NTCP0 models*

The  $SMp(x)$  function of [6] was derived from the well-known Triangular model (TM), as a result of including powers  $p_1$  and  $p_2$  ( $p_1$  and  $p_2 \geq 0$ ).

$$TM = \begin{cases} f_1(x; a, b, c) * MaxTM \\ f_2(x; a, b, c) * MaxTM \end{cases} \quad (2)$$

$$SMp = \begin{cases} f_1(x; a, b, c)^{p_1} * MaxSMp \\ f_2(x; a, b, c)^{p_2} * MaxSMp \end{cases} \quad (3)$$

where  $a$ ,  $b$  and  $c$  are TM and  $SMp$  parameters, and  $MaxTM$  and  $MaxSMp$  are the respective maximum values of the TM and  $SMp$  models.

The  $SMp(x)$  can play the role of some probability density functions and DPDs, such as normal distribution and BD. Also, this can generate the three types:  $SMp_1$ ,  $SMp_2$  and  $SMp_3$ . For example, NTCP0 Vs. D model of [4] is a type  $SMp_3$ , which has a 100%-deterministic region, a stochastic and a 0%-deterministic, respectively defined by the parameters  $TD_{min} \geq 0$  and  $TD_{max}$  as

$$SMp\ NTCP\ 0(D) = \left[ \frac{TDmax - D}{TDmax - TDmin} \right]^{pNO}$$

(4)

$TDmin$  — maximum value of  $D$  for  $NTCP0 = 100\%$ . ( $TDmin \geq 0$ );  $TDmax$  — minimum value of  $D$  for  $NTCP0 = 0\%$ ;  $pNO$  — power in this model.  $pNO > 0$ ;  $D$  —  $Dpres$  function of  $d$  for a constant  $n$ ; or function of  $n$  for a constant  $d$ . In  $D < TDmin$  and  $D > TDmax$ ,  $SMp\ NTCP0(D)$  is respectively equal to 100% and 0%.

The current NTCP models provide approaches of this metric; i.e., NTCP estimations. An experienced radiation team will be able to assume good NTCP ( $x_i$ ) distributions. This implies good NTCP0 estimations, too.

### ***The NTCP( $x_i$ ) DPD assumed***

The tumor control probability (TCP) is a metric related to cell kill in a determined tumor tissue. For this reason, one can estimate its value using a computational simulation based on its own probabilistic concept, as has been developed in [7]. Contrary to simulated TCP calculations, nowadays, the determination of NTCP0 by means of mathematical models or computational simulations for treatments with few or no data is very complicated or almost impossible. Front of these difficulties, there is an option of assuming NTCP( $x_i$ ) distributions using generators of DPDs, like BD. For choosing the BD parameter  $p$ , one should consider that:

- 1 — if  $p \ll 0.5$ , the NTCP0 is the event with maximum probability (EwMP);
- 2 — if  $p < 0.5$ , one of the complications is the EwMP, and  $NTCP0 \gg 0\%$ ; if  $p \approx 0.5$ , one of the complications is the EwMP, and  $NTCP0 > 0\%$ ;
- 3 — if  $p > 0.5$ , one of the complications is the EwMP, and  $NTCP0 \approx 0\%$ .

The Figure 2 illustrates a hypothetical example of a NTCP( $x_i$ ) DPD for describing or assuming the probabilities of late complications discussed in [8], and associated to BT treatment for prostate cancer. The  $NTCP0 = NTCP(0) = 24\%$ . This value increases if  $D$  or  $R0$  decreases, and vice versa, as a result of variations of  $d$  for a treatment with a constant  $n$ ; or variations of  $n$  for a constant  $d$ ; or variation of  $R0$ .







Figure 2. Hypothetical example of a  $NTCP(x_i)$  discrete probabilistic distribution for describing or assuming the probabilities of late complications associated with a BT treatment for prostate cancer. D — prescribed dose;  $NTC_0$  — no complication;  $NTC_1$  — leakage of urine;  $NTC_2$  — cancer of the bladder;  $NTC_3$  — cancer of the lower bowel;  $NTC_4$  — erection problems (impotence). The  $NTCP_0 = NTCP_0 = 24\%$  is represented by a x; and its value increases if D or R0 decreases, and vice versa, as is shown by the four arrows on the right side of the y-axis.

Figure 3 shows an example of an option of the Matlab application for an assumed NTCP distribution generated by the BD expression:  $BD(x;0.4,6)$  for a head & neck disease location.



**Figure 3.** An example of the third option of the NTCP0cal/NTCP0CalDr application for an assumed NTCP distribution generated by the BD expression:  $BD(x;0.4,6)$  for a head & neck disease location.

For selecting  $NTCP(x_i)$  and its correspondent  $x_i$ , the aspect contained in the Table 1, sub-region of the disease and other clinical and physical factors should be considered. The table is based on some QUANTEC studies.

**Table 1.** Late complications of the BT treatments for their correspondent disease location

Late complications	Disease location				
	Head and Neck	Breast	Chest	Abdomen	Pelvis
Radiation (Rad.) brain	[9]				
Rad. induced optic neuropathy	[10]				
Myelopathy	[11]	[11]	[11]	[11]	[11]
Sensorineural hearing loss	[12]				
Xerostomia	[13]				
Rad. larynx and pharynx complications	[14]				
Rad. lung		[15]	[15]		
Rad. heart		[16]	[16]		
Rad. esophagus		[17]	[17]	[17]	
Liver dysfunction		[18]	[18]	[18]	
Rad. stomach and small bowel				[19]	[19]
Rad. kidney				[20]	[20]
Genitourinary				[21]	[21]
Rad. rectal				[22]	[22]
Rad. penile bulb				[23]	[23]

### **Other aspects**

From revisions of studies related to use of NTCP in the evaluations of S-Es of the BT, we can say that:

- the majority of current NTCP models are DVH-based;
- the risk of toxicity is the way of evaluating the S-Es of radiation oncology treatments;

- NTCP is used mainly for evaluations of individual or principal complications or Endpoints of radiation treatments.

Nowadays, as described in [10], [17] and [24], the NTCP studies have been focused on separated OARs, or the principal late complications of a radiation treatment of an OAR; however, these treatments have various normal tissue complications; in other words, they have their own associated NTCP( $x_i$ ) DPDs.

The fractional radiation treatment has two independent variables: 1 — Number of fractions ( $n$ ); and 2 — Dose per fraction ( $d$ ). For this reason, the SMp NTCP0(D) could be expressed as SMp NTCP0( $d$ ) for a constant  $n$ ; or as SMp NTCP0( $n$ ) for a constant  $d$ .


Because of the difficulty of obtaining NTCP model parameters for different combinations of  $n$  and  $d$ , the equivalent dose of 2 Gy per fraction (EQD<sub>2</sub>) was derived. But it is very important to consider that EQD<sub>2</sub> establishes a cellular radiosensitivity equivalence, not a normal complication one.

As shown in the Figure 1, if SMp NTCP0(D) model parameters are not known for a determined combination of  $n$  and  $d$ , we suggest that a NTCP( $x_i$ ) DPD should be assumed using a binomial distribution. For example, in Figure 4 (f) the BD( $x$ ;5,0.54) can be assumed for describing the NTCP DPD of this figure.

Figure 4 illustrates a generic example for showing variations of the late complications and NTCP0 for a BT treatment of a constant number of fractions and six different dose per fraction values. We want to show with this figure that:

- 1 — Any specific BT treatment given to an homogeneous patient populations has an associated acute/late NTCP( $x_i$ ) DPD, where  $i=0:nc$  and  $nc$ : Number of complications; NTCP0 = NTCP(0) and TNTCP = 100% – NTCP0;
- 2 — For a treatment with a constant  $n$ , if  $d$  increases TNTCP increases, and NTCP0 decreases; i.e. the number of patients with late complications increases, and the number of those without complications decrease;
- 3 — Each NTCP( $x_i$ ) complication ( $I > 0$ ) has an independent behavior when  $d$  increases. For example: C1 decreases when  $d$  increases in CD; C3 keeps its value in A–D; and C2 increases in AB; and when D

increases as a result of increases of  $d$ , the  $NTCP(x_i)$  cannot be described with increasing functions, but these can describe TNTCP; and of course the decreasing functions of  $NTCP_0$ .



**Figure 4.** Illustrations of a generic example of a BT treatment with a constant number of fractions ( $n$ ) dose per fractions, and **(A)**: for dose per fraction  $d_1$ ; **(B)**:  $d_2$ ; **(C)**:  $d_3$ ; **(D)**:  $d_4$ ; **(E)**:  $d_5$  and **(F)**:  $d_6$ ; where  $d_1 < d_2 < d_3 < d_4 < d_5 < d_6$ . The treatment has associated five late complications (C1, C2, C3, C4 and Others). We graphically and numerically show the independent variations of each late complications, and  $NTCP_0$

The  $SMp\ NTCP_0(D)$  model does not require DVH values of the OARs, nor their derivations, such as the EUD ; but the prescribed dose. Contrary to our models, the widely used LKB, and relative seriality model are DVH-based.

Implementing NCTP0 in the BT will represent the following advantages compared to the current S-E evaluations:

- the SMp NTCP0(D) and SMp NTCP0(R0) models are mathematically less complex than the LKB NTCP(Deff), where *Deff*: Effective dose;
- contrary to other NTCP models, these models do not involve OARs nor complications with different grade of severity. According to the type of OAR, one should use the LKB or relative seriality;
- given these models are not DVH-based, calculations of: EUD, Deff, or Maximum dose (Dmax) are not required. This model uses only information of the treatment.

Some previously discussed aspects and others of [7] are probabilistic foundations of our NTCP0 applications, and show why its validation is a priori. The validation of the NTCP0 methodologies is a priori because these are wholly based on strong probabilistic foundations, such as the normal complications of the specific radiation oncology treatments, as stochastic processes of more than outcome, have their own NTCP( $x_i$ ) DPDs, where NTCP0 = NTCP(0).

## Conclusions

The LKB NTCP(Deff) model is the normal cumulative distribution function (NCDF). As a cumulative distribution function, the NCDF has a sigmoidal shape and should be used for calculating the probability  $P(Deff \leq x)$  if Deff follows a normal distribution. For this reason, its use is not wholly appropriated as a NTCP model.

The current NTCP models used for evaluating S-Es in the radiation treatments provide NTCP approaches. An experienced radiation oncology team can assume a good NTCP( $x_i$ ) DPD based on database. Although an NTCP distribution is generated, the team should be interested only in one value, NTCP0. The NTCP0 estimations will be corrected in the future when a major data are available.

Concerning the mathematical correlations, the NTCP0(D) and NTCP0(R0) models are three-parameter phenomenological, and given the number of parameters and type, it is very easy to fit whatever real data NTCP0 Vs. D or R0, whose radiobiological mean behaviors should be described with decreasing functions aimed at acceptable estimations of S-Es.

Given gathering a data that lets us reproduce real graphical representations is too difficult or impossible; we have developed a generic example based on strong radiobiological and probabilistic foundations.

## References

1. [1] Shiraishi Y, Hanada T, Ohashi T, et al. Normal Tissue Complication Probability (NTCP) Modeling of Late Rectal Bleeding Following Iodine-125 Prostate Brachytherapy Combined With External Beam Radiation Therapy. *Int J Radiat Oncol Biol Phys.* 2015; 93(3): E210, doi: [10.1016/j.ijrobp.2015.07.1080](https://doi.org/10.1016/j.ijrobp.2015.07.1080).
2. Prayongrat A, Kobashi K, Ito YM, et al. The normal tissue complication probability model-based approach considering uncertainties for the selective use of radiation modality in primary liver cancer patients. *Radiother Oncol.* 2019; 135: 100–106, doi: [10.1016/j.radonc.2019.03.003](https://doi.org/10.1016/j.radonc.2019.03.003), indexed in Pubmed: [31015154](https://pubmed.ncbi.nlm.nih.gov/31015154/).
3. Report of AAPM TG-166. The Use and QA of Biologically Related Models for Treatment Planning. 2012.
4. Frometa-Castillo T, Pyakuryal A, Wals-Zurita A, et al. Proposals of models for new formulations of the current complication-free cure (P+) and uncomplicated tumor control probability (UTCP) concepts, and total normal tissue complication probability of late complications. *Int J Radiat Biol.* 2020; 96(7): 847–850, doi: [10.1080/09553002.2020.1741722](https://doi.org/10.1080/09553002.2020.1741722), indexed in Pubmed: [32163306](https://pubmed.ncbi.nlm.nih.gov/32163306/).
5. Frometa-Castillo T, Pyakuryal A, Mesbahi A, et al. The mathematical, probabilistic and computational generators of discrete probabilistic distributions applied to medical physics. *Med Phys Int J.* 2020; 8(3): 491, doi: [498](https://doi.org/10.1016/j.medphys.2020.03.003).
6. Frometa-Castillo T. The  $SMP(x \text{ or } y; PX_{min}, X_{max}, ML, p_1, p_2, Max)$  a probabilistic distribution, or a probability density function of a random variable X, Chapter 48 of 16th International Conference on Information Technology-New Generations (ITNG 2019), Springer Nature Switzerland AG; 2018.
7. Frometa-Castillo T, Pyakuryal A, Narayanasamy G, et al. Computational simulator that calculates tumor control probability in a tumor heterogeneously irradiated for fractionated radiation oncology treatments. *Simulation.* 2021; 98(4): 285–294, doi: [10.1177/00375497211039460](https://doi.org/10.1177/00375497211039460).
8. Long term side effects of internal radiotherapy (brachytherapy) . <https://www.cancerresearchuk.org/about-cancer/prostate-cancer/treatment/radiotherapy/internal-radiotherapy/long-term-side-effects-internal-radiotherapy>.
9. Lawrence YR, Li XA, el Naqa I, et al. Radiation dose-volume effects in the brain. *Int J Radiat Oncol Biol Phys.* 2010; 76(3 Suppl): S20–S27, doi: [10.1016/j.ijrobp.2009.02.091](https://doi.org/10.1016/j.ijrobp.2009.02.091), indexed in Pubmed: [20171513](https://pubmed.ncbi.nlm.nih.gov/20171513/).
10. Mayo C, Martel MK, Marks LB, et al. Radiation dose-volume effects of optic nerves and chiasm. *Int J Radiat Oncol Biol Phys.* 2010; 76(3 Suppl): S28–S35, doi: [10.1016/j.ijrobp.2009.07.1753](https://doi.org/10.1016/j.ijrobp.2009.07.1753), indexed in Pubmed: [20171514](https://pubmed.ncbi.nlm.nih.gov/20171514/).
11. Kirkpatrick JP, van der Kogel AJ, Schultheiss TE. Radiation dose-volume effects in the spinal cord. *Int J Radiat Oncol Biol Phys.* 2010; 76(3 Suppl): S42–S49, doi: [10.1016/j.ijrobp.2009.04.095](https://doi.org/10.1016/j.ijrobp.2009.04.095), indexed in Pubmed: [20171517](https://pubmed.ncbi.nlm.nih.gov/20171517/).
12. Bhandare N, Jackson A, Eisbruch A, et al. Radiation Therapy and Hearing Loss. *Int J Radiat Oncol Biol Phys.* 2010; 76(3): S50–S57, doi: [10.1016/j.ijrobp.2009.04.096](https://doi.org/10.1016/j.ijrobp.2009.04.096).

13. Deasy JO, Moiseenko V, Marks L, et al. Radiotherapy dose-volume effects on salivary gland function. *Int J Radiat Oncol Biol Phys.* 2010; 76(3 Suppl): S58-S63, doi: [10.1016/j.ijrobp.2009.06.090](https://doi.org/10.1016/j.ijrobp.2009.06.090), indexed in Pubmed: [20171519](https://pubmed.ncbi.nlm.nih.gov/20171519/).
14. Rancati T, Schwarz M, Allen AM, et al. Radiation dose-volume effects in the larynx and pharynx. *Int J Radiat Oncol Biol Phys.* 2010; 76(3 Suppl): S64-S69, doi: [10.1016/j.ijrobp.2009.03.079](https://doi.org/10.1016/j.ijrobp.2009.03.079), indexed in Pubmed: [20171520](https://pubmed.ncbi.nlm.nih.gov/20171520/).
15. Marks LB, Bentzen SM, Deasy JO, et al. Radiation dose-volume effects in the lung. *Int J Radiat Oncol Biol Phys.* 2010; 76(3 Suppl): S70-S76, doi: [10.1016/j.ijrobp.2009.06.091](https://doi.org/10.1016/j.ijrobp.2009.06.091), indexed in Pubmed: [20171521](https://pubmed.ncbi.nlm.nih.gov/20171521/).
16. Gagliardi G, Constine LS, Moiseenko V, et al. Radiation dose-volume effects in the heart. *Int J Radiat Oncol Biol Phys.* 2010; 76(3): 77-85.
17. Werner-Wasik M, Yorke E, Nam J, et al. Radiation dose-volume effects in the esophagus. *Int J Radiat Oncol Biol Phys.* 2010; 76(3): S86-S93.
18. Pan CC, Kavanagh BD, Dawson LA, et al. Radiation-associated liver injury. *Int J Radiat Oncol Biol Phys.* 2010; 76(3 Suppl): S94-100, doi: [10.1016/j.ijrobp.2009.06.092](https://doi.org/10.1016/j.ijrobp.2009.06.092), indexed in Pubmed: [20171524](https://pubmed.ncbi.nlm.nih.gov/20171524/).
19. Kavanagh BD, Pan CC, Dawson LA, et al. Radiation dose-volume effects in the stomach and small bowel. *Int J Radiat Oncol Biol Phys.* 2010; 76(3 Suppl): S101-S107, doi: [10.1016/j.ijrobp.2009.05.071](https://doi.org/10.1016/j.ijrobp.2009.05.071), indexed in Pubmed: [20171503](https://pubmed.ncbi.nlm.nih.gov/20171503/).
20. Dawson LA, Kavanagh BD, Paulino AC, et al. Radiation-associated kidney injury. *Int J Radiat Oncol Biol Phys.* 2010; 76(3 Suppl): S108-S115, doi: [10.1016/j.ijrobp.2009.02.089](https://doi.org/10.1016/j.ijrobp.2009.02.089), indexed in Pubmed: [20171504](https://pubmed.ncbi.nlm.nih.gov/20171504/).
21. Viswanathan AN, Yorke ED, Marks LB, et al. Radiation dose-volume effects of the urinary bladder. *Int J Radiat Oncol Biol Phys.* 2010; 76(3 Suppl): S116-S122, doi: [10.1016/j.ijrobp.2009.02.090](https://doi.org/10.1016/j.ijrobp.2009.02.090), indexed in Pubmed: [20171505](https://pubmed.ncbi.nlm.nih.gov/20171505/).
22. Michalski JM, Gay H, Jackson A, et al. Radiation dose-volume effects in radiation-induced rectal injury. *Int J Radiat Oncol Biol Phys.* 2010; 76(3 Suppl): S123-S129, doi: [10.1016/j.ijrobp.2009.03.078](https://doi.org/10.1016/j.ijrobp.2009.03.078), indexed in Pubmed: [20171506](https://pubmed.ncbi.nlm.nih.gov/20171506/).
23. Roach M, Nam J, Gagliardi G, et al. Radiation dose-volume effects and the penile bulb. *Int J Radiat Oncol Biol Phys.* 2010; 76(3 Suppl): S130-S134, doi: [10.1016/j.ijrobp.2009.04.094](https://doi.org/10.1016/j.ijrobp.2009.04.094), indexed in Pubmed: [20171507](https://pubmed.ncbi.nlm.nih.gov/20171507/).
24. Luo R, Wu VWC, He B, et al. Development of a normal tissue complication probability (NTCP) model for radiation-induced hypothyroidism in nasopharyngeal carcinoma patients. *BMC Cancer.* 2018; 18(1): 575, doi: [10.1186/s12885-018-4348-z](https://doi.org/10.1186/s12885-018-4348-z), indexed in Pubmed: [29776390](https://pubmed.ncbi.nlm.nih.gov/29776390/).