

Potential Use of Immature Oocyte to Improve Fertility Preservation Outcome: A Narrative Review

by Batara Sirait

Submission date: 12-Apr-2022 11:57AM (UTC+0700)

Submission ID: 1808553873

File name: Batara_Immanuel_Sirait.pdf (820.21K)

Word count: 6394

Character count: 35654

Review Article

Potential Use of Immature Oocyte to Improve Fertility Preservation Outcome: A Narrative Review

Batara Sirait^{1,2,3}, Ahmad Aulia Jusuf^{1,4}, Budi Wiweko^{5,6,7}, Nining Handayani^{3,8}, Daniel Abidin Aubry³, R. Muharam^{6,7}

¹Doctoral program, Faculty of Medicine, University of Indonesia, ²Department of Obstetrics and Gynaecology, Faculty of Medicine Universitas Kristen Indonesia, ³Morula IVF Jakarta Clinic, ⁴Department of Histology, Faculty of Medicine, Universitas Indonesia, ⁵Human Reproductive, Infertility, and Family Planning Research Center, Indonesian Medical Education and Research Institutes, Faculty of Medicine, University of Indonesia, ⁶Department of Obstetrics and Gynecology, Division of Reproductive Endocrinology and Infertility, Faculty of Medicine, University of Indonesia, ⁷Yasmin Infertility Clinic, Dr. Cipto Mangunkusumo General Hospital, ⁸IRSI Research and Training Centre, Jakarta, Indonesia

ABSTRACT

Fertility preservation through gamete vitrification has become one of the critical strategies to secure a childbearing potential in patients who are diagnosed with cancer or risks of infertility. Preserving the gametes would prevent the deleterious effects of cancer drugs or radiotherapy exposure on the quality of the gametes. Furthermore, *in vitro* fertilisation of vitrified mature human oocytes has lately demonstrated promising results that are reflected in the increased survival rate of thawed oocytes and the resultant clinical pregnancy rate. However, limitations in the cryopreservation of mature oocytes of cancer patients persist. Ovarian stimulation protocols which comprise administering gonadotrophin-releasing hormones could aggravate cancer or delay essential cancer therapy. Considering such circumstances, vitrification of immature oocytes would become a rational option. While the vitrification procedure of mature oocytes has been established, the vitrification of immature oocytes remains controversial due to a low post-thaw *in vitro* maturation and fertilisation rate. Apparent cryoinjuries to the immature oocytes post thawing or warming have been observed in both human and animal model oocytes. An alternative strategy was therefore proposed to improve the effectiveness of utilising immature oocytes for fertility preservation by conducting the *in vitro* oocyte maturation process first before vitrification. This method has prevailed, especially in oncofertility patients. Although the success rate of the clinical outcomes remains low, this approach, in conjugation with proper counselling, might provide oncofertility patients with an opportunity to preserve their reproductive potential.

KEYWORDS: Cancer-related infertility, fertility preservation, immature oocytes, *in vitro* fertilisation, vitrification

INTRODUCTION

The first successful births of twin human babies from fertilisation of frozen-thawed pre-ovulatory mature oocyte were reported in Australia in 1986.^[1] The strategy of utilising vitrified mature oocytes in an *in vitro* fertilisation (IVF) programme, however, has endured a slow acceptance due to the low survival rate of the oocytes post thawing. Experimental research on oocyte cryopreservation indicates the difficulties of vitrifying a large single oocyte cell.^[2] The urgency to improve the success rate of oocyte cryopreservation


has risen in several countries such as Italy, Austria, Germany and Switzerland due to legislation that restricts embryo cryopreservation.^[3] A significant improvement in cancer prognosis after treatments in young adolescent patients has also validated the demand for an oocyte cryopreservation programme as means to preserve the reproductive potential of girls or women who are about

Address for correspondence: Dr. Batara Sirait, Department of Obstetrics and Gynaecology, Faculty of Medicine Universitas Kristen Indonesia, Mayjen Sutoyo Street Number 2, RT.9/RW.6, Cawang, Kramat Jati East Jakarta, 13630, Indonesia. E-mail: batarasirait@gmail.com

23

Received: 29-07-2021
Accepted: 07-02-2022

Revised: 02-02-2022
Published: 31-03-2022

Access this article online	
Quick Response Code: 	Website: www.jhrsonline.org
DOI: 10.4103/jhrs.jhrs_112_21	

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Sirait B, Jusuf AA, Wiweko B, Handayani N, Aubry DA, Muharam R. Potential use of immature oocyte to improve fertility preservation outcome: A narrative review. *J Hum Reprod Sci* 2022;15:3-11.

to undergo cancer therapies.^[4,5] Oocyte freezing as an alternative ²⁸ method to treat infertility was first approved in the UK by the Human Fertilisation ⁶ and Embryology Authority in 2000^[6] followed by the American Society for Reproductive Medicine in 2013.^[7]

Vitrification is preferably performed when the ⁵⁶ oocyte is at a mature stage, namely metaphase II (MII). To promote ¹⁰ *in vitro* maturation of oocytes, ovarian stimulation begins at day 2 or 3 of a menstrual cycle during the follicular phase. Exogenous gonadotropin hormone is administered once daily to support follicular growth until a minimum of two or three follicles has reached 18 mm. Oocyte maturation trigger utilising ³⁸ gonadotropin-releasing hormone agonist or human chorionic gonadotropin is then injected and the ovum pick-up procedure is commenced 36 h later.^[8] Nonetheless, the gonadotropin stimulation approach is unsuitable for certain patients with underlying conditions such as cancer. Women with breast cancer, for instance, are particularly sensitive to the elevation of serum oestradiol (oestrogen-sensitive tumour) and are therefore advised to avoid undergoing ovarian stimulation using the gonadotropin hormone.^[9] In addition to the risks of cancer aggravation, the time required for ovarian stimulation procedure could delay crucial cancer treatment.⁵⁷ Investigating the potential use of immature oocytes as an alternative option for fertility preservation therefore ² becomes admissible. Contrary to the collection of *in vivo* matured oocytes, immature oocytes can be ⁵⁷ retrieved conveniently at any stage of the ovarian cycle.^[10] More importantly, the necessity of the ovarian stimulation protocol could be bypassed. A case report published in ²³ 2022 demonstrated the convenient method of retrieving immature oocytes from antral follicles during a conservative surgery for ovarian cancer, indicating the feasibility and development prospects of such strategy.^[11]

Several investigations were carried out to establish an optimal fertility preservation strategy using immature oocytes that could assure the chance of bearing a child in a specific group of cancer patients.^[12-14] In the earlier decade of oocyte cryopreservation history, a low survival rate of cryopreserved mature oocyte post thawing has led to a postulation of a higher cryoinjury in resistance in germinal vesicle (GV) immature oocyte compared to the MII mature oocyte. At that point, GV was believed to be the ideal stage for vitrification rather than MII because of its lack of a microtubular spindle system. Moreover, chromosomes in GV are enclosed by a nuclear membrane, which was considered to reduce the risk of chromosome injuries and prevent the polyploidy or aneuploidy occurrence possibly induced by the extreme cooling condition

during vitrification.^[15] However, ⁴⁶ increasing evidence has suggested that the GV-stage oocytes are as vulnerable to cryoinjuries as the mature-stage oocytes.^[16-19] Some studies ³ have subsequently recommended performing an *in vitro* maturation (IVM) of the immature oocytes before vitrification,^[20-24] while others have opposed this idea.^[25-27] This literature provides a comprehensive review on the safety and current progress of vitrifying immature human oocytes for possible use of fertility preservation.

METHODS

The literature review was conducted using several search engines including Google Scholar and PubMed. Boolean search strategy (AND, OR, NOT) has been applied to identify the relevant articles using terms such as oocyte, egg, vitrification, freezing, slow freezing, cryopreservation, immature and fertility preservation. Keyword phrases included in the search were 'immature egg freezing', 'immature oocyte vitrification' and 'immature oocyte freezing' [Figure 1].

DISCUSSION

The clinical indications for women who could benefit from fertility cryopreservation using ⁴⁸ mature oocytes

Several factors including the cancer type and grade, urgency of the cancer treatment and marital status should be taken into consideration when opting for the fertility preservation programme. The clinical algorithm for female cancer patients aiming to retain their reproductive potential has been well defined.²⁵^[4,9,28] As previously described, gonadotropin ovarian stimulation may be contraindicated in certain cancer patients with hormonal-sensitive tumours such as desmoplastic tumours and breast cancer (especially in oestrogen receptor-positive type and breast adenocarcinoma).^[13,29] Administering exogenous gonadotropin during ovarian stimulation could induce an increased oestradiol level up to 15 times more than the natural cycle.^[30] Thus, oncologists may advise the patients to undergo oocyte retrieval without ovarian stimulation.^[12,13] Likewise, women who cannot delay their cancer treatments for a 2- to 6-week stimulation protocol could benefit from the retrieval and cryopreservation of immature oocytes.^[9]

Clinical evidence on the benefits of preserving fertility using immature oocytes was proven in several studies.^[10,12-14,29] A 2010 study demonstrated ⁶ a novel approach of collecting immature oocytes without ovarian stimulation in ³⁸ 20 infertile women diagnosed with breast cancer. The oocyte retrieval was performed under sedation 36 h following a 10,000 IU HCG injection.




Figure 1: Search strategies for narrative review

19 GV-stage oocytes were subsequently matured *in vitro* through a 24-h culture and the resultant mature oocytes were used for further treatment. The yield of the *in vitro* mature oocytes ranged between 1 and 22 and the median vitrified embryos was 4 with a range of 1–13. Of the 38 women, 18 opted for the IVM process followed by the oocyte vitrification while the remaining agreed to fertilise their mature oocytes and vitrify the resultant embryos.^[13]

47 An increased number of *in vitro* matured oocytes for vitrification was reported in two oncofertility women through a combination of immature oocyte aspiration at both the follicular and the luteal phase.^[29] Moreover, evident benefits of retrieving immature oocytes at the luteal phase were demonstrated in a woman aged 21 who was incapable of suspending her cancer treatment and was advised against receiving the gonadotropin therapy.^[29] Supporting the previous results, a prospective study on 248 breast cancer women has produced a similar IVM rate of immature oocytes aspirated at either the follicular or the luteal phase. The increasing yield of

mature oocytes post IVM has certainly generated an interest in the practice of such strategy as an urgent approach for fertility preservation.^[10]

Another plausible method of fertility preservation is through immature oocytes aspiration from excised ovarian tissue combining with an ovarian tissue o-banking. This strategy was applied in four women who were diagnosed with Hodgkin lymphoma, breast cancer and rectal cancer.^[12] Immature oocytes were collected from the excised ovarian tissue and were subjected to the IVM culture. The mean oocyte maturation rate post IVM was 79%. All of the oncofertility patients managed to acquire at least one mature oocyte for vitrification. Furthermore, a large retrospective cohort study in 2015 comprising 255 cancer patients has validated the safety and advantages of harvesting the ovarian tissue for immature oocyte collection as means to attain an increased total number of *in vitro* matured oocytes and fertilisation rate.^[14]

As the fertilisation of the *in vitro* matured oocytes was decided based on the marital status, patient preference or

age, most studies on fertility preservation using immature oocytes have heretofore reported the yield of the *in vitro* matured oocytes as the main outcome.^[10,12-14,29] Only few studies managed to describe the downstream IVF outcomes including the total number of embryos derived from the *in vitro* matured oocytes.^[13,14] Hourvitz *et al.* reported a mean number of vitrified embryos between 1.67 ± 0.56 and 3.39 ± 0.73 depending on the immature oocyte collection procedure.^[14] Although the results were encouraging, clear benefits of utilising immature oocytes to obtain embryos for further treatment are difficult to define due to the small sample sizes of the available reports. Therefore, detailed information regarding the current success rate of fertility preservation should be informed to the patients who wish to undergo the program.

Immature oocyte collection in IVF could also benefit women with polycystic ovary syndrome who are at risk of an ovarian hyperstimulation syndrome subsequent to ovarian stimulation.^[31] The fertility preservation programme would also cater to the increasingly modern trend of postponing childbearing due to social or non-medical reasons. A study in the UK provided several background and clinical characteristics of 26 women who underwent fertility cryopreservation. The mean age of the 27 women involved in the study was 36.7 years. They were highly educated, and half of the participants were professionally employed.^[32]

Competency of *in vitro* matured germinal vesicle or post germinal vesicle breakdown metaphase I oocytes: A lesson from a stimulated fresh *in vitro* fertilisation cycles

Clinical use of immature oocytes obtained during a stimulated fresh IVF cycle is disputable even without the vitrification processing. *In vitro* matured GV and MI oocytes lack the competency to improve the clinical pregnancy.^[33] Although a similar fertilisation rate was observed between the *in vivo* and *in vitro* matured oocytes, Shu *et al.* concluded that the clinical pregnancy and live birth rate of transferring embryos derived from the *in vitro* matured oocytes were unsatisfactory.^[33] A 2010 study has also shown the inconspicuous effectiveness of using immature oocytes derived from the stimulated IVF cycles. Two hundred sixty-three immature oocytes subjected to IVM were compared with their sibling *in vivo* matured oocytes ($n = 234$). Although both groups acquired comparable fertilisation rates, the developmental quality of the day 2 cleavage stage in the immature oocyte group was lower in regard to the blastomeric number and symmetry. Moreover, none of the 17 transferred embryos derived from the *in vitro* matured oocytes were

successfully implanted.^[34] Another study also observed a low clinical efficacy of the *in vitro* matured oocytes which did not culminate to a single clinical pregnancy in the five cases of embryo transfer.^[35]

Nonetheless, a promising utilisation of *in vitro* matured non-GV or germinal vesicle breakdown (GVBD) has recently been reported by Olid *et al.*^[36] IVM of GVBD oocytes in G-2™ US media (Vitrolife, Sweden) resulted in a 10% clinical pregnancy rate and a 5.6% live birth rate. Concordantly, an evidence on the clinical benefit of *in vitro* matured II-stage oocyte was recorded through one successful live birth of a healthy normal baby.^[37]

An IVF using *in vitro* matured oocytes is undeniably less favourable than the *in vivo* matured oocytes. However, performing IVM of non-GV immature oocytes is another option, particularly in cycles which yield a low number of mature oocytes. The possible reasons underlying the low developmental competence of *in vitro* matured non-GV stages were implied by Ferrer-Vaquero *et al.*^[38] This study highlighted the several differences in the cytoplasmic maturation between *in vivo* and *in vitro* matured oocytes. Alterations in the mitochondrial membrane potential, cluster number of endoplasmic reticulum and actin cytoskeleton were particularly observed in *in vitro* matured oocytes.

Should *in vitro* maturation be performed before or after vitrification?: A lesson from the use of immature oocytes in *in vitro* fertilisation to retain reproductive potential

Following the successful birth of a human female baby derived from a frozen-thawed *in vitro* matured immature oocyte in 1998 Georgia – America, cryopreservation of immature oocytes has been proposed as an alternative strategy to preserve the gametes.^[39] Over the past two decades, numerous studies have revealed the challenges of immature oocyte vitrification. Several limitations of the approach are well discussed in an existing review as follows: (i) even though the survival rates of the frozen-thawed mature and immature oocyte are similar, the capacity of the surviving immature oocytes to undergo maturation *in vitro* is considerably low; (ii) fertilisation capacity of *in vitro* matured oocytes is also lower compared to the *in vivo* matured oocytes; and (iii) vitrification has been shown to evoke injuries of the chromosomes and cytoplasmic organelles even in the GV stage, disclaiming the theories that were previously suggested. The irreversible injuries of the organelles could be accountable for the lack of maturation and fertilisation capacity of the vitrified immature oocytes.^[16,19] However, several contradictory

studies have proved that the outcomes of vitrifying oocytes at immature stage were not inferior to that of mature oocytes.^[25-27]

As presented in Table 1, most studies recommended performing IVM before vitrification to improve the clinical utilisation of the immature oocytes. Nonetheless, three studies suggested otherwise, proving that vitrifying oocytes before IVM or after were equally efficient. An in-depth evaluation is therefore still required to identify the factors which led to the different outcomes^[37] and to confidently determine whether vitrification of immature oocytes should be done before or after IVM. The current methodology for IVM is not sufficient enough to induce oocytes with a viability that is on par with the *in vivo* matured oocytes. Advancements in the IVM process might become an essential key to enhance the quality of *in vitro* matured oocytes and a stepping stone towards improvement in the fertility preservation programme.^[18]

A standard procedure for IVM is to culture the immature oocytes in media supplemented with human serum albumin and highly purified gonadotrophins for up to 30 h until a polar body extrusion is observed.^[40] Of note, the collection of immature oocytes from small antral follicles may slightly be more time – and technically – demanding than the collection of mature oocytes. A cell strainer with minute pores might be utilised to search for immature oocytes with minimal cumulus complex.^[40] No particular commercialised brand of media has been deemed more effective in promoting IVM.

Advancements for IVM of oocytes could only be achieved with a clear insight into the *in vivo* maturation process. The mechanism involves a series of complex signalling pathways that are regulated by specific factors expressed through different cells found within the cumulus-oocyte complex. These cells are connected by gap junctions and connexins that need to be maintained *in vitro* to preserve the intra- and intercellular communications. In the pursuit of mimicking an *in vitro* condition which complements the *in vivo* micro-environments, several pre-maturation enhancements of IVM^[7] have been attempted. Vanhoutte *et al.*^[41] designed a three-dimensional (3D) culture system utilising an extracellular matrix composed of collagen. Retrieved immature oocytes with intact cumulus complex were cultured on polymerised collagen supplied with IVM media and phosphodiesterase-3 inhibitor (meiotic arrester). Subsequently, the oocytes were recovered from the gel, washed and denuded for fertilisation through conventional IVF. The 3D culture managed to maintain the communication framework between cells found in the cumulus-oocyte complex that

in turn brought on more meiotically competent oocytes and improved embryo development compared to the conventional IVM method.

In addition to the pre-maturation culture, several studies have shown an enhanced efficacy of IVM by fine-tuning the media with paracrine oocyte maturation-promoting factors such as FSH and amphiregulin^[42] and oocyte-secreted factors such as bone morphogenetic protein 15 and growth differentiation factor 9.^[43] Nonetheless, while the promising outcomes of these experiments could bridge the gap in the effectiveness between an IVM-IVF cycle and a conventional IVF cycle, large prospective studies are obligatory before these findings could be translated to clinical practice. Additionally, methods to reduce the cryoinjuries during vitrification should also be pursued to accomplish an altogether viable strategy for fertility preservation.

The cellular and molecular causes of low fertility potential in immature oocytes after vitrification

Several studies have revealed the consequences of immature oocyte vitrification in humans^[44,45] and animal models^[46-49] on a cellular and molecular level. In a 1988 study conducted by Van Blerk *et al.*,^[13] mouse GV vitrification was shown to cause premature chromosome condensation, extrusion of chromatin fragmentation in the cytoplasm, microtubule defects, impaired relocation of mitochondria and protein synthesis alteration.^[46] An experimental study that examined the post-vitrification cycle cell status and chromosomal integrity of GV-stage rat oocytes also pointed out similar findings. Following a warming procedure, the GV-stage oocytes immediately entered the M-phase cell cycle but failed to maintain chromosomal integrity due to multiple chromatin fusion. The study also highlighted a remarkably low number of F-actin in both the cytoplasm and the cortical area and an irregular retraction of trans-zona pellucida post vitrification.^[47] Several defects of the cytoskeleton, aberrant density of the mitochondrial matrix and microtubule were also observed in other animal studies.^[48,49]

Utilising transmission electron microscope, frozen-thawed GV oocytes were shown to prematurely release peripheral cortical granules onto the subzonal space during IVM. Additionally, the number of cortical granules was reduced remarkably causing an increased density of the zona pellucida ultrastructure and subsequent zona pellucida hardening. Aggregates of small mitochondria-smooth endoplasmic reticulum (M-SER) were found to be randomly distributed throughout the cytoplasm.^[44] Furthermore, detrimental effects of cryoprotectant exposure on the chromosomes and microtubules of immature human

Table 1: Summary table of studies evaluated the various outcomes of vitrified immature oocyte before or after *in vitro* maturation

Study design	Year	Main objective	Sample size	Main outcome	Conclusion	References
<i>In vitro</i> experimental study	2009	To compare the efficacy of vitrification before and after IVM of immature oocytes	Immature human oocytes (n=472) Group 1 consisted of 219 GV that was vitrified before IVM Group 2 comprised 253 GV that was vitrified after IVM (n=79, while the remaining (n=99) were setting for control group)	A comparable survival rate post vitrification was observed between Group 1 and 2 (85.4% vs. 86.1%, respectively) However, the number of matured oocytes was significantly higher in Group 2 than in Group 1 A similar outcomes including fertilisation rate, cleavage and blastocyst rate were observed	IVM of GV-stage oocytes before vitrification yields better outcomes in terms of maturation rate	Cao et al. ^[20]
<i>In vitro</i> experimental study	2012	Comparing the effectiveness of IVM of immature oocytes before and after vitrification	184 immature (GV or MI) human oocytes were randomly allocated to Group 1 (vitrified after IVM, n=100) and Group 2 (vitrified before IVM, n=84)	While the survival rates were similar between Group 1 and 2 (86.9% and 84.5%), the maturation rate was significantly higher in Group 1 than Group 2 (46% vs. 23.8%) Fertilisation rate and embryo development potency between the Group 1 and 2 were comparable. However, no transferable blastocysts were obtained in both groups	A high number of <i>in vitro</i> matured oocytes survived the vitrification procedure because IVM was conducted before vitrification	Fasano et al. ^[21]
<i>In vitro</i> experimental study	2012	To investigate the best stage for vitrification of cumulus-free immature oocytes	Immature oocytes were obtained from 120 IVF patients 24 GV-stage oocytes were allocated to freezing prior to IVM (n=109) and after IVM (n=107)	A comparable survival rate between vitrified-GV before and after IVM was observed (69.7% vs. 70.5%, respectively) However, the low maturation rate (51.3% vs. 75.7%) and high spontaneous cleavage were observed in GV oocytes of vitrified before IVM	Vitrification of <i>in vitro</i> matured GV oocytes acquired more preferable outcomes than GV oocytes vitrified before IVM	Wang et al. ^[23]
Cross-sectional study	2013	Evaluating the maturation rate and viability of immature oocytes post warming	Infertile human oocytes A total of 53 fresh immature oocytes were subjected to fresh IVM as a control group (fIVM) and 50 immature oocytes were vitrified prior to IVM (vIVM)	Maturation rate of fIVM was superior over vIVM (88.7% vs. 56%, respectively, P<0.001)	Performing IVM prior to vitrification attained the maturation capacity of immature oocytes and the viability to further support embryonic development	Yazdanpanah et al. ^[24]
Experimental study	2016	Observing cryoinjuries on nuclear integrity such as spindle, DNA, as well as embryonic aneuploidy and ZPD after warming	Infertile human oocytes Thirty-one <i>in vivo</i> mature oocytes were used as control (Group A, with or without vitrification) Two hundred-twenty-six immature oocytes were allocated to Group B (IVM group before vitrification, n=113) and Group C (immature group, n=110)	Aneuploidy rate was comparable amongst groups (P<0.05). However, comet cells were significantly lower in Group A than Group B and C (P<0.05) Group C showed a lower occurrence and retardance value of spindle compared to Group A and B 45% cleavage rate was also lower than that of Group A and B (P<0.05)	According to post warming molecular analysis, the most suitable stages for oocyte vitrification are at the <i>in vivo</i> mature and <i>in vitro</i> matured stages	Song et al. ^[22]

Contd...

Table 1: Contd...

Study design	Year	Main objective	Sample size	Main outcome	Conclusion	References
<i>In vitro</i> experimental study	2011	Evaluating the effect of GV vitrification on the methylation profile followed by IVM	184 GV oocytes subjected to vitrification followed by IVM and 120 GV oocytes were matured <i>in vitro</i> as a control group	<i>In vitro</i> matured oocytes derived from vitrified-GV showed comparable methylation profiles as fresh <i>in vitro</i> matured GV oocytes	Oocyte vitrification at the GV stage did not impair the methylation profile	Al-Khtib <i>et al.</i> ^[25]
<i>In vitro</i> experimental study	2016	Assessing the effects of GV vitrification on nuclear configuration, DNA methylation as well as cytoplasmic maturation and maturation rate post warming	Infertile human oocytes A total of 284 oocytes were allocated into three groups <i>In vivo</i> mature oocyte group (MII as the control group, <i>n</i> =56) IVM group (vitro-MII, <i>n</i> =106) Immature vitrified group (cryo-MII, <i>n</i> =122)	The cryo-MII group was shown to have the highest abnormal spindle and chromosome configuration amongst the groups (<i>P</i> <0.05) However, global DNA methylation, mitochondria and cortical granules distribution (reflecting cytoplasmic maturation), as well as maturation rate post warming were similar amongst the three groups	Although GV vitrification could disrupt the configuration of spindle and chromosome, GV stage can preserve its viability and undergo IVM after warming	Liu <i>et al.</i> ^[27]
<i>In vitro</i> experimental study	2016	Investigation on whether oocyte vitrification delivers better results before or after IVM in regards to the survival, maturation rate and embryonic development	221 unused human GV from ICSI-IVF cycles were grouped Group 1 as the control group (GV oocytes were matured up to the MII stage without vitrification) Group 2 (GV-stage oocytes were vitrified and subsequently matured <i>in vitro</i> until reaching the MII stage) Group 3 (GV oocytes underwent IVM followed by a vitrification-thawing procedure)	The post warming survival rates between Group 2 and group 3 were similar (<i>P</i> =0.810) Group 2 showed the highest number of <i>in vitro</i> matured oocytes (83.7%) compared to Group 1 (63.4) and Group 3 (56.6%) Blastocyst formation rates were low in the overall three groups (<i>P</i> =0.432)	Vitrifying GV-stage oocytes resulted in a higher survival rate and maturation rate post warming	Molina <i>et al.</i> ^[26]

10

IVM=*In vitro* maturation, GV=Germinal vesicle, MI=Metaphase I, fIVM=Fresh IVM, vIVM=Vitrified IVM, ZPD: Zona pellucida density, ICSI=Intra-cytoplasmic sperm injection

oocytes were also apparent. An increased incidence of abnormal chromosomes and microtubules was exhibited in the *in vitro* matured GV oocytes.^[45]

CONCLUSION

Fertility preservation utilising immature oocytes could increase the availability of mature oocytes and consequently embryos for cryopreservation. Ineffectiveness of immature oocyte vitrification for fertility preservation persists on account of the vitrification injuries observed in frozen-thawed oocytes of both the animal models and humans. Alternatively, performing IVM before vitrification would increase the survival and fertilisation rate of the immature oocytes. Developing methods that could significantly improve IVM and reduce cryoinjuries during vitrification is imperative to deliver the clinical benefits of immature oocyte cryopreservation as means to preserve fertility.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Chen C. Pregnancy after human oocyte cryopreservation. *Lancet* 1986;1:884-6.
- Argyle CE, Harper JC, Davies MC. Oocyte cryopreservation: Where are we now? *Hum Reprod Update* 2016;22:440-9.
- Rienzi L, Gracia C, Maggiulli R, LaBarbera AR, Kaser DJ, Ubaldi FM, *et al.* Oocyte, embryo and blastocyst cryopreservation in ART: Systematic review and meta-analysis comparing slow-freezing versus vitrification to produce evidence for the development of global guidance. *Hum Reprod Update* 2017;23:139-55.
- Kim SY, Lee JR. Fertility preservation option in young women with ovarian cancer. *Future Oncol* 2016;12:1695-8.
- Siegel R, Ward E, Brawley O, Jemal A. *Cancer statistics, 2011*:

- The impact of eliminating socioeconomic and racial disparities on premature cancer deaths. *CA Cancer J Clin* 2011;61:212-36.
6. Wise J. UK lifts ban on frozen eggs. *BMJ* 2000;320:334.
 7. American Society for Reproductive Medicine and the Society for Assisted Reproductive Technology. Mature oocyte cryopreservation: A guideline. *Fertil Steril* 2013;99:37-43.
 8. Cobo A, Garcia-Velasco JA, Coello A, Domingo J, Pellicer A, Remohi J. Oocyte vitrification as an efficient option for elective fertility preservation. *Fertil Steril* 2016;105:755-64.e8.
 9. Rodriguez-Wallberg KA, Oktay K. Options on fertility preservation in female cancer patients. *Cancer Treat Rev* 2012;38:354-61.
 10. Grynberg M, Poulain M, Le Parco S, Sifer C, Fanchin R, Frydman N. Similar *in vitro* maturation rates of oocytes retrieved during the follicular or luteal phase offer flexible options for urgent fertility preservation in breast cancer patients. *Hum Reprod* 2016;31:623-9.
 11. Fadini R, Dal Canto M, Mignini Renzini M, Milani R, Fruscio R, Cantù MG, *et al.* Embryo transfer following *in vitro* maturation and cryopreservation of oocytes recovered from antral follicles during conservative surgery for ovarian cancer. *J Assist Reprod Genet* 2012;29:779-81.
 12. Huang JY, Tulandi T, Holzer H, Tan SL, Chian RC. Combining ovarian tissue cryobanking with retrieval of immature oocytes followed by *in vitro* maturation and vitrification: An additional strategy of fertility preservation. *Fertil Steril* 2008;89:567-72.
 13. Huang JY, Chian RC, Gilbert L, Fleischer D, Holzer H, Dermatas E, *et al.* Retrieval of immature oocytes from unstimulated ovaries followed by *in vitro* maturation and vitrification: A novel strategy of fertility preservation for breast cancer patients. *Am J Surg* 2010;200:177-83.
 14. Hourvitz A, Yerushalmi GM, Maman E, Raanani H, Elizur S, Brengauz M, *et al.* Combination of ovarian tissue harvesting and immature oocyte collection for fertility preservation increases preservation yield. *Reprod Biomed Online* 2015;31:497-505.
 15. Toth TL, Lanzendorf SE, Sandow BA, Veeck LL, Hassen WA, Hansen K, *et al.* Cryopreservation of human prophase I oocytes collected from unstimulated follicles. *Fertil Steril* 1994;61:1077-82.
 16. Lee JA, Sekhon L, Grunfeld L, Copperman AB. *In vitro* maturation of germinal vesicle and metaphase I eggs prior to cryopreservation optimizes reproductive potential in patients undergoing fertility preservation. *Curr Opin Obstet Gynecol* 2014;26:168-73.
 17. Brambillasca F, Guglielmo MC, Coticcchio G, Mignini Renzini M, Dal Canto M, Fadini R. The current challenges to efficient immature oocyte cryopreservation. *J Assist Reprod Genet* 2013;30:1531-9.
 18. Son WY, Henderson S, Cohen Y, Dahan M, Buckett W. Immature oocyte for fertility preservation. *Front Endocrinol (Lausanne)* 2019;10:464.
 19. Khalili MA, Shahedi A, Ashourzadeh S, Nottola SA, Macchiarelli G, Palmerini MG. Vitrification of human immature oocytes before and after *in vitro* maturation: A review. *J Assist Reprod Genet* 2017;34:1413-26.
 20. Cao Y, Xing Q, Zhang ZG, Wei ZL, Zhou P, Cong L. Cryopreservation of immature and *in vitro* matured human oocytes by vitrification. *Reprod Biomed Online* 2009;19:369-73.
 21. Fasano G, Demeestere I, Englert Y. *In-vitro* maturation of human oocytes: Before or after vitrification? *J Assist Reprod Genet* 2012;29:507-12.
 22. Song WY, Peng ZF, Chen XM, Jin HX, Yao GD, Shi SL, *et al.* Effects of vitrification on outcomes of *in vivo*-mature, *in vitro*-mature and immature human oocytes. *Cell Physiol Biochem* 2016;38:2053-62.
 23. Wang H, Racowsky C, Combelles CM. Is it best to cryopreserve human cumulus-free immature oocytes before or after *in vitro* maturation? *Cryobiology* 2012;65:79-87.
 24. Yazdanpanah F, Khalili MA, Eftekhari M, Karimi H. The effect of vitrification on maturation and viability capacities of immature human oocytes. *Arch Gynecol Obstet* 2013;288:439-44.
 25. Al-Khtib M, Perret A, Khoeiry R, Ibala-Romdhane S, Blachre T, Greze C, *et al.* Vitrification at the germinal vesicle stage does not affect the methylation profile of H19 and KCNQ1OT1 imprinting centers in human oocytes subsequently matured *in vitro*. *Fertil Steril* 2011;95:1955-60.
 26. Molina I, Gómez J, Balasch S, Pellicer N, Novella-Maestre E. Osmotic-shock produced by vitrification solutions improves immature human oocytes *in vitro* maturation. *Reprod Biol Endocrinol* 2016;14:27.
 27. Liu MH, Zhou WH, Chu DP, Fu L, Sha W, Li Y. Ultrastructural changes and methylation of human oocytes vitrified at the germinal vesicle stage and matured *in vitro* after thawing. *Gynecol Obstet Invest* 2016;82:252-61.
 28. Summezer M, Oktay K. Fertility reservation in female patients. *Hum Reprod Update* 2004;10:251-66.
 29. Demirtas E, Elizur SE, Holzer H, Gidoni Y, Son WY, Chian RC, *et al.* Immature oocyte retrieval in the luteal phase to preserve fertility in cancer patients. *Reprod Biomed Online* 2008;17:520-3.
 30. Polim A, Handayani N, Aprilliana T, Silvia R, Sirait B, Boediono A, *et al.* Association between estradiol levels and clinical outcomes of IVF cycles with single blastocyst embryo transfer. *Asian Pac J Reprod* 2021;10:49-55.
 31. Lim KS, Chae SJ, Choo CW, Ku YH, Lee HJ, Hur CY, *et al.* *In vitro* maturation: Clinical applications. *Clin Exp Reprod Med* 2013;40:143-7.
 32. Lavery S, Baldwin K, Mitchell H, Culley L, Hudson N. Oocyte cryopreservation for social reasons: Demographic profile and disposal intentions of UK users. *Reprod Biomed* 2015;31:239-45.
 33. Shu Y, Gebhardt J, Watt J, Lyon J, Dasig D, Behr B. Fertilization, embryo development, and clinical outcome of immature oocytes from stimulated intracytoplasmic sperm injection cycles. *Fertil Steril* 2007;87:1022-7.
 34. Reichman DE, Politch J, Ginsburg ES, Racowsky C. Extended *in vitro* maturation of immature oocytes from stimulated cycles: An analysis of fertilization potential, embryo development, and reproductive outcomes. *J Assist Reprod Genet* 2010;27:347-56.
 35. Shin SB, Cho JW, Lee SH, Yang KM, Lim CK, Lee HS. Fertilization and pregnancy potential of immature oocytes from stimulated intracytoplasmic sperm injection cycles. *Clin Exp Reprod Med* 2013;40:7-11.
 36. Martin-Palomino Olid N, García D, Rodríguez A, Vassena R. Could fertility clinics offer a sizable improvement of live birth rates by maturing post-GVBD oocytes *in vitro*? *J Assist Reprod Genet* 2019;36:1927-34.
 37. Vanhoutte L, De Sutter P, Van der Elst J, Dhont M. Clinical benefit of metaphase I oocytes. *Reprod Biol Endocrinol* 2005;3:71.
 38. Ferrer-Vaquero A, Barragán M, Rodríguez A, Vassena R. Altered cytoplasmic maturation in rescued *in vitro* matured oocytes. *Hum Reprod* 2019;34:1095-105.
 39. Tucker MJ, Wright G, Morton PC, Massey JB. Birth after cryopreservation of immature oocytes with subsequent *in vitro* maturation. *Fertil Steril* 1998;70:578-9.
 40. Son WY, Tan SL. Laboratory and embryological aspects

- of hCG-primed *in vitro* maturation cycles for patients with polycystic ovaries. *Hum Reprod Update* 2010;16:675-89.
41. Vanhoutte L, Nogueira D, Dumortier F, De Sutter P. Assessment of a new *in vitro* maturation system for mouse and human cumulus-enclosed oocytes: Three-dimensional prematuration culture in the presence of a phosphodiesterase 3-inhibitor. *Hum Reprod* 2009;24:1946-59.
 42. Sánchez F, Lolicato F, Romero S, De Vos M, Van Ranst H, Verheyen G, *et al.* An improved IVM method for cumulus-oocyte complexes from small follicles in polycystic ovary syndrome patients enhances oocyte competence and embryo yield. *Hum Reprod* 2017;32:2056-68.
 43. Hussein TS, Thompson JG, Gilchrist RB. Oocyte-secreted factors enhance oocyte developmental competence. *Dev Biol* 2006;296:514-21.
 44. Shahedi A, Hosseini A, Khalili MA, Norouzi M, Salehi M, Piriaei A, *et al.* The effect of vitrification on ultrastructure of human *in vitro* matured germinal vesicle oocytes. *Eur J Obstet Gynecol Reprod Biol* 2013;167:69-75.
 45. Park SE, Son WY, Lee SH, Lee KA, Ko JJ, Cha KY. Chromosome and spindle configurations of human oocytes matured *in vitro* after cryopreservation at the germinal vesicle stage. *Fertil Steril* 1997;68:920-6.
 46. Van Blerkom J. Maturation at high frequency of germinal-vesicle-stage mouse oocytes after cryopreservation: Alterations in cytoplasmic, nuclear, nucleolar and chromosomal structure and organization associated with vitrification. *Hum Reprod* 1989;4:883-98.
 47. Kim SS, Olsen R, Kim DD, Albertini DF. The impact of vitrification on immature oocyte cell cycle and cytoskeletal integrity in a rat model. *J Assist Reprod Genet* 2014;31:739-47.
 48. Wu C, Rui R, Dai J, Zhang C, Ju S, Xie B, *et al.* Effects of cryopreservation on the developmental competence, ultrastructure and cytoskeletal structure of porcine oocytes. *Mol Reprod Dev* 2006;73:1454-62.
 49. Turathum B, Saikhun K, Sangsuwan P, Kitiyanant Y. Effects of vitrification on nuclear maturation, ultrastructural changes and gene expression of canine oocytes. *Reprod Biol Endocrinol* 2010;8:70.

Potential Use of Immature Oocyte to Improve Fertility Preservation Outcome: A Narrative Review

ORIGINALITY REPORT

16%

SIMILARITY INDEX

10%

INTERNET SOURCES

14%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	taiwanoncologysociety.org.tw Internet Source	1%
2	"In Vitro Fertilization", Springer Science and Business Media LLC, 2019 Publication	1%
3	link.springer.com Internet Source	1%
4	www.frontiersin.org Internet Source	1%
5	"Abstract book of the 31 st ESHRE Annual Meeting, Lisbon, Portugal, 14–17 June 2015", Human Reproduction, 2015. Publication	<1%
6	"Textbook of Assisted Reproduction", Springer Science and Business Media LLC, 2020 Publication	<1%
7	epdf.pub Internet Source	<1%

8	Internet Source	<1 %
9	www.bioseek.eu Internet Source	<1 %
10	www.ecerm.org Internet Source	<1 %
11	www.repdevmed.org Internet Source	<1 %
12	Submitted to CSU, San Marcos Student Paper	<1 %
13	S. Samuel Kim, Rachel Olsen, Dojun David Kim, David F. Albertini. "The impact of vitrification on immature oocyte cell cycle and cytoskeletal integrity in a rat model", <i>Journal of Assisted Reproduction and Genetics</i> , 2014 Publication	<1 %
14	ijfs.ir Internet Source	<1 %
15	www.biorxiv.org Internet Source	<1 %
16	Ingrid Segers, Erlisa Bardhi, Ileana Mateizel, Ellen Van Moer, Rik Schots, Greta Verheyen, Herman Tournaye, Michel De Vos. "Live births following fertility preservation using in-vitro maturation of ovarian tissue oocytes", <i>Human Reproduction</i> , 2020	<1 %

17

Submitted to Udayana University

Student Paper

<1 %

18

"Cryopreservation of Mammalian Gametes and Embryos", Springer Science and Business Media LLC, 2017

Publication

<1 %

19

Jack Yu Jen Huang, Ri-Cheng Chian, Lucy Gilbert, David Fleischer et al. "Retrieval of immature oocytes from unstimulated ovaries followed by in vitro maturation and vitrification: A novel strategy of fertility preservation for breast cancer patients", The American Journal of Surgery, 2010

Publication

<1 %

20

Sonigo, Charlotte, Alice Seroka, Isabelle Cédrin-Durnerin, Nathalie Sermondade, Christophe Sifer, and Michael Grynberg. "History of ABVD alters the number of oocytes vitrified after in vitro maturation in fertility preservation candidates", Future Oncology, 2016.

Publication

<1 %

21

Submitted to University of New South Wales

Student Paper

<1 %

22

Xinyan Cao, Jingchun Li, Hailong Xue, Shiyong Wang, Weigang Zhao, Zhanyu Du, Yifeng Yang,

<1 %

Zhigang Yue. "Effect of vitrification on meiotic maturation, mitochondrial distribution and glutathione synthesis in immature silver fox cumulus oocyte complexes", *Theriogenology*, 2017

Publication

23

Geetha R. "Techniques for Cognitive Pathways from Natrinai", *International Research Journal of Tamil*, 2022

Publication

24

Wang, Haiyan, Catherine Racowsky, and Catherine M.H. Combelles. "Is it best to cryopreserve human cumulus-free immature oocytes before or after in vitro maturation?", *Cryobiology*, 2012.

Publication

25

www.tandfonline.com

Internet Source

26

N. Martin-Palomino Olid, D. García, A. Rodríguez, Rita Vassena. "Could fertility clinics offer a sizable improvement of live birth rates by maturing post-GVBD oocytes in vitro?", *Journal of Assisted Reproduction and Genetics*, 2019

Publication

27

Y. Liu, Y.X. Cao, Z.G. Zhang, Q. Xing. " Artificial oocyte activation and human failed-matured

<1 %

<1 %

<1 %

<1 %

<1 %

oocyte vitrification followed by maturation ", Zygote, 2011

Publication

28

obgyn.onlinelibrary.wiley.com

Internet Source

<1 %

29

repository.uki.ac.id

Internet Source

<1 %

30

www.ncbi.nlm.nih.gov

Internet Source

<1 %

31

"In vitro maturation: a committee opinion",
Fertility and Sterility, 2020

Publication

<1 %

32

Biennial Review of Infertility, 2009.

Publication

<1 %

33

Ghetler, Y.. "Human oocyte cryopreservation
and the fate of cortical granules", Fertility and
Sterility, 200607

Publication

<1 %

34

J. Laurincik. "Differences in pronucleus
formation and first cleavage following in vitro
fertilization between pig oocytes matured in
vivo and in vitro", Reproduction, 11/01/1994

Publication

<1 %

35

Julie Barberet, Fatima Barry, Cécile Choux,
Magali Guilleman, Sara Karoui, Raymond
Simonot, Céline Bruno, Patricia Fauque. "What

<1 %

impact does oocyte vitrification have on epigenetics and gene expression?", Clinical Epigenetics, 2020

Publication

36

Yuanyuan LIANG. "Cryopreservation of immature buffalo oocytes: Effects of cytochalasin B pretreatment on the efficiency of cryotop and solid surface vitrification methods : IMMATURED BUFFALO OOCYTE VITRIFICATION", Animal Science Journal, 04/2012

Publication

<1 %

37

www.createfertility.co.uk

Internet Source

<1 %

38

"Diminished Ovarian Reserve and Assisted Reproductive Technologies", Springer Science and Business Media LLC, 2020

Publication

<1 %

39

Eliza C. Curnow, John P. Ryan, Douglas M. Saunders, Eric S. Hayes. "Primate model of metaphase I oocyte in vitro maturation and the effects of a novel glutathione donor on maturation, fertilization, and blastocyst development", Fertility and Sterility, 2011

Publication

<1 %

40

Ester Polak de Fried M.D., Judith Notrica M.Sc., Mara Rubinstein M.D., Andrea Marazzi M.D., Mercedes Gómez Gonzalez M.Sc..

<1 %

"Pregnancy After Human Donor Oocyte Cryopreservation and Thawing in Association with Intracytoplasmic Sperm Injection in a Patient with Ovarian Failure", Fertility and Sterility, 1998

Publication

41

Ezgi Demirtas. "In-vitro Maturation of Human Oocytes", Reproductive Endocrinology and Infertility, 2010

Publication

42

Park, S.E.. "Chromosome and spindle configurations of human oocytes matured in vitro after cryopreservation at the germinal vesicle stage", Fertility and Sterility, 1997

Publication

43

Sylvie Lierman, Annelies Tolpe, Ilse De Croo, Stefanie De Gheselle et al. "Low feasibility of in vitro matured oocytes originating from cumulus complexes found during ovarian tissue preparation at the moment of gender confirmation surgery and during testosterone treatment for fertility preservation in transgender men", Fertility and Sterility, 2021

Publication

44

Wei Shi, Bo Xu, Li-Min Wu, Ren-Tao Jin, Hong-Bing Luan, Li-Hua Luo, Qing Zhu, Lars Johansson, Yu-Sheng Liu, Xian-Hong Tong. "Oocytes with a Dark Zona Pellucida

<1 %

<1 %

<1 %

<1 %

Demonstrate Lower Fertilization, Implantation and Clinical Pregnancy Rates in IVF/ICSI Cycles", PLoS ONE, 2014

Publication

45

Z.J. Chen. "Effects of sucrose concentration on the developmental potential of human frozen-thawed oocytes at different stages of maturity", Human Reproduction, 08/27/2004

Publication

<1 %

46

coek.info

Internet Source

<1 %

47

journals.plos.org

Internet Source

<1 %

48

uobrep.openrepository.com

Internet Source

<1 %

49

www.tjoddergisi.org

Internet Source

<1 %

50

zoologicalletters.biomedcentral.com

Internet Source

<1 %

51

Allen C. Schroeder. "Developmental capacity of mouse oocytes that undergo maturation in vitro: Effect of the hormonal state of the oocyte donor", Gamete Research, 09/1989

Publication

<1 %

52

Jack Huang, Ri-Cheng Chian, Seang Tan. "Ovarian Hyperstimulation Syndrome

<1 %

Prevention Strategies: In Vitro Maturation", Seminars in Reproductive Medicine, 2010

Publication

53

Mahboobeh Amoushahi, Karin Lykke-Hartmann. "Distinct Signaling Pathways Distinguish in vivo From in vitro Growth in Murine Ovarian Follicle Activation and Maturation", *Frontiers in Cell and Developmental Biology*, 2021

Publication

<1 %

54

Fausta Brambillasca, Maria Cristina Guglielmo, Giovanni Coticchio, Mario Mignini Renzini et al. "The current challenges to efficient immature oocyte cryopreservation", *Journal of Assisted Reproduction and Genetics*, 2013

Publication

<1 %

55

Irma Virant-Klun, Andrej Vogler. "In vitro maturation of oocytes from excised ovarian tissue in a patient with autoimmune ovarian insufficiency possibly associated with Epstein-Barr virus infection", *Reproductive Biology and Endocrinology*, 2018

Publication

<1 %

56

Methods in Molecular Biology, 2014.

Publication

<1 %

57

Michel De Vos. "Fertility preservation in women with cancer: in vitro maturation of

<1 %

oocytes", Expert Review of Quality of Life in Cancer Care, 2016

Publication

58

Weon-Young Son, Sara Henderson, Yoni Cohen, Michael Dahan, William Buckett.
"Immature Oocyte for Fertility Preservation",
Frontiers in Endocrinology, 2019

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On