



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

A Review of Non-Invasive Sampling in Wildlife Disease and Health Research

Citation for published version:

Schilling, A-K, Mazzamuto, MV & Romeo, C 2022, 'A Review of Non-Invasive Sampling in Wildlife Disease and Health Research: What's New?', *Animals*, vol. 12, no. 13, 1719, pp. 1-24.
<https://doi.org/10.3390/ani12131719>

Digital Object Identifier (DOI):

[10.3390/ani12131719](https://doi.org/10.3390/ani12131719)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Animals

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.



Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Review

A Review of Non-Invasive Sampling in Wildlife Disease and Health Research: What's New?

Anna-Katarina Schilling^{1,†}, Maria Vittoria Mazzamuto^{2,3,†} and Claudia Romeo^{4,*}

¹ Previously Royal (Dick) School of Veterinary Studies and Roslin Institute, University of Edinburgh, Easter Bush, Midlothian EH25 9RG, UK; annakatarinaschilling@gmail.com

² Haub School of Environment and Natural Resources, University of Wyoming, 1000 E. University Ave., Laramie, WY 82072, USA; mariavittoria.mazzamuto@uwyo.edu

³ Department of Theoretical and Applied Sciences, University of Insubria, Via J.H. Dunant 3, 21100 Varese, Italy

⁴ Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna (IZSLER), Via Bianchi 9, 25124 Brescia, Italy

* Correspondence: claudiarosa.romeo@izsler.it

† These authors contributed equally to this work.

Simple Summary: The interest in wildlife research has increased in the last decades as more scientists work within a One Health framework that regards human, livestock and wildlife health as connected entities. To minimise the impact of research on wildlife, collecting samples with as little disturbance of the animals as possible is important. In our review, we assess the use of so-called non-invasive sampling and summarise which samples can be used successfully when carrying out research on wildlife diseases and health status. Our results show that interest in minimally invasive sampling has steadily increased since the 2010s. Topics able to employ these methods include disease research, but also stress and other hormone assessments, pollution studies, and dietary studies. At the moment, such methods are mainly used to collect samples from land mammals, however, they can also be used in a wide range of other animals. Ever more capable analytical methods will allow for an even wider use of such “animal-friendly” sampling methods.

Abstract: In the last decades, wildlife diseases and the health status of animal populations have gained increasing attention from the scientific community as part of a One Health framework. Furthermore, the need for non-invasive sampling methods with a minimal impact on wildlife has become paramount in complying with modern ethical standards and regulations, and to collect high-quality and unbiased data. We analysed the publication trends on non-invasive sampling in wildlife health and disease research and offer a comprehensive review on the different samples that can be collected non-invasively. We retrieved 272 articles spanning from 1998 to 2021, with a rapid increase in number from 2010. Thirty-nine percent of the papers were focussed on diseases, 58% on other health-related topics, and 3% on both. Stress and other physiological parameters were the most addressed research topics, followed by viruses, helminths, and bacterial infections. Terrestrial mammals accounted for 75% of all publications, and faeces were the most widely used sample. Our review of the sampling materials and collection methods highlights that, although the use of some types of samples for specific applications is now consolidated, others are perhaps still underutilised and new technologies may offer future opportunities for an even wider use of non-invasively collected samples.

Keywords: non-invasive methods; wildlife species; wildlife pathogens; 3Rs; animal health; field sampling



Citation: Schilling, A.-K.;

Mazzamuto, M.V.; Romeo, C. A

Review of Non-Invasive Sampling in Wildlife Disease and Health Research: What's New? *Animals* **2022**, *12*, 1719. <https://doi.org/10.3390/ani12131719>

Academic Editor: Laila Darwich Soliva

Received: 30 May 2022

Accepted: 29 June 2022

Published: 2 July 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

In the last decades, awareness of the importance of studying and monitoring disease in wild animal populations has steadily grown. The need for a One Health approach

that integrates wildlife, human, and domestic animal health into a single framework is now widely recognised [1] and has gained further attention due to the recent SARS-CoV-2 pandemic [2]. In addition to their relevance as a source of pathogens that may spill over to humans and domestic animals, many wild species are currently threatened by emerging diseases (e.g., white-nose syndrome in bats [3], chytridiomycosis in amphibians [4], or *Chlamydia* infection in koalas [5]), making wildlife disease research paramount from a conservation perspective [6,7]. Additionally, the need for a more complete understanding of disease circulation in natural populations has sparked the interest of many researchers who are now investigating those intrinsic factors affecting host–pathogen dynamics such as the host physiological and immunological status [8–11].

In parallel to the growing interest in wildlife disease, attention towards animal welfare in research has been growing since the '60s, when Russell and Burch [12] proposed the 3Rs principle. It aims to replace animal use in research whenever possible, reduce the numbers of animals employed, and to refine the methods to limit their pain and distress. This principle has been incorporated in several legislations (e.g., [13]) and is now widely applied to laboratory animals. Ethical advances in wildlife research have been comparably slower, but the need for the development of non-invasive methods to safeguard animal welfare and avoid taking valuable animals from vulnerable populations is being advocated by several authors [14–16].

Wildlife disease and health surveys often rely on the opportunistic sampling of carcasses, and this is certainly an important resource for passive disease surveillance (e.g., [17–20] in the present Special Issue), but it can introduce relevant biases into epidemiological studies [21]. Random sampling of living individuals should therefore be preferred whenever researchers are interested in studying disease circulation in natural populations [22]. However, sampling living, free-ranging animals is a challenging endeavour, and biologists and veterinarians need to cope with several issues particular to the field that are not encountered when working with domestic or laboratory animals [23]. First and foremost, relevant constraints are related to logistics, as field work is often carried out in inaccessible locations, far away from laboratory facilities, making sample storage and transport a major concern. Another issue is related to capture and handling: capturing wild animals is normally work-demanding and indeed represents a source of distress for the animal, possibly altering physiological parameters [11] and posing ethical problems.

Hence, a sampling method for wildlife should ideally be non-invasive, possibly require no trapping and/or handling to limit distress (although for some species this might be unavoidable) and be feasible in field conditions. Thanks to the growing interest in both wildlife diseases and health, the number of studies proposing alternative approaches aimed at making wildlife sampling easier and less invasive has increased. Such new approaches either focus on alternative methods to collect and use traditional samples, or on the validation of new sample materials that may represent a reliable alternative to traditional samples that would mostly require invasive collection (e.g., blood). Developing non-invasive methods for wildlife studies is likely to always be a multiple-step process in which samples are initially collected post-mortem or during direct contact with the animal to be able to establish whether a certain parameter or pathogen can be detected reliably. It then needs to be established how a sufficient amount of material can be collected from a live animal without direct interference and causing as little distress as possible. Finally, a further step may be necessary to prove that the results from the samples collected post-mortem and in live animals are actually comparable.

Our aim was to (i) analyse the publication trends on non-invasive sampling methods in wildlife health research and (ii) offer a review on the different samples that can be collected from wildlife non-invasively, what they are useful for and how they can be collected to survey for different pathogens or to investigate the physiological and immunological parameters in wild animals. Our intent in this review was to mimic a wildlife researcher, not necessarily a veterinarian, and possibly new to sampling for disease and/or physiology,

starting to plan a new project involving field sampling, or are interested in using banked samples they have gained access to.

2. Materials and Methods

We searched for published literature on the Web of Science, Scopus, and Google Scholar platforms up until December 2021. Specifically, we used the following search string: (non-invasive OR noninvasive) AND (method* OR technique* OR tool* OR sampling) AND (wildlife OR “wild animals”) AND (disease OR pathogen* OR parasite* OR health). All search results were screened and those not relevant to the present synthesis were excluded. Additionally, we also used publications cited in the articles we found. Studies including post-mortem sampling were retained only when aimed at developing or validating non-invasive methods to be later applied on living individuals. Each paper was classified based on the parameters reported in Table 1.

Table 1. The categories used to classify articles included in the present review.

Topic	Type of Article	Host's Taxon	Material Collected	Collection Method ¹	Detection Method ²
Disease	Research ⁵	Amphibia	Faeces	Collection from habitat	Analytical chemistry
	Bacteria Method ⁶	Aves	Hair-feathers-skin	Hair trapping	Coprological method
	Disease Review	Fish	Imaging and remote sensing	New device	Coprological method combined ¹¹
	Ectoparasites	Mammalia	Invertebrates ⁸	Other animals ⁹	Imaging
	Endoparasites	Marine mammal	Saliva and other body fluids	Post-mortem ¹⁰	Immunoassay
	Fungi	Reptilia	Urine	Trapping and handling	Molecular method
	Helminths	Several ⁷	Several ⁷	Visual	Molecular method combined ¹²
	Protozoa			Several ⁷	Other ³
	Virus				Several ⁷
	Other ³				
Non-disease	Diet				
	Immunity				
	Physiology ⁴				
	Pollutants				
	Reproductive condition				
	Stress				
	Other ³				
Both					

¹ Method used to collect the sample; ² Method applied on the sample for pathogen detection and/or health parameters assessment; ³ Topic not included in any of the above categories (e.g., prion diseases); ⁴ Hormones other than stress steroids, metabolism, body temperature, and other physiological parameters; ⁵ Research articles applying non-invasive sampling methods; ⁶ Research articles defining, testing, or validating new non-invasive methods; ⁷ Two or more of the above categories included; ⁸ Blood-sucking invertebrates used to collect blood from target species; ⁹ for example, blood-sucking invertebrates, sniffer dogs; ¹⁰ roadkill, found carcasses, or hunted animals used for non-invasive method validation; ¹¹ Coprological method combined with immunoassay, molecular method or observation; ¹² Molecular method combined with microscopy or culture.

To assess the independence between some of the categories used to classify papers (Table 1), we performed a Pearson's chi squared independence test. We used Monte Carlo simulation to obtain *p*-values without assuming asymptotically normal behaviour from small cell count (degrees of freedom do not go into the equation at any stage of the computation, so there is no reason to report them). Specifically, we tested the independence between “Biological material collected” and “Host's taxon”, “Biological material collected” and “Topic”, “Detection method” and “Host's taxon”. For these analyses, we excluded papers that were classified as “Reviews” and records with NAs.

3. Results

Our search returned a total of 340 publications. After screening, we retained 272 papers spanning from 1998–2021 (median = 2016, Figure 1) and discarded the ones that were not

relevant to this review. Thirty-nine percent of the papers were focussed on diseases, 58% on other topics, and 3% on both (see Figure 1a for a breakdown of the number of papers per year). We also classified papers based on their type: research articles (38%), methodological articles (51%), and reviews (11%) (Figure 1b).




Figure 1. The number of articles on non-invasive sampling in wildlife published from 1998 to 2021 divided by general topic (a) and type (b). For a definition of categories, see Table 1.

Regarding the topic, “stress” was the one with the highest number of papers published ($n = 67$), followed by “physiology” ($n = 42$) and “virus” ($n = 29$). The other topics accounted for less than 10% each (Figure 2; Supplementary Materials Table S1).

The animal taxon most widely investigated was terrestrial mammals (Mammalia), accounting for 75% of all papers. The other taxa have been less studied: Aves 9%, Amphibia 6%, marine mammals 6%, Reptilia 3%, fish 1%, several taxa together <1%. Sixteen papers were classified as NA if the paper was a general review or methodological without a target species or taxon and were not included in the calculation of the percentage. Fifteen percent of the papers had captive wildlife as the host, mostly terrestrial mammals.

Faeces were used as the biological sample in 50% of both the disease and non-disease papers. The second most used biological material was saliva and other body fluids (14%) (Figure 3; Table S2).




Figure 2. The percentage of published articles per topic. For a definition of categories, see Table 1.




Figure 3. The percentage of published articles by collected material. For a definition of categories, see Table 1.

The methods mostly used to collect biological materials were “Collection from the habitat” (39%) and “Trapping and handling” (36%) (Figure S1, Table S3). Finally, the detection methods mostly used were “Immunoassay” (39%), “Molecular method” (23%) and “Analytical chemistry” (13%) (Figure S2, Table S4).

Our analysis showed that “Biological material collected” was not independent from “Host’s taxon” ($\chi^2 = 147.04$, $p < 0.001$). In fact, there was a positive association between “Saliva and other body fluids” and “Amphibia”: this contributed 38% to the total Chi-square score, accounting for most of the difference between the expected and observed values (Table S5). “Biological material collected” was also not independent from “Topic” ($\chi^2 = 249.08$, $p < 0.001$): the test showed that “hair-feathers-skin” and “pollutants” were positively associated (22% contribution to the total Chi-square score; (Table S6). We found independence between “Host’s taxon” and “Detection method” ($\chi^2 = 58.77$, $p = 0.136$).

4. Discussion

We reviewed the scientific literature published in the last 25 years aimed at implementing non-invasive methods in wildlife disease research. Our analysis of the retrieved publications showed that the number of published articles regarding non-invasive sampling methods has been steadily rising since 2010. This suggests an increased ethical awareness of researchers and a growing need to comply with the more advanced ethical standards requested by society. Most of the publications were methodological articles aimed at describing new methods for the non-invasive collection of samples, but many of these were later applied in research studies. Several reviews were also retrieved, further testifying that the interest in non-invasive methods is nowadays consolidated.

The growing interest in non-invasive methods might also be partially related to the mounting evidence on the impact of acute stress on the physiological parameters and the immune response. Avoiding trapping and handling becomes in some cases a data quality requirement on top of the ethical necessity. This is further reflected by our analysis of the publication topic: stress was by far the most studied topic among those using non-invasive methods, followed by other physiological parameters (e.g., other hormones, body temperature, etc.) and viral diseases. Despite this, a considerable number of articles still relied on captures for sample collection, but this is, in our opinion, hardly avoidable when dealing with certain animal species or sample types or when the identity of the sampled individual is relevant to the research question. However, we believe that nowadays, capture protocols, can—and must—be defined to be as minimally invasive as possible. Additionally, concerning wild birds, many countries have continuous ringing programs in place that may represent an optimal opportunity for sampling, without resorting to additional captures specific for the research aim.

Regarding the type of samples, faeces were by far the most frequently sampled material, followed by saliva and other body fluids and by hairs, feathers, or skin. This is at least partly due to the sheer number of articles on stress evaluation, as faeces are usually the material of choice when studying chronic stress in mammals, but it is also likely to be a consequence of the versatility of this biological material, which can be used for several purposes (see below). Likewise, immunoassays are usually employed to quantify hormones, or their metabolites and they were indeed the most frequent technique applied on samples in the retrieved papers. These were followed by molecular methods, which nowadays are a fundamental tool in pathogen detection, especially viruses and bacteria. Finally, the vast majority of the publications regarded terrestrial mammals, which is unsurprising, considering that such taxonomic bias is well-recognised and common to several fields of wildlife biology [24–27], disease ecology included [28,29].

Below, we will present an overview of the reviewed techniques and methods for collecting samples non-invasively from wildlife, both for disease and health or physiology surveys.

4.1. Faeces

Thanks to their versatility and ease of sampling, faeces are undoubtedly the most widely used material for the non-invasive monitoring of diseases and health in wildlife. This was confirmed by our analysis on the published literature where 126 articles using faecal samples were retrieved.

Regarding their use, faecal samples are of course essential for endoparasite investigations on living animals as they can be used to detect helminth eggs or larvae and coccidian oocysts that are shed with faeces. This is still mostly conducted through traditional copromicroscopy and morphological identification of the isolated specimens [30–33]; molecular approaches can also be applied to identify endoparasite life stages at the species level, since this cannot always be achieved through morphology alone. The extraction, amplification, and sequencing of parasite DNA using species-specific or group-specific primers can be carried out directly from faecal material (e.g., *Echinococcus multilocularis* in coyotes *Canis latrans* [34]), but most often faeces are first processed to allow for the isolation

of helminth eggs and larvae [35–37], or oocysts [38,39] to achieve better results. Moreover, constant advances in genomics and a greater affordability of high-throughput sequencing make metabarcoding a promising tool for the simultaneous screening of all parasitic DNA present in a faecal sample (reviewed in [40]).

These same molecular methods can also be employed to detect viral nucleic acids from faeces, avoiding invasive blood sampling. Faecal samples were used to screen for a wide variety of viruses belonging to several families (e.g., [41–43]), sometimes of zoonotic (e.g., Lassa virus in rodents [44]) or conservation relevance (e.g., peste des petits ruminants in wild ungulates [45]). Specific molecular analyses (i.e., protein misfolding cyclic amplification, PMCA) on faeces have also been used successfully to detect prion diseases such as chronic wasting disease, CWD, in ungulates [46]. Faeces are also routinely used to screen for pathogenic bacteria by standard microbiological techniques or molecular identification (*Clostridium perfringens* [47]; *Rickettsia felis* [48]; *Salmonella enterica* serovar. Enteritidis [49] and Typhimurium [50]; *Mycobacterium bovis* [51]), and nowadays, high-throughput metagenomic techniques allow us to characterise the whole gut microbial community composition at once including the pathogenic bacteria from a single faecal sample [52,53]. Of course, in a One Health context, faeces from wildlife can also be used to investigate antimicrobial resistance (AMR), either by screening for AMR genes through molecular methods, or by culture and phenotypical methods (e.g., [54] on *Staphylococcus aureus* in lemurs).

Other than for pathogen detection, faecal samples represent a source of data on the general health status of wildlife, as they can be used to assess several physiological parameters, the presence of pollutants, and the diet. The metabolites of steroid and thyroid hormones are excreted with faeces, allowing for their non-invasive quantification through specific immunoassays (reviewed in [55–57]) or, less frequently, through mass spectrometry [58]. It is important to note that the species-specific validation of hormonal immunoassays is crucial to obtain reliable results since hormone metabolism and excretion may greatly vary, even within the same genus [59]. However, glucocorticoid metabolites have already been used as a measure of chronic stress and assessed from faeces in a variety of mammalian species [60–67], birds [68–70] as well as reptiles [71]. Likewise, immunoassays on faeces allow for the quantification of the faecal metabolites of sexual steroids [72–75] and can be used, for instance, to monitor the reproductive cycles and pregnancy in endangered species (wombats species [76]; southern white rhinoceros *Ceratotherium simum simum* [77]; collared peccary *Pecari tajacu* [78]; tigers *Panthera tigris* [79]). Finally, faeces can also be used to study the diet of vertebrates (either by traditional microscopy or metagenomic techniques: [80–83]) and to screen for the presence of pollutants or toxic compounds by analytical chemistry [84,85].

For studies at the population-level or disease surveillance, fresh faeces can be collected directly from the habitat (e.g., from pastures, dwellings, latrines, or under nesting sites, depending on the species) without any direct contact with the animal (wild cats *Felis silvestris silvestris* [86]; badgers *Meles meles* [87]; Gentoos penguins *Pygoscelis papua* [88]; several Carnivora and Artiodactyla species in Brazil [89]; Amur leopards *Panthera pardus orientalis* [90]; wolverines *Gulo gulo* [42]). In the case of large mammals, GPS-telemetry might be useful to monitor animal movements and locate areas where faeces are most likely to be found, as it was conducted by Van der Goot and colleagues [77] on southern white rhinoceros or by Di Francesco and colleagues [91] on wolf (*Canis lupus*) packs. For flying or arboreal species (e.g., the lesser horseshoe bat *Rhinolophus hipposideros* [92]), plastic sheets can be placed under nests or roosting sites to facilitate sample collection.

However, when information at the individual-level is needed, researchers may need to resort to observation from a distance to collect fresh faeces as soon as an individual defecates (Upland geese *Chloephaga picta leucoptera* [69]; Barbary macaques *Macaca sylvanus* [93]; red deer *Cervus elaphus* [94]; North Pacific grey whales *Eschrichtius robustus* [75]) or to captures, especially in the case of small birds and mammals (southern pied babblers *Turdoides bicolor* [95]; Muridae and Tupaide in Borneo [96]; Namaqua rock mice *Micaelamys namaquensis* [65]; desert gerbils [66]). Serres-Corral and colleagues [67] applied an interest-

ing technique to individually mark faeces from lions (*Panthera leo*) held in captivity, making use of non-toxic coloured waxes administered with feed. However, applying this method on free-living individuals might prove more challenging.

Once collected, faecal samples may be generally kept safely at room temperature for a limited amount of time, but will then need to be stored appropriately, depending on the specific aim of the study. For example, when carrying out standard copromicroscopy to identify helminth eggs or larvae, to preserve their morphology, faeces are best stored dry at 4 °C and processed within a few days, while freezing is not recommended [97]. On the other hand, storage at −20 °C is needed in the case of genetic analyses or immunoassays on hormones [98]. Additionally, a certain sterility of the sample is sometimes required, for instance, Knutie and Gotanda [52] developed a cheap and easy-to-build device to collect faeces from birds in sterile conditions for microbiome studies. It is also best to avoid contamination with urine when analysing faeces for hormone metabolites because metabolites are also excreted with urine (see below), potentially leading to an overestimation of hormone levels [55,57,59]. Finally, it is worth mentioning that the extraction of nucleic acids from faeces can be challenging compared to other samples due to the presence of inhibiting substances; as a consequence for molecular analyses, specific extraction protocols are required [99,100]. Moreover, when using molecular tools to survey for parasite life stages, additional sample processing steps aimed at breaking egg walls or larvae such as magnetic beads or sonication are needed in order to be able to efficiently extract and amplify DNA (e.g., [101]).

4.2. Urine

Urine was used as the sample material in 19 publications included in this review, 14 of these were studies on primates. Beyond the standard clinical dipstick, urine can be used in the detection of pathogens and to assess the physiological parameters associated with stress, reproductive status, inflammation, and metabolic status.

Urine has been used to detect helminths, for example, the eggs of the nematode *Stephanurus dentatus* that parasitize the urinary tract of wild boars (*Sus scrofa*, [102]) and the antigens of *Taenia serialis* in the urine of gelada monkeys (*Theropitecus gelada*, [103]). Bacterial pathogens with a tropism for kidneys such as zoonotic *Leptospira* spp. have been successfully detected in urine as part of the monitoring and surveillance efforts [104]. The detection of some viruses (e.g., paramyxoviruses in African fruit bats) or antiviral antibodies against simian T-lymphotrophic virus type 1 (STLV-1) in chimpanzees (*Pan troglodytes*) is also possible [105–107]. Urine can also be used for hormonal assessment. For example, stress steroids and their metabolites excreted with urine have been successfully quantified through immunoassays in several mammalian and amphibian species [108–114]. Sexual steroid hormone levels (or their metabolites) can also be assessed from the urine and have thus been used to investigate the reproductive status in mammals (giant pandas *Ailuropoda melanoleuca* [115]; chimpanzees [116]) as well as amphibians (reviewed in [112]). Other hormones that can be isolated from urine are oxytocin [117] and thyroid hormones (reviewed in [56]). The cell-mediated immune response has been also monitored from urine in a range of primates by the quantification of neopterin, a biomarker that increases when an acute inflammation is present [114,118–123]. Metabolic state has also been assessed by measuring urine triiodothyronine in macaques [114], by determining C-peptide and ketone bodies in orangutans [124], and by assessing the nitrogen:creatinine ratio in wild moose [125,126].

In arboreal mammals, urine can often be collected from the ground or leaves immediately after spontaneous urination or by spreading plastic sheets under known roost or sleeping sites [103,105–108,114–121,123,124]. In cold climates, it may be possible to collect naturally frozen urine samples from the snow, however, the dilution effect of the snow needs to be taken into account for analysis [125,126]. If mammals are anaesthetised for other procedures, it is likely that they will urinate during recovery, so sample collection can be attempted in this phase [123]. In amphibians or small mammals, collection during

handling is often possible, either by stimulating urination by massaging the belly or by using small suction devices employing capillary forces, which are applied to the cloaca or end of the urethra [104,109,110,112,113]. After collection, the two main ways of storing urine samples are either immediate freezing, and afterwards limiting the freeze–thaw cycles to a minimum, or the use of filter papers. If fresh urine for neopterin analysis is to be kept for longer periods of time at room temperature, biocidal preservatives can be added to reduce its deterioration [118].

Similarly to faeces, for surveillance efforts in a population, the identification of individuals may not be that relevant; however, it remains important to be aware if there is a risk of contamination of the samples with urine from other cohabitating species than the target one [107]. Finally, the volume of urine samples is often limited and contamination with faeces and soil or exposure to sunlight can have an impact on the analytical results, depending on the component assessed [114,117–119].

4.3. Saliva and Other Body Fluids

The use of saliva and other body fluids (exhaled breath condensate (blow), breath, lacrimal fluid, skin mucus) was described in 34 papers included in this review. This sample category was widely used on amphibians (in 15 out of the 22 publications retrieved for the taxon), where skin mucus represents an important resource for the non-invasive monitoring of diseases and physiological parameters.

The potential uses of saliva are manifold and include stress monitoring, pathogen detection, and antibody detection. Stress monitoring is possible both by cortisol measurement [127,128] and by detecting a suppression in secretory Immunoglobulin A, an indicator of chronic stress [129], given that baseline values are established for the species and population under investigation. Both bacterial pathogens excreted orally such as *Mycobacterium bovis* in Australian brushtail possums and oral microbiomes can be assessed by extracting bacterial DNA and RNA from saliva [130,131]. A wide range of viruses can be isolated from saliva including economically relevant viruses such as foot-and-mouth disease virus [132,133], and classical and African swine fever [134,135] as well as viruses of conservation concern such as the elephant endotheliotropic herpesvirus [136]. All examples of virus isolation from saliva that we found in this review were conducted in mammals [132–139]. Antibody detection in saliva has, for example, been attempted for *M. bovis* antibodies in wild boar [140]. The collection of saliva samples can be carried out directly by rolling a cotton swab over the oral mucosa of a restrained animal [127,128,131,136,137], or indirectly by using artificial baited objects or natural objects from the animals' environment that they routinely interact with. An artificial object that can be attractive to a range of inquisitive mammalian species including, for example, suids and primates, are ropes covered in bait [133–135,137,140]. Baited salt licks [132], interactive enrichment toys [139], and WaxTags® [130] are other objects that encourage animals to lick them and leave saliva behind for collection. Natural objects include all items that a specific species tends to chew and discharge, for example, leaves in some primates [138]. When choosing between direct and indirect collection, it needs to be considered that while direct collection allows us to assign a sample to an individual with great certainty, it also requires this animal to be physically or chemically restrained or long-term habituated and trained for sampling [128]. Indirect sampling removes the need to handle an animal, but requires observation (camera or direct) of the collection object [138], or additional DNA assessment to establish the individual that a sample came from. Choosing a safe and effective object for saliva collection for a species requires detailed knowledge of its behaviour and preferences. To optimise sampling, pre-trials with different objects are often necessary. The risk of interactions of non-target species with the object, objects becoming unretrievable or being swallowed also need to be managed [130,135,137]. Some objects such as cotton ropes may contain natural PCR inhibitors that need to be considered when samples are processed [135,137]. Generally, in saliva samples, it needs to be considered that the oral microflora can be sensitive to

external conditions [131] and that low pathogen excretion can occur, potentially reducing the sensitivity of surveillance efforts using saliva [134].

Blow collection has been described for a range of marine mammals including, for example, humpback whales (*Megaptera novaeangliae*), harbour porpoises (*Phocoena phocoena*), and belugas (*Delphinapterus leucas*) [141–143]. It has been used for bacterial detection [144], respiratory microbiome analysis [145], and viral detection [142] as well as volatile organic compound analysis [146] and cortisol and the detection of other steroid hormones [141,143]. Blow can be collected in a variety of vials, bags, and dishes, either held in the hand if it is possible to get close to the animal or mounted on a pole to be able to work from a greater distance [141,142,146]. Petri dishes tend to allow for the collection of the largest volumes of fluid and can be readily covered in materials and membranes to enhance sampling based on the individual research question [141,143]. A factor to consider when planning to use blow samples is that the sample volume and dilution cannot be controlled, and that the confirmation of the presence of sufficient organic material for analysis may need to be included in any experimental design [141,143].

Breath collection in mammals has been trialled for use in indirect pathogen detection, as the presence of certain pathogens such as *M. bovis*, MAP or *Brucella spp.* can lead to characteristic changes in the composition of volatile organic compounds in breath [147]. As the collection is currently only possible using a complex mask that needs to be tightly fitted to the animal's nose, it can only be used in habituated and fixated individuals [147], which will limit its applicability to wildlife.

Lacrimal fluids have been assessed as a low impact sample to detect cortisol in harbour seals [148]. Animals need to be restrained for sampling and a cotton swab inserted into the conjunctival sac for 15 s. The cortisol concentration in lacrimal fluid did show good correlation with values measured in blood in the same animals, but the measurement range for this new sample type would need to be established for each species it is to be used on [148].

Skin mucus has been collected from amphibians [111,149–155] and fish [156–158]. In amphibians, the main focus of studies utilising skin mucus has been the detection of the fungal pathogen *Batrachochytrium dendrobatidis* [111,150,151,153–155]; however, wider investigations of the skin microbiome [149] and cortisol detection [152] are further uses for this material. The sensitivity of pathogen detection in skin mucus may be limited when the intensity of infection is low [151]. In fish, skin mucus is mainly used for stress assessments, both by isolating cortisol and its metabolites [157] and by assessing the oxidative stress using F2-isoprostanes [156,158]. Mucus collection has only been described during direct handling, either by swabbing or scraping of the skin [151,152,156,158]. This may in itself alter the cortisol excretion, both systemically and locally in the skin [152]. A further method for the minimally invasive assessment of steroid hormones in amphibians and fish is the immersion of caught individuals in a clean tank of water for a predetermined period of time, after which the animal is moved and excreted hormones can be extracted from the water [159–162]. These assays can also be used on environmental water samples; however, ambient cortisol and metabolites as well as steroid hormones may persist in the environment to varying degrees and limit the information that can be gathered about an individual in, for example, a pond [162].

Overall, saliva and other body-associated fluids have great potential both in health and welfare surveillance (microbiome studies, stress assessment) and in pathogen monitoring. However, being able to collect samples that can clearly be assigned to an individual animal without handling it remains a challenge as well as the standardisation of the volume and dilution of samples collected with more hand-off methods.

4.4. Hair, Feathers, and Skin

The collection and analysis of hair, feather, or skin samples were mentioned in 25 papers included in this review. These sample materials are the preferred media for

investigating contaminants; indeed, out of the 16 retrieved publications aimed at detecting pollutants or medicine residuals, 11 relied on either hair or feathers.

Hair samples are used for pollutant and mineral detection in a wide range of mammals, from bats to polar bears [163–168]. Similarly, hair can be used to detect medicine residuals [169]. Further uses are the determination of cortisol concentrations for stress monitoring [167,170–176] and the assessment of steroid hormone concentrations [177]. Some pathogens, for example, adenovirus and squirrelpox virus in Eurasian red squirrels (*Sciurus vulgaris*) [178] and protozoa *Leishmania infantum* can be isolated from hair samples [179,180]. A final use linked to wildlife health is the assessment of nitrogen, carbon, and sulphur composition in the hair to gain insights into the animals' diet including periods of negative energy balance [181,182]. Hair is therefore a very versatile and valuable source of information, however, while a collection is possible without direct contact with the animal, for example, by using hair traps, barbed wire, or collection from known rubbing trees [164,181], most studies included in this review gathered their samples either during direct handling of the physically or chemically restrained animal [168,170,171,177,178,181,182] or from carcasses during post-mortem examination [163–167,169,172,176,178–182]. This implies that further research into low-impact collection methods from live, free-roaming animals is necessary to use this sample type to its full potential. Once the sample is collected, analytical and diagnostic methods for minerals, pollutants, residuals, diet, and pathogens are usually well-established. It does need to be established for each species, though, whether the correlation between the blood levels and levels in hair is reliable, as it may be poor in some species [168]. With regard to pathogens, the location a pathogen takes after infection needs to be taken into consideration to determine whether an animal is infected with a certain pathogen or if the pathogen may just be present on its fur as a contaminant. To use hair samples for cortisol determination, it needs to be considered that in some species, age, pregnancy, colour, body region, sex, and season influence cortisol levels [170,174,175], and that local cortisol production in the hair follicle may occur, in which case the hair cortisol concentration may not offer information on the HPA-axis activity [170]. A final factor to consider when designing studies using hair samples is the number of hairs needed to produce reliable results, as too small a sample may result in false negatives, while too large a sample may cause inhibition during analysis [178].

For feathers, our search only flagged up publications using them for the detection of heavy metal pollution (As Cd, Cu, Hg, Ni, Pb, Se, Zn) or mercury alone (Hg) [183–187]. However, based on the experience with hair, it may eventually be possible to use feathers for the detection of other contaminants such as medicine residuals. While one study relied on feathers collected post-mortem [183], others have used the plucking of feathers during physical restraint and handling [184,187]. It has, however, been shown that meaningful results can also be achieved by collecting clean, freshly shed feathers from nest boxes, still allowing a certain level of information of which animal a feather came from, or at least narrowing the pool down to the breeding pair and their offspring [186]. Disturbance to the nest needs to be weighed against handling stress to determine which of the two collection methods actually has less impact on the species under investigation in each study.

Skin samples of marine mammals can be used for trace element detection (Al, V, Cr, Mn, Cu, Zn, As, Se, Rb, Sr, Mo, Cd, Pb), but only if a full biopsy is collected, which, depending on size, can actually be an invasive procedure [188]. Truly minimally invasive samples of dead superficial skin cells collected with combs can be used for ectoparasite detection, for example, *Antarctophthirus microchir* in American sea lions (*Otaria flavescens*) [189]. Skin scrapings are commonly used on several species for *Sarcoptes scabiei* detection. However, depending on the handler experience and technique used, they may reach a depth where they draw blood and are no longer truly minimally invasive. Fraser and colleagues [190] compared the efficacy of PCR on skin scrapings and on less-invasive skin swabs for *S. scabiei* detection in wombats (*Vombatus ursinus*). Although they showed that PCR has a higher sensitivity than standard microscopy for ectoparasite detection, they concluded that PCR

on swabs is less accurate [190]. For reptiles, the detection of mercury in skin scutes has been described [191]. It would be of interest to assess whether the shed skins of reptiles could be used with similarly reliable results.

4.5. Imaging and Remote Sensing

This sample material class included 22 articles that mostly studied the physiological parameters of wild mammals (terrestrial and marine), birds, or reptiles. However, a few studies have also used imaging, observation, and remote sensing to investigate the parasites and health conditions.

One of the tools used in most of the studies was infrared thermography (IRT). Thanks to the development of new and lower cost portable systems, IRT has become a non-invasive technique that allows for the precise measurement of infrared radiation, and hence spatial variation in body surface temperature, it can be used at less than one metre to investigate specific body regions, or from a distance up to hundreds of meters when just the detection of an animal in the wild is desired. The physiological parameter mostly recorded with IRT clearly was the body temperature of wildlife in the wild and in captivity. While some studies have focused on the study of thermoregulation mechanisms (e.g., [192–194]), others have used body temperature as a proxy for health, body condition, and the metabolic state of an animal (e.g., [195–197]). For example, Burns, McCafferty, and Kennedy [192] studied the nesting thermal biology of wild leatherback turtles (*Dermochelys coriacea*), avoiding any interference or stress in this reproductive phase thanks to the use of thermal imaging. Researchers were able to study the spatial and temporal variation in the body surface temperature through all phases of nesting behaviour, and to estimate the relationships between body temperatures and the environmental conditions of the beach. It is important to highlight that those studies using IRT need to consider some important parameters to record accurate temperatures and avoid errors (e.g., variation in emissivity, evaporative cooling, radiative heating of the coat) [198]. This was also highlighted by Horton and colleagues [193], who paired the technology of IRT with that of unoccupied aerial systems (drones) to document the blowhole and dorsal fin skin temperature, respiration rate, and heart rate of adult and calf humpback whales (*Megaptera novaeangliae*). Few attempts have been made to use IRT in the detection of disease in wild mammals [mange [199]; rabies [200]; foot-and-mouth disease [201]]. However, the latter two studies were carried out in a controlled environment in captivity. Arenas and colleagues [199] tested IRT to diagnose inflamed skin infected by sarcoptic mange (*Sarcoptes scabiei*) in wild Spanish ibex (*Capra pyrenaica*). Unfortunately, visual observation resulted in being more effective than IRT due to the low accuracy of the thermal image at distances greater than 100 m.

Since *S. scabiei* results in visible hair loss and affects a broad host range, another tool that has been successfully used to detect mange in wild canids are camera traps [202,203]. However, the distance at which the animal is detected by the camera, hence the resolution of the image, could be a limitation if this tool is to be used for early diagnosis to prevent epidemics of mange that could otherwise have high morbidity and mortality rates. Camera traps can be used to detect any health condition that affects the look of an animal in the wild. For example, Lacroux and colleagues [204] successfully investigated the occurrence of facial dysplasia in wild forest olive baboons (*Papio anubis*) in Uganda as a suspected alteration of embryonic development due to the use of pesticides in crops at the border of their habitat.

A more recent paper showed the use of a common survey tool for bats as a surveillance method for white-nose syndrome (WNS) [205]. Thanks to the collection of bat echolocation sound, the authors developed site-specific prediction models for bat activity accounting for seasonal and annual temperature variation prior to the known occurrence of WNS. These models were then used to monitor changes in bat activity that may signal the potential presence of WNS. While this seems to be an effective method to detect WNS, the authors highlight the need to collect baseline (pre-WNS) data across several years and sites and then look for bat activity in excess of the predicted range during winter months, followed

by bat activity below the predicted range during the following spring and summer. As a consequence, this approach cannot be applied to populations that might already be infected and is simple to apply only if passive autonomous recording units are deployed.

4.6. Other

A few papers have used other animals as either a detection or collection tool. This is the case of the sample class “Invertebrates”, where researchers have used blood sucking invertebrates to collect and then analyse the blood of the target wildlife species (six papers). It must be noted, however, that although these studies considered collecting blood samples through invertebrates as non-invasive, this method is lethal to the invertebrates used. Two studies used leeches to collect blood from mammals. Kvapil and colleagues [206] used medicinal leeches (*Hirudo medicinalis*) as an alternative to more complex and invasive methods for blood sampling for preventive medicine and epidemiological studies in zoo animals looking for antibodies to tick-borne encephalitic virus. Alfano and colleagues [207] collected leech-derived iDNA (terrestrial leeches: *Haemadipsa picta*, *H. zeylanica*) in the forests of the Malaysian Borneo to detect and identify known and novel mammalian viruses. In this study, iDNA was also paired with eDNA from waterholes to study wildlife viral diversity and to detect novel potentially zoonotic viruses prior to their emergence. Thomsen, Voigh, and colleagues [208,209] tested the use of kissing bugs (Reduviidae; Triatominae; Heteroptera) as a gentler system to collect blood to detect rabbit haemorrhagic disease virus in domestic rabbits [209] and in captive primates [208]. Hoffmann and colleagues [210] assessed the feasibility of the fly-based surveillance of wildlife infectious diseases, specifically for adenoviruses (family Adenoviridae), in wild-living vertebrates in a tropical rainforest in Côte d’Ivoire. They concluded that this method could probably be used to detect the genetic material of wildlife infectious agents causing wildlife mass mortality in pristine areas. However, they also highlighted that characterising the genetic diversity of wildlife infectious agents through fly-based monitoring may not be cost-efficient. Other invertebrates such as engorged ticks collected from wild hosts may be used to indirectly screen for pathogens, as was carried out, for instance, by Zechmeisterova and colleagues [211], who used engorged ticks to detect parasitic protozoa in the Iberian lizard (*Lacerta schreiberi*).

The detection of viral infections can also be conducted collecting samples from the environment where the targeted wildlife live. The study previously mentioned by Alfano and colleagues [207] paired the iDNA analyses with those of the eDNA collected from waterholes in Tanzania and Mongolia. The indirect transmission of beak and feather disease virus, which is of global concern and can cause lethal infections, was tested by Martens and colleagues [212] using nest swabs. They provided novel insights into the potential role of nest cavities and other fomites in the indirect transmission of this virus and possibly other pathogens, and offered a non-invasive method for the surveillance of pathogens in wild bird populations.

The use of odour-detection dogs was evaluated as the methodology to locate faecal samples that were then analysed to detect parasite infections. Detection dogs resulted in being more efficient than humans in detecting scat samples [213] and allowed for the collection of more than 600 cervid scats in the study by Teixeira and colleagues [39] in Brazil with the aim of detecting and genetically characterising the infections by *Cryptosporidium* in brockets (*Mazama gouazoubira*). Curry and colleagues [214] tested, for the first time, the use of sniffer dogs in the medical diagnoses of wildlife species. They investigated the reliability of a trained dog for pregnancy detection in polar bears *Ursus maritimus*. The dog had a reduced sensitivity in testing versus training and the authors discussed the possible causes. Moreover, the authors suggest that it is likely that many unique cases of condition are required to sufficiently train an odour-detection dog, which may be prohibitive in wildlife studies when sample sizes are liable to be limited.

4.7. Additional Considerations

We designed our search to mimic a wildlife researcher, possibly new to sampling for disease and/or physiology, starting to plan a new project involving field sampling or interested in using banked samples they have gained access to. In particular, minimally invasive samples are often collected from wildlife without a specific goal in mind and stored until it is possible to acquire funding or connect to colleagues with the right skills and equipment to analyse them further. This approach may seem alien to researchers used to working in laboratory-based settings; however, it is extremely important in wildlife research due to the often challenging field conditions that make every sampling occasion an opportunity to be seized. Therefore, we designed our search in the broadest way we could think of, attempting to capture as many publications presenting or discussing minimally invasive methods as possible. However, we soon realised that our search missed several publications that might have been of interest. For instance, even some of our own papers using non-invasive methods were not retrieved (e.g., [215,216]). Other articles deserving of attention were pointed out to us by peer-reviewers such as a study using ticks to survey for pathogens in wombats [217], another one describing the use of a cytology cell sampler to collect samples from lesions in marine mammals [218], or yet another describing the use of drones to collect blow from them [219]. We were also made aware of the opportunity to detect viruses and cortisol from bird feathers [220,221] or to use pellets from birds of prey or owls for microbiological studies [222,223]. This led us to reflect on why all of those articles had been missed and we concluded that in most cases, neither we nor our colleagues had made it sufficiently explicit that the research employed non-invasive methods (i.e., we had not included the term in the title, keywords or the abstract, and sometimes not even in the full text). This may reduce the opportunities to inform future projects and may slow down the process of employing minimally invasive methods to their full potential. We would therefore like to use this final paragraph to encourage the research community using non-invasive methods on wildlife to tag this clearly and make the information easily accessible to all those interested.

5. Conclusions

Our review of the literature addressing non-invasive methods for the disease and health monitoring of wildlife has highlighted the growing interest of the research community in sampling free-ranging animals non-invasively, with as little interference as possible. However, we believe that there is still much room for improvement, for instance, saliva and urine have been proven to be effective methods for the detection of some pathogens, but still appear underutilised, perhaps because collection can be more challenging compared to other mediums. However, our review highlighted that some specific methods and novel devices for their sampling have already been developed. Additionally, new opportunities for the development and implementation of non-invasive sampling may arise in the future following advances in rapidly evolving fields such as molecular biology or imaging and remote sensing technologies. Finally, we would like to stress again that, although trapping and handling might be still unavoidable in some cases, researchers should always employ rigorous protocols aimed at minimising distress for the animal. A responsible use of wildlife in research, in line with biodiversity conservation efforts, must make the further development of non-invasive sampling methods and a comprehensive use of all samples available for as many types of analyses as possible a priority. We sincerely hope that the trends for the increased use of non-invasive methods as identified in this review continues.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ani12131719/s1>. Table S1. Number and percentage of papers published per topic. Table S2. Number and percentage of papers published per biological material collected (21 NA removed). Figure S1. Percentage of published articles classified based on method used to collect the samples. Table S3. Number and percentage of papers published per biological material collection method (26 NA removed). Figure S2. Percentage of published articles classified based

on the detection method. Table S4. Number and percentage of papers published per detection method (18 NA removed). Table S5. Percentage contribution to the Chi-squared independence test biological material sample collected—Host's taxon. Table S6. Percentage contribution to Chi-squared independence test Topic—Biological material sample collected.

Author Contributions: Conceptualisation, C.R. and A.-K.S.; Methodology, C.R., A.-K.S. and M.V.M.; Formal analysis, M.V.M.; Data curation, C.R., A.-K.S. and M.V.M.; Writing—original draft preparation, C.R., A.-K.S. and M.V.M.; Writing—review & editing, C.R., A.-K.S. and M.V.M. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: No new data were created or analyzed in this study. Data sharing is not applicable to this article.

Acknowledgments: We thank the Editor and three anonymous reviewers for the careful and insightful review of our manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Destoumieux-Garzón, D.; Mavingui, P.; Boetsch, G.; Boissier, J.; Darriet, F.; Duboz, P.; Fritsch, C.; Giraudoux, P.; Le Roux, F.; Morand, S.; et al. The One Health Concept: 10 Years Old and a Long Road Ahead. *Front. Vet. Sci.* **2018**, *5*, 14. [[CrossRef](#)] [[PubMed](#)]
2. Watsa, M. Wildlife Disease Surveillance Focus Group Rigorous Wildlife Disease Surveillance. *Science* **2020**, *369*, 145–147. [[CrossRef](#)] [[PubMed](#)]
3. Blehert, D.S.; Hicks, A.C.; Behr, M.; Meteyer, C.U.; Berlowski-Zier, B.M.; Buckles, E.L.; Coleman, J.T.H.; Darling, S.R.; Gargas, A.; Niver, R.; et al. Bat White-Nose Syndrome: An Emerging Fungal Pathogen? *Science* **2009**, *323*, 227. [[CrossRef](#)] [[PubMed](#)]
4. Fisher, M.C.; Garner, T.W.J.; Walker, S.F. Global Emergence of *Batrachochytrium dendrobatidis* and *Amphibian chytridiomycosis* in Space, Time, and Host. *Annu. Rev. Microbiol.* **2009**, *63*, 291–310. [[CrossRef](#)]
5. Quigley, B.L.; Timms, P. Helping Koalas Battle Disease—Recent Advances in Chlamydia and Koala Retrovirus (KoRV) Disease Understanding and Treatment in Koalas. *FEMS Microbiol. Rev.* **2020**, *44*, 583–605. [[CrossRef](#)]
6. Daszak, P.; Cunningham, A.A.; Hyatt, A.D. Emerging Infectious Diseases of Wildlife—Threats to Biodiversity and Human Health. *Science* **2000**, *287*, 443–449. [[CrossRef](#)]
7. Thompson, R.C.A.; Lymbery, A.J.; Smith, A. Parasites, Emerging Disease and Wildlife Conservation. *Int. J. Parasitol.* **2010**, *40*, 1163–1170. [[CrossRef](#)]
8. Beldomenico, P.M.; Begon, M. Disease Spread, Susceptibility and Infection Intensity: Vicious Circles? *Trends Ecol. Evol.* **2010**, *25*, 21–27. [[CrossRef](#)]
9. Hawley, D.M.; Altizer, S.M. Disease Ecology Meets Ecological Immunology: Understanding the Links between Organismal Immunity and Infection Dynamics in Natural Populations. *Funct. Ecol.* **2011**, *25*, 48–60. [[CrossRef](#)]
10. Pedersen, A.B.; Babayan, S.A. Wild Immunology. *Mol. Ecol.* **2011**, *20*, 872–880. [[CrossRef](#)]
11. Hing, S.; Narayan, E.J.; Thompson, R.C.A.; Godfrey, S.S.; Hing, S.; Narayan, E.J.; Thompson, R.C.A.; Godfrey, S.S. The Relationship between Physiological Stress and Wildlife Disease: Consequences for Health and Conservation. *Wildl. Res.* **2016**, *43*, 51–60. [[CrossRef](#)]
12. Russell, W.M.S.; Burch, R.L. *The Principles of Humane Experimental Technique*; Universities Federation for Animal Welfare: Wheathampstead, UK, 1959.
13. The European Parliament; The Council of The European Union. Directive 2010/63/EU of the European Parliament and of the Council of 22 September 2010 on the Protection of Animals Used for Scientific Purposes Text with EEA Relevance. *Off. J. Eur. Union* **2010**, *276*, 32010L0063.
14. Lindsjö, J.; Fahlman, Å.; Törnqvist, E. Animal Welfare from Mouse to Moose—Implementing the Principles of the 3Rs in Wildlife Research. *J. Wildl. Dis.* **2016**, *52*, S65–S77. [[CrossRef](#)] [[PubMed](#)]
15. Field, K.A.; Paquet, P.C.; Artelle, K.; Proulx, G.; Brook, R.K.; Darimont, C.T. Publication Reform to Safeguard Wildlife from Researcher Harm. *PLoS Biol.* **2019**, *17*, e3000193. [[CrossRef](#)] [[PubMed](#)]
16. Zemanova, M.A. Towards More Compassionate Wildlife Research through the 3Rs Principles: Moving from Invasive to Non-Invasive Methods. *Wildl. Biol.* **2020**, *2020*, wlb.00607. [[CrossRef](#)]
17. Fusillo, R.; Romanucci, M.; Marcelli, M.; Massimini, M.; Della Salda, L. Health and Mortality Monitoring in Threatened Mammals: A First Post Mortem Study of Otters (*Lutra lutra* L.) in Italy. *Animals* **2022**, *12*, 609. [[CrossRef](#)]
18. Rohner, S.; Wohlsein, P.; Prenger-Berninghoff, E.; Ewers, C.; Waindok, P.; Strube, C.; Baechlein, C.; Becher, P.; Wilmes, D.; Rickerts, V.; et al. Pathological Findings in Eurasian Otters (*Lutra lutra*) Found Dead between 2015–2020 in Schleswig-Holstein, Germany. *Animals* **2022**, *12*, 59. [[CrossRef](#)]

19. Shuttleworth, C.M.; Everest, D.; Holmes, P.; Bell, S.; Cripps, R. An Opportunistic Assessment of the Impact of Squirrelpox Disease Outbreaks upon a Red Squirrel Population Sympatric with Grey Squirrels in Wales. *Animals* **2022**, *12*, 99. [[CrossRef](#)]
20. Vengušt, G.; Kuhar, U.; Jerina, K.; Švara, T.; Gombač, M.; Bandelj, P.; Vengušt, D.Ž. Passive Disease Surveillance of Alpine Chamois (*Rupicapra r. rupicapra*) in Slovenia between 2000 and 2020. *Animals* **2022**, *12*, 1119. [[CrossRef](#)]
21. Nusser, S.M.; Clark, W.R.; Otis, D.L.; Huang, L. Sampling Considerations for Disease Surveillance in Wildlife Populations. *J. Wildl. Manag.* **2008**, *72*, 52–60. [[CrossRef](#)]
22. Lachish, S.; Murray, K.A. The Certainty of Uncertainty: Potential Sources of Bias and Imprecision in Disease Ecology Studies. *Front. Vet. Sci.* **2018**, *5*, 90. [[CrossRef](#)] [[PubMed](#)]
23. Ryser-Degiorgis, M.-P. Wildlife Health Investigations: Needs, Challenges and Recommendations. *BMC Vet. Res.* **2013**, *9*, 223. [[CrossRef](#)] [[PubMed](#)]
24. Clark, J.A.; May, R.M. Taxonomic Bias in Conservation Research. *Science* **2002**, *297*, 191–192. [[CrossRef](#)] [[PubMed](#)]
25. Pyšek, P.; Richardson, D.M.; Pergl, J.; Jarošík, V.; Sixtová, Z.; Weber, E. Geographical and Taxonomic Biases in Invasion Ecology. *Trends Ecol. Evol.* **2008**, *23*, 237–244. [[CrossRef](#)]
26. Meyer, C.; Kreft, H.; Guralnick, R.; Jetz, W. Global Priorities for an Effective Information Basis of Biodiversity Distributions. *Nat. Commun.* **2015**, *6*, 8221. [[CrossRef](#)]
27. Rosenthal, M.F.; Gertler, M.; Hamilton, A.D.; Prasad, S.; Andrade, M.C.B. Taxonomic Bias in Animal Behaviour Publications. *Anim. Behav.* **2017**, *127*, 83–89. [[CrossRef](#)]
28. Han, B.A.; Kramer, A.M.; Drake, J.M. Global Patterns of Zoonotic Disease in Mammals. *Trends Parasitol.* **2016**, *32*, 565–577. [[CrossRef](#)]
29. Plourde, B.T.; Burgess, T.L.; Eskew, E.A.; Roth, T.M.; Stephenson, N.; Foley, J.E. Are Disease Reservoirs Special? Taxonomic and Life History Characteristics. *PLoS ONE* **2017**, *12*, e0180716. [[CrossRef](#)]
30. Hoby, S.; Schwarzenberger, F.; Doherr, M.G.; Robert, N.; Walzer, C. Steroid Hormone Related Male Biased Parasitism in Chamois, *Rupicapra Rupicapra Rupicapra*. *Vet. Parasitol.* **2006**, *138*, 337–348. [[CrossRef](#)]
31. Stringer, A.P.; Smith, D.; Kerley, G.I.H.; Linklater, W.L. Reducing Sampling Error in Faecal Egg Counts from Black Rhinoceros (*Diceros bicornis*). *Int. J. Parasitol. Parasites Wildl.* **2014**, *3*, 1–5. [[CrossRef](#)]
32. Catalano, S.; Symeou, A.; Marsh, K.J.; Borlase, A.; Léger, E.; Fall, C.B.; Sène, M.; Diouf, N.D.; Ianniello, D.; Cringoli, G.; et al. Mini-FLOTAC as an Alternative, Non-Invasive Diagnostic Tool for *Schistosoma Mansoni* and Other Trematode Infections in Wildlife Reservoirs. *Parasit. Vectors* **2019**, *12*, 439. [[CrossRef](#)] [[PubMed](#)]
33. Carrau, T.; Martínez-Carrasco, C.; Garijo, M.M.; Alonso, F.; de Ybáñez, R.R.; Tizzani, P. Evaluation of the Baermann–Wetzel Method for Detecting Lungworm Larvae in Wild Ruminants from Faecal Samples. *J. Helminthol.* **2021**, *95*. [[CrossRef](#)] [[PubMed](#)]
34. Liccioli, S.; Rogers, S.; Greco, C.; Kutz, S.J.; Chan, F.; Ruckstuhl, K.E.; Massolo, A. Assessing Individual Patterns of *Echinococcus Multilocularis* Infection in Urban Coyotes: Non-Invasive Genetic Sampling as an Epidemiological Tool. *J. Appl. Ecol.* **2015**, *52*, 434–442. [[CrossRef](#)]
35. Al-Sabi, M.N.S.; Deplazes, P.; Webster, P.; Willeßen, J.L.; Davidson, R.K.; Kapel, C.M.O. PCR Detection of *Angiostrongylus vasorum* in Faecal Samples of Dogs and Foxes. *Parasitol. Res.* **2010**, *107*, 135–140. [[CrossRef](#)]
36. Poglayen, G.; Gori, F.; Morandi, B.; Galuppi, R.; Fabbri, E.; Caniglia, R.; Milanese, P.; Galaverni, M.; Randi, E.; Marchesi, B.; et al. Italian Wolves (*Canis lupus italicus altobello*, 1921) and Molecular Detection of Taeniids in the Foreste Casentinesi National Park, Northern Italian Apennines. *Int. J. Parasitol. Parasites Wildl.* **2017**, *6*, 1–7. [[CrossRef](#)] [[PubMed](#)]
37. Solórzano-García, B.; Pérez-Ponce de León, G. Helminth Parasites of Howler and Spider Monkeys in Mexico: Insights into Molecular Diagnostic Methods and Their Importance for Zoonotic Diseases and Host Conservation. *Int. J. Parasitol. Parasites Wildl.* **2017**, *6*, 76–84. [[CrossRef](#)]
38. Bertram, M.R.; Hamer, G.L.; Snowden, K.F.; Hartup, B.K.; Hamer, S.A. Coccidian Parasites and Conservation Implications for the Endangered Whooping Crane (*Grus americana*). *PLoS ONE* **2015**, *10*, e0127679. [[CrossRef](#)]
39. Teixeira, W.F.P.; De Oliveira, M.L.; de Faria Peres, P.H.; Nagata, W.B.; Santana, B.N.; Oliveira, B.C.M.; Duarte, J.M.B.; Cardoso, T.C.; Lopes, W.D.Z.; Bresciani, K.D.S. *Cryptosporidium Parvum* in Brown Brocket (*Mazama gouazoubira*) from Brazil: First Report of the Subtype IIAA16G3R1 in Cervids. *Parasitol. Int.* **2021**, *80*, 102216. [[CrossRef](#)]
40. Aivelo, T.; Medlar, A. Opportunities and Challenges in Metabarcoding Approaches for Helminth Community Identification in Wild Mammals. *Parasitology* **2018**, *145*, 608–621. [[CrossRef](#)]
41. De Carvalho Ferreira, H.C.; Weesendorp, E.; Quak, S.; Stegeman, J.A.; Loeffen, W.L.A. Suitability of Faeces and Tissue Samples as a Basis for Non-Invasive Sampling for African Swine Fever in Wild Boar. *Vet. Microbiol.* **2014**, *172*, 449–454. [[CrossRef](#)]
42. Bando, R.A.; Bautista, J.; Lund, M.; Newkirk, E.; Squires, J.; Varsani, A.; Kraberger, S. Identification of Novel Circovirus and Anelloviruses from Wolverines Using a Non-Invasive Faecal Sampling Approach. *Infect. Genet. Evol.* **2021**, *93*, 104914. [[CrossRef](#)] [[PubMed](#)]
43. Hardmeier, I.; Aeberhard, N.; Qi, W.; Schoenbaechler, K.; Kraettli, H.; Hatt, J.-M.; Fraefel, C.; Kubacki, J. Metagenomic Analysis of Faecal and Tissue Samples from 18 Endemic Bat Species in Switzerland Revealed a Diverse Virus Composition Including Potentially Zoonotic Viruses. *PLoS ONE* **2021**, *16*, e0252534. [[CrossRef](#)] [[PubMed](#)]
44. Wood, R.; Bangura, U.; Mariën, J.; Douno, M.; Fichet-Calvet, E. Detection of Lassa Virus in Wild Rodent Feces: Implications for Lassa Fever Burden within Households in the Endemic Region of Faranah, Guinea. *One Health* **2021**, *13*, 100317. [[CrossRef](#)]

45. Bataille, A.; Kwiatek, O.; Belfkhi, S.; Mounier, L.; Parida, S.; Mahapatra, M.; Caron, A.; Chubwa, C.C.; Keyyu, J.; Kock, R.; et al. Optimization and Evaluation of a Non-Invasive Tool for Peste Des Petits Ruminants Surveillance and Control. *Sci. Rep.* **2019**, *9*. [[CrossRef](#)] [[PubMed](#)]
46. Cheng, Y.C.; Hannaoui, S.; John, T.R.; Dudas, S.; Czub, S.; Gilch, S. Early and Non-Invasive Detection of Chronic Wasting Disease Prions in Elk Feces by Real-Time Quaking Induced Conversion. *PLoS ONE* **2016**, *11*, e0166187. [[CrossRef](#)] [[PubMed](#)]
47. Fujita, S.; Kageyama, T. Polymerase Chain Reaction Detection of *Clostridium perfringens* in Feces from Captive and Wild Chimpanzees, Pan Troglodytes. *J. Med. Primatol.* **2007**, *36*, 25–32. [[CrossRef](#)]
48. Keita, A.K.; Socolovschi, C.; Ahuka-Mundeke, S.; Ratmanov, P.; Butel, C.; Ayouba, A.; Inogwabini, B.-I.; Muyembe-Tamfum, J.-J.; Mpoudi-Ngole, E.; Delaporte, E.; et al. Molecular Evidence for the Presence of *Rickettsia felis* in the Feces of Wild-Living African Apes. *PLoS ONE* **2013**, *8*, e54679. [[CrossRef](#)]
49. Smith, A.; Ismail, H.; Henton, M.; Keddy, K. GERMS-SA Surveillance Network Similarities between *Salmonella enteritidis* Isolated from Humans and Captive Wild Animals in South Africa. *J. Infect. Dev. Ctries.* **2014**, *8*, 1615–1619. [[CrossRef](#)]
50. Grant, D.; Todd, P.A.; Pennycott, T. Monitoring Wild Greenfinch (*Carduelis chloris*) for *Salmonella enterica typhimurium*. *Ecol. Res.* **2007**, *22*, 571–574. [[CrossRef](#)]
51. King, H.C.; Murphy, A.; James, P.; Travis, E.; Porter, D.; Sawyer, J.; Cork, J.; Delahay, R.J.; Gaze, W.; Courtenay, O.; et al. Performance of a Noninvasive Test for Detecting *Mycobacterium bovis* Shedding in European Badger (*Meles meles*) Populations. *J. Clin. Microbiol.* **2015**, *53*, 2316–2323. [[CrossRef](#)]
52. Knutie, S.A.; Gotanda, K.M. A Non-Invasive Method to Collect Fecal Samples from Wild Birds for Microbiome Studies. *Microb. Ecol.* **2018**, *76*, 851–855. [[CrossRef](#)] [[PubMed](#)]
53. Wolf, J.F.; Kriss, K.D.; MacAulay, K.M.; Munro, K.; Patterson, B.R.; Shafer, A.B.A. Gut Microbiome Composition Predicts Summer Core Range Size in Two Divergent Ungulates. *FEMS Microbiol. Ecol.* **2021**, *97*, fiab048. [[CrossRef](#)] [[PubMed](#)]
54. Schaumburg, F.; Mugisha, L.; Kappeller, P.; Fichtel, C.; Köck, R.; Köndgen, S.; Becker, K.; Boesch, C.; Peters, G.; Leendertz, F. Evaluation of Non-Invasive Biological Samples to Monitor *Staphylococcus aureus* Colonization in Great Apes and Lemurs. *PLoS ONE* **2013**, *8*, e78046. [[CrossRef](#)]
55. Palme, R.; Touma, C.; Arias, N.; Dominchin, M.F.; Lepschy, M. Steroid Extraction: Get the Best out of Faecal Samples. *Wien. Tierärztl. Monatsschrift* **2013**, *9*, 238–246.
56. Behringer, V.; Deimel, C.; Hohmann, G.; Negrey, J.; Schaeb, F.S.; Deschner, T. Applications for Non-Invasive Thyroid Hormone Measurements in Mammalian Ecology, Growth, and Maintenance. *Horm. Behav.* **2018**, *105*, 66–85. [[CrossRef](#)]
57. Sheriff, M.J.; Dantzer, B.; Delehanty, B.; Palme, R.; Boonstra, R. Measuring Stress in Wildlife: Techniques for Quantifying Glucocorticoids. *Oecologia* **2011**, *166*, 869–887. [[CrossRef](#)]
58. Gesquiere, L.R.; Ziegler, T.E.; Chen, P.A.; Epstein, K.A.; Alberts, S.C.; Altmann, J. Measuring Fecal Testosterone in Females and Fecal Estrogens in Males: Comparison of RIA and LC/MS/MS Methods for Wild Baboons (*Papio cynocephalus*). *Gen. Comp. Endocrinol.* **2014**, *204*, 141–149. [[CrossRef](#)]
59. Touma, C.; Palme, R. Measuring Fecal Glucocorticoid Metabolites in Mammals and Birds: The Importance of Validation. *Ann. N. Y. Acad. Sci.* **2005**, *1046*, 54–74. [[CrossRef](#)]
60. Laver, P.N.; Ganswindt, A.; Ganswindt, S.B.; Alexander, K.A. Non-Invasive Monitoring of Glucocorticoid Metabolites in Banded Mongooses (*Mungos mungo*) in Response to Physiological and Biological Challenges. *Gen. Comp. Endocrinol.* **2012**, *179*, 178–183. [[CrossRef](#)]
61. Shutt, K.; Setchell, J.M.; Heistermann, M. Non-Invasive Monitoring of Physiological Stress in the Western Lowland Gorilla (*Gorilla gorilla gorilla*): Validation of a Fecal Glucocorticoid Assay and Methods for Practical Application in the Field. *Gen. Comp. Endocrinol.* **2012**, *179*, 167–177. [[CrossRef](#)]
62. Ludwig, C.; Wachter, B.; Silinski-Mehr, S.; Ganswindt, A.; Bertschinger, H.; Hofer, H.; Dehnhard, M. Characterisation and Validation of an Enzyme-Immunoassay for the Non-Invasive Assessment of Faecal Glucocorticoid Metabolites in Cheetahs (*Acinonyx jubatus*). *Gen. Comp. Endocrinol.* **2013**, *180*, 15–23. [[CrossRef](#)] [[PubMed](#)]
63. Rangel-Negrin, A.; Coyohua-Fuentes, A.; Chavira, R.; Canales-Espinosa, D.; Dias, P. Primates Living Outside Protected Habitats Are More Stressed: The Case of Black Howler Monkeys in the Yucatan Peninsula. *PLoS ONE* **2014**, *9*. [[CrossRef](#)]
64. Narayan, E. Physiological Stress Levels in Wild Koala Sub-Populations Facing Anthropogenic Induced Environmental Trauma and Disease. *Sci. Rep.* **2019**, *9*, 6031. [[CrossRef](#)]
65. Ramahlo, M.; Chimimba, C.; Pirk, C.; Ganswindt, A. Non-Invasive Monitoring of Adrenocortical Activity in Free-Ranging Namaqua Rock Mice *Micaelamys namaquensis* from South Africa in Response to Anthropogenic Land Use and Season. *Wildl. Biol.* **2019**, *2019*, 1–6. [[CrossRef](#)]
66. Navarro-Castilla, Á.; Garrido, M.; Hawlena, H.; Barja, I. Non-Invasive Monitoring of Adrenocortical Activity in Three Sympatric Desert Gerbil Species. *Animals* **2021**, *11*, 75. [[CrossRef](#)] [[PubMed](#)]
67. Serres-Corral, P.; Fernandez-Bellon, H.; Padilla-Sole, P.; Carbajal, A.; Lopez-Bejar, M. Evaluation of Fecal Glucocorticoid Metabolite Levels in Response to a Change in Social and Handling Conditions in African Lions (*Panthera leo bleyenberghi*). *Animals* **2021**, *11*, 1877. [[CrossRef](#)] [[PubMed](#)]
68. Baltic, M.; Jenni-Eiermann, S.; Arlettaz, R.; Palme, R. A Noninvasive Technique to Evaluate Human-Generated Stress in the Black Grouse. *Ann. N. Y. Acad. Sci.* **2005**, *1046*, 81–95. [[CrossRef](#)]

69. Gladbach, A.; Gladbach, D.J.; Koch, M.; Kuchar, A.; Möstl, E.; Quillfeldt, P. Can Faecal Glucocorticoid Metabolites Be Used to Monitor Body Condition in Wild Upland Geese *Chloephaga picta leucoptera*? *Behav. Ecol. Sociobiol.* **2011**, *65*, 1491–1498. [[CrossRef](#)]
70. Ozella, L.; Anfossi, L.; Di Nardo, F.; Pessani, D. Non-Invasive Monitoring of Adrenocortical Activity in Captive African Penguin (*Spheniscus demersus*) by Measuring Faecal Glucocorticoid Metabolites. *Gen. Comp. Endocrinol.* **2015**, *224*, 104–112. [[CrossRef](#)]
71. Ganswindt, S.B.; Myburgh, J.G.; Cameron, E.Z.; Ganswindt, A. Non-Invasive Assessment of Adrenocortical Function in Captive Nile Crocodiles (*Crocodylus niloticus*). *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **2014**, *177*, 11–17. [[CrossRef](#)]
72. Bautista, L.M.; Silván, G.; Cáceres, S.; Martínez-Fernández, L.; Bravo, C.; Illera, J.C.; Alonso, J.C.; Blanco, G. Faecal Sexual Steroids in Sex Typing and Endocrine Status of Great Bustards. *Eur. J. Wildl. Res.* **2013**, *59*, 815–822. [[CrossRef](#)]
73. Pribbenow, S.; Wachter, B.; Ludwig, C.; Weigold, A.; Dehnhard, M. Validation of an Enzyme-Immunoassay for the Non-Invasive Monitoring of Faecal Testosterone Metabolites in Male Cheetahs (*Acinonyx jubatus*). *Gen. Comp. Endocrinol.* **2016**, *228*, 40–47. [[CrossRef](#)] [[PubMed](#)]
74. Khonmee, J.; Rojanasthien, S.; Thitaram, C.; Sumretprasong, J.; Aunsusin, A.; Chaisongkram, C.; Songsasen, N. Non-Invasive Endocrine Monitoring Indicates Seasonal Variations in Gonadal Hormone Metabolites in Dholes (*Cuon alpinus*). *Conserv. Physiol.* **2017**, *5*, cox001. [[CrossRef](#)]
75. Lemos, L.S.; Olsen, A.; Smith, A.; Chandler, T.E.; Larson, S.; Hunt, K.; Torres, L.G. Assessment of Fecal Steroid and Thyroid Hormone Metabolites in Eastern North Pacific Gray Whales. *Conserv. Physiol.* **2020**, *8*, coaa110. [[CrossRef](#)] [[PubMed](#)]
76. Paris, M.C.J.; White, A.; Reiss, A.; West, M.; Schwarzenberger, F. Faecal Progesterone Metabolites and Behavioural Observations for the Non-Invasive Assessment of Oestrous Cycles in the Common Wombat (*Vombatus ursinus*) and the Southern Hairy-Nosed Wombat (*Lasiorchinus latifrons*). *Anim. Reprod. Sci.* **2002**, *72*, 245–257. [[CrossRef](#)]
77. Van der Goot, A.; Dalerum, F.; Ganswindt, A.; Martin, G.; Millar, R.; Paris, M. Faecal Progestagen Profiles in Wild Southern White Rhinoceros (*Ceratotherium simum simum*). *Afr. Zool.* **2013**, *48*, 143–151. [[CrossRef](#)]
78. Mayor, P.; Guimaraes, D.A.; da Silva, J.; Jori, F.; Lopez-Bejar, M. Reproductive Monitoring of Collared Peccary Females (*Pecari tajacu*) by Analysis of Fecal Progesterone Metabolites. *Theriogenology* **2019**, *134*, 11–17. [[CrossRef](#)]
79. Mondol, S.; Booth, R.K.; Wasser, S.K. Fecal Stress, Nutrition and Reproductive Hormones for Monitoring Environmental Impacts on Tigers (*Panthera tigris*). *Conserv. Physiol.* **2020**, *8*, coz091. [[CrossRef](#)]
80. Seamster, V.; Waits, L.; Macko, S.; Shugart, H. Coyote (*Canis latrans*) Mammalian Prey Diet Shifts in Response to Seasonal Vegetation Change. *Isotopes Environ. Health Stud.* **2014**, *50*, 343–360. [[CrossRef](#)]
81. Monterroso, P.; Godinho, R.; Oliveira, T.; Ferreras, P.; Kelly, M.J.; Morin, D.J.; Waits, L.P.; Alves, P.C.; Mills, L.S. Feeding Ecological Knowledge: The Underutilised Power of Faecal DNA Approaches for Carnivore Diet Analysis. *Mammal Rev.* **2019**, *49*, 97–112. [[CrossRef](#)]
82. O'Rourke, D.R.; Mangan, M.T.; Mangan, K.E.; Bokulich, N.A.; MacManes, M.D.; Foster, J.T. Lord of the Diptera (and Moths and a Spider): Molecular Diet Analyses and Foraging Ecology of Indiana Bats in Illinois. *Front. Ecol. Evol.* **2021**, *9*, 12. [[CrossRef](#)]
83. Ando, H.; Mukai, H.; Komura, T.; Dewi, T.; Ando, M.; Isagi, Y. Methodological Trends and Perspectives of Animal Dietary Studies by Noninvasive Fecal DNA Metabarcoding. *Environ. DNA* **2020**, *2*, 391–406. [[CrossRef](#)]
84. Sage, M.; Fourel, I.; Cœurdassier, M.; Barrat, J.; Berny, P.; Giraudoux, P. Determination of Bromadiolone Residues in Fox Faeces by LC/ESI-MS in Relationship with Toxicological Data and Clinical Signs after Repeated Exposure. *Environ. Res.* **2010**, *110*, 664–674. [[CrossRef](#)] [[PubMed](#)]
85. Webster, A.B.; Callealta, F.J.; Ganswindt, A.; Bennett, N.C. A Non-Invasive Assessment of Essential Trace Element Utilization at Different Trophic Levels in African Wildlife. *J. Environ. Manage.* **2021**, *293*, 112820. [[CrossRef](#)] [[PubMed](#)]
86. Napoli, E.; Anile, S.; Arrabito, C.; Scornavacca, D.; Mazzamuto, M.V.; Gaglio, G.; Otranto, D.; Giannetto, S.; Brianti, E. Survey on Parasitic Infections in Wildcat (*Felis silvestris silvestris schreber, 1777*) by Scat Collection. *Parasitol. Res.* **2016**, *115*, 255–261. [[CrossRef](#)]
87. Stewart, L.; Tort, N.; Meakin, P.; Argudo, J.; Nzuma, R.; Reid, N.; Delahay, R.; Ashford, R.; Montgomery, W.; Grant, I. Development of a Novel Immunochromatographic Lateral Flow Assay Specific for *Mycobacterium bovis* Cells and Its Application in Combination with Immunomagnetic Separation to Test Badger Faeces. *BMC Vet. Res.* **2017**, *13*. [[CrossRef](#)]
88. Lynch, M.A.; Youngflesh, C.; Agha, N.H.; Ottinger, M.A.; Lynch, H.J. Tourism and Stress Hormone Measures in Gentoo Penguins on the Antarctic Peninsula. *Polar Biol.* **2019**, *42*, 1299–1306. [[CrossRef](#)]
89. Dib, L.V.; Palmer, J.P.S.; de Souza Carvalho Class, C.; Pinheiro, J.L.; Ramos, R.C.F.; dos Santos, C.R.; Fonseca, A.B.M.; Rodríguez-Castro, K.G.; Gonçalves, C.F.; Galetti, P.M.; et al. Non-Invasive Sampling in Itatiaia National Park, Brazil: Wild Mammal Parasite Detection. *BMC Vet. Res.* **2020**, *16*, 295. [[CrossRef](#)]
90. Hou, Z.; Peng, Z.; Ning, Y.; Liu, D.; Chai, H.; Jiang, G. An Initial Coprological Survey of Parasitic Fauna in the Wild Amur Leopard (*Panthera pardus orientalis*). *Integr. Zool.* **2020**, *15*, 375–384. [[CrossRef](#)]
91. Di Francesco, C.E.; Smoglica, C.; Paoletti, B.; Angelucci, S.; Innocenti, M.; Antonucci, A.; Di Domenico, G.; Marsilio, F. Detection of Selected Pathogens in Apennine Wolf (*Canis lupus italicus*) by a Non-Invasive GPS-Based Telemetry Sampling of Two Packs from Majella National Park, Italy. *Eur. J. Wildl. Res.* **2019**, *65*, 84. [[CrossRef](#)]
92. Afonso, E.; Tournant, P.; Foltête, J.-C.; Giraudoux, P.; Baurand, P.-E.; Roué, S.; Canella, V.; Vey, D.; Scheifler, R. Is the Lesser Horseshoe Bat (*Rhinolophus hipposideros*) Exposed to Causes That May Have Contributed to Its Decline? A Non-Invasive Approach. *Glob. Ecol. Conserv.* **2016**, *8*, 123–137. [[CrossRef](#)]
93. Maréchal, L.; Semples, S.; Majolo, B.; MacLarnon, A. Assessing the Effects of Tourist Provisioning on the Health of Wild Barbary Macaques in Morocco. *PLoS ONE* **2016**, *11*, e0155920. [[CrossRef](#)]

94. Albery, G.F.; Kenyon, F.; Morris, A.; Morris, S.; Nussey, D.H.; Pemberton, J.M. Seasonality of Helminth Infection in Wild Red Deer Varies between Individuals and between Parasite Taxa. *Parasitology* **2018**, *145*, 1410–1420. [[CrossRef](#)]
95. Jepsen, E.M.; Ganswindt, A.; Ngcamphalala, C.A.; Bourne, A.R.; Ridley, A.R.; McKechnie, A.E. Non-Invasive Monitoring of Physiological Stress in an Afrotropical Arid-Zone Passerine Bird, the Southern Pied Babbler. *Gen. Comp. Endocrinol.* **2019**, *276*, 60–68. [[CrossRef](#)]
96. Wells, K.; Smales, L.R.; Kalko, E.K.V.; Pfeiffer, M. Impact of Rain-Forest Logging on Helminth Assemblages in Small Mammals (*Muridae*, *Tupaïidae*) from Borneo. *J. Trop. Ecol.* **2007**, *23*, 35–43. [[CrossRef](#)]
97. Foreyt, W.J. *Veterinary Parasitology Reference Manual*; John Wiley & Sons: Hoboken, NJ, USA, 2013; ISBN 9781118682265.
98. Biswas, S.; Bhatt, S.; Paul, S.; Modi, S.; Ghosh, T.; Habib, B.; Nigam, P.; Talukdar, G.; Pandav, B.; Mondol, S. A Practical Faeces Collection Protocol for Multidisciplinary Research in Wildlife Science. *Curr. Sci.* **2019**, *116*, 537803. [[CrossRef](#)]
99. Marrero, P.; Fregel, R.; Cabrera, V.M.; Nogales, M. Extraction of High-Quality Host DNA from Feces and Regurgitated Seeds: A Useful Tool for Vertebrate Ecological Studies. *Biol. Res.* **2009**, *42*, 147–151. [[CrossRef](#)]
100. Ramón-Laca, A.; Soriano, L.; Gleeson, D.; Godoy, J.A. A Simple and Effective Method for Obtaining Mammal DNA from Faeces. *Wildl. Biol.* **2015**, *21*, wlb.00855. [[CrossRef](#)]
101. Santa, M.A.; Pastran, S.; Klein, C.; Ruckstuhl, K.; Massolo, A. Evaluation of an Automated Magnetic Bead-Based DNA Extraction and Real-Time PCR in Fecal Samples as a Pre-Screening Test for Detection of *Echinococcus multilocularis* and *Echinococcus canadensis* in Coyotes. *Parasitol. Res.* **2019**, *118*, 119–125. [[CrossRef](#)]
102. Fernández-Vizcaino, E.; Martínez-Carrasco, C.; Moratal, S.; Barroso, P.; Vicente, J. Detection of *Stephanurus dentatus* in Wild Boar Urine Using Different Parasitological Techniques. *Int. J. Parasitol. Parasites Wildl.* **2021**, *15*, 31–34. [[CrossRef](#)]
103. Schneider-Crease, I.; Griffin, R.H.; Gomery, M.A.; Dorny, P.; Noh, J.C.; Handali, S.; Chastain, H.M.; Wilkins, P.P.; Nunn, C.L.; Snyder-Mackler, N.; et al. Identifying Wildlife Reservoirs of Neglected Taeniid Tapeworms: Non-Invasive Diagnosis of Endemic *Taenia serialis* Infection in a Wild Primate Population. *PLoS Negl. Trop. Dis.* **2017**, *11*, e0005709. [[CrossRef](#)] [[PubMed](#)]
104. Seidlova, V.; Nemcova, M.; Pikula, J.; Bartonička, T.; Ghazaryan, A.; Heger, T.; Kokurewicz, T.; Orlov, O.L.; Patra, S.; Píacek, V.; et al. Urinary Shedding of Leptospire in Palearctic Bats. *Transbound. Emerg. Dis.* **2021**, *68*, 3089–3095. [[CrossRef](#)]
105. Leendertz, F.H.; Boesch, C.; Ellerbrok, H.; Rietschel, W.; Couacy-Hymann, E.; Pauli, G. Non-Invasive Testing Reveals a High Prevalence of Simian T-Lymphotropic Virus Type 1 Antibodies in Wild Adult Chimpanzees of the Tai National Park, Côte d'Ivoire. *J. Gen. Virol.* **2004**, *85*, 3305–3312. [[CrossRef](#)] [[PubMed](#)]
106. Baker, K.S.; Tachedjian, M.; Barr, J.; Marsh, G.A.; Todd, S.; Cramer, G.; Cramer, S.; Smith, I.; Holmes, C.E.G.; Suu-Ire, R.; et al. Achimota Pararubulavirus 3: A New Bat-Derived Paramyxovirus of the Genus Pararubulavirus. *Viruses* **2020**, *12*, 1236. [[CrossRef](#)] [[PubMed](#)]
107. Baker, K.S.; Todd, S.; Marsh, G.; Fernandez-Loras, A.; Suu-Ire, R.; Wood, J.L.N.; Wang, L.F.; Murcia, P.R.; Cunningham, A.A.Y. 2012 Co-Circulation of Diverse Paramyxoviruses in an Urban African Fruit Bat Population. *J. Gen. Virol.* **2012**, *93*, 850–856. [[CrossRef](#)]
108. Muller, M.N.; Wrangham, R.W. Dominance, Cortisol and Stress in Wild Chimpanzees (*Pan troglodytes schweinfurthii*). *Behav. Ecol. Sociobiol.* **2004**, *55*, 332–340. [[CrossRef](#)]
109. Narayan, E.; Molinia, F.; Christi, K.; Morley, C.; Cockrem, J.; Narayan, E.; Molinia, F.; Christi, K.; Morley, C.; Cockrem, J. Urinary Corticosterone Metabolite Responses to Capture, and Annual Patterns of Urinary Corticosterone in Wild and Captive Endangered Fijian Ground Frogs (*Platymantis vitiana*). *Aust. J. Zool.* **2010**, *58*, 189–197. [[CrossRef](#)]
110. Narayan, E.; Hero, J.-M. Urinary Corticosterone Responses and Haematological Stress Indicators in the Endangered Fijian Ground Frog (*Platymantis vitiana*) during Transportation and Captivity. *Aust. J. Zool.* **2011**, *59*, 79. [[CrossRef](#)]
111. Kindermann, C.; Narayan, E.J.; Hero, J.-M. Urinary Corticosterone Metabolites and Chytridiomycosis Disease Prevalence in a Free-Living Population of Male Stony Creek Frogs (*Litoria wilcoxii*). *Comp. Biochem. Physiol. A. Mol. Integr. Physiol.* **2012**, *162*, 171–176. [[CrossRef](#)]
112. Narayan, E.J. Non-Invasive Reproductive and Stress Endocrinology in Amphibian Conservation Physiology. *Conserv. Physiol.* **2013**, *1*, cot011. [[CrossRef](#)]
113. Narayan, E.J.; Gramapurohit, N.P. Sexual Dimorphism in Baseline Urinary Corticosterone Metabolites and Their Association with Body-Condition Indices in a Peri-Urban Population of the Common Asian Toad (*Duttaphrynus melanostictus*). *Comp. Biochem. Physiol. A. Mol. Integr. Physiol.* **2016**, *191*, 174–179. [[CrossRef](#)]
114. Sadoughi, B.; Girard-Buttoz, C.; Engelhardt, A.; Heistermann, M.; Ostner, J. Non-Invasive Assessment of Metabolic Responses to Food Restriction Using Urinary Triiodothyronine and Cortisol Measurement in Macaques. *Gen. Comp. Endocrinol.* **2021**, *306*. [[CrossRef](#)]
115. Kinoshita, K.; Miyazaki, M.; Morita, H.; Vassileva, M.; Tang, C.; Li, D.; Ishikawa, O.; Kusunoki, H.; Tsenkova, R. Spectral Pattern of Urinary Water as a Biomarker of Estrus in the Giant Panda. *Sci. Rep.* **2012**, *2*, 856. [[CrossRef](#)]
116. Mouri, K.; Shimizu, K. Stability of Chimpanzee (*Pan troglodytes*) Urinary Reproductive Hormones during Long-Term Preservation on Filter Paper. *Primates* **2021**, *62*, 289–296. [[CrossRef](#)]
117. Moscovice, L.R.; Ziegler, T.E. Peripheral Oxytocin in Female Baboons Relates to Estrous State and Maintenance of Sexual Consortships. *Horm. Behav.* **2012**, *62*, 592–597. [[CrossRef](#)]
118. Heistermann, M.; Higham, J.P. Urinary Neopterin, a Non-Invasive Marker of Mammalian Cellular Immune Activation, Is Highly Stable under Field Conditions. *Sci. Rep.* **2015**, *5*, 16308. [[CrossRef](#)]
119. Behringer, V.; Stevens, J.; Leendertz, F.; Hohmann, G.; Deschner, T. Validation of a Method for the Assessment of Urinary Neopterin Levels to Monitor Health Status in Non-Human-Primate Species. *Front. Physiol.* **2017**, *8*, 51. [[CrossRef](#)]

120. Löhrich, T.; Behringer, V.; Wittig, R.M.; Deschner, T.; Leendertz, F.H. The Use of Neopterin as a Noninvasive Marker in Monitoring Diseases in Wild Chimpanzees. *EcoHealth* **2018**, *15*, 792–803. [[CrossRef](#)]
121. Wu, D.F.; Behringer, V.; Wittig, R.M.; Leendertz, F.H.; Deschner, T. Urinary Neopterin Levels Increase and Predict Survival during a Respiratory Outbreak in Wild Chimpanzees (Taï National Park, Côte d'Ivoire). *Sci. Rep.* **2018**, *8*, 13346. [[CrossRef](#)]
122. Löhrich, T. Health Monitoring in Great Apes: The Use of Neopterin as a Non-Invasive Marker in Monitoring Diseases in Wild Chimpanzees (*Pan troglodytes verus*). Ph.D. Thesis, Freien Universität Berlin, Berlin, Germany, 2020. [[CrossRef](#)]
123. Sacco, A.J.; Mayhew, J.A.; Watsa, M.; Erkenswick, G.; Binder, A.K. Detection of Neopterin in the Urine of Captive and Wild Platyrrhines. *BMC Zool.* **2020**, *5*, 2. [[CrossRef](#)]
124. Emery Thompson, M.; Knott, C.D. Urinary C-Peptide of Insulin as a Non-Invasive Marker of Energy Balance in Wild Orangutans. *Horm. Behav.* **2008**, *53*, 526–535. [[CrossRef](#)]
125. Peterson, R.O.; Hoy, S.R.; Vucetich, L.M.; Vucetich, J.A.; Kraus, V.B.; Huebner, J.L. Urinary Biomarkers to Non-Invasively Evaluate Health in Wild Moose (*Alces alces*). *Osteoarthr. Cartil.* **2018**, *26*, S196–S197. [[CrossRef](#)]
126. Rosenblatt, E.; DeBow, J.; Blouin, J.; Donovan, T.; Murdoch, J.; Creel, S.; Rogers, W.; Gieder, K.; Fortin, N.; Alexander, C. Juvenile Moose Stress and Nutrition Dynamics Related to Winter Ticks, Landscape Characteristics, Climate-Mediated Factors and Survival. *Conserv. Physiol.* **2021**, *9*. [[CrossRef](#)]
127. Millspaugh, J.J.; Washburn, B.E.; Milanick, M.A.; Beringer, J.; Hansen, L.P.; Meyer, T.M. Non-Invasive Techniques for Stress Assessment in White-Tailed Deer. *Wildl. Soc. Bull. 1973–2006* **2002**, *30*, 899–907.
128. Menargues, A.; Urios, V.; Mauri, M. Welfare Assessment of Captive Asian Elephants (*Elephas maximus*) and Indian Rhinoceros (*Rhinoceros unicornis*) Using Salivary Cortisol Measurement. *Anim. Welf.* **2008**, *17*, 305–312.
129. Staley, M.; Conners, M.G.; Hall, K.; Miller, L.J. Linking Stress and Immunity: Immunoglobulin A as a Non-Invasive Physiological Biomarker in Animal Welfare Studies. *Horm. Behav.* **2018**, *102*, 55–68. [[CrossRef](#)]
130. Emami-Khoyi, A.; Agnew, T.W.; Adair, M.G.; Murphy, E.C.; Benmazouz, I.; Monsanto, D.M.; Parbhu, S.P.; Main, D.C.; Le Roux, R.; Golla, T.R.; et al. A New Non-Invasive Method for Collecting DNA From Small Mammals in the Field, and Its Application in Simultaneous Vector and Disease Monitoring in Brushtail Possums. *Front. Environ. Sci.* **2021**, *9*, 318. [[CrossRef](#)]
131. Filek, K.; Trotta, A.; Gračan, R.; Di Bello, A.; Corrente, M.; Bosak, S. Characterization of Oral and Cloacal Microbial Communities of Wild and Rehabilitated Loggerhead Sea Turtles (*Caretta caretta*). *Anim. Microbiome* **2021**, *3*, 59. [[CrossRef](#)]
132. Menegotto, A. *Potential Role of Telemetry and Non-Invasive Sampling in Wild Boar in the Surveillance and the Control of Foot-and-Mouth Disease in Bulgaria*; Università di Padova: Padua, Italy, 2013.
133. Mouchantat, S.; Haas, B.; Böhle, W.; Globig, A.; Lange, E.; Mettenleiter, T.C.; Depner, K. Proof of Principle: Non-Invasive Sampling for Early Detection of Foot-and-Mouth Disease Virus Infection in Wild Boar using a Rope-in-a-Bait Sampling Technique. *Vet. Microbiol.* **2014**, *172*, 329–333. [[CrossRef](#)]
134. Mouchantat, S.; Globig, A.; Böhle, W.; Petrov, A.; Strebelow, H.-G.; Mettenleiter, T.C.; Depner, K. Novel Rope-Based Sampling of Classical Swine Fever Shedding in a Group of Wild Boar Showing Low Contagiosity upon Experimental Infection with a Classical Swine Fever Field Strain of Genotype 2.3. *Vet. Microbiol.* **2014**, *170*, 425–429. [[CrossRef](#)]
135. Männistö, H.E. Collection of Oral Fluid Samples from Wild Boar in the Field Conditions to Detect African Swine Fever Virus (ASFV). Ph.D. Thesis, Estonian University of Life Sciences, Tartu, Estonia, 2018.
136. Common, S.M.; Yun, Y.; Silva-Fletcher, A.; Thitaram, C.; Janyamethakul, T.; Khammesri, S.; Molenaar, F.M. Developing a Non-Invasive Method of Detecting Elephant Endotheliotropic Herpesvirus Infections Using Faecal Samples. *Vet. Rec.* **2021**, *190*, e833. [[CrossRef](#)]
137. Evans, T.S.; Barry, P.A.; Gilardi, K.V.; Goldstein, T.; Deere, J.D.; Fike, J.; Yee, J.; Ssebide, B.J.; Karmacharya, D.; Cranfield, M.R.; et al. Optimization of a Novel Non-Invasive Oral Sampling Technique for Zoonotic Pathogen Surveillance in Nonhuman Primates. *PLoS Negl. Trop. Dis.* **2015**, *9*, e0003813. [[CrossRef](#)]
138. Smiley Evans, T.; Gilardi, K.V.K.; Barry, P.A.; Ssebide, B.J.; Kinani, J.F.; Nizeyimana, F.; Noheri, J.B.; Byarugaba, D.K.; Mudakikwa, A.; Cranfield, M.R.; et al. Detection of Viruses Using Discarded Plants from Wild Mountain Gorillas and Golden Monkeys. *Am. J. Primatol.* **2016**, *78*, 1222–1234. [[CrossRef](#)]
139. Seeber, P.A.; Soilemetzidou, S.E.; East, M.L.; Walzer, C.; Greenwood, A.D. Equine Behavioral Enrichment Toys as Tools for Non-Invasive Recovery of Viral and Host DNA. *Zoo Biol.* **2017**, *36*, 341–344. [[CrossRef](#)]
140. Barasona, J.A.; Barroso-Arévalo, S.; Rivera, B.; Gortázar, C.; Sánchez-Vizcaíno, J.M. Detection of Antibodies against *Mycobacterium bovis* in Oral Fluid from Eurasian Wild Boar. *Pathogens* **2020**, *9*, 242. [[CrossRef](#)]
141. Thompson, L.A.; Spoon, T.R.; Goertz, C.E.C.; Hobbs, R.C.; Romano, T.A. Blow Collection as a Non-Invasive Method for Measuring Cortisol in the Beluga (*Delphinapterus leucas*). *PLoS ONE* **2014**, *9*, e114062. [[CrossRef](#)]
142. Groch, K.R.; Blazquez, D.N.H.; Marcondes, M.C.C.; Santos, J.; Colosio, A.; Díaz Delgado, J.; Catão-Dias, J.L. Cetacean Morbillivirus in Humpback Whales' Exhaled Breath. *Transbound. Emerg. Dis.* **2021**, *68*, 1736–1743. [[CrossRef](#)]
143. Reckendorf, A.; Schmicke, M.; Bunskoek, P.; Hansen, K.; Thybo, M.; Strube, C.; Siebert, U. Is Harbor Porpoise (*Phocoena phocoena*) Exhaled Breath Sampling Suitable for Hormonal Assessments? *Animals* **2021**, *11*, 907. [[CrossRef](#)]
144. Acevedo-Whitehouse, K.; Rocha-Gosselin, A.; Gendron, D. A Novel Non-Invasive Tool for Disease Surveillance of Free-Ranging Whales and Its Relevance to Conservation Programs. *Anim. Conserv.* **2010**, *13*, 217–225. [[CrossRef](#)]
145. Lima, N.; Rogers, T.; Acevedo-Whitehouse, K.; Brown, M.V. Temporal Stability and Species Specificity in Bacteria Associated with the Bottlenose Dolphins Respiratory System. *Environ. Microbiol. Rep.* **2012**, *4*, 89–96. [[CrossRef](#)]

146. Cumeras, R.; Cheung, W.H.K.; Gulland, F.; Goley, D.; Davis, C.E. Chemical Analysis of Whale Breath Volatiles: A Case Study for Non-Invasive Field Health Diagnostics of Marine Mammals. *Metabolites* **2014**, *4*, 790–806. [[CrossRef](#)] [[PubMed](#)]
147. Peled, N.; Ionescu, R.; Nol, P.; Barash, O.; McCollum, M.; VerCauteren, K.; Koslow, M.; Stahl, R.; Rhyan, J.; Haick, H. Detection of Volatile Organic Compounds in Cattle Naturally Infected with *Mycobacterium bovis*. *Sens. Actuators B Chem.* **2012**, *171–172*, 588–594. [[CrossRef](#)]
148. Gundlach, N.H.; Piechotta, M.; Siebert, U. Is Lachrymal Fluid a Potential Method for Cortisol Measurement in Wild Harbor Seals? A Pilot Study. *Int. J. Vet. Med. Res. Rep.* **2016**, *1–10*. [[CrossRef](#)]
149. Woodhams, D.; Brandt, H.; Baumgartner, S.; Kielgast, J.; Kupfer, E.; Tobler, U.; Davis, L.; Schmidt, B.; Bel, C.; Hodel, S.; et al. Interacting Symbionts and Immunity in the Amphibian Skin Mucosome Predict Disease Risk and Probiotic Effectiveness. *PLoS ONE* **2014**, *9*, e96375. [[CrossRef](#)]
150. Zhu, W.; Fan, L.; Soto-Azat, C.; Yan, S.; Gao, X.; Liu, X.; Wang, S.; Liu, C.; Yang, X.; Li, Y. Filling a Gap in the Distribution of *Batrachochytrium dendrobatidis*: Evidence in Amphibians from Northern China. *Dis. Aquat. Organ.* **2016**, *118*, 259–265. [[CrossRef](#)]
151. DiRenzo, G.V.; Campbell Grant, E.H.; Longo, A.V.; Che-Castaldo, C.; Zamudio, K.R.; Lips, K.R. Imperfect Pathogen Detection from Non-Invasive Skin Swabs Biases Disease Inference. *Methods Ecol. Evol.* **2018**, *9*, 380–389. [[CrossRef](#)]
152. Santymire, R.M.; Manjerovic, M.B.; Sacerdote-Velat, A. A Novel Method for the Measurement of Glucocorticoids in Dermal Secretions of Amphibians. *Conserv. Physiol.* **2018**, *6*, coy008. [[CrossRef](#)]
153. Smith, H.; Pasmans, F.; Dhaenens, M.; Deforce, D.; Bonte, D.; Verheyen, K.; Lens, L.; Martel, A. Skin Mucosome Activity as an Indicator of *Batrachochytrium salamandrivorans* Susceptibility in Salamanders. *PLoS ONE* **2018**, *13*. [[CrossRef](#)]
154. Ghosh, P.N.; Verster, R.; Sewell, T.R.; O'Hanlon, S.J.; Brookes, L.M.; Rieux, A.; Garner, T.W.J.; Weldon, C.; Fisher, M.C. Discriminating Lineages of *Batrachochytrium dendrobatidis* Using Quantitative PCR. *Mol. Ecol. Resour.* **2021**, *21*, 1452–1459. [[CrossRef](#)]
155. Rothstein, A.; Byrne, A.; Knapp, R.; Briggs, C.; Voyles, J.; Richards-Zawacki, C.; Rosenblum, E. Divergent Regional Evolutionary Histories of a Devastating Global Amphibian Pathogen. *Proc. R. Soc. B-Biol. Sci.* **2021**, *288*, 20210782. [[CrossRef](#)]
156. Dzul-Caamal, R.; Olivares-Rubio, H.F.; Salazar-Coria, L.; Rocha-Gómez, M.A.; Vega-López, A. Multivariate Analysis of Biochemical Responses using Non-Invasive Methods to Evaluate the Health Status of the Endangered Blackfin Goodeid (*Girardinichthys viviparus*). *Ecol. Indic.* **2016**, *60*, 1118–1129. [[CrossRef](#)]
157. Fernández-Alacid, L.; Sanahuja, I.; Ordóñez-Grande, B.; Sánchez-Nuño, S.; Viscor, G.; Gisbert, E.; Herrera, M.; Ibarz, A. Skin Mucus Metabolites in Response to Physiological Challenges: A Valuable Non-Invasive Method to Study Teleost Marine Species. *Sci. Total Environ.* **2018**, *644*, 1323–1335. [[CrossRef](#)] [[PubMed](#)]
158. Bulloch, P.; Schur, S.; Muthumuni, D.; Xia, Z.; Johnson, W.; Chu, M.; Palace, V.; Su, G.; Letcher, R.; Tomy, G.T. F2-Isoprostanes in Fish Mucus: A New, Non-Invasive Method for Analyzing a Biomarker of Oxidative Stress. *Chemosphere* **2020**, *239*, 124797. [[CrossRef](#)] [[PubMed](#)]
159. Zuberi, A.; Ali, S.; Brown, C. A Non-Invasive Assay for Monitoring Stress Responses: A Comparison between Wild and Captive-Reared Rainbowfish (*Melanotania duboulayi*). *Aquaculture* **2011**, *321*, 267–272. [[CrossRef](#)]
160. Gabor, C.R.; Fisher, M.C.; Bosch, J. A Non-Invasive Stress Assay Shows That Tadpole Populations Infected with *Batrachochytrium dendrobatidis* Have Elevated Corticosterone Levels. *PLoS ONE* **2013**, *8*, e56054. [[CrossRef](#)] [[PubMed](#)]
161. Baugh, A.T.; Bastien, B.; Still, M.B.; Stowell, N. Validation of Water-Borne Steroid Hormones in a Tropical Frog (*Physalaemus pustulosus*). *Gen. Comp. Endocrinol.* **2018**, *261*, 67–80. [[CrossRef](#)]
162. Tornabene, B.; Hossack, B.; Crespi, E.; Breuner, C. Evaluating Corticosterone as a Biomarker for Amphibians Exposed to Increased Salinity and Ambient Corticosterone. *Conserv. Physiol.* **2021**, *9*, 1–15. [[CrossRef](#)]
163. Sobańska, M.A. Wild Boar Hair (*Sus scrofa*) as a Non-Invasive Indicator of Mercury Pollution. *Sci. Total Environ.* **2005**, *339*, 81–88. [[CrossRef](#)]
164. Bechshoft, T.; Derocher, A.E.; Richardson, E.; Mislan, P.; Lunn, N.J.; Sonne, C.; Dietz, R.; Janz, D.M.; St. Louis, V.L. Mercury and Cortisol in Western Hudson Bay Polar Bear Hair. *Ecotoxicology* **2015**, *24*, 1315–1321. [[CrossRef](#)]
165. Hernout, B.V.; McClean, C.J.; Arnold, K.E.; Walls, M.; Baxter, M.; Boxall, A.B.A. Fur: A Non-Invasive Approach to Monitor Metal Exposure in Bats. *Chemosphere* **2016**, *147*, 376–381. [[CrossRef](#)]
166. Treu, G.; Krone, O.; Unnsteinsdóttir, E.R.; Greenwood, A.D.; Czirják, G.Á. Correlations between Hair and Tissue Mercury Concentrations in Icelandic Arctic Foxes (*Vulpes lagopus*). *Sci. Total Environ.* **2018**, *619–620*, 1589–1598. [[CrossRef](#)] [[PubMed](#)]
167. Montillo, M.; Caslini, C.; Peric, T.; Prandi, A.; Netto, P.; Tubaro, F.; Pedrotti, L.; Bianchi, A.; Mattiello, S. Analysis of 19 Minerals and Cortisol in Red Deer Hair in Two Different Areas of the Stelvio National Park: A Preliminary Study. *Animals* **2019**, *9*, 492. [[CrossRef](#)] [[PubMed](#)]
168. Stoklasová, L.; Váhala, J.; Hejčmanová, P. Minerals in the Blood, Hair, and Faeces of the Critically Endangered Western Derby Eland under Human Care in Two Wildlife Reserves in Senegal. *Biol. Trace Elem. Res.* **2020**, *195*, 105–109. [[CrossRef](#)] [[PubMed](#)]
169. Richards, N.; Cook, G.; Simpson, V.; Hall, S.; Harrison, N.; Scott, K. Qualitative Detection of the NSAIDs Diclofenac and Ibuprofen in the Hair of Eurasian Otters (*Lutra lutra*) Occupying UK Waterways with GC-MS. *Eur. J. Wildl. Res.* **2011**, *57*, 1107–1114. [[CrossRef](#)]
170. Keckeis, K.; Lepschy, M.; Schöpfer, H.; Moser, L.; Troxler, J.; Palme, R. Hair Cortisol: A Parameter of Chronic Stress? Insights from a Radiometabolism Study in Guinea Pigs. *J. Comp. Physiol. B* **2012**, *182*, 985–996. [[CrossRef](#)]

171. Terwissen, C.V.; Mastromonaco, G.F.; Murray, D.L. Influence of Adrenocorticotrophin Hormone Challenge and External Factors (Age, Sex, and Body Region) on Hair Cortisol Concentration in Canada Lynx (*Lynx canadensis*). *Gen. Comp. Endocrinol.* **2013**, *194*, 162–167. [[CrossRef](#)]
172. Caslini, C.; Comin, A.; Peric, T.; Prandi, A.; Pedrotti, L.; Mattiello, S. Use of Hair Cortisol Analysis for Comparing Population Status in Wild Red Deer (*Cervus elaphus*) Living in Areas with Different Characteristics. *Eur. J. Wildl. Res.* **2016**, *62*, 713–723. [[CrossRef](#)]
173. Charalambous, R.; Narayan, E. Cortisol Measurement in Koala (*Phascolarctos cinereus*) Fur. *JOVE-J. Vis. Exp.* **2019**. [[CrossRef](#)]
174. Crill, C.; Janz, D.M.; Kusch, J.M.; Santymire, R.M.; Heyer, G.P.; Shury, T.K.; Lane, J.E. Investigation of the Utility of Feces and Hair as Non-Invasive Measures of Glucocorticoids in Wild Black-Tailed Prairie Dogs (*Cynomys ludovicianus*). *Gen. Comp. Endocrinol.* **2019**, *275*, 15–24. [[CrossRef](#)]
175. Heimbürge, S.; Kanitz, E.; Otten, W. The Use of Hair Cortisol for the Assessment of Stress in Animals. *Gen. Comp. Endocrinol.* **2019**, *270*, 10–17. [[CrossRef](#)]
176. Potratz, E.J.; Brown, J.S.; Gallo, T.; Anchor, C.; Santymire, R.M. Effects of Demography and Urbanization on Stress and Body Condition in Urban White-Tailed Deer. *Urban Ecosyst.* **2019**, *22*, 807–816. [[CrossRef](#)]
177. Cattet, M.; Stenhouse, G.B.; Boulanger, J.; Janz, D.M.; Kapronczai, L.; Swenson, J.E.; Zedrosser, A. Can Concentrations of Steroid Hormones in Brown Bear Hair Reveal Age Class? *Conserv. Physiol.* **2018**, *6*, coy001. [[CrossRef](#)] [[PubMed](#)]
178. Everest, D.; Tolhurst-Cherriman, D.; Davies, H.; Dastjerdi, A.; Ashton, A.; Blackett, T.; Meredith, A.; Milne, E.; Mill, A.; Shuttleworth, C. Assessing a Potential Non-Invasive Method for Viral Diagnostic Purposes in European Squirrels. *HYSTRIX—Ital. J. Mammal.* **2019**, *30*, 44–50. [[CrossRef](#)]
179. Muñoz-Madrid, R.; Belinchón-Lorenzo, S.; Iniesta, V.; Fernández-Cotrina, J.; Parejo, J.C.; Serrano, F.J.; Monroy, I.; Baz, V.; Gómez-Luque, A.; Gómez-Nieto, L.C. First Detection of *Leishmania infantum* Kinetoplast DNA in Hair of Wild Mammals: Application of QPCR Method to Determine Potential Parasite Reservoirs. *Acta Trop.* **2013**, *128*, 706–709. [[CrossRef](#)]
180. Ortega, M.V.; Moreno, I.; Domínguez, M.; de la Cruz, M.L.; Martín, A.B.; Rodríguez-Bertos, A.; López, R.; Navarro, A.; González, S.; Mazariegos, M.; et al. Application of a Specific Quantitative Real-Time PCR (QPCR) to Identify *Leishmania infantum* DNA in Spleen, Skin and Hair Samples of Wild Leporidae. *Vet. Parasitol.* **2017**, *243*, 92–99. [[CrossRef](#)]
181. Hopkins, J.B.; Whittington, J.; Clevenger, A.P.; Sawaya, M.A.; St. Clair, C.C. Stable Isotopes Reveal Rail-Associated Behavior in a Threatened Carnivore. *Isotopes Environ. Health Stud.* **2014**, *50*, 322–331. [[CrossRef](#)]
182. Rysava, K.; McGill, R.; Matthiopoulos, J.; Hopcraft, J. Re-Constructing Nutritional History of Serengeti Wildebeest from Stable Isotopes in Tail Hair: Seasonal Starvation Patterns in an Obligate Grazer. *Rapid Commun. Mass Spectrom.* **2016**, *30*, 1461–1468. [[CrossRef](#)]
183. Espín, S.; Martínez-López, E.; Gómez-Ramírez, P.; María-Mojica, P.; García-Fernández, A.J. Razorbills (*Alca torda*) as Bioindicators of Mercury Pollution in the Southwestern Mediterranean. *Mar. Pollut. Bull.* **2012**, *64*, 2461–2470. [[CrossRef](#)]
184. Costa, R.A.; Eeva, T.; Eira, C.; Vaqueiro, J.; Vingada, J.V. Assessing Heavy Metal Pollution Using Great Tits (*Parus major*): Feathers and Excrements from Nestlings and Adults. *Environ. Monit. Assess.* **2013**, *185*, 5339–5344. [[CrossRef](#)]
185. Di Marzio, A.; Gómez-Ramírez, P.; Barbar, F.; Lambertucci, S.A.; García-Fernández, A.J.; Martínez-López, E. Mercury in the Feathers of Bird Scavengers from Two Areas of Patagonia (Argentina) under the Influence of Different Anthropogenic Activities: A Preliminary Study. *Environ. Sci. Pollut. Res.* **2018**, *25*, 13906–13915. [[CrossRef](#)]
186. Liu, Q.; Chen, Y.; Maltby, L.; Zheng, Y.; Ma, Q. Trace Elements Exposure of Endangered Crested Ibis (*Nipponia nippon*) under in Situ and Ex Situ Conservations. *Environ. Pollut.* **2019**, *253*, 800–810. [[CrossRef](#)] [[PubMed](#)]
187. Sanchez, M.; Flavier, M.; Paller, V.; Rebancos, C.; Alcantara, A.; Sanchez, R.; Pelegrina, D. Presence of Heavy Metals in Feathers of Avian Species and in Soils of Barangay Ipil-Calancan Bay, Sta. Cruz, Marinduque Island, Philippines. *PERTANIKA J. Trop. Agric. Sci.* **2019**, *42*, 1009–1021.
188. Bryan, C.E.; Christopher, S.J.; Balmer, B.C.; Wells, R.S. Establishing Baseline Levels of Trace Elements in Blood and Skin of Bottlenose Dolphins in Sarasota Bay, Florida: Implications for Non-Invasive Monitoring. *Sci. Total Environ.* **2007**, *388*, 325–342. [[CrossRef](#)] [[PubMed](#)]
189. Ebmer, D.; Navarrete, M.J.; Muñoz, P.; Flores, L.M.; Gärtner, U.; Taubert, A.; Hermosilla, C. Antartophthirus Microchir Infestation in Synanthropic South American Sea Lion (*Otaria flavescens*) Males Diagnosed by a Novel Non-Invasive Method. *Parasitol. Res.* **2019**, *118*, 1353–1361. [[CrossRef](#)]
190. Fraser, T.A.; Martin, A.; Polkinghorne, A.; Carver, S. Comparative Diagnostics Reveals PCR Assays on Skin Scrapings Is the Most Reliable Method to Detect Sarcoptes Scabiei Infestations. *Vet. Parasitol.* **2018**, *251*, 119–124. [[CrossRef](#)]
191. Schneider, L.; Eggins, S.; Maher, W.; Vogt, R.C.; Krikowa, F.; Kinsley, L.; Eggins, S.M.; Da Silveira, R. An Evaluation of the Use of Reptile Dermal Scutes as a Non-Invasive Method to Monitor Mercury Concentrations in the Environment. *Chemosphere* **2015**, *119*, 163–170. [[CrossRef](#)]
192. Burns, T.J.; McCafferty, D.J.; Kennedy, M.W. Core and Body Surface Temperatures of Nesting Leatherback Turtles (*Dermochelys coriacea*). *J. Therm. Biol.* **2015**, *51*, 15–22. [[CrossRef](#)]
193. Horton, T.W.; Hauser, N.; Cassel, S.; Klaus, K.F.; Fettermann, T.; Key, N. Doctor Drone: Non-Invasive Measurement of Humpback Whale Vital Signs Using Unoccupied Aerial System Infrared Thermography. *Front. Mar. Sci.* **2019**, *6*, 466. [[CrossRef](#)]
194. Valle, C.; Grijalva, C.; Calle, P.; Munoz-Perez, J.; Quezada, G.; Vera, C.; Lewbart, G. Methods of Body Temperature Assessment in *Conolophus subcristatus*, *Conolophus pallidus* (Galapagos Land Iguanas), and *Amblyrhynchus cristatus* X C. *Subcristatus* Hybrid. *PEERJ* **2019**, *7*. [[CrossRef](#)]

195. Jerem, P.; Jenni-Eiermann, S.; Herborn, K.; McKeegan, D.; McCafferty, D.J.; Nager, R.G. Eye Region Surface Temperature Reflects Both Energy Reserves and Circulating Glucocorticoids in a Wild Bird. *Sci. Rep.* **2018**, *8*, 1907. [[CrossRef](#)]
196. Mellish, J.; Nienaber, J.; Polasek, L.; Horning, M. Beneath the Surface: Profiling Blubber Depth in Pinnipeds with Infrared Imaging. *J. Therm. Biol.* **2013**, *38*, 10–13. [[CrossRef](#)] [[PubMed](#)]
197. Narayan, E.; Perakis, A.; Meikle, W. Using Thermal Imaging to Monitor Body Temperature of Koalas (*Phascolarctos cinereus*) in A Zoo Setting. *Animals* **2019**, *9*, 1094. [[CrossRef](#)] [[PubMed](#)]
198. Mccafferty, D.J. The Value of Infrared Thermography for Research on Mammals: Previous Applications and Future Directions. *Mammal Rev.* **2007**, *37*, 207–223. [[CrossRef](#)]
199. Arenas, A.J.; Gómez, F.; Salas, R.; Carrasco, P.; Borge, C.; Maldonado, A.; O'Brien, D.J.; Martínez-Moreno, F.J. An Evaluation of the Application of Infrared Thermal Imaging to the Tele-Diagnosis of Sarcoptic Mange in the Spanish Ibex (*Capra pyrenaica*). *Vet. Parasitol.* **2002**, *109*, 111–117. [[CrossRef](#)]
200. Dunbar, M.R.; MacCarthy, K.A. Use of Infrared Thermography to Detect Signs of Rabies Infection in Raccoons (*Procyon lotor*). *J. Zoo Wildl. Med.* **2006**, *37*, 518–523. [[CrossRef](#)]
201. Dunbar, M.R.; Johnson, S.R.; Rhyan, J.C.; McCollum, M. Use of Infrared Thermography to Detect Thermographic Changes in Mule Deer (*Odocoileus hemionus*) Experimentally Infected with Foot-and-Mouth Disease. *J. Zoo Wildl. Med.* **2009**, *40*, 296–301. [[CrossRef](#)]
202. Galaverni, M.; Palumbo, D.; Fabbri, E.; Caniglia, R.; Greco, C.; Randi, E. Monitoring Wolves (*Canis lupus*) by Non-Invasive Genetics and Camera Trapping: A Small-Scale Pilot Study. *Eur. J. Wildl. Res.* **2012**, *58*, 47–58. [[CrossRef](#)]
203. Murray, M.H.; Fidino, M.; Lehrer, E.W.; Simonis, J.L.; Magle, S.B. A Multi-State Occupancy Model to Non-Invasively Monitor Visible Signs of Wildlife Health with Camera Traps That Accounts for Image Quality. *J. Anim. Ecol.* **2021**, *90*, 1973–1984. [[CrossRef](#)]
204. Lacroux, C.; Guma, N.; Krief, S. Facial Dysplasia in Wild Forest Olive Baboons (*Papio anubis*) in Sebitoli, Kibale National Park, Uganda: Use of Camera Traps to Detect Health Defects. *J. Med. Primatol.* **2019**, *48*, 143–153. [[CrossRef](#)]
205. Hicks, L.L.; Schwab, N.A.; Homyack, J.A.; Jones, J.E.; Maxell, B.A.; Burkholder, B.O. A Statistical Approach to White-Nose Syndrome Surveillance Monitoring Using Acoustic Data. *PLoS ONE* **2020**, *15*, e0241052. [[CrossRef](#)]
206. Kvapil, P.; Kastelic, M.; Jež, N.; Sedlák, K.; Kašpárková, N.; Jelovšek, M.; Avšič-Županc, T.; Bártová, E.; Račnik, J. Detection of Antibodies against Tick-Borne Encephalitis Virus in Zoo Animals Using Non-Invasive Blood Sampling with Medicinal Leeches (*Hirudo medicinalis*). *Pathogens* **2021**, *10*, 952. [[CrossRef](#)] [[PubMed](#)]
207. Alfano, N.; Dayaram, A.; Axtner, J.; Tsangaras, K.; Kampmann, M.-L.; Mohamed, A.; Wong, S.T.; Gilbert, M.T.P.; Wilting, A.; Greenwood, A.D. Non-Invasive Surveys of Mammalian Viruses Using Environmental DNA. *Methods Ecol. Evol.* **2021**, *12*, 1941–1952. [[CrossRef](#)]
208. Thomsen, R.; Voigt, C.C. Non-Invasive Blood Sampling from Primates using Laboratory-Bred Blood-Sucking Bugs (*Dipetalogaster maximus*; *Reduviidae*, *Heteroptera*). *Primates* **2006**, *47*, 397–400. [[CrossRef](#)] [[PubMed](#)]
209. Voigt, C.C.; Peschel, U.; Wibbelt, G.; Frölich, K. An Alternative, Less Invasive Blood Sample Collection Technique for Serologic Studies Utilizing Triatomine Bugs (*Heteroptera*; *Insecta*). *J. Wildl. Dis.* **2006**, *42*, 466–469. [[CrossRef](#)]
210. Hoffmann, C.; Stockhausen, M.; Merkel, K.; Calvignac-Spencer, S.; Leendertz, F.H. Assessing the Feasibility of Fly Based Surveillance of Wildlife Infectious Diseases. *Sci. Rep.* **2016**, *6*, 37952. [[CrossRef](#)]
211. Zechmeisterová, K.; de Bellocq, J.G.; Široký, P. Diversity of Karyolysus and Schellackia from the Iberian Lizard *Lacerta schreiberi* with Sequence Data from Engorged Ticks. *Parasitology* **2019**, *146*, 1690–1698. [[CrossRef](#)]
212. Martens, J.M.; Stokes, H.S.; Berg, M.L.; Walder, K.; Raidal, S.R.; Magrath, M.J.L.; Bennett, A.T.D. A Non-Invasive Method to Assess Environmental Contamination with Avian Pathogens: Beak and Feather Disease Virus (BFDV) Detection in Nest Boxes. *PeerJ* **2020**, *8*, e9211. [[CrossRef](#)]
213. De Oliveira, M.L.; Norris, D.; Ramírez, J.F.M.; de Peres, P.H.F.; Galetti, M.; Duarte, J.M.B. Dogs Can Detect Scat Samples More Efficiently than Humans: An Experiment in a Continuous Atlantic Forest Remnant. *Zool. Curitiba* **2012**, *29*, 183–186. [[CrossRef](#)]
214. Curry, E.; Skogen, M.; Roth, T. Evaluation of an Odour Detection Dog for Non-Invasive Pregnancy Diagnosis in Polar Bears (*Ursus maritimus*): Considerations for Training Sniffer Dogs for Biomedical Investigations in Wildlife Species. *J. Zoo Aquar. Res.* **2021**, *9*, 1–7. [[CrossRef](#)]
215. Lurz, P.W.W.; Krauze-Gryz, D.; Gryz, J.; Meredith, A.; Schilling, A.-K.; Thain, C.; Heller, E. Invisible Threats to Native Mammals—Mercury Levels in Three Eurasian Red Squirrel Populations. *Hystrix Ital. J. Mammal.* **2017**, *28*, 280–283. [[CrossRef](#)]
216. Romeo, C.; Piscitelli, A.P.; Santicchia, F.; Martinoli, A.; Ferrari, N.; Wauters, L.A. Invading Parasites: Spillover of an Alien Nematode Reduces Survival in a Native Species. *Biol. Invasions* **2021**, *23*, 3847–3857. [[CrossRef](#)]
217. Beard, D.; Stannard, H.J.; Old, J.M. Morphological Identification of Ticks and Molecular Detection of Tick-Borne Pathogens from Bare-Nosed Wombats (*Vombatus ursinus*). *Parasit. Vectors* **2021**, *14*, 60. [[CrossRef](#)] [[PubMed](#)]
218. Segura-Göthlin, S.; Fernández, A.; Arbelo, M.; Felipe-Jiménez, I.; Colom-Rivero, A.; Almunia, J.; Sierra, E. The Validation of a Non-Invasive Skin Sampling Device for Detecting Cetacean Poxvirus. *Animals* **2021**, *11*, 2814. [[CrossRef](#)]
219. Apprill, A.; Miller, C.A.; Moore, M.J.; Durban, J.W.; Fearnbach, H.; Barrett-Lennard, L.G. Extensive Core Microbiome in Drone-Captured Whale Blow Supports a Framework for Health Monitoring. *mSystems* **2017**, *2*, e00119-17. [[CrossRef](#)] [[PubMed](#)]
220. Lebarbenchon, C.; Poulson, R.; Shannon, K.; Slagter, J.; Slusher, M.J.; Wilcox, B.R.; Berdeen, J.; Knutsen, G.A.; Cardona, C.J.; Stallknecht, D.E. Isolation of Influenza A Viruses from Wild Ducks and Feathers in Minnesota (2010–2011). *Avian Dis.* **2013**, *57*, 677–680. [[CrossRef](#)] [[PubMed](#)]

221. Frongia, G.N.; Peric, T.; Leoni, G.; Satta, V.; Berlinguer, F.; Muzzeddu, M.; Prandi, A.; Naitana, S.; Comin, A. Assessment of Cortisol and DHEA Concentrations in Griffon Vulture (*Gyps fulvus*) Feathers to Evaluate its Allostatic Load. *Ann. Anim. Sci.* **2020**, *20*, 85–96. [[CrossRef](#)]
222. Dipineto, L.; Bossa, L.M.D.L.; Pace, A.; Russo, T.P.; Gargiulo, A.; Ciccarelli, F.; Raia, P.; Caputo, V.; Fioretti, A. Microbiological Survey of Birds of Prey Pellets. *Comp. Immunol. Microbiol. Infect. Dis.* **2015**, *41*, 49–53. [[CrossRef](#)]
223. Nguyen, Q. The Isolation of Presumptive Pathogens from the Pellets of Seven North American Owl Species. *Merge* **2019**, *3*, 1–21.