

PAPER • OPEN ACCESS

## Transition neutrino magnetic moments in CEvNS

To cite this article: Patrick D. Bolton *et al* 2021 *J. Phys.: Conf. Ser.* **2156** 012218

View the [article online](#) for updates and enhancements.

### You may also like

- [Comparative Analysis of Gender Differences in Environmental Values of Chinese People](#)  
Ningning Wu, Shuai Wang, Xiao Liang et al.
- [Community cooling infrastructure from waste heat among diverse building types in Rourkela Steel Township, India](#)  
R Sen, E Teitelbaum and F Meggers
- [The comparison of velaroidal shell structures of square plane load bearing properties](#)  
E Tupikova and M Berdiev



*Benefit from connecting  
with your community*

## ECS Membership = Connection

**ECS membership connects you to the electrochemical community:**

- Facilitate your research and discovery through ECS meetings which convene scientists from around the world;
- Access professional support through your lifetime career;
- Open up mentorship opportunities across the stages of your career;
- Build relationships that nurture partnership, teamwork—and success!

**Join ECS!**      **Visit [electrochem.org/join](https://electrochem.org/join)**



# Transition neutrino magnetic moments in $CE\nu NS$

Patrick D. Bolton<sup>1,2</sup>, Frank F. Deppisch<sup>1</sup>, Kåre Fridell<sup>\*</sup><sup>3</sup>, Julia Harz<sup>3</sup>, Chandan Hati<sup>3</sup> and Suchita Kulkarni<sup>4</sup>

<sup>1</sup>Department of Physics and Astronomy, University College London, London WC1E 6BT, United Kingdom

<sup>2</sup>SISSA, International School for Advanced Studies, INFN, Sezione di Trieste, Via Bonomea 265, I-34136 Trieste, Italy

<sup>3</sup>Physik Department T70, Technische Universität München, James-Frank-Straße 1, D-85748 Garching, Germany

<sup>4</sup>Institute of Physics, NAWI Graz, University of Graz, Universitätsplatz 5, A-8010 Graz, Austria

E-mail: kare.fridell@tum.de

**Abstract.** Coherent Elastic Neutrino Nucleus Scattering ( $CE\nu NS$ ) is a novel technique to look for new physics beyond the Standard Model. We study the prospects of probing a transition magnetic moment in  $CE\nu NS$  experiments. Showing the NUCLEUS experiment as an example, we demonstrate that properties of a potential sterile neutrino can be deduced.

## 1. Introduction

Coherent neutrino nucleus scattering ( $CE\nu NS$ ) was observed for the first time at the COHERENT experiment in 2017 [1]. One of the major difficulties in observing such scatterings is the smallness of the nuclear recoil that is induced, with recoil energies being in the keV range. The near-future experiment NUCLEUS [2, 3], located at the CHOOZ nuclear site in France, aims to reach a sensitivity to nuclear recoils in the  $\mathcal{O}(10)$  eV range.

After the first observation of  $CE\nu NS$ , the novelty in  $CE\nu NS$  experiments has shifted from confirming the existence of this process, which is predicted by the Standard Model (SM), to using it as an observable to constrain neutrino oscillation parameters [4, 5] or search for New Physics (NP) beyond the SM (BSM) [6–11]. A possible BSM scenario that could induce a signal in  $CE\nu NS$  experiments is the existence of an enhanced neutrino magnetic moment. This magnetic moment, connecting neutrinos with photons, can provide an additional interaction channel between neutrinos and nuclei, thereby enhancing the  $CE\nu NS$  signal.


One further possibility is that one of the fermions participating in the magnetic moment vertex is sterile neutrino  $N$ , which can be of either Dirac or Majorana nature, rather than an active SM neutrino  $\nu_{\alpha L}$ . Such a vertex can be realized by the Lagrangian

$$\mathcal{L} \supset \mu_{\nu N}^{\alpha} \bar{\nu}_{\alpha L} \sigma_{\mu\nu} P_R N F^{\mu\nu} + \text{h.c.}, \quad (1)$$

where  $F^{\mu\nu} = \partial^{\mu} A^{\nu} - \partial^{\nu} A^{\mu}$  is the field strength tensor corresponding to the electromagnetic field  $A^{\mu}$ . The terms in Eq. (1) correspond to dimension 5 effective field theory (EFT) operators, that can be UV-completed in for example renormalisable inverse see-saw models, for both Dirac and Majorana active neutrinos [12, 13].

\*Speaker





**Figure 1. Left:** Neutrino transition magnetic moment vertex. **Right:** Primakoff upscattering of a neutrino  $\nu$  to a sterile neutrino  $N$  via a nucleus  $A$ , with subsequent decay of the sterile neutrino to an active neutrino and a photon.

The evidence for non-zero neutrino masses, coming from neutrino oscillation experiments [14], motivates the search for sterile neutrinos that could take part in the active neutrino mass generation mechanism. In case the sterile neutrino is a Majorana particle, the transition magnetic moment vertex will also contribute to a radiative neutrino Majorana mass term [6], possibly explaining the origin of active neutrino masses.

## 2. Radiative $\text{CE}\nu\text{NS}$

The transition neutrino magnetic moment, shown in Fig. 1 (left), can lead to Primakoff upscattering in  $\text{CE}\nu\text{NS}$  events, where an incoming active neutrino transitions into a sterile neutrino in its interaction with the nucleus. Assuming that the sterile neutrino is heavy, it can decay back into an active neutrino and an additional photon via the same vertex that produced the initial upscattering, as depicted in Fig. 1 (right).


In Ref. [6], we proposed a new type of experimental search, where the possibility to detect a final state photon is present in  $\text{CE}\nu\text{NS}$  experiments. This could allow for efficient background rejection, while simultaneously providing a new way to search for sterile neutrinos. In such a set-up, a photon detector is placed a distance  $l_0$  downstream of a  $\text{CE}\nu\text{NS}$  target. The sterile neutrinos that decay via the transition magnetic moment before reaching the photon detector will emit a photon that can be detected. This requirement of decaying within the experimental set-up can be accounted for in the effective width of the sterile neutrino  $\Gamma_N \propto 1/l_0$ , such that longer distances  $l_0$  generally lead to a larger proportion of  $N$  being able to provide a signal [6]. If the nuclear recoil in the upscattering is also recorded, the photon- and nuclear recoil signals are coincident and can provide a clear signature for these types of radiative  $\text{CE}\nu\text{NS}$  events.

## 3. Constraints

There are numerous constraints on the neutrino transition magnetic moment coupling. This coupling is flavour dependent, and therefore different experiments probe different flavour couplings (depending on the source of neutrinos in terms of their flavour). For experiments that use neutrinos from pion decays, the most stringent constraints are provided for the  $\mu$ -flavour coupling. Astrophysical observations generally constrain flavour-universally, and most stringently so in the low- $m_N$  part of the parameter space.

The NUCLEUS experiment uses a nuclear reactor as the source of neutrinos, where electron-flavour antineutrinos are emitted in the  $\beta$ -decays of fission products. Therefore, the relevant constraints to consider for NUCLEUS are those on the electron-flavour neutrino transition magnetic moment coupling  $\mu_{\nu N}^e$ . As shown in Fig. 2, the parameter space of this coupling is constrained by both astrophysical observations and laboratory experiments.

The reach of the near-future NUCLEUS experiment for the radiative  $\text{CE}\nu\text{NS}$  mode with an active neutrino final state, in the case where final state photon detection is possible and 10g



**Figure 2.** Parameter space of the electron flavour neutrino transition magnetic dipole moment  $\mu_{\nu N}^e$  as a function of the sterile neutrino mass  $m_N$ . The reach of the NUCLEUS experiment for the radiative  $CE\nu NS$  mode is shown in red solid and dot-dashed lines for the near-future 10g detector and black solid and dot-dashed lines for the far-future 10 kg upgrade, where solid lines refer to an active neutrino final state and dot-dashed lines correspond to another sterile fermion  $N'$  in the final state, assuming a 3 year run time [6]. Existing constraints are shown in coloured areas. Astrophysical constraints come from Borexino, SN1987A, and BBN [15], while laboratory constraints come from LSND, LEP [16], and COHERENT [6]. Future projections are shown in dashed lines for DUNE near detector [17] and NUCLEUS for Primakoff upscattering [6]. All constraints are given at 90% C.L. unless otherwise stated.

of target material is used, is within the region excluded by the COHERENT experiment in the parameter space of  $\mu_{\nu N}^e$  and  $m_N$ , assuming a 3 year run time for NUCLEUS [6]. A possible far-future upgrade of NUCLEUS, consisting of 1kg of target material, would have a much greater reach in the parameter space than the near-future 10g experiment. However, also the region that would be explored by the upgrade is excluded by the COHERENT experiment.

The exclusion line in Fig. 2 corresponding to the COHERENT experiment assumes that Primakoff upscattering via a neutrino transition magnetic moment would cause a deviation from the standard  $CE\nu NS$  signal, and since such a deviation is not seen, a part of the relevant parameter space is excluded [6]. Therefore, the process that corresponds to the exclusion line from COHERENT includes one instance of the neutrino transition magnetic moment, while the exclusion line from NUCLEUS, corresponding to the radiative  $CE\nu NS$  mode, includes two instances. Therefore, prime objective in the search for a radiative  $CE\nu NS$  mode is not to constrain new regions of parameter space, but rather to find the details of a possible signal coming from Primakoff upscattering. As seen in Ref. [6], the potential Primakoff upscattering exclusion limit from the near- and far-future NUCLEUS is well within new regions of parameter space. Furthermore, in the radiative mode, it is possible that the decay of  $N$  contains final state fermions  $N'$  other than the active electron-flavour neutrino, possibly enhancing the experimental reach of the radiative mode further.

#### 4. Conclusion

One possible method to search for NP in the neutrino sector is via  $CE\nu NS$  experiments, where the sterile neutrino could be produced in Primakoff upscattering events. The improved sensitivity

in the near-future CE $\nu$ NS experiment NUCLEUS will allow sterile neutrinos with MeV-scale masses to be probed in regions of parameter space that are so far unconstrained [6].

The final state photon in radiative CE $\nu$ NS events could in principle be detected if a photon detector is placed downstream of the CE $\nu$ NS target [6]. A coincident signal of a final state photon and a nuclear recoil in the CE $\nu$ NS target could reduce the background, and help to identify the process producing the signals.

The potential that CE $\nu$ NS experiments have in providing more details about the properties of neutrinos is largely still untapped, due mainly to the novelty of such experiments. Therefore, the future of CE $\nu$ NS is very exciting, and the next generation of experiments will be crucial in the development of this relatively new search mode.

## Acknowledgements

The authors are grateful to the NUCLEUS collaboration, in particular to B. Mauri, E. Mazzucato, C. Nones, J. Rothe, T. Lasserre, F. Reindl, R. Strauss, M. Vignati for many useful discussions regarding the NUCLEUS experiment. K. F., J. H. and C. H. acknowledge support from the DFG Emmy Noether Grant No. HA 8555/1-1. K. F. acknowledges support from the DFG Collaborative Research Centre “Neutrinos and Dark Matter in Astro- and Particle Physics” (SFB 1258). J. H. acknowledges discussions at the Aspen Center for Physics, supported by National Science Foundation grant PHY-1607611. P. D. B. and F. F. D. acknowledge support from the UK Science and Technology Facilities Council (STFC) via the Consolidated Grants ST/P00072X/1 and ST/T000880/1. P. D. B. has received support from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 860881-HIDDeN. S. K. is supported by the Austrian Academy of Sciences Elise-Richter grant project number V592-N27.

## References

- [1] D. Akimov *et al.* [COHERENT], *Science* **357** (2017) no.6356, 1123-1126 doi:10.1126/science.aao0990 [arXiv:1708.01294 [nucl-ex]].
- [2] G. Angloher *et al.* [NUCLEUS], *Eur. Phys. J. C* **79** (2019) no.12, 1018 doi:10.1140/epjc/s10052-019-7454-4 [arXiv:1905.10258 [physics.ins-det]].
- [3] J. Rothe *et al.* [NUCLEUS], *J. Low Temp. Phys.* **199** (2019) no.1-2, 433-440 doi:10.1007/s10909-019-02283-7
- [4] P. Coloma and T. Schwetz, *Phys. Rev. D* **94** (2016) no.5, 055005 [erratum: *Phys. Rev. D* **95** (2017) no.7, 079903] doi:10.1103/PhysRevD.94.055005 [arXiv:1604.05772 [hep-ph]].
- [5] P. B. Denton, J. Gehrlein and R. Pestes, *Phys. Rev. Lett.* **126** (2021) no.5, 051801 doi:10.1103/PhysRevLett.126.051801 [arXiv:2008.01110 [hep-ph]].
- [6] P. D. Bolton, F. F. Deppisch, K. Fridell, J. Harz, C. Hati and S. Kulkarni, [arXiv:2110.02233 [hep-ph]].
- [7] P. Coloma, P. A. N. Machado, I. Martínez-Soler and I. M. Shoemaker, *Phys. Rev. Lett.* **119** (2017) no.20, 201804 doi:10.1103/PhysRevLett.119.201804 [arXiv:1707.08573 [hep-ph]].
- [8] O. G. Miranda, D. K. Papoulias, M. Tórtola and J. W. F. Valle, *JHEP* **07** (2019), 103 doi:10.1007/JHEP07(2019)103 [arXiv:1905.03750 [hep-ph]].
- [9] P. S. Bhupal Dev, K. S. Babu, P. B. Denton, P. A. N. Machado, C. A. Argüelles, J. L. Barrow, S. S. Chatterjee, M. C. Chen, A. de Gouvêa and B. Dutta, *et al.* *SciPost Phys. Proc.* **2** (2019), 001 doi:10.21468/SciPostPhysProc.2.001 [arXiv:1907.00991 [hep-ph]].
- [10] P. Coloma, I. Esteban, M. C. Gonzalez-Garcia and M. Maltoni, *JHEP* **02** (2020), 023 doi:10.1007/JHEP02(2020)023 [arXiv:1911.09109 [hep-ph]].
- [11] A. N. Khan, D. W. McKay and W. Rodejohann, *Phys. Rev. D* **104** (2021) no.1, 015019 doi:10.1103/PhysRevD.104.015019 [arXiv:2104.00425 [hep-ph]].
- [12] R. N. Mohapatra, *Phys. Rev. Lett.* **56** (1986), 561-563 doi:10.1103/PhysRevLett.56.561
- [13] R. N. Mohapatra and J. W. F. Valle, *Phys. Rev. D* **34** (1986), 1642 doi:10.1103/PhysRevD.34.1642
- [14] P.A. Zyla *et al.* [Particle Data Group], *PTEP* **2020** (2020) no.8, 083C01 doi:10.1093/ptep/ptaa104
- [15] V. Brdar, A. Greljo, J. Kopp and T. Opferkuch, *JCAP* **01** (2021), 039 doi:10.1088/1475-7516/2021/01/039 [arXiv:2007.15563 [hep-ph]].
- [16] G. Magill, R. Plestid, M. Pospelov and Y. D. Tsai, *Phys. Rev. D* **98** (2018) no.11, 115015 doi:10.1103/PhysRevD.98.115015 [arXiv:1803.03262 [hep-ph]].
- [17] T. Schwetz, A. Zhou and J. Y. Zhu, *JHEP* **21** (2020), 200 doi:10.1007/JHEP07(2021)200 [arXiv:2105.09699 [hep-ph]].