

Weierstraß-Institut
für Angewandte Analysis und Stochastik
Leibniz-Institut im Forschungsverbund Berlin e. V.

Preprint

ISSN 2198-5855

**Stability of the solution set of quasi-variational inequalities and
optimal control**

Amal Alphonse¹, Michael Hintermüller^{1,2}, Carlos N. Rautenberg^{1,2}

submitted: March 20, 2019

¹ Weierstrass Institute
Mohrenstr. 39
10117 Berlin
Germany
E-Mail: amal.alphonse@wias-berlin.de
michael.hintermueller@wias-berlin.de
carlos.rautenberg@wias-berlin.de

² Humboldt-Universität zu Berlin
Unter den Linden 6
10099 Berlin
Germany
E-Mail: hint@math.hu-berlin.de
carlos.rautenberg@math.hu-berlin.de

No. 2582
Berlin 2019



2010 *Mathematics Subject Classification.* 47J20, 49J40, 49M15, 65J15, 65K10.

Key words and phrases. Quasi-variational inequality, optimal control, multivalued solution maps.

This research was carried out in the framework of MATHEON supported by the Einstein Foundation Berlin within the ECMath projects OT1, SE5 and SE19 as well as the Excellence Cluster Math⁺ Berlin within project AA4-3, EF3-5, the transition project OT6, and the MATHEON project A-AP24. The authors further acknowledge the support of the DFG through the DFG-SPP 1962: Priority Programme Non-smooth and Complementarity-based Distributed Parameter Systems: Simulation and Hierarchical Optimization within Projects 10 and 11.

Edited by
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)
Leibniz-Institut im Forschungsverbund Berlin e. V.
Mohrenstraße 39
10117 Berlin
Germany

Fax: +49 30 20372-303
E-Mail: preprint@wias-berlin.de
World Wide Web: <http://www.wias-berlin.de/>

Stability of the solution set of quasi-variational inequalities and optimal control

Amal Alphonse, Michael Hintermüller, Carlos N. Rautenberg

Abstract

For a class of quasivariational inequalities (QVIs) of obstacle-type the stability of its solution set and associated optimal control problems are considered. These optimal control problems are non-standard in the sense that they involve an objective with set-valued arguments. The approach to study the solution stability is based on perturbations of minimal and maximal elements to the solution set of the QVI with respect to monotonic perturbations of the forcing term. It is shown that different assumptions are required for studying decreasing and increasing perturbations and that the optimization problem of interest is well-posed.

1 Introduction

Quasi-variational inequalities (QVIs) are powerful mathematical models for the description of complex physical phenomena. Such models arise in many scientific areas including superconductivity ([39, 49, 50, 11, 14, 45, 32, 31, 30]), continuum mechanics ([23]), impulse control problems ([16, 17, 15, 18]), growth of sandpiles ([12, 13, 14, 43, 44, 46, 47]), and the formation of networks of lakes and rivers ([13, 44, 46]), among others.

In general, QVIs are nonlinear, nonconvex, and nonsmooth problems with non-unique (i.e., set-valued) solutions. In physical models like the growth of sandpiles or the determination of the magnetic field in superconductors, each of these solutions fulfills physical laws confirming that they are not artifacts of the mathematical formulation (compare the results in [11, 12, 13, 14, 43, 6]). In some cases, like the QVI arising in impulse control problems, extremals of the solution set can be determined, in the sense that there exist minimal and maximal elements of the solution set which are related to the value functional [16].

The mathematical treatment of QVIs entails several possible directions. In addition to the “order” approach followed in this paper, at least two more are worth mentioning. In some cases, the QVI can be expressed as a generalized equation, and hence a particular instance of a more general problem class; see, e.g., [35, 36] and also [24, 34, 25]. In problems involving constraints on derivatives, special forms of constraint regularization that modify the original partial differential operator may be suitable, see [49, 40, 8, 9]. For details on these and further approaches, we refer the reader to [4].

Given the complexity of QVIs, their optimal control represents a task which is yet even more complex than the study of the QVI itself. While without any structural properties of the solution set the treatment of the control problem appears very hard if not impossible, solution properties such as the availability of extremal elements provide useful starting points for the successful analysis of the control problem and characterizations of its solutions. For this purpose, the study of the stability of minimal and maximal elements of the solution set with respect to perturbations of the forcing term represents a fundamental analytical step for the subsequent study of the control problem. Concerning the latter considered in

infinite dimensions, we note that the literature is rather scarce; we refer to [2, 22, 21, 42] for some of the very few contributions. Finite dimensional cases have been studied in [41] and the references therein. On the other hand, the study of optimal control problems for variational inequalities (VIs) has been the subject of a number of recent studies; see, e.g., [28, 33, 29, 53] and the references therein. We note here that—to the best of our knowledge—the study of the stability of minimal and maximal solutions of QVIs and the optimal control thereof, with both being focus topics of this work, have not yet been treated in the literature. We further note that the stability of the solution set is also of relevance in identification problems involving QVIs; see [27].

The paper is organized as follows. In section 2 we introduce the optimal control problem associated to the QVI of interest, and we provide the mathematical foundation of the structure of spaces under consideration and their associated ordering. Additionally, in section 3 we study two classes of applications associated to impulse control problems and to QVIs arising as the coupling of VIs and nonlinear partial differential equations (PDEs), respectively. In section 4, we discuss the fundamental results due to Tartar that determine the existence of minimal and maximal solutions of the QVIs of interest. Abstract stability results from the operator theoretic point of view are the subject of section 5, along with an example exploring limitations. In section 6, we study minimal and maximal solutions under perturbations of the forcing term from below and from above. The paper ends in section 7 which studies the well-posedness of the control problem for the QVI.

Notation

Throughout the paper we assume that Ω is an open subset of \mathbb{R}^N , and $L^p(\Omega)$ for $1 \leq p \leq \infty$ denotes the usual Lebesgue space. For $\nu > 0$, we define

$$L_\nu^\infty(\Omega) := \{z \in L^\infty(\Omega) : z(x) \geq \nu \text{ for almost all (f.a.a.) } x \in \Omega\}.$$

Additionally, $H_0^1(\Omega)$ and $H^1(\Omega)$ denote the usual Sobolev spaces; see [1].

For a Banach space X we write $\|\cdot\|_X$ for a norm on X and X' for the topological dual of X with $\langle \cdot, \cdot \rangle_{X', X}$ the associated duality pairing, respectively. For a sequence $\{z_n\}_{n \in \mathbb{N}}$ in X we denote its strong convergence to $z \in X$ by “ $z_n \rightarrow z$ ” and weak convergence by “ $z_n \rightharpoonup z$ ”. Further, for two Banach spaces X_1 and X_2 , we write $\mathcal{L}(X_1, X_2)$ for the space of bounded linear operators from X_1 to X_2 .

2 A class of optimization problems with QVI constraints

2.1 Preliminaries

Let (V, H, V') be a Gelfand triple of Hilbert spaces, i.e., $V \hookrightarrow H \hookrightarrow V'$, where the embedding $V \hookrightarrow H$ is dense and continuous, H is identified with H' , and the embedding $H \hookrightarrow V'$ is dense and continuous as well (see [26] and also, e.g., [20]). Also, from now on we use $\langle \cdot, \cdot \rangle := \langle \cdot, \cdot \rangle_{V', V}$ and (\cdot, \cdot) for the inner product in H .

Let further $H^+ \subset H$ be a closed convex cone satisfying $H^+ = \{v \in H : (v, y) \geq 0 \text{ for all } y \in H^+\}$. Note that H^+ defines the cone of non-negative elements inducing the vector ordering: $x \leq y$ if and only if $y - x \in H^+$. Given $x \in H$, let x^+ denote the orthogonal projection of x onto H^+ , and $x^- := x - x^+$ the one onto $H^- := -H^+$. Clearly, one has the decomposition $x = x^+ - x^- \in$

$H^+ - H^+$ for every $x \in H$, and $(x^+, x^-) = 0$. Further, the infimum and supremum of two elements $x, y \in H$ are defined as $\sup(x, y) := x + (y - x)^+$ and $\inf(x, y) := x - (x - y)^+$, respectively. The supremum of an arbitrary subset of H that is bounded (in the order) above is also properly defined since H is Dedekind complete. In other words: For a set $\{x_i\}_{i \in J}$ where J is completely ordered and bounded from above, we have that $\{x_i\}_{i \in J}$ is a generalized Cauchy sequence in H (see [7, Chapter 15, §15.2, Proposition 1]), and then Dedekind completeness follows (see [3, Chapter 4, Theorem 4.9 and Corollary 4.10]). This additionally implies that norm convergence preserves order, i.e., if $z_n \leq y_n$ for every $n \in \mathbb{N}$ and $z_n \rightarrow z$ and $y_n \rightarrow y$ both in H , then $z \leq y$. Also, we write $z_n \downarrow z$ in H if $z_n \geq z_{n+1}$ for all $n \in \mathbb{N}$ and $z_n \rightarrow z$ in H , and analogously for $z_n \uparrow z$. Further, we have that if the sequence $\{z_n\}$ is non-increasing (non-decreasing) and bounded from below (above) in the sense of the order, then there exists $z \in H$ for which $z_n \downarrow z$ ($z_n \uparrow z$) in H . Now, concerning V we assume that $y \in V$ implies $y^+ \in V$, and that $(\cdot)^+ : V \rightarrow V$ is a bounded operator, i.e., we have $M > 0$ with $\|y^+\|_V \leq M\|y\|_V$ for all $y \in V$.

Given $x, y \in H$ such that $x \leq y$, we define the closed “interval” with x and y as its respective endpoints by $[x, y] := \{z \in H : x \leq z \text{ and } z \leq y\}$. Furthermore, we write $[y, +\infty)$ and $(-\infty, y]$ instead of $\{z \in H : z \geq y\}$ and $\{z \in H : z \leq y\}$, respectively.

Next we get more specific with respect to V and H . In fact, both are assumed to be spaces of maps $h : \Omega \rightarrow \mathbb{R}$ over some open set $\Omega \subset \mathbb{R}^N$ with the following dense and continuous embedding: $L^\infty(\Omega) \hookrightarrow H$ such that $L^\infty(\Omega) \hookrightarrow V'$, as well. Our prototypical example for this setting is $V := H_0^1(\Omega)$ and $H := L^2(\Omega)$ with $H^+ := L_+^2(\Omega)$, the closed convex cone of non-negative maps in $L^2(\Omega)$ with “ $v \leq w$ ” for $v, w \in H$ iff $v(x) \leq w(x)$ almost everywhere (a.e.) on Ω . Here, we have $v^+(x) := \max\{v(x), 0\}$ for $x \in \Omega$.

Let $A : V \rightarrow V'$ be a (possibly nonlinear) operator that is

(-) homogenous of order one, i.e., $A(tu) = tA(u)$ for all $u \in V, t > 0$;

(-) Lipschitz continuous, i.e., there exists $C > 0$ such that

$$\|A(u) - A(v)\|_{V'} \leq C\|u - v\|_V, \quad \text{for all } u, v \in V;$$

(-) strongly monotone, i.e., there exists $c > 0$ such that

$$\langle A(u) - A(v), u - v \rangle \geq c\|u - v\|_V^2, \quad \text{for all } u, v \in V;$$

(-) T-monotone, i.e.,

$$\langle A(u) - A(v), (u - v)^+ \rangle \geq 0, \quad \text{for all } u, v \in V,$$

where equality holds if and only if $(u - v)^+ = 0$.

A well-known example for A in the case $V = H_0^1(\Omega)$ (or $V = H^1(\Omega)$) and $H = L^2(\Omega)$ is given by the elliptic partial differential operator

$$\langle Av, w \rangle = \sum_{i,j} \int_{\Omega} a_{ij}(x) \frac{\partial v}{\partial x_j} \frac{\partial w}{\partial x_i} dx + \sum_i \int_{\Omega} a_i(x) \frac{\partial v}{\partial x_i} w + \int_{\Omega} a_0(x) vw dx, \quad (1)$$

under suitable assumptions on a_{ij}, a_i and a_0 such as, e.g., $a_{ij}, a_0 \in L^\infty(\Omega), a_i \equiv 0, \sum a_{ij}(x) \xi_j \xi_i \geq c|\xi|^2$ for all $\xi = \{\xi_i\} \in \mathbb{R}^N$, and $a_0(x) \geq \epsilon > 0$ f.a.a. $x \in \Omega$.

For the definition of the constraint set of the QVI we need a map Φ with the following properties: There exist $\underline{y}, \bar{y} \in H$ such that $\underline{y} \leq \bar{y}$ for which $\Phi : [\underline{y}, \bar{y}] \rightarrow H^+ \cup \{+\infty\}$, and Φ is increasing in $[\underline{y}, \bar{y}]$. The latter yields that if $v, w \in [\underline{y}, \bar{y}]$ and $v \leq w$ then $\Phi(v) \leq \Phi(w)$. Further properties of Φ will be specified below.

Next, we define the set-valued map $\mathbf{K} : H^+ \rightarrow 2^V$ as

$$\mathbf{K}(\psi) := \{v \in V : v \leq \psi\}. \quad (2)$$

Note that $\mathbf{K}(\psi) \subset V$ is non-empty, closed and convex. We also set $\mathbf{K}(+\infty) := V$.

2.2 Problem formulations

The QVI problem of interest is the following one.

Problem (P_{QVI}) : Let $f \in V'$ be given.

$$\text{Find } y \in \mathbf{K}(\Phi(y)) : \langle A(y) - f, v - y \rangle \geq 0, \quad \forall v \in \mathbf{K}(\Phi(y)). \quad (P_{\text{QVI}})$$

This problem admits (in general) multiple solutions due to the non-convexity resulting from $y \mapsto \mathbf{K}(\Phi(y))$. Let $\mathbf{Q}(f)$ denote the associated solution set.

In applications, one is typically interested in confining the solution set $\mathbf{Q}(f)$ to a certain interval $[\underline{y}, \bar{y}]$ for some given $\underline{y}, \bar{y} \in H$. This can be done by considering f a control force and by solving the following optimal control problem:

Problem (\mathbb{P}) :

$$\begin{aligned} & \text{minimize } J(\mathbf{O}, f) := J_1(T_{\text{sup}}(\mathbf{O}), T_{\text{inf}}(\mathbf{O})) + J_2(f) \text{ over } (\mathbf{O}, f) \in 2^H \times U, \\ & \text{subject to } f \in U_{\text{ad}}, \\ & y \in \mathbf{O}, \quad \mathbf{O} = \{z \in V : z \text{ solves } P_{\text{QVI}}\}. \end{aligned} \quad (\mathbb{P})$$

Here $U_{\text{ad}} \subset U \subset V'$ is the set of admissible controls. Moreover, $J_1 : H \times H \rightarrow \mathbb{R}$ and $J_2 : U \rightarrow \mathbb{R}$, and for $\underline{y}, \bar{y} \in H$ we define the set-valued map

$$T_{\text{sup}}(\mathbf{O}) := \begin{cases} \sup_{z \in \mathbf{O} \cap [\underline{y}, \bar{y}]} z, & \mathbf{O} \cap [\underline{y}, \bar{y}] \neq \emptyset, \\ \underline{y}, & \text{otherwise,} \end{cases}$$

and analogously

$$T_{\text{inf}}(\mathbf{O}) := \begin{cases} \inf_{z \in \mathbf{O} \cap [\underline{y}, \bar{y}]} z, & \mathbf{O} \cap [\underline{y}, \bar{y}] \neq \emptyset, \\ \bar{y}, & \text{otherwise.} \end{cases}$$

Problems of type (\mathbb{P}) have not yet been considered in the literature and pose several formidable challenges. For instance, the proof of existence of a solution is highly delicate due to the dependence $y \mapsto \mathbf{K}(\Phi(y))$ and the fact that $y = y(f)$. As a consequence, the direct method of the calculus of variations is only applicable if certain convergence properties of that constraint set can be guaranteed. Another delicacy is related to the potential set-valuedness of the solution of the QVI in the constraint system of (\mathbb{P}). This fact requires to identify a suitable selection mechanism such as the one identifying the maximal or minimal solution, respectively, if available at all. We note, however, that in the special case where $T_{\text{inf}}(\mathbf{Q}(f))$ and $T_{\text{sup}}(\mathbf{Q}(f))$ also belong to $\mathbf{Q}(f)$, they are the minimal and maximal solution, respectively, to (P_{QVI}) in $V \cap [\underline{y}, \bar{y}]$. Then the proof of existence of solutions to (\mathbb{P}) reduces to a stability result for this minimal and maximal solution to the QVI of interest.

3 Examples of application

Our work here is motivated by the following two applications. The first one is associated to QVIs that result from coupling a variational inequality (VI) to a nonlinear partial differential equation (PDE). Such models have recently been studied in connection with thermoforming; see [5]. The other problem class is given by QVIs arising in impulse control as pioneered by Bensoussan and Lions. We briefly describe both problem types in the sequel.

3.1 QVIs arising from coupling VIs and nonlinear PDEs

Consider the following class of compliant obstacle problems where the obstacle is given implicitly by solving a PDE, thus coupling a VI and a PDE. It consists in finding $(y, \Phi, z) \in V \times H \times W$ such that

$$y \leq \Phi, \quad \langle A(y) - f, y - v \rangle \leq 0, \quad \forall v \in V : v \leq \Phi, \quad (3)$$

$$\langle Bz + G(\Phi, y) - g, w \rangle = 0 \quad \forall w \in W, \quad (4)$$

$$\Phi = Lz, \quad \text{in } H. \quad (5)$$

Here, $V \hookrightarrow W \hookrightarrow H \hookrightarrow W' \hookrightarrow V'$, $f, g \in H^+$, $G : H \times V \rightarrow H$ is continuous and bounded, i.e., for some $M_G > 0$, $\|G(\Phi, y)\|_H \leq M_G(\|\Phi\|_H + \|y\|_V)$, for all $(\Phi, y) \in H \times V$. Further, $L : W \rightarrow H$ is an increasing linear continuous map. Additionally, $B \in \mathcal{L}(W, W')$ is strongly monotone and satisfies $\langle Bz^+, z^- \rangle \leq 0$ for all $z \in W$.

Under mild conditions, the above problem can be cast into the form of (P_{QVI}) as follows. Let $v \in V$, and consider the problem of finding $z \in W$ such that

$$\langle Bz + G(\phi, v) - g, w \rangle = 0 \quad \forall w \in W, \quad (6)$$

$$\phi = Lz, \quad \text{in } H. \quad (7)$$

Assuming that for each $v \in V$, $z \mapsto G(Lz, v)$ is monotone, one can show the existence of a unique solution $z(v) \in W$ of (6)–(7). Now set $\Phi(v) := \phi$. Suppose additionally that $\langle G(Lz, y), z^- \rangle \leq 0$ for all $z \in W$ and $y \in V \cap H^+$ so that $z(v) \geq 0$ and $\Phi(v) = Lz(v) \geq 0$ for all v for each $v \in V$, and that if $v_1 \leq v_2$ then

$$(G(Lv, v_1) - G(Lw, v_2), (v - w)^+) > 0,$$

for all w, v with $(v - w)^+ \neq 0$. Then $z(v_1) \leq z(v_2)$ which implies for $\Phi(v) := Lz(v)$ that $v_1 \leq v_2$ implies $\Phi(v_1) \leq \Phi(v_2)$, as L is increasing. This finally shows that (3)–(5) has the form (P_{QVI}) .

In view of controlling the outcome of a stationary industrial process one is clearly interested in forcing the solution set $\mathbf{Q}(f)$ to be a singleton which is close to a pre-specified desired state y_d . This can be modelled as follows.

$$\text{minimize } \frac{1}{2} \int_{\Omega} |T_{\text{sup}}(\mathbf{Q}(f)) - T_{\text{inf}}(\mathbf{Q}(f))|^2 + \frac{1}{2} \int_{\Omega} |y_d - T_{\text{inf}}(\mathbf{Q}(f))|^2 + \frac{\lambda}{2} |f|_U^2 \quad (8)$$

$$\text{subject to } 0 < \nu \leq f \leq F, \quad f \in U,$$

for given $\lambda, \nu, F > 0$. Here, U denotes the underlying control space. Note that the first term in the objective aims at minimizing the distance between the maximal and minimal solution targeting single-valued $\mathbf{Q}(f)$, the second term aims at tracking y_d , whereas the last term associates an $(U-)$ average cost of λ to the control action. Notice that the smaller λ the cheaper the cost of the control gets and the smaller one expects the first two terms in the objective. Clearly, (8) fits the form of (\mathbb{P}) .

Example 1. A possible setting for this problem class satisfying all assumptions invoked so far is given by $V = H_0^1(\Omega)$, $H = L^2(\Omega)$, $W = H^1(\Omega)$, with

$$\begin{aligned} \langle Ay, z \rangle &= \sum_{i,j} \int_{\Omega} a_{ij}(x) \frac{\partial y}{\partial x_j} \frac{\partial z}{\partial x_i} dx + \int_{\Omega} a_0(x) y z dx, \quad \forall y, z \in V, \\ \langle Bv, w \rangle &= \sum_{i,j} \int_{\Omega} b_{ij}(x) \frac{\partial v}{\partial x_j} \frac{\partial w}{\partial x_i} dx + \int_{\Omega} b_0(x) v w dx, \quad \forall v, w \in W, \end{aligned}$$

$a_{ij}, b_{ij}, a_0, b_0 \in L^\infty(\Omega)$, $\sum a_{ij}(x) \xi_j \xi_i \geq c |\xi|^2$ and $\sum b_{ij}(x) \xi_j \xi_i \geq c |\xi|^2$ for all $\xi = \{\xi_i\} \in \mathbb{R}^N$ and some $c > 0$ and $a_0(x) \geq 0$ and $b_0(x) \geq \epsilon > 0$ f.a.a. $x \in \Omega$. Additionally, for $y \geq 0$

$$G(\Phi, y) = (\Phi - y)^+, \quad \text{and} \quad (Lz)(x) = k(x)z(x)$$

with $k \in L^\infty(\Omega)^+$. Further, $U = \mathbb{R}^M$ for some $M \in \mathbb{N}$, where $f = \sum_{m=1}^M f_m \chi_{\Omega_m}$, $f_m \in \mathbb{R}$ and $\Omega_m \subset \Omega$ for each m , and $\|f\|_U := \|\{f_m\}\|_{\mathbb{R}^M}$. In this setting, $\Phi : \{y \in H : 0 \leq y\} \rightarrow H^+$ is non-decreasing.

3.2 Impulse control

We consider impulse control problems (see [17]) for the following stochastic differential equation

$$du = b(u)dt + \sigma(u)dw(t), \quad u(0) = x \in \mathbb{R}^N,$$

where $b, \sigma : \mathbb{R}^N \rightarrow \mathbb{R}^N$ are Lipschitz functions whose regularity will be specified later. Let $a_{ij} := \sigma_i \sigma_j^\top / 2$.

The control is carried on instances $0 \leq \theta_1 \leq \theta_2 \leq \dots$, and the system is forced from $y(\theta_n^-)$ to $y(\theta_n^-) + \xi_n$ on the instance θ_n . The energy associated to the process is given by the expected value

$$\mathbb{E} \left\{ \int_0^{\tau_x} f(u(t)) e^{-\int_0^t a_0(u(s)) ds} dt + \sum_n (k + c_0(\xi_n)) e^{-\int_0^{\theta_n} a_0(u(s)) ds} \chi_{\theta_n < \infty} \right\} =: \mathcal{H}(f, w, x),$$

where $w = \{(\theta_n, \xi_n)\}_{n=1}^\infty$, and with $\tau_x := \inf\{t : u(t^-) \notin \Omega \text{ or } u(t) \notin \Omega\}$, for some open $\Omega \subset \mathbb{R}^N$. In this setting, f uniquely determines the value function

$$\min_{w \in \mathcal{W}} \mathcal{H}(f, w, x),$$

which represents the cost of the optimal control associated to the initial condition x and the cost function f . Here, \mathcal{W} is the set of all possible instances and jumps $\{(\theta_n, \xi_n)\}_{n=1}^\infty$. The optimization of the above quantity via choosing f turns out to be of interest. Indeed, in specific applications f determines the value of a certain stock or energy type per unit of time. The goal is then to solve

$$\text{minimize} \int_{\Omega} \left(s - \min_{w \in \mathcal{W}} \mathcal{H}(f, w, x) \right)^2 dx + \frac{\lambda}{2} |f|_U^2 \quad \text{subject to } f \in U_{\text{ad}}, \quad (9)$$

where $U_{\text{ad}} \subset U$ is the set of admissible functions f , $|f|_U^2$ represents cost of the choice of f , $\lambda > 0$ is a weight, and $s \geq 0$ is a desired average cost that could be zero.

3.2.1 Bounded case

We consider Ω bounded with a sufficiently smooth boundary, with $V = H_0^1(\Omega)$, $H = L^2(\Omega)$, and $H^+ = L_+^2(\Omega)$, where A is of the type (1) with

$$\begin{aligned} a_{ij} = a_{ji} &\in W^{1,\infty}(\mathbb{R}^N), \quad 1 \leq i, j \leq N, \quad \sum a_{ij} \xi_i \xi_j \geq \alpha |\xi|^2, \quad \alpha > 0, \forall \xi \in \mathbb{R}^N, \\ a_i, a_0 &\in L^\infty(\mathbb{R}^N), \quad b_i = -a_i + \sum_j \frac{\partial a_{ij}}{\partial x_j} \in W^{1,\infty}(\mathbb{R}^N), \\ a_0(x) &\geq r > 0, \quad \text{f.a.a. } x \in \Omega. \end{aligned}$$

Consider

$$(\Phi y)(x) := k + \operatorname{ess\,inf}_{x+\xi \in \bar{\Omega}} (c_0(\xi) + y(x + \xi)),$$

where $c_0 \in C(\mathbb{R}_+^N, \mathbb{R})$ is such that $c_0(0) = 0$ is sub-linear and non-decreasing, with $f \in L^p(\Omega)$ with $p > N$ and $f \geq 0$.

In this setting one can show that the solution set $\mathbf{Q}(f)$ of (P_{QVI}) is a singleton, $\mathbf{Q}(f) = \{y^*\}$, and y^* determines the value function of the impulse control problem of interest (see [17]), i.e.,

$$y^*(x) = \min_{w \in \mathcal{W}} \mathcal{H}(f, w, x),$$

a.e. for $x \in \Omega$. Hence, problem (9) amounts to controlling the solution to the quasi-variational inequality and is, thus, of the form (\mathbb{P}) .

3.2.2 Unbounded case

Let $\omega(x) := \exp(-\mu\sqrt{1+|x|^2})$ for $x \in \mathbb{R}^N$, and consider the weighted spaces $V = H^1(\mathbb{R}^N, \omega)$, and $H = L^2(\mathbb{R}^N, \omega)$ with $H^+ = L_+^2(\mathbb{R}^N, \omega)$ the usual cone of non-negative maps. In particular, $L^2(\mathbb{R}^N, \omega)$ is the space of (equivalence classes of) measurable functions $h : \mathbb{R}^N \rightarrow \mathbb{R}$ for which $|h|_{L^2(\mathbb{R}^N, \omega)}^2 := \int_{\mathbb{R}^N} |h(x)|^2 \omega(x)^2 dx < +\infty$, and $H^1(\mathbb{R}^N, \omega)$ is the space of (equivalence classes of) functions $g : \mathbb{R}^N \rightarrow \mathbb{R}$ for which g and its weak gradient ∇g belong to $L^2(\mathbb{R}^N, \omega)$ and $L^2(\mathbb{R}^N, \omega)^N$, respectively.

The operator $A : V \rightarrow V'$ is given by

$$\langle Av, y \rangle = \sum_{i,j} \int_{\mathbb{R}^N} a_{ij} \frac{\partial v}{\partial x_j} \frac{\partial y}{\partial x_i} \omega^2 dx + \sum_i \int_{\mathbb{R}^N} a_i \frac{\partial v}{\partial x_j} y \omega^2 dx + a_0 \int_{\mathbb{R}^N} v y \omega^2 dx,$$

with a_{ij}, a_i, b_i as in Section 3.2.1, but with $a_0(x) = r$ for all x and a real $r > 0$.

Define

$$U := \{f : \mathbb{R}^N \rightarrow \mathbb{R} : \text{measurable and } 0 \leq f(x) \leq C(1 + |x|^s), \quad x \in \mathbb{R}^N, s \geq 0\},$$

and the map Φ by

$$(\Phi y)(z) := k + \operatorname{ess\,inf}_{\xi \geq 0} (c_0(\xi) + y(z + \xi)),$$

where $k > 0$, and $c_0 \in C(\mathbb{R}_+^N, \mathbb{R})$, with $c_0(0) = 0$, is sub-linear, non-decreasing with $\lim_{|\xi| \rightarrow +\infty} c_0(\xi) = +\infty$ and for which $c_0(\xi) \leq a|\xi|^\gamma$ for some $a, \gamma > 0$.

In this scenario, the set of solutions $\mathbf{Q}(f)$ of (P_{QVI}) is *not* a singleton, and both $T_{\inf}(\mathbf{Q}(f))$ and $T_{\sup}(\mathbf{Q}(f))$ have probabilistic interpretations associated to the value function in impulse control. In particular,

$$T_{\inf}(\mathbf{Q}(f))(x) = \min_{w \in \mathcal{W}} \mathcal{H}(f, w, x),$$

i.e., $T_{\inf}(\mathbf{Q}(f))$ is the value function associated with the initial impulse control problem. Then (9) has the form of (\mathbb{P}) for appropriate choices of J_1 and J_2 .

4 Increasing maps and QVI solutions

This section is strongly related to a result due to Tartar [51]; see also [7]. Upon converting (P_{QVI}) into a fixed-point equation, the corresponding approach yields the existence of a solution for an increasing fixed-point map under very mild assumptions. We note that the technique is analogous to the one by Kolodner and Birkhoff; see [38, 10, 19].

We start by recalling Tartar's result (compare [7, Chapter 15, §15.2]) which rests on increasing maps. In this vein, we call $T : H \rightarrow H$ increasing iff $v \leq w$ implies $T(v) \leq T(w)$.

Theorem 2 (BIRKHOFF-TARTAR). *Suppose $T : H \rightarrow H$ is an increasing map and let \underline{y} be a sub-solution and \bar{y} be a super-solution of the map T , that is:*

$$\underline{y} \leq T(\underline{y}) \quad \text{and} \quad T(\bar{y}) \leq \bar{y}.$$

If $\underline{y} \leq \bar{y}$, then the set of fixed points of the map T in the interval $[\underline{y}, \bar{y}]$ is non-empty and has a smallest and a largest element. $y \in$

We apply the above result to (P_{QVI}) and first need to introduce the following VI.

Problem (P_{VI}) : Let $\psi \in H^+$, $f \in V'$ be given.

$$\text{Find } y \in \mathbf{K}(\psi) : \langle Ay - f, v - y \rangle \geq 0, \quad \forall v \in \mathbf{K}(\psi). \quad (P_{VI})$$

The solution to (P_{VI}) can be proven to be unique by standard methods. For $(f, \psi) \in (V', H^+)$, we denote the unique solution to (P_{VI}) as $S(f, \psi)$. Before we can make use of Theorem 2, we state the following property of the map $(f, \psi) \mapsto S(f, \psi)$. Its proof can be found on [48].

Proposition 1. *Let $f_1, f_2 \in V'$ and $\psi_1, \psi_2 \in H^+$ be such that $f_1 \leq f_2$ in V' and $\psi_1 \leq \psi_2$. Then it holds that $S(f_1, \psi_1) \leq S(f_2, \psi_2)$.*

We note that in the above result $f_1 \leq f_2$ in V' is well-defined, since V inherits the order in H , so that $f_1 \leq f_2$ iff $\langle f_2 - f_1, v \rangle \geq 0$ for all $v \in V$ such that $v \geq 0$. Further observe that the case $\psi = +\infty$ is also allowed, where $S(f, +\infty)$ denotes the solution of the unconstrained problem

$$\text{Find } y \in V \text{ such that } \langle A(y), v \rangle = \langle f, v \rangle, \quad \text{for all } v \in V.$$

This implies that

$$S(f, \psi) \leq S(f, +\infty), \quad \forall f \in V', \psi \in H^+.$$

In order to apply the Birkhoff-Tartar Theorem to the QVI problem of interest, we need to identify a proper interval $[\underline{y}, \bar{y}]$, with \underline{y} a sub-solution and \bar{y} a super-solution of the map $y \mapsto S(f, \Phi(y))$. In our case, we choose $\underline{y} = 0$, since we infer from Proposition 1 that

$$0 = S(0, \Phi(0)) \leq S(f, \Phi(0)),$$

for any $f \geq 0$ in V' . On the other hand, we assume that $f \in U_{\text{ad}} \subset V'$ is bounded from above (in the V' -order) by some F . Then let $\bar{y} = S(F, +\infty)$, for which

$$S(f, \Phi(\bar{y})) \leq S(F, +\infty) = \bar{y}.$$

This leads to the following result.

Theorem 3 (TARTAR). *Let $U_{\text{ad}} \subset \{f \in V' : 0 \leq f \leq F\}$ for some $0 \leq F \in V'$. Then, there are \underline{y}, \bar{y} such that for each $f \in U_{\text{ad}}$, the set of fixed points of the map $y \mapsto S(f, \Phi(y))$ in the interval $[\underline{y}, \bar{y}]$ is non-empty and contains a smallest and a largest element, i.e., there are fixed points y_{\min}^* and y_{\max}^* in V such that*

$$\mathbf{Q}(f) \cap [\underline{y}, \bar{y}] = \mathbf{Q}(f) \cap [y_{\min}^*, y_{\max}^*] \neq \emptyset.$$

In light of Theorems 2 and 3, there exist operators \mathbf{m} and \mathbf{M} , which map an increasing map on the interval $[\underline{y}, \bar{y}]$ to its minimal and maximal fixed points, respectively; insofar that sub- and super-solutions \underline{y} and \bar{y} exist.

We fix some notation now. In the case of a general increasing map T , with sub- and super-solutions \underline{y} and \bar{y} , respectively, we denote by $\mathbf{m}(T)$ and $\mathbf{M}(T)$ its minimal and maximal fixed points in some interval $[\underline{y}, \bar{y}]$. When the map T is given by $y \mapsto S(f, \Phi(y))$ for some f , we write $\mathbf{m}(f)$ and $\mathbf{M}(f)$. In particular, it follows that if $\mathbf{Q}(f)$ is the set of solutions of (P_{QVI}) , then

$$T_{\text{sup}}(\mathbf{Q}(f)) = \mathbf{M}(f), \quad \text{and} \quad T_{\text{inf}}(\mathbf{Q}(f)) = \mathbf{m}(f),$$

where $T_{\text{sup}}, T_{\text{inf}}$ are given in (\mathbb{P}) .

For an operator T as in Theorem 2, the fixed points $\mathbf{m}(T)$ and $\mathbf{M}(T)$ are determined (see [7] for a proof) by the maximal and minimal elements of the sets $Z(T)$ and $\tilde{Z}(T)$, respectively, where

$$\begin{aligned} Z(T) &= \{x \in X(T) : x \leq y \text{ for all } y \in Y(T)\}, \\ \tilde{Z}(T) &= \{y \in Y(T) : x \leq y \text{ for all } x \in X(T)\}, \end{aligned}$$

and

$$\begin{aligned} X(T) &= \{x \in H : x \in [\underline{y}, \bar{y}] \text{ and } x \leq T(x)\}, \\ Y(T) &= \{x \in H : x \in [\underline{y}, \bar{y}] \text{ and } x \geq T(x)\}. \end{aligned}$$

In the following section, we use this setting for $\mathbf{m}(T)$ and $\mathbf{M}(T)$ to establish stability results. We also provide an equivalent definition that is exploited subsequently.

5 Stability results

For the existence of optimal controls for our problem of interest, we need to study the stability of the maps $f \mapsto \mathbf{m}(f)$ and $f \mapsto \mathbf{M}(f)$. In the general case of an increasing map T , we now prove that $\mathbf{m}(T)$ and $\mathbf{M}(T)$ are stable from below and above, respectively, provided T has certain complete continuity properties.

Theorem 4. *Let $T, R_n, U_n : H \rightarrow V \subset H$ be increasing mappings with $n \in \mathbb{N}$. Assume further:*

- (i) *$T : V \rightarrow V$ is completely continuous with respect to monotone sequences, i.e., if $v_n \rightarrow v$ in V and $v_n \leq v_{n+1}$ (or $v_n \geq v_{n+1}$) for all $n \in \mathbb{N}$, then $T(v_n) \rightarrow T(v)$ in V .*

(ii) Sets of fixed points of T, R_n, U_n (assuming they exist) are uniformly bounded in V with respect to $n \in \mathbb{N}$, and that

$$\underline{y} \leq R_n(v) \leq R_{n+1}(v) \leq T(v) \leq U_{n+1}(v) \leq U_n(v) \leq \bar{y}, \quad \forall v \in [\underline{y}, \bar{y}], n \in \mathbb{N},$$

for some \underline{y} and \bar{y} in V .

(iii) If $\{v_n\}$ and $\{w_n\}$ are bounded sequences in V such that $v_n \leq v_{n+1} \leq \bar{y}$ and $w_n \geq w_{n+1} \geq \underline{y}$, then

$$\lim_{n \rightarrow \infty} \|R_n(v_n) - T(v_n)\|_V = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \|U_n(w_n) - T(w_n)\|_V = 0.$$

Let \mathbf{m} and \mathbf{M} be the operators that take an increasing map with sub- and supersolutions $[\underline{y}, \bar{y}]$ into the minimal and maximal solutions of Theorem 2, respectively. Then

$$\mathbf{m}(R_n) \rightarrow \mathbf{m}(T) \text{ in } V, \quad \text{and} \quad \mathbf{M}(U_n) \rightarrow \mathbf{M}(T) \text{ in } V,$$

and

$$\mathbf{m}(R_n) \uparrow \mathbf{m}(T) \text{ in } H, \quad \text{and} \quad \mathbf{M}(U_n) \downarrow \mathbf{M}(T) \text{ in } H,$$

as $n \rightarrow \infty$, respectively.

Proof. First note that since $\underline{y} \leq R_n(v) \leq T(v) \leq U_n(v) \leq \bar{y}$, the operators \mathbf{m} and \mathbf{M} are well defined on T, R_n and U_n for each $n \in \mathbb{N}$ since each of these maps is increasing with the same sub- and supersolutions. We introduce the sets

$$\begin{aligned} X(T) &= \{x \in H : x \in [\underline{y}, \bar{y}] \text{ and } x \leq T(x)\}, \\ Y(T) &= \{x \in H : x \in [\underline{y}, \bar{y}] \text{ and } x \geq T(x)\}, \\ Z(T) &= \{x \in X(T) : x \leq y \text{ for all } y \in Y(T)\}; \end{aligned}$$

and similarly for each R_n and U_n , $n \in \mathbb{N}$.

Since $R_n(v) \leq R_{n+1}(v) \leq T(v)$ for all $v \in [\underline{y}, \bar{y}]$ it follows that

$$X(R_n) \subset X(T) \text{ and } Y(T) \subset Y(R_n), \text{ and hence } Z(R_n) \subset Z(T), \quad (10)$$

and also

$$X(R_n) \subset X(R_{n+1}) \text{ and } Y(R_{n+1}) \subset Y(R_n), \text{ and hence } Z(R_n) \subset Z(R_{n+1}). \quad (11)$$

Clearly $Z(R_n)$ and $Z(T)$ are not empty, since \underline{y} belongs to either of them. Following the proof of Tartar's Theorem (compare [7]) we observe that $\mathbf{m}(R_n)$ and $\mathbf{m}(T)$ correspond to the maximal elements of $Z(R_n)$ and $Z(T)$, respectively. Consequently, it follows from (10) and (11) that

$$\mathbf{m}(R_n) \leq \mathbf{m}(R_{n+1}) \leq \mathbf{m}(T), \quad \forall n \in \mathbb{N}. \quad (12)$$

Hence, $\{\mathbf{m}(R_n)\}$ is a monotonically increasing sequence which is bounded from above (for the ordering ' \leq '), which implies that $\mathbf{m}(R_n) \rightarrow \hat{y}$ in H , for some $\hat{y} \in H$. We also know that the sets of fixed points of the maps are uniformly bounded in V . Therefore, we infer $\mathbf{m}(R_n) \rightarrow \hat{y}$ in V , that the

sequence is non-decreasing, and hence $T(\mathbf{m}(R_n)) \rightarrow T(\hat{y})$ in V . Since $\mathbf{m}(R_n) = R_n(\mathbf{m}(R_n))$ and

$$\lim_{j \rightarrow \infty} \|R_n(\mathbf{m}(R_n)) - T(\mathbf{m}(R_n))\|_V = 0,$$

we have $R_n(\mathbf{m}(R_n)) \rightarrow T(\hat{y})$. Therefore, $\mathbf{m}(R_n) \rightarrow T(\hat{y})$, but since $\mathbf{m}(R_n) \rightarrow \hat{y}$ it follows that $\mathbf{m}(R_n) \rightarrow \hat{y}$ in V , respectively, where \hat{y} is a fixed point of T .

Since $\mathbf{m}(R_n) \leq \mathbf{m}(T)$ for all n , we have $\hat{y} \leq \mathbf{m}(T)$. However, $\mathbf{m}(T)$ is the minimal fixed point of T , and therefore $\hat{y} = \mathbf{m}(T)$. Summarizing we have

$$\mathbf{m}(R_n) \rightarrow \mathbf{m}(T) \text{ in } V \quad \text{and} \quad \mathbf{m}(R_n) \uparrow \mathbf{m}(T) \text{ in } H.$$

Now we consider the upper bound. We define

$$\tilde{Z}(T) := \{y \in Y(T) : x \leq y \text{ for all } x \in X(T)\},$$

and analogously for U_n , $n \in \mathbb{N}$. Since $T(v) \leq U_{n+1}(v) \leq U_n(v)$ for all $v \in [\underline{y}, \bar{y}]$ and $n \in \mathbb{N}$ it follows that

$$X(T) \subset X(U_n) \text{ and } Y(U_n) \subset Y(T), \text{ and hence } \tilde{Z}(U_n) \subset \tilde{Z}(T), \quad (13)$$

and also

$$X(U_{n+1}) \subset X(U_n) \text{ and } Y(U_n) \subset Y(U_{n+1}) \text{ hence } \tilde{Z}(U_n) \subset \tilde{Z}(U_{n+1}). \quad (14)$$

Clearly, $\bar{y} \in \tilde{Z}(T)$, $\tilde{Z}(U_n)$ and then, as before, we apply Zorn's Lemma (with the reverse order) to find minimal elements $\mathbf{M}(T)$ and $\mathbf{M}(U_n)$, such that

$$\mathbf{M}(T) \leq \mathbf{M}(U_{n+1}) \leq \mathbf{M}(U_n) \leq \bar{y}.$$

Then, $\{-\mathbf{M}(U_n)\}$ is a monotonically increasing sequence which is bounded above for the ordering ' \leq '. This implies that $\mathbf{M}(U_n) \rightarrow \check{y}$ in H for some $\check{y} \in H$. Since $\{\mathbf{M}(U_n)\}$ is also uniformly bounded in V , we have $\mathbf{M}(U_n) \rightarrow \check{y}$ and this latter sequence is also non-increasing. Therefore, we infer $T(\mathbf{M}(U_n)) \rightarrow T(\check{y})$. Since $\mathbf{M}(U_n) = U_n(\mathbf{M}(U_n))$ and

$$\lim_{j \rightarrow \infty} \|U_n(\mathbf{M}(U_n)) - T(\mathbf{M}(U_n))\|_V = 0,$$

we get $U_n(\mathbf{M}(U_n)) \rightarrow T(\check{y})$ and $\mathbf{M}(U_n) \rightarrow \check{y}$, both in V , where $\check{y} = T(\check{y})$.

As in the previous case, since $\mathbf{M}(T) \leq \mathbf{M}(U_n)$, we have that $\mathbf{M}(T) \leq \check{y}$. However, $\mathbf{M}(T)$ is the maximal fixed point to T , and therefore $\check{y} = \mathbf{M}(T)$. Hence, we have

$$\mathbf{M}(U_n) \rightarrow \mathbf{M}(T) \text{ in } V \quad \text{and} \quad \mathbf{M}(U_n) \downarrow \mathbf{M}(T) \text{ in } H,$$

which ends the proof. □

This result is sharp regarding lower and upper approximations, as it is generally not possible to obtain $\mathbf{M}(R_n) \rightarrow \mathbf{M}(T)$ and $\mathbf{m}(U_n) \rightarrow \mathbf{m}(T)$. We illustrate this fact by means of the following one dimensional example.

Example 5.1. Let $T : [0, 1] \rightarrow [0, 1]$ be defined as

$$T(v) = \begin{cases} a, & 0 \leq v < a; \\ v, & a \leq v < b; \\ b, & b \leq v \leq 1. \end{cases}$$

with $0 < a < b < 1$ and where $\mathbf{m}(T) = a$ and $\mathbf{M}(T) = b$ and

$$R_n(v) = \begin{cases} a, & 0 \leq v < \frac{1}{n}, \\ T(v - \frac{1}{n}), & \frac{1}{n} \leq v \leq 1, \end{cases} \quad U_n(v) = \begin{cases} T(v + \frac{1}{n}), & 0 \leq v < 1 - \frac{1}{n}, \\ b, & 1 - \frac{1}{n} \leq v \leq 1. \end{cases}$$

Suppose that $n > N$ such that $\frac{1}{N} \leq a$ and $b \leq 1 - \frac{1}{N}$. Then, all the assumptions of the previous theorem hold, but $\mathbf{m}(R_n) = \mathbf{M}(R_n) = a$ and $\mathbf{m}(U_n) = \mathbf{M}(U_n) = b$ and hence $a = \mathbf{M}(R_n) \rightarrow \mathbf{M}(T) = b$ and $b = \mathbf{m}(U_n) \rightarrow \mathbf{m}(T) = a$ only hold for $a = b$, a contradiction.

Although, as observed in the previous example, a general approximation theorem (under the hypotheses of Theorem 4) for minimal and maximal fixed points seems elusive, we establish such a result for the specific case of the QVIs of interest. In order to achieve this, we first determine an equivalent definition of \mathbf{m} and \mathbf{M} but from slightly different means as in the Birkhoff-Tartar Theorem (see [7]).

Lemma 1. Let $T : H \rightarrow H$ be an increasing map with sub-solution \underline{y} and super-solution \bar{y} such that $\underline{y} \leq \bar{y}$. Then $\mathbf{m}(T)$, the maximal element of $Z(T)$, can also be defined as the maximal element of the set $Z^\bullet(T)$, which is defined as follows

$$\begin{aligned} X(T) &= \{x \in H : x \in [\underline{y}, \bar{y}] \text{ and } x \leq T(x)\}, \\ Y^\bullet(T) &= \{x \in H : x \in [\underline{y}, +\infty) \text{ and } x \geq T(x)\}, \\ Z^\bullet(T) &= \{x \in X(T) : x \leq y \text{ for all } y \in Y^\bullet(T)\}. \end{aligned}$$

Similarly, $\mathbf{M}(T)$, the minimal element of $\tilde{Z}(T)$, can also be defined as the minimal element of the set $\tilde{Z}^\bullet(T)$, defined as

$$\begin{aligned} X^\bullet(T) &= \{x \in H : x \in (-\infty, \bar{y}] \text{ and } x \leq T(x)\}, \\ Y(T) &= \{x \in H : x \in [\underline{y}, \bar{y}] \text{ and } x \geq T(x)\}, \\ \tilde{Z}^\bullet(T) &= \{y \in Y(T) : x \leq y \text{ for all } x \in X^\bullet(T)\}. \end{aligned}$$

Proof. We begin by noting that $\mathbf{m}(T)$ is the maximal element of $Z(T)$, and $\mathbf{M}(T)$ is the minimal element of $\tilde{Z}(T)$, as shown in the proof of the Birkhoff-Tartar Theorem (see [7, Chapter 15, §15.2]).

Since $\underline{y} \in Z^\bullet(T)$ and $Z^\bullet(T) \subset [\underline{y}, \bar{y}]$, $Z^\bullet(T)$ is nonempty and bounded in H , we may apply Zorn's Lemma (see [7]). Let $x^* \in Z^\bullet(T)$ be the maximal element of $Z^\bullet(T)$. It follows from $Y(T) \subset Y^\bullet(T)$ that $Z^\bullet(T) \subset Z(T)$. Therefore

$$x^* \leq \mathbf{m}(T),$$

where $\mathbf{m}(T)$ is the maximal element of $Z(T)$ and the minimum fixed point of T in $[\underline{y}, \bar{y}]$.

Since $x^* \in Z^\bullet(T)$, it follows by definition that $x^* \in X(T)$. Hence, we have $\underline{y} \leq x^* \leq \bar{y}$ and $x^* \leq T(x^*)$. Also, since T is an increasing map, it holds that $\underline{y} \leq T(\underline{y}) \leq T(x^*) \leq T(\bar{y}) \leq \bar{y}$ and $T(x^*) \leq T(T(x^*))$, i.e., $T(x^*) \in X(T)$. Furthermore, if $y \in Y^\bullet(T)$, then $x^* \leq y$. Hence, $T(x^*) \leq T(y) \leq y$, i.e., $T(x^*) \in Z^\bullet(T)$. However, $x^* \in Z^\bullet(T)$ is maximal, i.e., $T(x^*) \leq x^*$. Consequently, $x^* = T(x^*)$ and $x^* \in [\underline{y}, \bar{y}]$. Finally, $\mathbf{m}(T)$ is the minimal fixed point of T in $[\underline{y}, \bar{y}]$ so that

$$\mathbf{m}(T) \leq x^*.$$

Noting that $\tilde{Z}^\bullet(T) \subset [\underline{y}, \bar{y}]$ and $\bar{y} \in \tilde{Z}^\bullet(T)$, we can once again apply Zorn's Lemma (with the reversed order). Let x^* be the minimal element of $\tilde{Z}^\bullet(T)$. We have that $X(T) \subset X^\bullet(T)$ which implies $\tilde{Z}^\bullet(T) \subset \tilde{Z}(T)$. Therefore, it holds that

$$\mathbf{M}(T) \leq x^*,$$

where $\mathbf{M}(T)$ is the minimum element of $\tilde{Z}(T)$ and the maximum fixed point of T in $[\underline{y}, \bar{y}]$.

Since $x^* \in \tilde{Z}^\bullet(T)$, we have by definition that $x^* \in Y(T)$, i.e., $\underline{y} \leq x^* \leq \bar{y}$ and $T(x^*) \leq x^*$. Furthermore, the map T is increasing and therefore $\underline{y} \leq T(\underline{y}) \leq T(x^*) \leq T(\bar{y}) \leq \bar{y}$ and $T(T(x^*)) \leq T(x^*)$, i.e., $T(x^*) \in Y(T)$.

For an arbitrary $x \in X^\bullet(T)$, we have $x \leq x^*$ and $x \leq T(x) \leq T(x^*)$, i.e., $T(x^*) \in \tilde{Z}^\bullet(T)$. As x^* was the minimal element of $\tilde{Z}^\bullet(T)$, it follows that $x^* \leq T(x^*)$, yielding $T(x^*) = x^*$. However, $\mathbf{M}(T)$ is the maximal fixed point of T on $[\underline{y}, \bar{y}]$, so that

$$x^* \leq \mathbf{M}(T),$$

which completes the proof. \square

6 Monotone perturbations

We now prove a series of lemmas that are instrumental in establishing Theorem 5 in the subsequent section. The latter is a form of stability result for perturbations of the operators \mathbf{m} and \mathbf{M} . More specifically, it turns out that the minimal and maximal solutions of the map $y \mapsto S(f, \Phi(y))$ are stable in the norm of H with respect to perturbations in $L^\infty(\Omega) \hookrightarrow V'$ of the forcing term (under certain assumptions on Φ), i.e., if $\{f_n\}$ is in $L^\infty_\nu(\Omega)$ and $f_n \rightarrow f^*$ in $L^\infty(\Omega)$, then

$$\mathbf{m}(f_n) \rightarrow \mathbf{m}(f^*) \quad \text{and} \quad \mathbf{M}(f_n) \rightarrow \mathbf{M}(f^*) \quad \text{in } H.$$

The strategy of the proof consists in considering the cases of increasing and decreasing sequences of $\{f_n\}$ separately and then combine both cases to obtain the final result. This strategy is due to the different nature of these cases as indicated in Theorem 4 and Example 5.1. It can also be corroborated by the different structural hypotheses of Lemma 1, 2, 3 and 4. As expected, stability results associated to one-sided perturbations are more amenable than general ones.

In this section, all sequences of forcing terms $\{f_n\}$ are assumed to satisfy $0 \leq f_n \leq F$ for all $n \in \mathbb{N}$ and some $F \in V'$ such $F \geq 0$. Further, we consider the interval $[\underline{y}, \bar{y}]$, with $\underline{y} = 0$, and $\bar{y} \in V$ such that

$$\langle A(\bar{y}), v \rangle = \langle F, v \rangle, \quad \forall v \in V. \quad (15)$$

For any f with $0 \leq f \leq F$, we observe that

$$0 \leq S(f, \Phi(0)) \quad \text{and} \quad S(f, \Phi(\bar{y})) \leq S(F, +\infty) = \bar{y}.$$

Hence, we denote by $\mathbf{m}(f)$ and $\mathbf{M}(f)$ the minimal and maximal fixed points of the map $y \mapsto S(f, \Phi(y)) = S(f, y)$, respectively, on the interval $[\underline{y}, \bar{y}] = [0, A^{-1}(F)]$. Note that $\mathbf{m}(f)$ and $\mathbf{M}(f)$ are well defined according to Theorem 3.

In the following lemma, we start by considering the behavior of $\{\mathbf{m}(f_n)\}$ for non-increasing sequences $\{f_n\}$.

Lemma 1 (NON-INCREASING SEQUENCES OF \mathbf{m}). *Suppose that the following hold true:*

- (i) *The sequence $\{f_n\}$ in $L^\infty_\nu(\Omega)$ is non-increasing and $\lim_{n \rightarrow \infty} f_n = f^*$ in $L^\infty(\Omega)$ for some $f^* \in L^\infty_\nu(\Omega)$.*

(ii) The upper bound mapping Φ satisfies

$$\lambda\Phi(y) \geq \Phi(\lambda y), \quad \text{for all } \lambda \geq 1, y \in V \cap H^+,$$

and if $\{v_n\}$ is bounded in V and $v_n \downarrow v$ in H , then $\Phi(v_n) \rightarrow \Phi(v)$ in H .

Then, it follows that

$$\mathbf{m}(f_n) \downarrow \mathbf{m}(f^*) \text{ in } H, \quad \text{and} \quad \mathbf{m}(f_n) \rightarrow \mathbf{m}(f^*) \text{ in } V. \quad (16)$$

Proof. The proof is split into several steps for the sake of clarity.

Step 1: We start by showing: *If a sequence $\{z_n\}$ satisfies $z_n \rightarrow z^*$ in V , for some z^* , and is non-increasing and non-negative: $z_n \geq z_{n+1} \geq 0$ for all $n \in \mathbb{N}$, then it holds that $S(f_n, \Phi(z_n)) \rightarrow S(f^*, \Phi(z^*))$ in V .* We follow closely the ideas in [52] and include the proof here for the sake of completeness. Note first that the non-increasing nature of the sequence implies also that $z_n \downarrow z^*$ in H .

By our hypothesis on Φ , we have that $\Phi(z_n) \rightarrow \Phi(z^*)$ in H and $\Phi(z_n) \geq \Phi(z_{n+1})$, which implies that $\mathbf{K}(\Phi(z_n)) \supset \mathbf{K}(\Phi(z_{n+1}))$ and $\mathbf{K}(\Phi(z_n)) \supset \mathbf{K}(\Phi(z^*))$.

Since $\mathbf{K}(\Phi(z^*))$ is non-empty (note that $z^* \geq 0$ and $\Phi(z^*) \geq 0$), we have $0 \in \mathbf{K}(\Phi(z^*))$ and $0 \in \mathbf{K}(\Phi(z_n))$ for all $n \in \mathbb{N}$. Let $w_n := S(f_n, \Phi(z_n))$ and note that by Proposition 1 the associated sequence is decreasing and bounded from below (in the ordering), so that $w_n \downarrow w^*$ in H for some $w^* \in H$.

By definition $\langle Aw_n - f_n, v - w_n \rangle \geq 0$ for all $v \in \mathbf{K}(\Phi(z_n))$, and then, using the uniform monotonicity of A , we have

$$c\|w_n\|_V^2 \leq \langle Aw_n, w_n \rangle \leq \langle f_n, w_n \rangle. \quad (17)$$

Also, $\langle f_n, w_n \rangle \leq (\|f_n\|_{V'})\|w_n\|_V$ and $\|f_n\|_{V'} \leq C\|f_n\|_{L^\infty(\Omega)} < \infty$. Therefore, $\{w_n\}$ is bounded in V , and hence for some subsequence $w_{n_k} \rightarrow w^*$ in V (for the same w^* as before). But since $w_n \downarrow w^*$ in H , we get $w_n \rightarrow w^*$ in V .

Since $\Phi(z_n) \rightarrow \Phi(z^*)$ in H and $w_n \leq \Phi(z_n)$, we conclude

$$w^* \leq \Phi(z^*), \quad \text{i.e.,} \quad w^* \in \mathbf{K}(\Phi(z^*)).$$

By Minty's Lemma (see [48]) applied to the VI arising from $w_n = S(f_n, \Phi(z_n))$, we obtain

$$\langle Av - f_n, v - w_n \rangle \geq 0, \quad \forall v \in \mathbf{K}(\Phi(z_n)),$$

and in particular for all $v \in \mathbf{K}(\Phi(z^*)) \subset \mathbf{K}(\Phi(z_n))$. As $f_n \rightarrow f^*$ in $L^\infty(\Omega)$ (and hence in V'), $(v - w_n) \rightarrow (v - w^*)$ in V and $(v - w_n) \rightarrow (v - w^*)$ in H , we have

$$\lim_{k \rightarrow \infty} \langle Av - f_{n_k}, v - w_{n_k} \rangle = \langle Av - f^*, v - w^* \rangle \geq 0, \quad \forall v \in \mathbf{K}(\Phi(z^*)).$$

Additionally, since $w^* \in \mathbf{K}(\Phi(z^*))$, Minty's Lemma implies

$$\langle Aw^* - f^*, v - w^* \rangle \geq 0, \quad \forall v \in \mathbf{K}(\Phi(z^*)),$$

i.e., $w^* = S(f^*, \Phi(z^*))$.

Given that $w_n \rightarrow w^*$ in V and $\|f_n\|_{V'} \leq C\|f_n\|_{L^\infty(\Omega)} < \infty$, by

$$c\|w_n - w^*\|_V^2 \leq \langle Aw_n - Aw^*, w_n - w^* \rangle \leq \langle f_n + Aw^*, w^* - w_n \rangle,$$

we have $w_n \rightarrow w^*$ in V . That is,

$$S(f_n, \Phi(z_n)) \rightarrow S(f^*, \Phi(z^*)) \text{ in } V. \quad (18)$$

Before we continue with the next step of the proof, we define for $f \in V'$ the set-valued mappings

$$\begin{aligned} X(f) &= \{x \in H : \underline{y} \leq x \leq \bar{y} \text{ and } x \leq S(f, \Phi(x))\}, \\ Y^\bullet(f) &= \{x \in H : \underline{y} \leq x \text{ and } x \geq S(f, \Phi(x))\}, \\ Z^\bullet(f) &= \{x \in X(f) : x \leq y \text{ for all } y \in Y^\bullet(f)\}. \end{aligned}$$

Step 2: Let $\{z_n\}$ be the sequence of Step 1, i.e., $z_n \rightarrow z^$ in V that is also non-increasing in the sense $z_n \geq z_{n+1} \geq 0$ for all n . If $z_n \in Z^\bullet(f_n)$, then $z^* \in Z^\bullet(f^*)$.*

Since $f_n \in L^\infty_\nu(\Omega)$ with $f_n \geq f_{n+1}$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} f_n = f^* \geq \nu > 0$ in $L^\infty(\Omega)$, we have that $f^* \leq f_n$ for all $n \in \mathbb{N}$. Hence, $S(f^*, \Phi(x)) \leq S(f_n, \Phi(x))$ for all $n \in \mathbb{N}$ and we obtain the inequalities

$$X(f^*) \subset X(f_n) \text{ and } Y^\bullet(f_n) \subset Y^\bullet(f^*), \text{ and hence } Z^\bullet(f^*) \subset Z^\bullet(f_n). \quad (19)$$

Let $z_n \in Z^\bullet(f_n)$, then $z_n \in X(f_n)$, i.e., $\underline{y} \leq z_n \leq \bar{y}$ and $z_n \leq S(f_n, \Phi(z_n))$. Therefore, we have

$$\underline{y} \leq z^* \leq \bar{y} \text{ and } z^* \leq S(f^*, \Phi(z^*)), \text{ and hence } z^* \in X(f^*). \quad (20)$$

Let $y \in Y^\bullet(f^*)$ be arbitrary and consider $y_n := \lambda_n y$, with $\lambda_n := \|f_n/f\|_{L^\infty(\Omega)}$. Since $\lambda_n \downarrow 1$ (recall $f_n \rightarrow f$ in $L^\infty(\Omega)$ and $f_n, f \in L^\infty_\nu(\Omega)$ for all $n \in \mathbb{N}$), we infer $\underline{y} \leq \lambda_n \underline{y} \leq \lambda_n y = y_n$. Also, $\lambda_n f \geq f_n$ and by the structural assumption over Φ , we have $\lambda_n \Phi(y) \geq \Phi(\lambda_n y)$. Furthermore, we obtain the following chain of inequalities

$$\lambda_n y \geq \lambda_n S(f, \Phi(y)) = S(\lambda_n f, \lambda_n \Phi(y)) \geq S(f_n, \Phi(\lambda_n y))$$

i.e., $y_n \in Y^\bullet(f_n)$ and $y_n \rightarrow y$ in $L^\infty(\Omega)$.

Now, we have that $z_n \in Z^\bullet(f_n)$ and $z_n(x) \rightarrow z^*(x)$ a.e. in Ω and $z^* \in X(f^*)$, and for each $n \in \mathbb{N}$ we have $z_n \leq \tilde{y}$ for all $\tilde{y} \in Y^\bullet(f_n)$. Choosing $\tilde{y} = \lambda_n y$ as in the previous paragraph with $y \in Y^\bullet(f^*)$ arbitrary, we have that $z_n \leq \lambda_n y$. Henceforth, $z^* \leq y$ for all $y \in Y^\bullet(f^*)$, i.e., $z^* \in Z^\bullet(f^*)$.

Step 3. The minimal solutions $\mathbf{m}(f_n)$ and $\mathbf{m}(f^*)$ are well defined as the maximal elements of $Z^\bullet(f_n)$ and $Z^\bullet(f^*)$, respectively. It follows immediately from (19) that $\mathbf{m}(f^*) \leq \mathbf{m}(f_n)$, and by the same argument used to derive (19), we have that $0 \leq \mathbf{m}(f_{n+1}) \leq \mathbf{m}(f_n)$. Denote $z_n = \mathbf{m}(f_n)$, since $z_n = S(f_n, \Phi(z_n))$ and $0 \in \mathbf{K}(\Phi(z_n))$, a standard monotonicity argument gives $\|z_n\|_V \leq M < \infty$. Hence z_n is bounded in V , non-increasing and bounded below in order, and $z_n \in Z^\bullet(f_n)$. The monotone behaviour in addition to the boundedness implies that $z_n \rightarrow z^*$ in V , by Step 2 we have that $z^* \in Z^\bullet(f^*)$. Since $z_n = S(f_n, \Phi(z_n))$, by Step 1, we have that $z_n \rightarrow z^*$ in V , $z^* = S(f^*, \Phi(z^*))$, i.e., z^* is a fixed point of the map $z \mapsto S(f^*, \Phi(z))$ and hence $\mathbf{m}(f^*) \leq z^*$. From the definition of $Z^\bullet(f^*)$ we infer that $z^* \leq y$ for all $y \in Y^\bullet(f^*)$, and we readily observe $\mathbf{m}(f^*) \in Y^\bullet(f^*)$, so that $z^* \leq \mathbf{m}(f^*)$, i.e., $\mathbf{m}(f^*) = z^*$. \square

A fundamental step in the previous lemma utilizes that

$$S(f_n, \Phi(z_n)) \rightarrow S(f^*, \Phi(z^*)), \quad \text{in } V$$

when $f_n \rightarrow f$ in V' . A sufficient condition for this to hold true is related to the Mosco convergence (see [48]) of $\{\mathbf{K}(\Phi(z_n))\}$ towards $\mathbf{K}(\Phi(z^*))$:

Definition 1 (MOSCO CONVERGENCE). *Let \mathbf{K} and \mathbf{K}_n , for each $n \in \mathbb{N}$, be non-empty, closed and convex subsets of V . Then the sequence $\{\mathbf{K}_n\}$ is said to converge to \mathbf{K} in the sense of Mosco as $n \rightarrow \infty$, denoted by $\mathbf{K}_n \xrightarrow{M} \mathbf{K}$, if the following two conditions are fulfilled:*

- (i) *For each $w \in \mathbf{K}$, there exists $\{w_{n'}\}$ such that $w_{n'} \in \mathbf{K}_{n'}$ for $n' \in \mathbb{N}' \subset \mathbb{N}$ and $w_{n'} \rightarrow w$ in V .*
- (ii) *If $w_n \in \mathbf{K}_n$ and $w_n \rightarrow w$ in V along a subsequence, then $w \in \mathbf{K}$.*

Mosco convergence of unilaterally constrained sets is equivalent (in the case when the obstacles are quasi-continuous and V a certain Sobolev space) to convergence of the obstacles in the sense of the capacity (which might be cumbersome to prove beyond rather simple examples). It is also well-known that the convergence of the obstacles in the sense of $L^\infty(\Omega)$ is a sufficient condition for Mosco convergence, although this might be rather a strong assumption for some applications. In the previous case, we are able to avoid that strong assumption rather elegantly by assuming only the H convergence of the obstacles. In the next case, for non-increasing sequences, the $L^\infty(\Omega)$ or V convergence might be avoided by using an argument of Toyoizumi (see [52]) by using geometrical considerations of the obstacles. For this matter, we consider the following assumption.

Assumption 1. *If $v_n \rightarrow v$ in V , then Φ satisfies one of the following:*

- (a) *$\Phi(v_n) \rightarrow \Phi(v)$ in $L^\infty(\Omega)$, or $\Phi(v_n) \rightarrow \Phi(v)$ in V .*
- (b) *$\Phi(v_n) \rightarrow \Phi(v)$ in H and if $v \in V \cap H^+$, then $\Phi(v) \in V$ and $\mathcal{Q}\Phi(v) \geq 0$ in V , for some strongly monotone $\mathcal{Q} \in \mathcal{L}(V, V')$, such that $\langle \mathcal{Q}v^-, v^+ \rangle \leq 0$ for all $v \in V$.*

With the above definition in mind we are now in the position to provide the stability result for minimal solutions and for non-decreasing sequences of forcing terms.

Lemma 2 (NON-DECREASING SEQUENCES FOR \mathbf{m}). *Suppose the following:*

- (i) *The sequence $\{f_n\}$ in H^+ is non-decreasing and $\lim_{n \rightarrow \infty} f_n = f^*$ in H for some $f^* \in H$.*
- (ii) *The upper bound mapping Φ satisfies Assumption 1.*

Then, the following hold true:

$$\mathbf{m}(f_n) \uparrow \mathbf{m}(f^*) \text{ in } H, \quad \text{and} \quad \mathbf{m}(f_n) \rightarrow \mathbf{m}(f^*) \text{ in } V.$$

Proof. We use the result of Theorem 4 with $R_n(v) := S(f_n, \Phi(v))$ and $S(v) := S(f^*, \Phi(v))$. The classical continuity result for $f \mapsto S(f, \Phi(y))$ (see [37]) states:

$$\|S(f^*, \Phi(y)) - S(f_n, \Phi(y))\|_V \leq \frac{1}{c} \|f^* - f_n\|_{V'}. \quad (21)$$

Since $f_n \rightarrow f^*$ in V' as $n \rightarrow \infty$, we have $S(f_n, \Phi(y)) \rightarrow S(f^*, \Phi(y))$ in V , uniformly on bounded sets for y . Additionally, by the usual monotonicity argument and using $v = 0$ as a test function, we obtain that $\|S(f, \Phi(y))\|_V \leq \frac{1}{c}\|f\|_{V'}$ which implies that the set of fixed points of the maps $y \mapsto S(f_n, \Phi(y))$, for $n \in \mathbb{N}$, and $y \mapsto S(f^*, \Phi(y))$ is uniformly bounded. Since $S(f_n, \Phi(y)) \leq S(f_{n+1}, \Phi(y)) \leq S(f^*, \Phi(y))$ we are only left to prove that

$$\lim_{n \rightarrow \infty} S(f_n, \Phi(v_n)) = S(f^*, \Phi(v)) \text{ in } V. \quad (22)$$

This will be achieved by proving Mosco convergence of the associated constraints.

Now we consider the two possible cases for Φ based on Assumption 1:

- (a) Suppose $\{v_n\}$ in V satisfies $v_n \rightarrow v$ in V and $\Phi(v_n) \rightarrow \Phi(v)$ in $L^\infty(\Omega)$. It is well-known that the latter implies $\mathbf{K}(\Phi(v_n)) \rightarrow \mathbf{K}(\Phi(v))$ in the sense of Mosco (see for example [48]).

Suppose that $\Phi(v_n) \rightarrow \Phi(v)$ in V . Let $w \leq \Phi(v)$, and consider $w_n = w - \Phi(v) + \Phi(v_n)$. Then, $w_n \leq \Phi(v_n)$ and also $w_n \rightarrow w$ in V , i.e., (i) in Definition 1 holds. Furthermore, if $y_n \leq \Phi(v_n)$ and $y_n \rightarrow y$ in V , then by Mazur's lemma it follows that $y \leq \Phi(v)$ which proves (ii) in Definition 1.

- (b) Suppose that $\{v_n\}$ in V satisfies $0 \leq v_n \leq v_{n+1}$ for all n , and $v_n \rightarrow v$ in V . Then $0 \leq \Phi(v_n) \leq \Phi(v_{n+1})$ since Φ is increasing and $\Phi(v_n) \rightarrow \Phi(v)$ in H by hypothesis. Hence, if $y_n \leq \Phi(v_n)$ and $y_n \rightarrow y$ in V , then by Mazur's lemma it follows that $y \leq \Phi(v)$, which proves (ii) in Definition 1.

In order to prove (i) in Definition 1, we now follow a modification of the argument in [52]. Let $w \in V$ such that $w \leq \Phi(v)$ and w_n be defined by

$$\langle r_n \mathcal{Q}w_n + w_n, v \rangle = \langle \phi_n, v \rangle, \text{ for all } v \in V, \quad (23)$$

where $r_n := \|\phi_n - w\|_H$ and $\phi_n := \min(w, \Phi(v_n))$, and note that $\phi_n \rightarrow w$ in H and $w \in V$. Then, we can prove that $w_n \rightarrow w$ in V . Since \mathcal{Q} is linear, bounded, and $\langle \mathcal{Q}v, v \rangle \geq c\|v\|_V^2$ for all $v \in V$, from the definition of w_n we have

$$\begin{aligned} r_n c \|w_n - w\|_V^2 + \|w_n - w\|_H^2 &\leq \langle (r_n \mathcal{Q} + I)(w_n - w), w_n - w \rangle \\ &\leq \langle \phi_n - w, w_n - w \rangle - r_n \langle \mathcal{Q}w, w_n - w \rangle \\ &\leq r_n (C_p + \|\mathcal{Q}w\|_{V'}) \|w_n - w\|_V, \end{aligned} \quad (24)$$

where C_p is the constant for the embedding $V \hookrightarrow H$. This implies that, $\{w_n\}$ is bounded in V , so that $w_n \rightarrow w^*$ (along a subsequence) for some $w^* \in V$. By taking the limit in (23), it is shown that $w^* = w$ and that $w_n \rightarrow w^*$ in V not only along a subsequence. It further follows that $w_n \rightarrow w$ in H , and since from (24) we observe

$$r_n c \|w_n - w\|_V^2 + \|w_n - w\|_H^2 \leq r_n (\|w_n - w\|_H + \langle \mathcal{Q}w, w - w_n \rangle), \quad (25)$$

we have that $w_n \rightarrow w$ in V .

Now we prove that $w_n \leq \Phi(v_n)$. Consider $v = (w_n - \Phi(v_n))^+$ and subtract $\langle r_n \mathcal{Q}\Phi(v_n) + \Phi(v_n), v \rangle$ from both sides of (23). Then, we get

$$\begin{aligned} r_n \langle \mathcal{Q}(w_n - \Phi(v_n)), (w_n - \Phi(v_n))^+ \rangle + \|(w_n - \Phi(v_n))^+\|_H^2 = \\ - r_n \langle \mathcal{Q}\Phi(v_n), (w_n - \Phi(v_n))^+ \rangle + (\min(w, \Phi(v_n)) - \Phi(v_n), (w_n - \Phi(v_n))^+). \end{aligned}$$

Note that $\min(w, \Phi(v_n)) - \Phi(v_n) \leq 0$ and by assumption $Q\Phi(v_n) \geq 0$. Therefore the right hand side is less or equal to zero. Additionally, since Q is linear, $\langle Qv^-, v^+ \rangle \leq 0$, and $\langle Qv, v \rangle \geq c\|v\|_V^2$ for all $v \in V$, we observe that

$$\begin{aligned} r_n c \| (w_n - \Phi(v_n))^+ \|_V^2 + \| (w_n - \Phi(v_n))^+ \|_H^2 \leq \\ r_n \langle Q(w_n - \Phi(v_n))^+, (w_n - \Phi(v_n))^+ \rangle + \| (w_n - \Phi(v_n))^+ \|_H^2 \leq 0. \end{aligned}$$

This yields $w_n \leq \Phi(v_n)$, i.e., (i) in Definition 1 holds

This completes the proof. \square

Lemma 1 and 2 are associated to non-increasing and non-decreasing sequences of minimal solutions. In the following we establish Lemma 3 and 4 that deal with the analogous results but for maximal solutions.

Lemma 3 (NON-INCREASING SEQUENCES FOR \mathbf{M}). *Suppose the following:*

- (i) *The sequence $\{f_n\}$ in H^+ is non-increasing and $\lim_{n \rightarrow \infty} f_n = f^*$ in H for some $f^* \in H$.*
- (ii) *The upper bound mapping Φ satisfies that if $\{v_n\}$ is bounded in V , $v_n \downarrow v$ in H , then $\Phi(v_n) \rightarrow \Phi(v)$ in H .*

Then, we have

$$\mathbf{M}(f_n) \downarrow \mathbf{M}(f^*) \text{ in } H, \quad \text{and} \quad \mathbf{M}(f_n) \rightarrow \mathbf{M}(f^*) \text{ in } V. \quad (26)$$

Proof. As obtained in the proof of Lemma 2, we have that $S(f_n, \Phi(y)) \rightarrow S(f^*, \Phi(y))$ in V and that the set of fixed points of the maps $y \mapsto S(f_n, \Phi(y))$, for $n \in \mathbb{N}$, and $y \mapsto S(f^*, \Phi(y))$ are uniformly bounded in V .

Let $\{v_n\}$ be such that $v_n \rightarrow v$ in V and $v_n \geq v_{n+1} \geq 0$ for all n , then $v_n \rightarrow v$ in H , $\Phi(v_n) \geq \Phi(v_{n+1}) \geq 0$ and $\Phi(v_n) \rightarrow \Phi(v)$ in H . Note that $f_n \rightarrow f^*$ in H is enough for step 1 of the proof of Lemma 1 to hold, i.e.,

$$\lim_{n \rightarrow \infty} S(f_n, \Phi(v_n)) = S(f, \Phi(v)), \quad \text{in } V.$$

Therefore applying Theorem 4 to $T_n(v) := S(f_n, \Phi(v))$ and $S(v) := S(f^*, \Phi(v))$, we obtain that (26) holds true. \square

Lemma 4 (NON-DECREASING SEQUENCES FOR \mathbf{M}). *Suppose the following:*

- (i) *The sequence $\{f_n\}$ in $L^\infty_\nu(\Omega)$ is non-decreasing and $\lim_{n \rightarrow \infty} f_n = f^*$ in $L^\infty(\Omega)$ for some f^* .*
- (ii) *The upper bound mapping Φ satisfies*

$$\lambda \Phi(y) \leq \Phi(\lambda y), \quad \text{for all } 0 < \lambda < 1, y \in V \cap H^+,$$

and Assumption 1.

Then, we have

$$\mathbf{M}(f_n) \uparrow \mathbf{M}(f^*) \text{ in } H, \quad \text{and} \quad \mathbf{M}(f_n) \rightarrow \mathbf{M}(f^*) \text{ in } V. \quad (27)$$

Proof. For f define the set-valued mappings

$$\begin{aligned} X^\bullet(f) &= \{x \in H : x \leq \bar{y} \text{ and } x \leq S(f, \Phi(x))\}, \\ Y(f) &= \{x \in H : \underline{y} \leq x \leq \bar{y} \text{ and } x \geq S(f, \Phi(x))\}, \\ \tilde{Z}^\bullet(f) &= \{y \in Y(f) : x \leq y \text{ for all } x \in X^\bullet(f)\}. \end{aligned}$$

If $\{z_n\}$ satisfies $z_n \rightarrow z^*$ in V for some $z^* \in V$ and is non-decreasing, i.e., $z_n \leq z_{n+1}$ for all $n \in \mathbb{N}$, then

$$S(f_n, \Phi(z_n)) \rightarrow S(f^*, \Phi(z^*)), \quad \text{in } V, \quad (28)$$

as proven in Lemma 2. We now show that if $z_n \in \tilde{Z}^\bullet(f_n)$, for all $n \in \mathbb{N}$, then $z^* \in \tilde{Z}^\bullet(f^*)$.

Since $f_n \leq f_{n+1}$, for all $n \in \mathbb{N}$, and $\lim_{n \rightarrow \infty} f_n = f^*$ in $L^\infty(\Omega)$, we have that $f_n \leq f^*$ and $S(f_n, \Phi(x)) \leq S(f^*, \Phi(x))$, for all $n \in \mathbb{N}$. Therefore,

$$X^\bullet(f_n) \subset X^\bullet(f^*) \text{ and } Y(f^*) \subset Y(f_n), \text{ and hence } \tilde{Z}^\bullet(f^*) \subset \tilde{Z}^\bullet(f_n). \quad (29)$$

Also, $z_n \in \tilde{Z}^\bullet(f_n)$ and hence $z_n \in Y(f_n)$, i.e., $\underline{y} \leq z_n \leq \bar{y}$ and $z_n \geq S(f_n, \Phi(z_n))$. Therefore, by (28) and since $z_n \rightarrow z^*$ in H (note that $z_n \leq z_{n+1} \leq \bar{y}$) we observe that

$$\underline{y} \leq z^* \leq \bar{y} \text{ and } z^* \geq S(f^*, \Phi(z^*)), \text{ and hence } z^* \in Y(f^*). \quad (30)$$

Let $x \in X^\bullet(f^*)$ be arbitrary and consider $x_n := \lambda_n x$, with $\lambda_n := \text{essinf} |f_n/f^*|$ which yields $\lambda_n \uparrow 1$. Indeed, since $f_n \leq f_{n+1}$, we have $|f_n/f^*| \leq |f_{n+1}/f^*| \leq 1$ and

$$\left| 1 - \text{essinf} \left| \frac{f_n}{f^*} \right| \right| = \text{essinf} \left| 1 - \left| \frac{f_n}{f^*} \right| \right| \leq \frac{\|f^* - f_n\|_{L^\infty(\Omega)}}{\nu},$$

where we have used that $f^* \in L^\infty_\nu(\Omega)$, and the result follows from the assumed convergence $f_n \rightarrow f$ in $L^\infty(\Omega)$.

Therefore, $x_n = \lambda_n x \leq \lambda_n \bar{y} \leq \bar{y}$, $\lambda_n f \leq f_n$, and by the structural assumption on Φ , we have $\lambda_n \Phi(y) \leq \Phi(\lambda_n y)$. Furthermore, we obtain the following chain of inequalities:

$$\lambda_n x \leq \lambda_n S(f, \Phi(x)) = S(\lambda_n f, \lambda_n \Phi(x)) \leq S(f_n, \Phi(\lambda_n x)),$$

i.e., $x_n \in X^\bullet(f_n)$ and $x_n \rightarrow x$ in H .

Since $z_n \in \tilde{Z}^\bullet(f_n)$, we have $x_n \leq z_n$ given the fact that $x_n \in X^\bullet(f_n)$. Additionally, along a subsequence we have that $x_n \rightarrow x$ and $z_n \rightarrow z$ in H so that $x \leq z^*$. However, $x \in X^\bullet(f^*)$ was arbitrary and hence, by (30), $z^* \in \tilde{Z}^\bullet(f^*)$.

Since $\mathbf{M}(f_n)$ and $\mathbf{M}(f^*)$ are well-defined as the minimal elements of $\tilde{Z}^\bullet(f_n)$ and $\tilde{Z}^\bullet(f^*)$, respectively, it follows immediately from (29) that $\mathbf{M}(f_n) \leq \mathbf{M}(f^*)$, and furthermore, we have that $\mathbf{M}(f_n) \leq \mathbf{M}(f_{n+1})$. Denoting $z_n = \mathbf{M}(f_n)$, we have $z_n = S(f_n, \Phi(z_n))$, and since $0 \in \mathbf{K}(\Phi(z_n))$, a monotonicity argument gives $\|z_n\|_V \leq \frac{1}{c} \|f_n\|_{V'} \leq \frac{1}{c} \|F\|_{V'} < \infty$. Hence, z_n is bounded in V , non-decreasing in order and $z_n \in \tilde{Z}^\bullet(f_n)$. Therefore, by the above paragraphs, we have that $z_n = \mathbf{M}(f_n) \rightarrow z^*$ in V and $z^* \in \tilde{Z}^\bullet(f^*)$ and additionally, since $z_n = S(f_n, \Phi(z_n))$, by (28), we have that $z_n \rightarrow z^*$ in V , $z^* = S(f^*, \Phi(z^*))$, i.e., z^* is a fixed point of the map $z \mapsto S(f^*, \Phi(z))$ and hence $z^* \leq \mathbf{M}(f^*)$. By definition of $\tilde{Z}^\bullet(f^*)$, we have that $x \leq z^*$ for all $x \in X^\bullet(f^*)$ and we readily observe $\mathbf{M}(f^*) \in X^\bullet(f^*)$, so that $\mathbf{M}(f^*) \leq z^*$, i.e., $\mathbf{M}(f^*) = z^*$. \square

7 Non-monotone perturbations and problem (\mathbb{P})

We are now in the position to establish our fundamental result concerning the behavior of the maps $f \mapsto \mathbf{m}(f)$ and $f \mapsto \mathbf{M}(f)$. Although the hypotheses of lemmas 1, 2, 3 and 4 seem to be quite diverse, when considering the intersection in the following theorem, the assumptions are simplified. As in the previous section we assume that $0 \leq f_n \leq F$ for any sequence $\{f_n\}$ and that $[\underline{y}, \bar{y}] = [0, A^{-1}F]$.

Theorem 5. *Let $\{f_n\}$ in $L^\infty_\nu(\Omega)$ be such that $\lim f_n = f^*$ in $L^\infty(\Omega)$ for some f^* , and suppose that the upper bound mapping $\Phi : V \cap H^+ \rightarrow H^+$ satisfies Assumption 1. Then the following hold true:*

(i) *If $\lambda\Phi(y) \geq \Phi(\lambda y)$ for any $\lambda > 1$ and any $y \in V \cap H^+$, we have*

$$\mathbf{m}(f_n) \rightarrow \mathbf{m}(f^*) \text{ in } H, \quad \text{and} \quad \mathbf{m}(f_n) \rightharpoonup \mathbf{m}(f^*) \text{ in } V. \quad (31)$$

(ii) *If $\lambda\Phi(y) \leq \Phi(\lambda y)$ for any $0 < \lambda < 1$ and any $y \in V \cap H^+$, we have*

$$\mathbf{M}(f_n) \rightarrow \mathbf{M}(f^*) \text{ in } H, \quad \text{and} \quad \mathbf{M}(f_n) \rightharpoonup \mathbf{M}(f^*) \text{ in } V. \quad (32)$$

Proof. Define $\hat{f}_n := \inf_{m \geq n} f_m$ and $\check{f}_n := \sup_{m \geq n} f_m$, so that $0 \leq \nu \leq \hat{f}_n \leq \hat{f}_{n+1} \leq F$, $F \geq \check{f}_n \geq \check{f}_{n+1} \geq \nu > 0$ for all $n \in \mathbb{N}$, and also $\lim_{n \rightarrow \infty} \hat{f}_n = \lim_{n \rightarrow \infty} \check{f}_n = f^*$ in $L^\infty(\Omega)$. Since $0 \leq \hat{f}_n \leq f_n \leq \check{f}_n \leq F$ and the map $H \ni y \mapsto S(f, \Phi(y))$ is increasing for any $f \in V'$, we have that $\mathbf{m}(\hat{f}_n)$, $\mathbf{m}(f_n)$, $\mathbf{m}(\check{f}_n)$ and $\mathbf{m}(f^*)$ as well as $\mathbf{M}(\hat{f}_n)$, $\mathbf{M}(f_n)$, $\mathbf{M}(\check{f}_n)$ and $\mathbf{M}(f^*)$ are well defined (note that $0 \leq f^* \leq F$), respectively. Moreover, we have that

$$0 \leq S(\hat{f}_n, \Phi(y)) \leq S(f_n, \Phi(y)) \leq S(\check{f}_n, \Phi(y)) \leq \bar{y} \quad \forall y \in [0, \bar{y}], n \in \mathbb{N}.$$

Hence from the inclusions (10) and (11), we obtain

$$0 \leq \mathbf{m}(\hat{f}_n) \leq \mathbf{m}(f_n) \leq \mathbf{m}(\check{f}_n) \leq \bar{y}, \quad \forall n \in \mathbb{N}, \quad (33)$$

and from the inclusions (13) and (14) that

$$0 \leq \mathbf{M}(\hat{f}_n) \leq \mathbf{M}(f_n) \leq \mathbf{M}(\check{f}_n) \leq \bar{y}, \quad \forall n \in \mathbb{N}. \quad (34)$$

Then, by lemmas 1, 2, 3 and 4 we have that $\mathbf{m}(\hat{f}_n) \rightarrow \mathbf{m}(f^*)$, $\mathbf{m}(\check{f}_n) \rightarrow \mathbf{m}(f^*)$, $\mathbf{M}(\hat{f}_n) \rightarrow \mathbf{M}(f^*)$ and $\mathbf{M}(\check{f}_n) \rightarrow \mathbf{M}(f^*)$, all in V and H . Hence, we find

$$\mathbf{m}(f_n) \rightarrow \mathbf{m}(f^*) \text{ in } H \quad \text{and} \quad \mathbf{M}(f_n) \rightarrow \mathbf{M}(f^*) \text{ in } H$$

by (33) and (34). Since $\{\mathbf{m}(f_n)\}$ and $\{\mathbf{M}(f_n)\}$ are bounded in V , they are also weakly convergent (along a subsequence) to $\mathbf{m}(f^*)$ and $\mathbf{M}(f^*)$, respectively. However, since the entire sequences $\{\mathbf{m}(f_n)\}$ and $\{\mathbf{M}(f_n)\}$ strongly converge in H , it further follows that they converge weakly (not only along a subsequence) in V . Hence (31) and (32) hold true. \square

With the aid of the previous theorem we can now formulate the result that proves the well-posedness of (\mathbb{P}) . We assume that $U \subset L^\infty(\Omega)$ and in particular that

$$U_{\text{ad}} \subset \{f \in L^\infty_\nu(\Omega) : f \leq F\},$$

for some $F \in V'$. As in previous sections $\underline{y} = 0$ and $\bar{y} = A^{-1}F$, so that $\mathbf{m}(f)$ and $\mathbf{M}(f)$ are defined as the minimal and maximal solutions, respectively, of the QVI in (P_{QVI}) . Hence, the reduced version of (\mathbb{P}) is given by

$$\begin{aligned} & \text{minimize } J_1(\mathbf{m}(f), \mathbf{M}(f)) + J_2(f), \\ & \text{subject to } f \in U_{\text{ad}}. \end{aligned} \tag{\tilde{\mathbb{P}}}$$

The well posedness of $(\tilde{\mathbb{P}})$ (and hence of (\mathbb{P})) is now shown in the following result.

Theorem 6. *Suppose that*

(i) $J_1 : V \times V \rightarrow \mathbb{R}$ *is weakly lower semicontinuous,*

(ii) $J_2 : L^\infty(\Omega) \rightarrow \mathbb{R}$ *is continuous,*

and both are bounded from below. In addition suppose that for each $\alpha > 0$ the set

$$\{f \in U_{\text{ad}} : J_2(f) \leq \alpha\},$$

is sequentially compact in $L^\infty(\Omega)$, and that Φ satisfies the assumptions of Theorem 5. Then, problem $(\tilde{\mathbb{P}})$, and hence of problem (\mathbb{P}) , admits a solution.

Proof. Given Theorem 5, the proof is just an application of the direct method of the calculus of variations. \square

7.1 Applications

We finally return to the applications considered earlier in the paper.

7.1.1 QVIs arising by coupling VIs and PDEs

We consider the problem class as described in section 3.1 and study conditions on G, B , and L to establish stability of minimum and maximum solutions to the QVI of interest. Recall that Φ in this setting is defined as $\Phi(y) = Lz(y)$ where $z(y)$ solves

$$\langle Bz + G(Lz, y) - g, w \rangle = 0 \quad \forall w \in W,$$

for $y \in V$.

Proposition 1. *Under the assumptions of section 3.1 suppose that if $\lambda \geq 1$ and $v \in H^+ \cap V$, then for all $z_1, z_2 \in V$, it holds true that*

$$(\lambda G(Lz_2, v) - G(Lz_1, \lambda v), (z_1 - \lambda z_2)^+) \leq 0.$$

Then, we have $\lambda \Phi(v) \geq \Phi(\lambda v)$.

Analogously, suppose that if $\lambda \in (0, 1)$ and $v \in H^+ \cap W$, then for all $z_1, z_2 \in V$, it holds true that

$$(G(Lz_2, \lambda v) - \lambda G(Lz_1, v), (\lambda z_1 - z_2)^+) \leq 0.$$

Then, it holds that $\lambda \Phi(y) \leq \Phi(\lambda y)$.

Proof. Let $w = z(\lambda v) - \lambda z(v)$ for $\lambda \geq 1$ and $v \in H^+ \cap V$. Since B is coercive and $\langle Bw^-, w^+ \rangle \leq 0$ we observe that

$$c|w^+|_W^2 \leq \langle Bw, w^+ \rangle = (\lambda G(Lz(v), v) - G(Lz(\lambda v), \lambda v), w^+) \leq 0,$$

i.e., $z(\lambda v) - \lambda z(v) \leq 0$, so that $Lz(\lambda v) - L(\lambda z(v)) \leq 0$ given that L preserves order. Hence, it follows that $\lambda\Phi(v) \geq \Phi(\lambda v)$.

Similarly, consider $w = \lambda z(v) - z(\lambda v)$ for $0 < \lambda < 1$ and $v \in H^+ \cap V$. Then,

$$c|w^+|_W^2 \leq \langle Bw, w^+ \rangle = (G(Lz(\lambda v) - \lambda G(Lz(v), v), \lambda v), w^+) \leq 0,$$

i.e., $\lambda z(v) - z(\lambda v) \leq 0$, so that $\lambda Lz(v) - Lz(\lambda v) \leq 0$ and hence $\lambda\Phi(v) \leq \Phi(\lambda v)$. \square

Note that the problem given in Example 1 satisfies the assumptions of the above proposition. Additionally, if the solution to $By = h$ satisfies $|y|_{H^2(\Omega)} \leq M|h|_{L^2(\Omega)}$ with M independent of h , then for dimensions $N = 1, 2, 3$ it is direct to infer that $\Phi : H_0^1(\Omega) \rightarrow L^\infty(\Omega)$ is completely continuous via Sobolev embeddings. Hence, all hypothesis of Theorem 5 are met, and the minimum and maximum solutions are stable for perturbations of f in $L^\infty(\Omega)$. Finally, if

$$\{f \in U : 0 < \nu \leq f \leq F \text{ and } \|f\|_U \leq \alpha\},$$

is sequentially compact in $L^\infty(\Omega)$ for each $\alpha > 0$, we have that Problem 8 has a solution. Further note that this last compactness assumption is satisfied for Example 1.

7.1.2 The impulse control problems

The previous can be directly applied to the impulse control problem in the bounded case. Let $\Omega = (0, 1)$. Then, we have that $V = H_0^1(\Omega)$ compactly embeds into $C(\overline{\Omega})$, and hence it follows that for

$$(\Phi y)(x) = k + \operatorname{ess\,inf}_{x+\xi \in \overline{\Omega}} (c_0(\xi) + y(x + \xi)),$$

with $k > 0$ and c_0 continuous, we have that if $v_n \rightharpoonup v$ in V , then $\Phi(v_n) \rightarrow \Phi(v)$ in $C(\overline{\Omega}) \subset L^\infty(\Omega)$. Hence, Φ satisfies Assumption 1. Furthermore, it follows that $\lambda\Phi(y) \geq \Phi(\lambda y)$ for any $\lambda > 1$ and any $y \in V \cap H^+$.

Consider $U = H^1(\Omega)$ and $U_{\text{ad}} := \{f \in U : 0 < \nu \leq f \leq F\}$ for some $F \in H^1(\Omega)^*$, $J_1(a, b) = \int_\Omega (A - a(x))^2 dx$, and $J_2(f) := \frac{\lambda}{2}|f|_U^2$. It follows that $\{f \in U_{\text{ad}} : J_2(f) \leq \alpha\}$ is sequentially compact in $L^\infty(\Omega)$ for each $\alpha > 0$ and that problem $(\tilde{\mathbb{P}})$ (which is the reduced version of problem (9)) has a solution by Theorem 6.

8 Conclusion

We have developed a theoretical framework for the study of optimal control problems with QVI constraints. Specifically, the reduced optimization problem of interest involves minimal and maximal points of the solution set to the QVI. The existence question reduces to the stability of two operators \mathbf{m} and \mathbf{M} , that relate the solution set of the QVI to its minimal and maximal elements, respectively. Stability of such maps was developed for monotonic and non-monotonic perturbations, and we have applied such results to applications involving QVIs arising from impulse control problems and problems involving VIs coupled with nonlinear PDEs.

References

- [1] R. A. Adams and J. J. F. Fournier. *Sobolev Spaces*. Academic Press, second edition, 2003.
- [2] S. Adly, M. t. Bergounioux, and M. Ait Mansour. Optimal control of a quasi-variational obstacle problem. *J. Global Optim.*, 47(3):421–435, 2010.
- [3] C. D. Aliprantis and O. Burkinshaw. *Positive operators*, volume 119. Springer Science & Business Media, 2006.
- [4] A. Alphonse, M. Hintermüller, and Rautenberg. Recent trends and views on elliptic quasi-variational inequalities. *WIAS Preprint No. 2518*, 2018.
Available at <http://dx.doi.org/10.20347/WIAS.PREPRINT.2518>.
- [5] A. Alphonse, M. Hintermüller, and C. N. Rautenberg. Directional differentiability for elliptic quasi-variational inequalities of obstacle type. *Calc. Var. Partial Differential Equations*, 58(1):58:39, 2019.
- [6] H. Antil and C. N. Rautenberg. Fractional elliptic quasi-variational inequalities: theory and numerics. *Interfaces Free Bound.*, 20(1):1–24, 2018.
- [7] J.-P. Aubin. *Mathematical methods of game and economic theory*. North-Holland, 1979.
- [8] A. Azevedo, F. Miranda, and L. Santos. Variational and quasivariational inequalities with first order constraints. *J. Math. Anal. Appl.*, 397(2):738–756, 2013.
- [9] A. Azevedo, F. Miranda, and L. Santos. *Stationary Quasivariational Inequalities with Gradient Constraint and Nonhomogeneous Boundary Conditions*, pages 95–112. Springer Berlin Heidelberg, Berlin, Heidelberg, 2014.
- [10] C. Baiocchi and A. Capelo. *Variational and Quasivariational Inequalities*. Wiley-Interscience, 1984.
- [11] J. W. Barrett and L. Prigozhin. A quasi-variational inequality problem in superconductivity. *Math. Models Methods Appl. Sci.*, 20(5):679–706, 2010.
- [12] J. W. Barrett and L. Prigozhin. A quasi-variational inequality problem arising in the modeling of growing sandpiles. *ESAIM Math. Model. Numer. Anal.*, 47(4):1133–1165, 2013.
- [13] J. W. Barrett and L. Prigozhin. Lakes and rivers in the landscape: a quasi-variational inequality approach. *Interfaces Free Bound.*, 16(2):269–296, 2014.
- [14] J. W. Barrett and L. Prigozhin. Sandpiles and superconductors: nonconforming linear finite element approximations for mixed formulations of quasi-variational inequalities. *IMA J. Numer. Anal.*, 35(1):1–38, 2015.
- [15] A. Bensoussan. *Stochastic Control by Functional Analysis Methods*. North-Holland, 1982.
- [16] A. Bensoussan and J.-L. Lions. Contrôle impulsionnel et inéquations quasi-variationnelles d'évolutions. *C. R. Acad. Sci. Paris*, 276:1333–1338, 1974.
- [17] A. Bensoussan and J.-L. Lions. *Contrôle impulsionnel et inéquations quasi variationnelles*, volume 11 of *Méthodes Mathématiques de l'Informatique [Mathematical Methods of Information Science]*. Gauthier-Villars, Paris, 1982.

- [18] A. Bensoussan and J.-L. Lions. *Impulse Control and Quasi-Variational Inequalities*. Gauthier-Villars, 1984.
- [19] G. Birkhoff. *Lattice Theory*, volume 25 of *Amer. Math. Soc. Colloquium Publ.* American Mathematical Society, Providence, 1961.
- [20] Z. Denkowski, S. Migórski, and N. S. Papageorgiou. *Introduction to Nonlinear Analysis: Applications*. Kluwer, 2003.
- [21] H. Dietrich. Über Probleme der optimalen Steuerung mit einer Quasivariationsungleichung. In *Operations Research Proceedings 1999 (Magdeburg)*, pages 111–116. Springer, Berlin, 2000.
- [22] H. Dietrich. Optimal control problems for certain quasivariational inequalities. *Optimization*, 49(1-2):67–93, 2001. In celebration of Prof. Dr. Alfred Göpfert 65th birthday.
- [23] A. Friedman. *Variational Principles and Free-Boundary Problems*. Wiley-Interscience, 1982.
- [24] T. Fukao and N. Kenmochi. Abstract theory of variational inequalities with Lagrange multipliers and application to nonlinear PDEs. *Math. Bohem.*, 139(2):391–399, 2014.
- [25] T. Fukao and N. Kenmochi. Quasi-variational inequality approach to heat convection problems with temperature dependent velocity constraint. *Discrete Contin. Dyn. Syst.*, 35(6):2523–2538, 2015.
- [26] I. M. Gel'fand and N. Y. Vilenkin. *Generalized functions. Vol. 4: Applications of harmonic analysis*. Translated by Amiel Feinstein. Academic Press, New York - London, 1964, 1964.
- [27] J. Gwinner, B. Jadamba, A. A. Khan, and M. Sama. Identification in variational and quasi-variational inequalities. *J. Convex Anal.*, 25(2):545–569, 2018.
- [28] R. Herzog, C. Meyer, and G. Wachsmuth. B- and strong stationarity for optimal control of static plasticity with hardening. *SIAM J. Optim.*, 23(1):321–352, 2013.
- [29] M. Hintermüller, B. S. Mordukhovich, and T. M. Surowiec. Several approaches for the derivation of stationarity conditions for elliptic MPECs with upper-level control constraints. *Math. Program.*, 146(1-2, Ser. A):555–582, 2014.
- [30] M. Hintermüller and C. N. Rautenberg. A sequential minimization technique for elliptic quasi-variational inequalities with gradient constraints. *SIAM J. Optim.*, 22(4):1224–1257, 2012.
- [31] M. Hintermüller and C. N. Rautenberg. Parabolic quasi-variational inequalities with gradient-type constraints. *SIAM J. Optim.*, 23(4):2090–2123, 2013.
- [32] M. Hintermüller and C. N. Rautenberg. On the uniqueness and numerical approximation of solutions to certain parabolic quasi-variational inequalities. *Port. Math.*, 74(1):1–35, 2017.
- [33] M. Hintermüller and T. Surowiec. First-order optimality conditions for elliptic mathematical programs with equilibrium constraints via variational analysis. *SIAM J. Optim.*, 21(4):1561–1593, 2011.
- [34] A. Kadoya, N. Kenmochi, and M. Niezgodka. Quasi-variational inequalities in economic growth models with technological development. *Adv. Math. Sci. Appl.*, 24(1):185–214, 2014.

- [35] R. Kano, Y. Murase, and N. Kenmochi. Nonlinear evolution equations generated by subdifferentials with nonlocal constraints. In *Nonlocal and abstract parabolic equations and their applications*, volume 86 of *Banach Center Publ.*, pages 175–194. Polish Acad. Sci. Inst. Math., Warsaw, 2009.
- [36] N. Kenmochi. Parabolic quasi-variational diffusion problems with gradient constraints. *Discrete Contin. Dyn. Syst. Ser. S*, 6(2):423–438, 2013.
- [37] D. Kinderlehrer and G. Stampacchia. *An Introduction to Variational Inequalities and Their Applications*. SIAM, 2000.
- [38] I. I. Kolodner. On completeness of partially ordered sets and fixpoint theorems for isotone mappings. *Amer. Math. Monthly*, 75:48–49, 1968.
- [39] M. Kunze and J. Rodrigues. An elliptic quasi-variational inequality with gradient constraints and some of its applications. *Mathematical Methods in the Applied Sciences*, 23:897–908, 2000.
- [40] F. Miranda, J.-F. Rodrigues, and L. Santos. On a p -curl system arising in electromagnetism. *Discrete Contin. Dyn. Syst. Ser. S*, 5(3):605–629, 2012.
- [41] B. S. Mordukhovich and J. V. Outrata. Coderivative analysis of quasi-variational inequalities with applications to stability and optimization. *SIAM J. Optim.*, 18(2):389–412, 2007.
- [42] Y. Murase, A. Kadoya, and N. Kenmochi. Optimal control problems for quasi-variational inequalities and its numerical approximation. *Discrete Contin. Dyn. Syst.*, (Dynamical systems, differential equations and applications. 8th AIMS Conference. Suppl. Vol. II):1101–1110, 2011.
- [43] L. Prigozhin. Quasivariational inequality describing the shape of a poured pile. *Zhurnal Vichislitel'noy Matematiki i Matematicheskoy Fiziki*, 7:1072–1080, 1986.
- [44] L. Prigozhin. Sandpiles and river networks: extended systems with non-local interactions. *Phys. Rev. E*, 49:1161–1167, 1994.
- [45] L. Prigozhin. On the Bean critical-state model in superconductivity. *European Journal of Applied Mathematics*, 7:237–247, 1996.
- [46] L. Prigozhin. Sandpiles, river networks, and type-ii superconductors. *Free Boundary Problems News*, 10:2–4, 1996.
- [47] L. Prigozhin and J. Barret. A quasi-variational inequality problem arising in the modeling of growing sandpiles. <http://www.cs.bgu.ac.il/leonid/sandqvi2012.pdf>.
- [48] J. F. Rodrigues. *Obstacle Problems in Mathematical Physics*. North-Holland, 1987.
- [49] J. F. Rodrigues and L. Santos. A parabolic quasi-variational inequality arising in a superconductivity model. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 29(1):153–169, 2000.
- [50] J. F. Rodrigues and L. Santos. Quasivariational solutions for first order quasilinear equations with gradient constraint. *Arch. Ration. Mech. Anal.*, 205(2):493–514, 2012.
- [51] L. Tartar. Inéquations quasi variationnelles abstraites. *C. R. Ac. Sci. Paris*, 178:1193–1196, 1974.
- [52] H. Toyozumi. Continuous dependence on obstacles in variational inequalities. *Funkcialaj Ekvacioj*, 34:103–115, 1991.

- [53] G. Wachsmuth. Towards M-stationarity for optimal control of the obstacle problem with control constraints. *SIAM J. Control Optim.*, 54(2):964–986, 2016.