

Weierstraß-Institut
für Angewandte Analysis und Stochastik
Leibniz-Institut im Forschungsverbund Berlin e. V.

Preprint

ISSN 2198-5855

**A phase field approach for optimal boundary control of damage
processes in two-dimensional viscoelastic media**

M. Hassan Farshbaf-Shaker¹, Christian Heinemann¹

submitted: February 2, 2015

¹ Weierstrass Institute
Mohrenstr. 39
10117 Berlin
Germany
E-Mail: MohammadHassanFarshbaf.Shaker@wias-berlin.de
Christian.Heinemann@wias-berlin.de

No. 2074

Berlin 2015



2010 *Mathematics Subject Classification.* 35A01, 35A02, 35D35, 35M33, 35M87, 35Q74, 49J20, 74A45, 74D10, 74F99, 74H20, 74H25, 74P99.

Key words and phrases. damage processes, phase field model, viscoelasticity, nonlinear parabolic inclusions, well-posedness, optimal control.

This work is partially supported by the Research Center Matheon in Berlin (Germany).

Edited by
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)
Leibniz-Institut im Forschungsverbund Berlin e. V.
Mohrenstraße 39
10117 Berlin
Germany

Fax: +49 30 20372-303
E-Mail: preprint@wias-berlin.de
World Wide Web: <http://www.wias-berlin.de/>

Abstract

In this work we investigate a phase field model for damage processes in two-dimensional viscoelastic media with nonhomogeneous Neumann data describing external boundary forces. In the first part we establish global-in-time existence, uniqueness, a priori estimates and continuous dependence of strong solutions on the data. The main difficulty is caused by the irreversibility as well as boundedness of the phase field variable which results in a doubly constrained PDE system. In the last part we consider an optimal control problem where a cost functional penalizes maximal deviations from prescribed damage profiles. The goal is to minimize the cost functional with respect to exterior forces acting on the boundary which play the role of the control variable in the considered model. To this end, we prove existence of minimizers and study a family of “local” approximations via adapted cost functionals.

1 Introduction

Damage phenomena in elastically deformable solids and their analytical studies have received a lot of attention in the mathematical literature, e.g., [4, 8, 12, 14, 16, 21, 27, 29, 31, 38]. Especially models which employ a phase field approach and incorporate higher order terms were focused in some recent works. In that case, an internal variable indicates the degree of structural integrity and, depending on the material and the scaling, may be defined as the volume or the surface density of microvoids or microcracks, respectively, as pointed out in [25]. This approach has also been utilized for approximations of surface discontinuities occurring in the displacement field of fracture models and turned out to be very useful for numerical implementations (see [1, 5, 14, 35]).

One of the main difficulty for a rigorous mathematical investigation of the underlying PDE systems is that the damage variable is usually doubly constrained: The variable is forced to be monotonically decreasing in time (irreversibility) and bounded in the unit interval. This kind of non-smooth evolution had motivated different concepts of weak solutions and regularization techniques in the literature (cf., e.g., [3, 21, 29]). However, to the authors’ best knowledge, a global-in-time well-posedness result for strong solutions with inhomogeneous boundary data was left open. Together with sufficiently strong a priori estimates such a result could be exploited to study optimal control problems typically arising in engineering problems focused on resistance against damage and failure. The following model problems with boundary control illustrate some practical examples:

- Suppose that a workpiece is exposed to external forces during an experiment and that certain parameters related to those forces can be controlled. A control problem could be to choose optimal parameters in order to prevent further damage in the material.
- Related to the first scenario we might be interested in calculating additional forces not to prevent but to redirect crack spreading to non-critical components of the structure and to avoid complete failure.

- Another problem might be the determination of external forces in order to deliberately induce a damage progression. For instance, it might be desirable to separate certain parts of the workpiece in industrial processes.

By now, to the authors' best knowledge, the mathematical contributions addressing those and related problems are inspired by the pioneering work [7] and employ fracture models to control the energy release rate of a single crack in a quasi-stationary setting by optimal shape design techniques, fibers or applied forces (see [19, 20, 26, 36] for more details). The cracks are explicitly modeled by non-smooth domains with or without non-penetration conditions for the deformation. A main issue consists in determining optimal forces or inclusions in the solid in order to cease crack propagation or to release as much energy as possible.

In this paper we would like to advance a different approach for such control problems by utilizing a phase field model for damage. The kind of model under consideration was motivated by FRÉMOND and NEDJAR in [13] and is stated below. Under certain structural assumptions we are able to investigate well-posedness of strong solutions and existence of optimal boundary controls for a coupled evolutionary system describing damage processes in viscoelastic materials in two spatial dimensions. A virtue of our approach is that we are able to control the damage progression on the whole domain of interest even at all times during the evolution if desired.

In the first part of this paper we study existence and then, for constant viscosity \mathbb{D} , well-posedness of the following PDE problem:

For a given time interval $(0, T)$ and reference configuration Ω with boundary Γ and outer unit normal ν , find (u, χ) such that

$$u_{tt} - \operatorname{div}(\mathbb{C}(\chi)\varepsilon(u) + \mathbb{D}(\chi)\varepsilon(u_t)) = \ell \quad \text{in } \Omega \times (0, T), \quad (1a)$$

$$\chi_t - \Delta\chi_t - \Delta\chi + \xi + \frac{1}{2}\mathbb{C}'(\chi)\varepsilon(u) : \varepsilon(u) + f'(\chi) = 0 \quad \text{in } \Omega \times (0, T) \quad (1b)$$

subject to the subgradient/pointwise constraints

$$\xi \in \partial I_{(-\infty, 0]}(\chi_t) \quad \text{in } \Omega \times (0, T), \quad (2a)$$

$$\chi \in [0, 1] \quad \text{in } \Omega \times (0, T) \quad (2b)$$

and the initial-boundary conditions

$$u(0) = u^0, \quad u_t(0) = v^0, \quad \chi(0) = \chi^0 \quad \text{in } \Omega, \quad (3a)$$

$$(\mathbb{C}(\chi)\varepsilon(u) + \mathbb{D}(\chi)\varepsilon(u_t)) \cdot \nu = b \quad \text{on } \Gamma \times (0, T), \quad (3b)$$

$$\nabla(\chi + \chi_t) \cdot \nu = 0 \quad \text{on } \Gamma \times (0, T). \quad (3c)$$

Equation (1a) describes the balance of forces in the workpiece according to the Kelvin-Voigt rheology. The displacement field is denoted by u , the external volume forces by ℓ , the linearized strain tensor by $\varepsilon(u) = \frac{1}{2}(\nabla u + (\nabla u)^T)$ and the stress tensor by $\sigma = \mathbb{C}(\chi)\varepsilon(u) + \mathbb{D}(\chi)\varepsilon(u_t)$. The first summand of σ contains the elastic contribution whereas the second summand models viscous effects. The coefficient \mathbb{C} designates the fourth-order damage-dependent stiffness tensor and \mathbb{D} the viscosity tensor. The second equation (1b) specifies the parabolic evolution law for the propagation of damage described by the variable χ under the constraints (2a) and (2b), where the subdifferential of the indicator function $I_{(-\infty, 0]} : \mathbb{R} \rightarrow \mathbb{R} \cup \{\infty\}$ is given by

$$\partial I_{(-\infty, 0]}(\chi_t) = \begin{cases} \{0\} & \text{if } \chi_t < 0, \\ [0, \infty) & \text{if } \chi_t = 0, \\ \emptyset & \text{if } \chi_t > 0. \end{cases}$$

The Laplacians $\Delta\chi$ and $\Delta\chi_t$ model diffusive effects of χ and χ_t and have a regularizing effect from the mathematical point of view (see the remark after the proof of Theorem 2.11). For a mechanical motivation of system (1)-(3) by means of balance laws and constitutive relations we refer to [9, 10, 13]. We would like to give the following interpretation for the subgradient constraint (2a):

By introducing the free energy F to system (1) as

$$F(u, \chi) = \int_{\Omega} \left(\frac{1}{2} |\nabla\chi|^2 + \frac{1}{2} \mathbb{C}(\chi) \varepsilon(u) : \varepsilon(u) + f(\chi) \right) dx, \quad (4)$$

we may rewrite (1b) as

$$\xi = -\chi_t + \Delta\chi_t - d_{\chi}F(u, \chi).$$

By virtue of the complementarity formulation for (2a), i.e.

$$\chi_t \leq 0, \quad \xi \cdot \chi_t = 0, \quad \xi \geq 0,$$

the evolution law (1b) reads as

$$\begin{aligned} (\mathbb{1} - \Delta)\chi_t &= -d_{\chi}F(u, \chi) && \text{if } -d_{\chi}F(u, \chi) \leq 0, \\ \chi_t &= 0 && \text{if } -d_{\chi}F(u, \chi) > 0, \end{aligned}$$

i.e., χ is governed by a gradient flow with respect to χ in the H^1 -norm whenever the driving force $-d_{\chi}F(u, \chi)$ is non-positive and $\chi_t = 0$ otherwise.

The second part in this paper is devoted to an optimal control problem. A cost functional \mathcal{J} will measure the maximal deviation of the damage variable χ from given prescribed damage profiles at the final time T and/or at all times in $[0, T]$ ($\lambda_Q, \lambda_{\Omega}, \lambda_{\Sigma} \geq 0$):

$$\mathcal{J}(\chi, b) := \frac{\lambda_Q}{2} \|\chi - \chi_Q\|_{L^{\infty}(\Omega \times (0, T))} + \frac{\lambda_{\Omega}}{2} \|\chi(T) - \chi_T\|_{L^{\infty}(\Omega \times (0, T))} + \frac{\lambda_{\Sigma}}{2} \|b\|_{L^2(\Gamma \times (0, T); \mathbb{R}^n)}^2. \quad (5)$$

A minimizer (χ, b) of \mathcal{J} under the constraint that χ solves system (1)-(3) for some displacement u and boundary data b indicates an evolution which approximates χ_Q and/or χ_T best in the sense of \mathcal{J} .

In the following we summarize the main results of our paper:

- In Theorem 2.11 we will prove existence of strong solutions for system (1)-(3) and for a so-called β -approximation in two spatial dimensions. In the latter case we replace the subgradient ξ in (2a) by a smooth approximation $\xi_{\beta}(\chi_t)$ with $\beta > 0$. On the one hand this enables us to perform the a priori estimates in Lemma 2.9, while, on the other hand, the β -approximation might be helpful for further studies such as optimality systems, numerical implementations etc. We emphasize that the existence analysis constitutes the main part of this paper and strongly relies on the two-dimensional *Ladyzhenskaya's inequality* originally devised for the 2D Navier-Stokes equations (see [23] and the calculation (38)).
- Continuous dependence on the data $(u^0, v^0, \chi^0, b, \ell)$ and, in particular, uniqueness of strong solutions for system (1)-(3) are proven in Theorem 2.12 (see also Corollary 2.13) under the assumption of constant viscosity \mathbb{D} . We also establish a priori estimates for the solutions in Corollary 2.14. These results allow us to define the solution operator and constitutes the fundament for the considered optimal control problem.

- Theorem 3.6 reveals existence to an optimal control problem where the cost functional penalizes deviations of the damage variable from given damage profiles in the L^∞ -norm (see (5)). The strong solutions of system (1)-(3) will be controlled via external boundary forces. We prove existence of optimal controls by using the β -approximation in the proof of Theorem 2.11 to define a family of optimal control problems. The minimizers or the optimal controls of the family of β -approximating control problems converge in a limit process (along a subsequence as $\beta \downarrow 0$) to an optimal control of the original control problem. In other words, we show that optimal controls for the family of β -approximating control problems are for some $\beta > 0$ likely to be “close” to optimal controls for the original control problem. It is natural to ask if the reverse holds, i.e., whether every optimal control for the original control problem can be approximated by a sequence of optimal controls of the β -approximating control problems. Unfortunately, we will not be able to prove such a “global result” that applies to all optimal controls for the original control problem. The reason for that lies on the non-convexity of the optimal control problems (both the original one and the β -approximating control problems) and consequently on the non-uniqueness of the optimal controls. However, a “local” result can be established by introducing so-called adapted optimal control problems in Theorem 3.10.

Let us recall some already established results in the mathematical literature of phase field models for damage/gradient-of-damage models:

- Local-in-time well-posedness of strong solutions for damage-elasticity systems with scalar-valued displacements and homogeneous Dirichlet conditions for the displacements is proven in [3, 4] and in [11, 12] for one dimensional models. The authors firstly replace the constraint (2b) by a second subgradient $\zeta \in \partial I_{[0,1]}(\chi)$ where ζ also occurs in the damage law. Then the local-in-time character guarantees the pointwise constraint (2b) and the vanishing of ζ since starting from an initial damage profile which satisfies $\chi^0 \in [\eta, 1]$ pointwise in Ω for a constant $\eta \in (0, 1]$ implies $\chi \in [\hat{\eta}, 1]$ pointwise in $\Omega \times (0, \hat{t})$ for further small constants $\hat{t}, \hat{\eta} > 0$ provided continuity of χ .

In contrast to this approach, we will ensure the constraint (2b) via a carefully chosen time-discretization scheme (see Definition 2.5 (iii)). The main advantage is that (2b) is carried over to global-in-time solutions. This is a crucial consequence since uniqueness of the damage-elasticity system with two subgradients is considered as an open problem (see [37, Remark 2.18]).

- Rate-independent gradient-of-damage models are explored in [29] and in various subsequent papers, e.g., [28, 30]. The rate-independence requires that the term χ_t in (1b) should be neglected which is a reasonable assumption if the damage progression is on a faster timescale than acting of the external forces. The authors considered non-smooth domains and employed weak notions referred to as *energetic formulation* in order to prove existence of solutions. The degenerating case where the material may loose all its elastic properties due to heavy damage is also studied. Further cases involving nonlinear r -Laplacians with $r > 1$ or even $r = 1$ instead of the classical Laplacian in (1b) are investigated in [31, 41], where also higher temporal regularity is shown.
- A weak notion for rate-dependent damage models coupled with Cahn-Hilliard equations was introduced in [16] for quasi-static balance of forces and in [17] with inertial effects and without the viscosity term in (1a). Existence of weak solutions is proven there for non-smooth domains and mixed-boundary conditions for the displacements whereas uniqueness is left open.

- A well-posedness result for damage models with (nonlocal) high-order s -Laplacian is shown in [21] (see also [22] for vanishing viscosity results in non-smooth settings). Among other results in that paper, the authors have proven existence of solutions in an energetic formulation and uniqueness in special situations. Concerning uniqueness in the case of $n \in \{2, 3\}$ dimensions, the s -Laplacian is assumed to be of higher order than the classical Laplacian.
- Coupled thermoviscoelastic and isothermal damage models incorporating p -Laplacian operators are analyzed in [37] (see also [38] for the full heat equation including all dissipative terms and [18] for damage-dependent heat expansion coefficients). In those works homogeneous Dirichlet boundary conditions for the displacements are assumed. Uniqueness is shown in the isothermal case by adopting $p > n$ and by dropping the irreversibility constraint (2a). Existence results for corresponding rate-independent thermoviscoelastic damage models are proven in the recent paper [24].

Structure of the paper

Section 2 is devoted to the well-posedness problem of system (1)-(3). We state the precise assumptions in Subsection 2.1 and introduce time-discretized and β -regularized approximations of (1)-(3) in Subsection 2.2. The existence proofs are carried out in Subsection 2.3 firstly for the time-discretized and then, by a limit analysis, for the time-continuous versions. In the final part of that section, i.e. in Subsection 2.4, we prove continuous dependence on the initial-boundary data. Then, equipped with the well-posedness result, we state the announced optimal control problem in Section 3. We prove existence of optimal controls via β -regularization in Subsection 3.1 and their approximation by means of an adapted cost functional in Subsection 3.2.

2 Analysis of the evolution inclusions

The approach presented in this work combines two different approximation techniques to obtain existence of solutions for system (1)-(3): semi-implicit time-discretization and regularization of the subgradient ξ in (2a). At first we will tackle the existence problem for the time-discrete and regularized system in Lemma 2.7. By passing the discretization fineness to 0, solutions of a time-continuous regularized system are obtained in Theorem 2.11 (i). In the final step, a further limit passage leads to solutions of the desired limit system (see Theorem 2.11 (ii)). Then, we conclude this section in Theorem 2.12 with a uniqueness and continuous dependence result.

2.1 Assumptions and notation

Throughout this work, we adopt the following assumptions:

- (A1) $\Omega \subseteq \mathbb{R}^n$ with $n \in \{1, 2\}$ is a bounded C^2 -domain. The boundary is denoted by Γ and the outer unit normal by ν .
- (A2) The damage-dependent stiffness tensor satisfies $\mathbb{C}(\cdot) = \mathbf{c}(\cdot)\mathbf{C}$, where the coefficient function \mathbf{c} is assumed to be in $\mathbf{c} \in C^{1,1}([0, 1]; \mathbb{R})$ and assumed to satisfy

$$\mathbf{c}(x) \geq 0 \quad \text{for all } x \in [0, 1].$$

Furthermore, we assume a convex-concave decomposition $\mathbf{c} = \mathbf{c}_1 + \mathbf{c}_2$ with convex $\mathbf{c}_1 \in C^1([0, 1]; \mathbb{R})$ and concave $\mathbf{c}_2 \in C^1([0, 1]; \mathbb{R})$ such that

$$\mathbf{c}'_1(0) \leq 0, \quad \mathbf{c}'_1(1) \geq \max_{s \in [0, 1]} -\mathbf{c}'_2(s) \quad (6a)$$

$$\mathbf{c}'_2(x) \leq 0 \quad \text{for all } x \in [0, 1]. \quad (6b)$$

The 4th order stiffness tensor $\mathbf{C} \in \mathcal{L}(\mathbb{R}_{\text{sym}}^{n \times n}; \mathbb{R}_{\text{sym}}^{n \times n})$ is assumed to be symmetric and positive definite, i.e.

$$\mathbf{C}_{ijkl} = \mathbf{C}_{jilk} = \mathbf{C}_{lkij} \text{ and } e : \mathbf{C}e \geq \eta |e|^2 \text{ for all } e \in \mathbb{R}_{\text{sym}}^{n \times n}$$

with constant $\eta > 0$.

- (A3) The damage-dependent viscosity tensor satisfies $\mathbb{D}(\cdot) = \mathbf{d}(\cdot)\mathbf{D}$, where the coefficient function \mathbf{d} satisfies $\mathbf{d} \in C^1([0, 1]; \mathbb{R})$ and

$$\mathbf{d}(x) \geq \eta > 0 \text{ for all } x \in [0, 1] \text{ and fixed } \eta > 0. \quad (7)$$

The 4th order tensor \mathbf{D} is given by $\mathbf{D} = \mu\mathbf{C}$, where $\mu > 0$ is a constant.

- (A4) The damage-dependent potential function f is assumed to be in $f \in C^{1,1}([0, 1]; \mathbb{R})$ and assumed to satisfy

$$f'(0) \leq 0 \text{ and } f'(1) \geq 0. \quad (8)$$

Remark 2.1 (i) A convex-concave decomposition $\mathbf{c} = \mathbf{c}_1 + \mathbf{c}_2$ in (A2) may be given by

$$\begin{aligned} \mathbf{c}_1(r) &:= \mathbf{c}(0) + \mathbf{c}'(0)r + \int_0^r \left(\int_0^s \max\{\mathbf{c}''(\tau), 0\} d\tau \right) ds, \\ \mathbf{c}_2(r) &:= \int_0^r \left(\int_0^s \min\{\mathbf{c}''(\tau), 0\} d\tau \right) ds, \end{aligned}$$

where \mathbf{c}_1 has to satisfy (6a). Note that condition (6b) is automatically satisfied here. In particular, if we choose a convex function for \mathbf{c} the conditions in (6) reduce to $\mathbf{c}'(0) \leq 0$ and $\mathbf{c}'(1) \geq 0$ as in (A4).

- (ii) The growth assumptions (6) and (8) will guarantee that the values of the damage variable are in the unit interval. To this end we will adapt the argumentation in [21, Proposition 5.5] to our situation. In that work the non-negativity of the damage variable is ensured via certain growth assumptions on the coefficient functions. Then, by taking the irreversibility constraint into account, i.e. $\chi_t \leq 0$, the authors infer $\chi \in [0, 1]$ in $\Omega \times (0, T)$ provided that the initial value satisfies $\chi^0 \in [0, 1]$ in Ω .

However, in our case, the irreversibility condition $\chi_t \leq 0$ is not be ensured in the β -regularization of system (1)-(3) (see Definition 2.5 (ii) and (iii)). Thus beside $\chi \geq 0$ we also need to show $\chi \leq 1$.

- (iii) The non-degeneracy condition (7) prevents the material from complete damage, i.e., even the maximal damaged parts (the region with $\chi = 0$) exhibit small viscous properties.
- (iv) The assumption $\mathbf{D} = \mu\mathbf{C}$ in (A3) is needed in the proof of Lemma 2.7 in step 2 in order to perform a regularity argument based on a transformation. It has already been employed in the mathematical literature (see [18, 38]).

Example 2.2 *The following example for c and f is covered by Assumptions (A2) and (A4):*

$$c(x) = x^2 \quad \text{and} \quad f(x) = \frac{1}{4\varepsilon}(1-x)^2$$

with constant $\varepsilon > 0$. With this choice, the energy functional (4) when replacing $\frac{1}{2}|\nabla\chi|^2$ by $\varepsilon|\nabla\chi|^2$ yields the (generalized) Ambrosio-Tortorelli approximation of the corresponding brittle fracture model (see [1]).

For later use, we define the solution space $\mathcal{U} \times \mathcal{X}$, where \mathcal{U} denotes the space of the displacements and \mathcal{X} the space of the damage evolutions given by

$$\begin{aligned} \mathcal{U} &:= H^1(0, T; H^2(\Omega; \mathbb{R}^n)) \cap W^{1,\infty}(0, T; H^1(\Omega; \mathbb{R}^n)) \cap H^2(0, T; L^2(\Omega; \mathbb{R}^n)), \\ \mathcal{X} &:= H^1(0, T; H^2(\Omega)). \end{aligned}$$

The space of boundary controls \mathcal{B} is defined as

$$\mathcal{B} := L^2(0, T; H^{1/2}(\Gamma; \mathbb{R}^n)) \cap H^1(0, T; L^2(\Gamma; \mathbb{R}^n)).$$

We also introduce the sets for brevity

$$Q := \Omega \times (0, T), \quad \Sigma := \Gamma \times (0, T).$$

Finally, let us mention that we make frequently use of the standard Young's inequality

$$ab \leq \delta a^2 + \frac{1}{4\delta} b^2 \quad \text{for all } a, b \in \mathbb{R} \text{ and all } \delta > 0$$

where $\delta > 0$ will be chosen when necessary and we write $C_\delta := \frac{1}{4\delta}$. Moreover, the symbols C , D , η and δ will denote positive constants throughout this work.

2.2 Notion of solutions

Let us consider two approximations of system (1)-(3): a regularized version where the indicator function $I_{(-\infty, 0]}$ in (2a) is replaced by a suitable smooth function I_β , $\beta \in (0, 1)$, and a time-discretized version of the regularized system. To this end, we introduce the following regularization:

Definition 2.3 (β -regularization) *Let the family of functions $\{I_\beta\}_{\beta \in (0, 1)} \subseteq C^{1,1}(\mathbb{R})$ denote a regularization of the indicator function $I_{(-\infty, 0]}$ in the following sense:*

- (i) $I_{\beta_1} \leq I_{\beta_2}$ pointwise in \mathbb{R} for every $\beta_1, \beta_2 \in (0, 1)$ with $\beta_1 \geq \beta_2$,
- (ii) $I_\beta \uparrow \infty$ pointwise in $[0, \infty)$ as $\beta \downarrow 0$,
- (iii) $I_\beta(x) = 0$ for all $x \leq 0$ and all $\beta \in (0, 1)$,
- (iv) $I_\beta''(x) \geq 0$ for a.e. $x \in \mathbb{R}$ and all $\beta \in (0, 1)$.

We may also write $\xi_\beta := I_\beta'$ in the following.

Remark 2.4 In particular, we may choose the Moreau-Yosida approximation given by (see [39, Lemma 5.17])

$$I_\beta(x) = \inf_{y \in \mathbb{R}} \left(\frac{|x-y|^2}{2\beta} + I_{(-\infty, 0]}(y) \right) = \begin{cases} 0 & \text{if } x \leq 0, \\ \frac{1}{2\beta}x^2 & \text{if } x > 0. \end{cases}$$

Let us mention that also C^∞ -approximations may be chosen for $\{I_\beta\}$ especially in view of optimality systems for optimal control problems (see [32, Chapter 5]).

Definition 2.5 (Strong solutions) For system (1)-(3) and their approximations we introduce the following notion of solutions:

(i) **Time-continuous limit system** ($\tau = 0, \beta = 0$).

Let the data $(u^0, v^0, \chi^0, b, \ell)$ be given. A solution of the time-continuous limit system is a pair of functions $(u, \chi) \in \mathcal{U} \times \mathcal{X}$ satisfying (1)-(3) in an a.e. sense and for a subgradient $\xi \in L^2(Q)$.

(ii) **Time-continuous β -regularized system** ($\tau = 0, \beta > 0$).

Let the data $(u^0, v^0, \chi^0, b, \ell)$ be given. A solution of the time-continuous β -regularized system is a pair of functions $(u, \chi) \in \mathcal{U} \times \mathcal{X}$ with $u(0) = u^0$, $\partial_t u(0) = v^0$ and $\chi(0) = \chi^0$ such that

$$u_{tt} - \operatorname{div}(\mathbb{C}(\chi)\varepsilon(u) + \mathbb{D}(\chi)\varepsilon(u_t)) = \ell \quad \text{a.e. in } Q, \quad (9a)$$

$$\chi_t - \Delta \chi_t - \Delta \chi + \xi_\beta(\chi_t) + \frac{1}{2}\mathbb{C}'(\chi)\varepsilon(u) : \varepsilon(u) + f'(\chi) = 0 \quad \text{a.e. in } Q, \quad (9b)$$

$$\chi \in [0, 1] \quad \text{a.e. in } Q, \quad (9c)$$

$$(\mathbb{C}(\chi)\varepsilon(u) + \mathbb{D}(\chi)\varepsilon(u_t)) \cdot \nu = b \quad \text{a.e. on } \Sigma, \quad (9d)$$

$$\nabla(\chi + \chi_t) \cdot \nu = 0 \quad \text{a.e. on } \Sigma. \quad (9e)$$

(iii) **Time-discrete β -regularized system** ($\tau, \beta > 0$).

Let $\{0, \tau, 2\tau, \dots, T\}$ denote an equidistant partition of $[0, T]$ with discretization fineness $\tau := T/M$ and $M \in \mathbb{N}$. Furthermore, let the data (u^0, u^{-1}, χ^0) , $\{b^k\}_{k=0, \dots, M}$ as well as $\{\ell^k\}_{k=0, \dots, M}$ be given. A solution of the time-discrete β -regularized system is a sequence $\{u^k, \chi^k\}_{k=0, \dots, M}$ of functions $u^k \in H^2(\Omega; \mathbb{R}^n)$ and $\chi^k \in H^2(\Omega)$ such that

$$\frac{u^k - 2u^{k-1} + u^{k-2}}{\tau^2} - \operatorname{div} \left(\mathbb{C}(\chi^k)\varepsilon(u^k) + \mathbb{D}(\chi^k)\varepsilon\left(\frac{u^k - u^{k-1}}{\tau}\right) \right) = \ell^k \quad \text{a.e. in } \Omega, \quad (10a)$$

$$\begin{aligned} \frac{\chi^k - \chi^{k-1}}{\tau} - \Delta \frac{\chi^k - \chi^{k-1}}{\tau} - \Delta \chi^k + \xi_\beta\left(\frac{\chi^k - \chi^{k-1}}{\tau}\right) \\ + \frac{1}{2}(\mathbf{c}'_1(\chi^k) + \mathbf{c}'_2(\chi^{k-1}))\mathbb{C}\varepsilon(u^{k-1}) : \varepsilon(u^{k-1}) + f'(\chi^k) = 0 \end{aligned} \quad \text{a.e. in } \Omega, \quad (10b)$$

$$\chi^k \in [0, 1] \quad \text{a.e. in } \Omega, \quad (10c)$$

$$\left(\mathbb{C}(\chi^k)\varepsilon(u^k) + \mathbb{D}(\chi^k)\varepsilon\left(\frac{u^k - u^{k-1}}{\tau}\right) \right) \cdot \nu = b^k \quad \text{a.e. on } \Gamma, \quad (10d)$$

$$\nabla\left(\chi^k + \frac{\chi^k - \chi^{k-1}}{\tau}\right) \cdot \nu = 0 \quad \text{a.e. on } \Gamma \quad (10e)$$

for all $k = 1, \dots, M$, where $\mathbf{c} = \mathbf{c}_1 + \mathbf{c}_2$ denotes the convex-concave decomposition from (A2).

Remark 2.6 If we assume $\nabla\chi^0 \cdot \nu = 0$ a.e. on Γ we even obtain

$$\nabla\chi \cdot \nu = \nabla\chi_t \cdot \nu = 0 \quad \text{a.e. on } \Sigma \quad (11)$$

instead of (3c) or (9e) and for all $k = 1, \dots, M$

$$\nabla\chi^k \cdot \nu = \nabla \frac{\chi^k - \chi^{k-1}}{\tau} \cdot \nu = 0 \quad \text{a.e. on } \Gamma \quad (12)$$

instead of (10e).

2.3 Existence of solutions

2.3.1 Existence for the time-discrete regularized system

At first we are going to show existence of time-discrete solution according to Definition 2.5 (iii). Let $\tau > 0$ and $\beta > 0$. To enhance readability, we will mostly omit the subscripts τ and β in $u_{\tau,\beta}^k$ and $\chi_{\tau,\beta}^k$.

Lemma 2.7 *Let the data $u^0, v^0 \in H^2(\Omega; \mathbb{R}^n)$, $\chi^0 \in H^2(\Omega)$ with $\chi^0 \in [0, 1]$ a.e. in Ω , $b^k \in H^{1/2}(\Gamma; \mathbb{R}^n)$ and $\ell^k \in L^2(\Omega; \mathbb{R}^n)$ for $k = 0, \dots, M$ be given. Then, there exists a strong solution $\{u^k, \chi^k\}_{k=0, \dots, M}$ of the time-discrete system in the sense of Definition 2.5 (iii).*

Proof. Starting from the initial values (u^0, u^{-1}, χ^0) with $u^{-1} := u^0 - \tau v^0$ we are going to construct $\{u^k, \chi^k\}_{k=0, \dots, M}$ by a recursive procedure. To this end, we decouple the discrete PDE problem into two distinct elliptic problems such that χ^k is obtained from χ^{k-1} and u^{k-1} , while u^k is gained from u^{k-1} , u^{k-2} , χ^k , b^k and ℓ^k .

Step 1: establishing equation (10b), (10c) and (10e)

We extend the function c_1 to $\tilde{c}_1 \in C^1(\mathbb{R}; \mathbb{R})$ in the following way

$$\tilde{c}_1(x) := \begin{cases} c_1(0) + xc_1'(0) & \text{if } x < 0, \\ c_1(x) & \text{if } x \in [0, 1], \\ c_1(1) + (x-1)c_1'(1) & \text{if } x > 1, \end{cases} \quad (13)$$

For convenience, we write c_1 instead of \tilde{c}_1 . The function f is extended in the same fashion as above. Note that due to Assumptions (A2) and (A4) we have

$$c_1(0) \leq c_1(x) \quad \text{for all } x \leq 0, \quad (14a)$$

$$c_1(1) + c_2'(z) \leq c_1(x) + c_2'(z)x \quad \text{for all } x \geq 1, z \in [0, 1], \quad (14b)$$

$$f(0) \leq f(x) \quad \text{for all } x \leq 0, \quad (14c)$$

$$f(1) \leq f(x) \quad \text{for all } x \geq 1. \quad (14d)$$

Let us give a proof for (14b): Consider the elementary estimate

$$-c_2'(z)(x-1) \leq \max_{y \in [0,1]} (-c_2'(y))(x-1)$$

holding for all $x \geq 1$ and all $z \in [0, 1]$. This is equivalent to

$$\mathbf{c}_1(1) + \mathbf{c}'_2(z) \leq \mathbf{c}_1(1) + \max_{y \in [0,1]} (-\mathbf{c}'_2(y))(x-1) + \mathbf{c}'_2(z)x$$

Now, taking the condition $\max_{y \in [0,1]} (-\mathbf{c}'_2(y)) \leq \mathbf{c}'_1(1)$ from (6a) into account yields the claim

$$\mathbf{c}_1(1) + \mathbf{c}'_2(z) \leq \underbrace{\mathbf{c}_1(1) + \mathbf{c}'_1(1)(x-1)}_{=\mathbf{c}_1(x)} + \mathbf{c}'_2(z)x.$$

Let us define the functional $\mathcal{F} : H^1(\Omega) \rightarrow \mathbb{R}$ by (note that $\chi^{k-1} \in [0, 1]$ a.e. in Ω)

$$\begin{aligned} \mathcal{F}(\chi) &:= \int_{\Omega} \left(\frac{1}{2} |\nabla \chi|^2 + \frac{1}{2} \mathbf{c}_1(\chi) \mathbf{C}\varepsilon(u^{k-1}) : \varepsilon(u^{k-1}) + \frac{1}{2} \mathbf{c}'_2(\chi^{k-1}) \chi \mathbf{C}\varepsilon(u^{k-1}) : \varepsilon(u^{k-1}) \right) dx \\ &\quad + \int_{\Omega} \left(f(\chi) + I_{\beta}((\chi - \chi^{k-1})\tau^{-1}) \right) dx + \frac{1}{2\tau} \int_{\Omega} |(\chi - \chi^{k-1})\tau^{-1}|^2 dx \\ &\quad + \frac{1}{2\tau} \int_{\Omega} |\nabla(\chi - \chi^{k-1})\tau^{-1}|^2 dx \end{aligned}$$

By the direct method in the calculus of variations, we obtain the existence of a minimizer of \mathcal{F} , which will be denoted by χ_m . Next, we show that the pointwise truncated function $\chi_m^{\#} := \min\{\chi_m^+, 1\}$ with $\chi_m^+ := \max\{\chi_m, 0\}$ is also a minimizer. The proof below extends the arguments used in [21, Proposition 5.5] (cf. Remark 2.1).

First of all, observe

$$\frac{1}{2} |\nabla \chi_m^{\#}|^2 \leq \frac{1}{2} |\nabla \chi_m|^2, \quad \left| \frac{\chi_m^{\#} - \chi^{k-1}}{\tau} \right|^2 \leq \left| \frac{\chi_m - \chi^{k-1}}{\tau} \right|^2 \quad (15)$$

and, by using (14c)-(14d),

$$f(\chi_m^{\#}) \leq f(\chi_m). \quad (16)$$

Due to $\chi^{k-1} \in [0, 1]$ and $I_{\beta}(x) = 0$ for $x \leq 0$ and the monotone increasingness of I_{β} , it follows

$$I_{\beta}\left(\frac{\chi_m^{\#} - \chi^{k-1}}{\tau}\right) \leq I_{\beta}\left(\frac{\chi_m - \chi^{k-1}}{\tau}\right). \quad (17)$$

and by (14a)

$$\mathbf{c}_1(\chi_m^+) \mathbf{C}\varepsilon(u^{k-1}) : \varepsilon(u^{k-1}) \leq \mathbf{c}_1(\chi_m) \mathbf{C}\varepsilon(u^{k-1}) : \varepsilon(u^{k-1}). \quad (18)$$

Because of $\mathbf{c}'_2(\chi^{k-1}) \leq 0$ (due to (6b) and $\chi^{k-1} \in [0, 1]$), we directly obtain

$$\mathbf{c}'_2(\chi^{k-1}) \chi_m^+ \mathbf{C}\varepsilon(u^{k-1}) : \varepsilon(u^{k-1}) \leq \mathbf{c}'_2(\chi^{k-1}) \chi_m \mathbf{C}\varepsilon(u^{k-1}) : \varepsilon(u^{k-1}). \quad (19)$$

Adding (18) and (19) yields

$$\begin{aligned} &(\mathbf{c}_1(\chi_m^+) + \mathbf{c}'_2(\chi^{k-1}) \chi_m^+) \mathbf{C}\varepsilon(u^{k-1}) : \varepsilon(u^{k-1}) \\ &\leq (\mathbf{c}_1(\chi_m) + \mathbf{c}'_2(\chi^{k-1}) \chi_m) \mathbf{C}\varepsilon(u^{k-1}) : \varepsilon(u^{k-1}). \end{aligned} \quad (20)$$

Moreover, from (14b) we infer

$$\begin{aligned} & (\mathbf{c}_1(\chi_m^\#) + \mathbf{c}_2(\chi^{k-1})\chi_m^\#)\mathbf{C}\varepsilon(u^{k-1}) : \varepsilon(u^{k-1}) \\ & \leq (\mathbf{c}_1(\chi_m^+) + \mathbf{c}_2(\chi^{k-1})\chi_m^+)\mathbf{C}\varepsilon(u^{k-1}) : \varepsilon(u^{k-1}). \end{aligned} \quad (21)$$

Combining the estimates (20) and (21), we have

$$\begin{aligned} & (\mathbf{c}_1(\chi_m^\#) + \mathbf{c}_2(\chi^{k-1})\chi_m^\#)\mathbf{C}\varepsilon(u^{k-1}) : \varepsilon(u^{k-1}) \\ & \leq (\mathbf{c}_1(\chi_m) + \mathbf{c}_2(\chi^{k-1})\chi_m)\mathbf{C}\varepsilon(u^{k-1}) : \varepsilon(u^{k-1}). \end{aligned} \quad (22)$$

Taking (15), (16), (17) and (22) into account, we end up with $\mathcal{F}(\chi_m^\#) \leq \mathcal{F}(\chi_m)$, i.e., $\chi_m^\#$ is also a minimizer of \mathcal{F} .

The Euler-Lagrange equation for the minimizer yields (10b) in a weak form by setting $\chi^k := \chi_m^\#$. By noticing that (10b) is a elliptic equation for χ^k with right hand side in $L^2(\Omega)$, we conclude $\chi^k \in H^2(\Omega)$ by elliptic regularity results for homogeneous Neumann problems (see, e.g. [15, Theorem 2.4.2.7] and remember that Γ is a C^2 -boundary by Assumption (A1)).

Step 2: establishing equation (10a)

Given the functions $\chi^k \in H^2(\Omega)$, $u^{k-1}, u^{k-2} \in H^2(\Omega; \mathbb{R}^n)$, $b^k \in H^{1/2}(\Gamma; \mathbb{R}^n)$ and $\ell^k \in L^2(\Omega; \mathbb{R}^n)$, we obtain a unique weak solution $u^k \in H^1(\Omega; \mathbb{R}^n)$ of the linear elliptic system (10a) via the well-known Lax-Milgram theorem (remember the assumption $\mathbf{D} = \mu\mathbf{C}$ from (A3)):

$$\begin{aligned} & \int_{\Omega} \left((\tau^2\mathbf{c}(\chi^k) + \tau\mathbf{d}(\chi^k)\mu)\mathbf{C}\varepsilon(u) : \varepsilon(\zeta) + u \cdot \zeta \right) dx \\ & = \int_{\Omega} \left(\tau^2\ell^k - \tau \operatorname{div}(\mathbf{d}(\chi^k)\mathbf{D}\varepsilon(u^{k-1})) + 2u^{k-1} - u^{k-2} \right) \cdot \zeta dx + \int_{\Gamma} b^k \cdot \zeta dx \end{aligned} \quad (23)$$

holding for all $\zeta \in H^1(\Omega; \mathbb{R}^n)$.

Now we use a modification of the regularity argument in [18, Proof of Lemma 4.1] and make use of the C^2 -regularity of Γ (see (A1)):

If we consider the test-function $\zeta = (\tau^2\mathbf{c}(\chi^k) + \tau\mathbf{d}(\chi^k)\mu)^{-1}\varphi$ where $\varphi \in H^1(\Omega; \mathbb{R}^n)$ is another test-function (remember that $\mathbf{c} \geq 0$ and $\mathbf{d} \geq \eta > 0$, see (A2)-(A3)) the linear elliptic system (23) rewrites to

$$\mathbf{a}(u^k, \varphi) = \langle \mathbf{q}, \varphi \rangle_{H^1}, \quad \varphi \in H^1(\Omega; \mathbb{R}^n). \quad (24)$$

with the bilinear form

$$\mathbf{a}(u, \varphi) := \int_{\Omega} \mathbf{C}\varepsilon(u) : \varepsilon(\varphi) dx$$

and the right hand side $\mathbf{q} \in H^1(\Omega; \mathbb{R}^n)'$ given by

$$\langle \mathbf{q}, \varphi \rangle_{H^1} := \int_{\Omega} R \cdot \varphi dx + \int_{\Gamma} N \cdot \varphi dx,$$

where R and N are defined as

$$R := \frac{\tau^2\mathbf{c}'(\chi^k) + \tau\mu\mathbf{d}'(\chi^k)}{\tau^2\mathbf{c}(\chi^k) + \tau\mathbf{d}(\chi^k)\mu} \nabla\chi^k \cdot \mathbf{C}\varepsilon(u^k) + \frac{\tau^2\ell^k - \tau \operatorname{div}(\mathbf{d}(\chi^k)\mathbf{D}\varepsilon(u^{k-1})) - u^k + 2u^{k-1} - u^{k-2}}{\tau^2\mathbf{c}(\chi^k) + \tau\mathbf{d}(\chi^k)\mu},$$

$$N := \frac{b^k}{\tau^2 c(\chi^k) + \tau d(\chi^k) \mu}.$$

Note that $N \in H^{1/2}(\Gamma; \mathbb{R}^n)$ and $R \in L^p(\Omega; \mathbb{R}^n)$ for all $p \in (1, 2)$, since $\varepsilon(u^k) \in L^2(\Omega; \mathbb{R}^{n \times n})$ and $\nabla \chi^k \in L^q(\Omega; \mathbb{R}^n)$ for all $q \in \mathbb{N}$.

In particular, $\mathbf{q} \in H^{2-s}(\Omega; \mathbb{R}^n)'$ for all $s \in (1, 3/2)$. We gain $u^k \in H^s(\Omega; \mathbb{R}^n)$ by applying the lower Sobolev H^s -regularity result from [6, Theorem 3.4.5 (ii)]. This, in turn, implies $\varepsilon(u^k) \in L^{2^*}(\Omega; \mathbb{R}^n)$ with the *fractional critical exponent* given in this case by $2^* = \frac{2n}{n-(s-1)2} > 2$ (see, e.g., [33, Theorem 6.7]). We obtain $R \in L^2(\Omega; \mathbb{R}^n)$. The H^2 -regularity result [6, Theorem 3.4.1] applied to the linear elliptic system (24) shows $u^k \in H^2(\Omega; \mathbb{R}^n)$. Thus (10a) is shown. \square

2.3.2 Existence result for the time-continuous system

The aim of this section is to provide existence of strong solutions in the sense of Definition 2.5 (i) and (ii). To this end, several a priori estimates for the time-discrete solutions will be established. The estimates will be used for the time-continuous limit analysis and for the optimal control problem in Section 3.

We assume that the initial data (u^0, v^0, χ^0) satisfy

$$u^0 \in H^2(\Omega; \mathbb{R}^n), \tag{25a}$$

$$v^0 \in H^1(\Omega; \mathbb{R}^n), \tag{25b}$$

$$\chi^0 \in H^2(\Omega) \text{ with } \chi^0 \in [0, 1] \text{ a.e. in } \Omega \text{ and } \nabla \chi^0 \cdot \nu = 0 \text{ a.e. on } \Gamma \tag{25c}$$

and the external forces (b, ℓ) are assumed to be in the following spaces:

$$b \in L^2(0, T; H^{1/2}(\Gamma; \mathbb{R}^n)) \cap H^1(0, T; L^2(\Gamma; \mathbb{R}^n)), \quad \ell \in L^2(0, T; L^2(\Omega; \mathbb{R}^n)). \tag{26}$$

For the moment, let us consider some approximations

$$\begin{aligned} \{v_\lambda^0\}_{\lambda \in (0,1)} &\subseteq H^2(\Omega; \mathbb{R}^n), \\ \{b_\lambda\}_{\lambda \in (0,1)} &\subseteq C^{1,1}(0, T; H^{1/2}(\Gamma; \mathbb{R}^n)), \\ \{\ell_\lambda\}_{\lambda \in (0,1)} &\subseteq C^{0,1}(0, T; L^2(\Omega; \mathbb{R}^n)) \end{aligned}$$

of the the initial velocity v^0 and the external forces b and ℓ such that (e.g. construction via convolution)

$$v_\lambda^0 \rightarrow v^0 \text{ strongly in } H^1(\Omega; \mathbb{R}^n), \tag{27a}$$

$$b_\lambda \rightarrow b \text{ strongly in } L^2(0, T; H^{1/2}(\Gamma; \mathbb{R}^n)) \cap H^1(0, T; L^2(\Gamma; \mathbb{R}^n)), \tag{27b}$$

$$\ell_\lambda \rightarrow \ell \text{ strongly in } L^2(0, T; L^2(\Omega; \mathbb{R}^n)) \tag{27c}$$

as $\lambda \downarrow 0$. Let us define the time-discretizations $b_{\tau,\lambda}^k$ and $\ell_{\tau,\lambda}^k$ by

$$b_{\tau,\lambda}^k := b_\lambda(\tau k), \quad \ell_{\tau,\lambda}^k := \ell_\lambda(\tau k).$$

For a sequence $\{h^k\}_{k=0,\dots,M}$ where $h^k \in \{u_{\tau,\beta}^k, \chi_{\tau,\beta}^k, b_{\tau,\lambda}^k, \ell_{\tau,\lambda}^k\}$, we define the piecewise constant and linear interpolation as

$$\left. \begin{aligned} \bar{h}(t) &:= h^k, & \underline{h}(t) &:= h^{k-1}, & \underline{\underline{h}}(t) &:= h^{k-2}, \\ h(t) &:= \frac{t - (k-1)\tau}{\tau} h^k + \frac{k\tau - t}{\tau} h^{k-1} \end{aligned} \right\} \quad \text{for } t \in ((k-1)\tau, k\tau]. \quad (28)$$

The left-continuous and right-continuous piecewise constant interpolation for a given time point t is denoted by

$$\begin{aligned} \bar{t}_\tau &:= \tau k & \text{for } \tau(k-1) < t \leq \tau k, \\ \underline{t}_\tau &:= \tau k & \text{for } \tau k \leq t < \tau(k+1). \end{aligned}$$

For notational convenience, we define the time-discrete velocity field and their interpolations by

$$v_{\tau,\beta}^k := \frac{u_{\tau,\beta}^k - u_{\tau,\beta}^{k-1}}{\tau} \quad \text{for } k = 0, \dots, M \quad \text{and} \quad \bar{v}_{\tau,\beta}, \underline{v}_{\tau,\beta}, v_{\tau,\beta} \quad \text{by (28)}. \quad (29)$$

As a first result, we prove convergence of the discretizations of the given data.

Lemma 2.8 *There exist subsequences $\tau_k \downarrow 0$ and $\lambda_k \downarrow 0$ as $k \uparrow \infty$ such that*

$$\begin{aligned} v_{\tau_k}^0 &\rightarrow v^0 \quad \text{strongly in } H^1(\Omega; \mathbb{R}^n), \\ b_{\tau_k, \lambda_k} &\rightarrow b \quad \text{strongly in } L^2(0, T; H^{1/2}(\Gamma; \mathbb{R}^n)) \cap H^1(0, T; L^2(\Gamma; \mathbb{R}^n)), \\ \bar{\ell}_{\tau_k, \lambda_k} &\rightarrow \ell \quad \text{strongly in } L^2(0, T; L^2(\Omega; \mathbb{R}^n)) \end{aligned}$$

as $k \uparrow \infty$. For readers' convenience we set $b_{\tau_k} := b_{\tau_k, \lambda_k}$ and $\bar{\ell}_{\tau_k} := \bar{\ell}_{\tau_k, \lambda_k}$ and omit the subscript k . Then the statement above reads as $v_\tau^0 \rightarrow v^0$, $b_\tau \rightarrow b$ and $\bar{\ell}_\tau \rightarrow \ell$ as $\tau \downarrow 0$.

Proof. For every fixed $\lambda > 0$, we find

$$b_{\tau,\lambda} \rightarrow b_\lambda \quad \text{strongly in } L^2(0, T; H^{1/2}(\Gamma; \mathbb{R}^n)) \cap H^1(0, T; L^2(\Gamma; \mathbb{R}^n)), \quad (30a)$$

$$\bar{\ell}_{\tau,\lambda} \rightarrow \ell_\lambda \quad \text{strongly in } L^2(0, T; L^2(\Omega; \mathbb{R}^n)) \quad (30b)$$

as $\tau \downarrow 0$. Indeed, the first convergence in (30a) follows by exploiting the Lipschitz continuity of $b_\lambda \in C^{0,1}(0, T; H^{1/2}(\Gamma; \mathbb{R}^n))$. Property (30b) can be proven with a similar argument. The convergence $b_{\tau,\lambda} \rightarrow b_\lambda$ in the $H^1(0, T; L^2(\Gamma; \mathbb{R}^n))$ -norm follows by the fundamental theorem of calculus for X -valued functions where $X := L^2(\Gamma; \mathbb{R}^n)$ and by the Lipschitz continuity of $\partial_t b_\lambda \in C^{0,1}(0, T; X)$:

$$\begin{aligned} \int_0^T \left\| \partial_t b_{\tau,\lambda}(t) - \partial_t b_\lambda(t) \right\|_X^2 dt &= \int_0^T \left\| \frac{b_\lambda(\bar{t}_\tau) - b_\lambda(\underline{t}_\tau)}{\tau} - \partial_t b_\lambda(t) \right\|_X^2 dt \\ &= \int_0^T \left\| \frac{1}{\tau} \int_{\underline{t}_\tau}^{\bar{t}_\tau} (\partial_t b_\lambda(s) - \partial_t b_\lambda(t)) ds \right\|_X^2 dt \\ &\leq \int_0^T \left(\frac{1}{\tau} \int_{\underline{t}_\tau}^{\bar{t}_\tau} \|\partial_t b_\lambda(s) - \partial_t b_\lambda(t)\|_X ds \right)^2 dt \\ &\leq C \int_0^T \left(\frac{1}{\tau} \int_{\underline{t}_\tau}^{\bar{t}_\tau} |s - t| ds \right)^2 dt \end{aligned}$$

$$\leq CT\tau^2.$$

The claim follows by using the convergences (27), (30) and an “ $\varepsilon/2$ ”-argument. \square

Lemma 2.9 (A priori estimates for the time-discrete system) *The following a priori estimates hold for strong solutions of the time-discrete system given in Definition 2.5 (iii) (recall that (29) implies $\bar{v}_{\tau,\beta} = \partial_t u_{\tau,\beta}$):*

(i) *First a priori estimate:*

There exists a constant $C > 0$ which continuously depends on

$$C = C(\|u^0\|_{H^1}, \|v^0\|_{L^2}, \|\chi^0\|_{H^1}, \|b\|_{L^2(0,T;L^2(\Gamma;\mathbb{R}^n))}, \|\ell\|_{L^2(0,T;L^2)}) \quad (31)$$

such that for all $\tau, \beta > 0$

$$\begin{aligned} \|u_{\tau,\beta}\|_{H^1(0,T;H^1)\cap W^{1,\infty}(0,T;L^2)} &\leq C, & \|\chi_{\tau,\beta}\|_{H^1(0,T;H^1)} &\leq C, \\ \|\underline{u}_{\tau,\beta}\|_{L^\infty(0,T;H^1)} &\leq C, & \|\underline{\chi}_{\tau,\beta}\|_{L^\infty(0,T;H^1)} &\leq C, \\ \|\bar{u}_{\tau,\beta}\|_{L^\infty(0,T;H^1)} &\leq C, & \|\bar{\chi}_{\tau,\beta}\|_{L^\infty(0,T;H^1)} &\leq C. \end{aligned}$$

(ii) *Second a priori estimate:*

There exists a constant $D > 0$ which continuously depends on

$$D = D(\|u^0\|_{H^2}, \|v^0\|_{H^1}, \|\chi^0\|_{H^2}, \|b\|_{L^2(0,T;H^{1/2}(\Gamma;\mathbb{R}^n))\cap H^1(0,T;L^2(\Gamma;\mathbb{R}^n))}, \|\ell\|_{L^2(0,T;L^2)})$$

such that for all $\tau, \beta > 0$

$$\begin{aligned} \|u_{\tau,\beta}\|_{H^1(0,T;H^2)\cap W^{1,\infty}(0,T;H^1)} &\leq D, & \|\chi_{\tau,\beta}\|_{H^1(0,T;H^2)} &\leq D, \\ \|\underline{u}_{\tau,\beta}\|_{L^\infty(0,T;H^2)} &\leq D, & \|\underline{\chi}_{\tau,\beta}\|_{L^\infty(0,T;H^2)} &\leq D, \\ \|\bar{u}_{\tau,\beta}\|_{L^\infty(0,T;H^2)} &\leq D, & \|\bar{\chi}_{\tau,\beta}\|_{L^\infty(0,T;H^2)} &\leq D, \\ \|v_{\tau,\beta}\|_{L^2(0,T;H^2)\cap L^\infty(0,T;H^1)\cap H^1(0,T;L^2)} &\leq D, & \|\xi_\beta(\partial_t \chi_{\tau,\beta})\|_{L^2(0,T;L^2)} &\leq D. \end{aligned}$$

Proof. We will omit the subscript τ and β in the time-discrete solutions.

To (i): In the following, we make use of a combined convex-concave estimate for: A convexity estimate for c_1 and concavity estimate for c_2 yield:

$$\begin{aligned} c_1(\chi^{k-1}) - c_1(\chi^k) &\geq c'_1(\chi^k)(\chi^{k-1} - \chi^k) \\ c_2(\chi^{k-1}) - c_2(\chi^k) &\geq c'_2(\chi^{k-1})(\chi^{k-1} - \chi^k). \end{aligned}$$

Adding them shows

$$\begin{aligned} c(\chi^{k-1}) - c(\chi^k) &= (c_1(\chi^{k-1}) - c_1(\chi^k)) + (c_2(\chi^{k-1}) - c_2(\chi^k)) \\ &\geq (c'_1(\chi^k) + c'_2(\chi^{k-1}))(\chi^{k-1} - \chi^k). \end{aligned}$$

By using this combined estimate and the positivity of \mathbf{C} , it holds

$$\begin{aligned}
& c(\chi^k) \mathbf{C} \varepsilon(u^k) : \varepsilon(u^k - u^{k-1}) \\
&= \frac{1}{2} c(\chi^k) \left(\mathbf{C} \varepsilon(u^k) : \varepsilon(u^k) - \mathbf{C} \varepsilon(u^{k-1}) : \varepsilon(u^{k-1}) + \mathbf{C} \varepsilon(u^k - u^{k-1}) : \varepsilon(u^k - u^{k-1}) \right) \\
&\geq \frac{c(\chi^k)}{2} \mathbf{C} \varepsilon(u^k) : \varepsilon(u^k) - \frac{c(\chi^{k-1})}{2} \mathbf{C} \varepsilon(u^{k-1}) : \varepsilon(u^{k-1}) \\
&\quad + \frac{1}{2} (c(\chi^{k-1}) - c(\chi^k)) \mathbf{C} \varepsilon(u^{k-1}) : \varepsilon(u^{k-1}) \\
&\geq \frac{c(\chi^k)}{2} \mathbf{C} \varepsilon(u^k) : \varepsilon(u^{k-1}) - \frac{c(\chi^{k-1})}{2} \mathbf{C} \varepsilon(u^{k-1}) : \varepsilon(u^{k-1}) \\
&\quad + \frac{1}{2} (c'_1(\chi^k) + c'_2(\chi^{k-1})) (\chi^{k-1} - \chi^k) \mathbf{C} \varepsilon(u^{k-1}) : \varepsilon(u^{k-1}). \tag{32}
\end{aligned}$$

Now, by testing equation (10a) with $u^k - u^{k-1}$, integrating over Ω , summing over the time index $k = 1, \dots, \bar{t}_\tau/\tau$, integrating by parts and using (10d), we obtain (remember that $v^k = (u^k - u^{k-1})/\tau$)

$$\begin{aligned}
& \sum_{k=1}^{\bar{t}_\tau/\tau} \int_{\Omega} (v^k - v^{k-1}) v^k \, dx + \sum_{k=1}^{\bar{t}_\tau/\tau} \int_{\Omega} c(\chi^k) \mathbf{C} \varepsilon(u^k) : \varepsilon(u^k - u^{k-1}) \, dx \\
&+ \int_0^{\bar{t}_\tau} \int_{\Omega} d(\chi^k) \mathbf{D} \varepsilon(\bar{v}) : \varepsilon(\bar{v}) \, dx \, ds \\
&= \int_0^{\bar{t}_\tau} \int_{\Omega} \bar{\ell} \cdot \bar{v} \, dx \, ds + \int_0^{\bar{t}_\tau} \int_{\Gamma} \bar{b} \cdot \bar{v} \, dx \, ds.
\end{aligned}$$

Applying elementary estimates including the convex-concave estimate (32), Korn's and Young's inequality and the trace theorem $H^1(\Omega; \mathbb{R}^n) \hookrightarrow L^2(\Gamma; \mathbb{R}^n)$ yield ($\eta, \delta, C_\delta > 0$ are constants)

$$\begin{aligned}
& \frac{1}{2} \|\bar{v}(t)\|_{L^2}^2 - \frac{1}{2} \|v^0\|_{L^2}^2 + \underbrace{\int_{\Omega} \frac{c(\bar{\chi}(t))}{2} \mathbf{C} \varepsilon(\bar{u}(t)) : \varepsilon(\bar{u}(t)) \, dx}_{\geq 0 \text{ by using (A2)}} - \int_{\Omega} \frac{c(\chi^0)}{2} \mathbf{C} \varepsilon(u^0) : \varepsilon(u^0) \, dx \\
&+ \int_0^{\bar{t}_\tau} \int_{\Omega} \frac{1}{2} (c'_1(\bar{\chi}) + c'_2(\underline{\chi})) (-\partial_t \chi) \mathbf{C} \varepsilon(\underline{u}) : \varepsilon(\underline{u}) \, dx \, ds + \eta \|\varepsilon(\bar{v})\|_{L^2(0, \bar{t}_\tau; L^2)}^2 \\
&\leq C_\delta \|\bar{\ell}\|_{L^2(0, \bar{t}_\tau; L^2)}^2 + C_\delta \|\bar{b}\|_{L^2(0, \bar{t}_\tau; L^2(\Gamma; \mathbb{R}^n))}^2 + \delta \|\bar{v}\|_{L^2(0, \bar{t}_\tau; H^1)}^2. \tag{33}
\end{aligned}$$

Testing equation (10b) with $\chi^k - \chi^{k-1}$, integrating over Ω , summing over the time index $k = 1, \dots, \bar{t}_\tau/\tau$, integrating by parts and using (10e), we obtain

$$\begin{aligned}
& \|\partial_t \chi\|_{L^2(0, \bar{t}_\tau; L^2)}^2 + \|\nabla \partial_t \chi\|_{L^2(0, \bar{t}_\tau; L^2)}^2 + \sum_{k=1}^{\bar{t}_\tau/\tau} \int_{\Omega} \nabla \chi^k \cdot \nabla (\chi^k - \chi^{k-1}) \, dx \\
&+ \sum_{k=1}^{\bar{t}_\tau/\tau} \int_{\Omega} \xi_\beta \left(\frac{\chi^k - \chi^{k-1}}{\tau} \right) (\chi^k - \chi^{k-1}) \, dx + \int_0^{\bar{t}_\tau} \int_{\Omega} \frac{1}{2} (c'_1(\bar{\chi}) + c'_2(\underline{\chi})) \partial_t \chi \mathbf{C} \varepsilon(\underline{u}) : \varepsilon(\underline{u}) \, dx \, ds \\
&+ \int_0^{\bar{t}_\tau} \int_{\Omega} f'(\bar{\chi}) \partial_t \chi \, dx \, ds
\end{aligned}$$

= 0

By using the monotonicity of ξ_β (see Definition 2.3), we get $\xi_\beta\left(\frac{\chi^k - \chi^{k-1}}{\tau}\right)(\chi^k - \chi^{k-1}) \geq 0$. Together with elementary convexity estimates and Young's inequality, we find

$$\begin{aligned} & \|\partial_t \chi\|_{L^2(0, \bar{t}_\tau; L^2)}^2 + \|\nabla \partial_t \chi\|_{L^2(0, \bar{t}_\tau; L^2)}^2 + \frac{1}{2} \|\nabla \bar{\chi}(t)\|_{L^2}^2 - \frac{1}{2} \|\nabla \chi^0\|_{L^2}^2 \\ & + \int_0^{\bar{t}_\tau} \int_\Omega \frac{1}{2} (c'_1(\bar{\chi}) + c'_2(\underline{\chi})) \partial_t \chi \mathbf{C} \varepsilon(\underline{u}) : \varepsilon(\underline{u}) \, dx \, ds \\ & \leq C_\delta \|f'(\bar{\chi})\|_{L^2(0, \bar{t}_\tau; L^2)}^2 + \delta \|\partial_t \chi\|_{L^2(0, \bar{t}_\tau; L^2)}^2. \end{aligned} \quad (34)$$

Adding (34) and (33), we see that the term

$$\int_0^{\bar{t}_\tau} \int_\Omega \frac{1}{2} (c'_1(\bar{\chi}) + c'_2(\underline{\chi})) \partial_t \chi \mathbf{C} \varepsilon(\underline{u}) : \varepsilon(\underline{u}) \, dx \, ds$$

cancels out in the calculations and we obtain

$$\begin{aligned} & \|\bar{v}(t)\|_{L^2}^2 + \|\nabla \bar{\chi}(t)\|_{L^2}^2 + \|\varepsilon(\bar{v})\|_{L^2(0, \bar{t}_\tau; L^2)}^2 + \|\partial_t \chi\|_{L^2(0, \bar{t}_\tau; L^2)}^2 + \|\nabla \partial_t \chi\|_{L^2(0, \bar{t}_\tau; L^2)}^2 \\ & \leq C_\delta \left(\|u^0\|_{H^1}^2 + \|v^0\|_{L^2}^2 + \|\chi^0\|_{H^1}^2 + \|\bar{\ell}\|_{L^2(0, \bar{t}_\tau; L^2)}^2 + \|\bar{b}\|_{L^2(0, \bar{t}_\tau; L^2(\Gamma; \mathbb{R}^n))}^2 + \|f'(\bar{\chi})\|_{L^2(0, \bar{t}_\tau; L^2)}^2 \right) \\ & + \delta \left(\|\bar{v}\|_{L^2(0, \bar{t}_\tau; H^1)}^2 + \|\partial_t \chi\|_{L^2(0, \bar{t}_\tau; L^2)}^2 \right). \end{aligned} \quad (35)$$

Korn's inequality yields

$$\|\varepsilon(\bar{v})\|_{L^2(0, \bar{t}_\tau; L^2)}^2 \geq \frac{1}{C} \|\bar{v}\|_{L^2(0, \bar{t}_\tau; H^1)}^2 - \|\bar{v}\|_{L^2(0, \bar{t}_\tau; L^2)}^2.$$

Now, choosing $\delta > 0$ small and noticing $\bar{\chi} \in [0, 1]$ a.e. in $\Omega \times (0, T)$, we obtain from (35)

$$\begin{aligned} & \|\bar{v}(t)\|_{L^2}^2 + \|\bar{\chi}(t)\|_{H^1}^2 + \|\bar{v}\|_{L^2(0, \bar{t}_\tau; H^1)}^2 + \|\partial_t \chi\|_{L^2(0, \bar{t}_\tau; H^1)}^2 \\ & \leq C \left(\|u^0\|_{H^1}^2 + \|v^0\|_{L^2}^2 + \|\chi^0\|_{H^1}^2 + \|\bar{\ell}\|_{L^2(0, \bar{t}_\tau; L^2)}^2 + \|\bar{b}\|_{L^2(0, \bar{t}_\tau; L^2(\Gamma; \mathbb{R}^n))}^2 + 1 \right) \\ & + C \int_0^{\bar{t}_\tau} \|\bar{v}\|_{L^2}^2 \, ds. \end{aligned}$$

We end up with the desired estimates in (i) by using a discrete version of Gronwall's lemma and

$$\|\bar{u}(t)\|_{H^1}^2 = \left\| u^0 + \int_0^{\bar{t}_\tau} \bar{v} \, ds \right\|_{H^1}^2 \leq C (\|u^0\|_{H^1}^2 + \|\bar{v}\|_{L^2(0, \bar{t}_\tau; H^1)}^2)$$

afterwards.

To (ii) – local-in-time result:

At first we are going to show the a priori estimates in (ii) for small time. In the next step global-in-time estimates will be derived.

Testing equation (10a) with $-\tau \operatorname{div} \left(c(\chi^k) \mathbf{C} \varepsilon(u^k) + d(\chi^k) \mathbf{D} \varepsilon \left(\frac{u^k - u^{k-1}}{\tau} \right) \right)$, integrating over Ω in space and summing over the time index $k = 1, \dots, \bar{t}_\tau/\tau$, we may write the result in the following way

$$\underbrace{\int_0^{\bar{t}_\tau} \int_\Omega -\partial_t v \cdot \operatorname{div} (c(\bar{\chi}) \mathbf{C} \varepsilon(\bar{u}) + d(\bar{\chi}) \mathbf{D} \varepsilon(\bar{v})) \, dx \, ds}_{=: T_1}$$

$$\begin{aligned}
& + \int_0^{\bar{t}_\tau} \int_\Omega \frac{1}{2} \left| \operatorname{div}(\mathbf{c}(\bar{\chi}) \mathbf{C}\varepsilon(\bar{u}) + \mathbf{d}(\bar{\chi}) \mathbf{D}\varepsilon(\bar{v})) \right|^2 dx ds \\
& + \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega \frac{1}{2} \left| \mathbf{c}'(\bar{\chi}) \nabla \bar{\chi} \cdot \mathbf{C}\varepsilon(\bar{u}) + \mathbf{c}(\bar{\chi}) \operatorname{div}(\mathbf{C}\varepsilon(\bar{u})) + \mathbf{d}'(\bar{\chi}) \nabla \bar{\chi} \cdot \mathbf{D}\varepsilon(\bar{v}) + \mathbf{d}(\bar{\chi}) \operatorname{div}(\mathbf{D}\varepsilon(\bar{v})) \right|^2}_{=:T_2} dx ds \\
& = \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega -\bar{\ell} \cdot \operatorname{div}(\mathbf{c}(\bar{\chi}) \mathbf{C}\varepsilon(\bar{u}) + \mathbf{d}(\bar{\chi}) \mathbf{D}\varepsilon(\bar{v}))}_{\leq C_\delta \|\bar{\ell}\|_{L^2(L^2)}^2 + \delta \|\operatorname{div}(\mathbf{c}(\bar{\chi}) \mathbf{C}\varepsilon(\bar{u}) + \mathbf{d}(\bar{\chi}) \mathbf{D}\varepsilon(\bar{v}))\|_{L^2(L^2)}^2} dx ds. \tag{36}
\end{aligned}$$

Note that the second summand and the third summand, i.e. T_2 , are identical. The splitting will simplify the calculations.

Testing equation (10b) with $-\Delta(\chi^k - \chi^{k-1})$, integrating over Ω in space and summing over $k = 1, \dots, \bar{t}_\tau/\tau$, we obtain

$$\begin{aligned}
& - \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega \partial_t \chi \Delta \partial_t \chi dx ds}_{=\|\nabla \partial_t \chi\|_{L^2(L^2)}^2 \text{ by using (12)}} + \underbrace{\|\Delta \partial_t \chi\|_{L^2(0, \bar{t}_\tau; L^2)}^2 + \sum_{k=1}^{\bar{t}_\tau/\tau} \int_\Omega \Delta \chi^k (\Delta \chi^k - \Delta \chi^{k-1}) dx}_{\geq \frac{1}{2} \|\Delta \bar{\chi}(t)\|_{L^2}^2 - \frac{1}{2} \|\Delta \chi^0\|_{L^2}^2} \\
& - \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega \xi_\beta(\partial_t \chi) \Delta \partial_t \chi dx ds}_{=:T_3} - \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega \frac{1}{2} (\mathbf{c}'_1(\bar{\chi}) + \mathbf{c}'_2(\underline{\chi})) \mathbf{C}\varepsilon(\underline{u}) : \varepsilon(\underline{u}) \Delta \partial_t \chi dx ds}_{=:T_4} \\
& - \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega f'(\bar{\chi}) \Delta \partial_t \chi dx ds}_{=:T_5} \\
& = 0. \tag{37}
\end{aligned}$$

In the following, we are going to estimate T_1, \dots, T_7 and conclude the claimed a priori estimates thereafter:

– To (T_1) : Integration by parts in space yields

$$\begin{aligned}
T_1 & = \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega \varepsilon(\partial_t v) : \mathbf{c}(\bar{\chi}) \mathbf{C}\varepsilon(\bar{u}) dx ds}_{=:T_1^{(1)}} + \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega \varepsilon(\partial_t v) : \mathbf{d}(\bar{\chi}) \mathbf{D}\varepsilon(\bar{v}) dx ds}_{=:T_1^{(2)}} \\
& - \underbrace{\int_0^{\bar{t}_\tau} \int_\Gamma \partial_t v \cdot ((\mathbf{c}(\bar{\chi}) \mathbf{C}\varepsilon(\bar{u}) + \mathbf{d}(\bar{\chi}) \mathbf{D}\varepsilon(\bar{v})) \cdot \nu) dx ds}_{=:T_1^{(3)}}
\end{aligned}$$

Note that we have no compensating $\partial_t v$ -term on the left-hand side of (36). To circumvent this problem we rewrite the term $T_1^{(1)}$ by using the discrete integration by parts formula in time

$$\sum_{k=1}^N \tau \frac{a^k - a^{k-1}}{\tau} b^k = a^N b^N - a^0 b^0 - \sum_{k=1}^N \tau a^{k-1} \frac{b^k - b^{k-1}}{\tau}.$$

Together with the Lipschitz continuity of \mathbf{c} we find:

$$\begin{aligned}
T_1^{(1)} &= - \int_0^{\bar{t}_\tau} \int_\Omega \varepsilon(\underline{v}) : \frac{\mathbf{c}(\bar{\chi})\mathbf{C}\varepsilon(\bar{u}) - \mathbf{c}(\underline{\chi})\mathbf{C}\varepsilon(\underline{u})}{\tau} dx ds \\
&\quad + \int_\Omega \varepsilon(\bar{v}(t)) : \mathbf{c}(\bar{\chi}(t))\mathbf{C}\varepsilon(\bar{u}(t)) dx - \int_\Omega \varepsilon(v^0) : \mathbf{c}(\chi^0)\mathbf{C}\varepsilon(u^0) dx \\
&= - \int_0^{\bar{t}_\tau} \int_\Omega \varepsilon(\underline{v}) : (\mathbf{c}(\bar{\chi}) - \mathbf{c}(\underline{\chi}))\tau^{-1}\mathbf{C}\varepsilon(\bar{u}) dx ds - \int_0^{\bar{t}_\tau} \int_\Omega \varepsilon(\underline{v}) : \mathbf{c}(\underline{\chi})\mathbf{C}\varepsilon(\bar{v}) dx ds \\
&\quad + \int_\Omega \varepsilon(\bar{v}(t)) : \mathbf{c}(\bar{\chi}(t))\mathbf{C}\varepsilon(\bar{u}(t)) dx - \int_\Omega \varepsilon(v^0) : \mathbf{c}(\chi^0)\mathbf{C}\varepsilon(u^0) dx \\
&\geq - C(\|\mathbf{c}\|_{L^\infty} + \|\mathbf{c}'\|_{L^\infty}) \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega |\varepsilon(\underline{v})| (|\partial_t \chi| |\varepsilon(\bar{u})| + |\varepsilon(\bar{v})|) dx ds}_{=:T_1^{(1,1)}} \\
&\quad - C\|\mathbf{c}\|_{L^\infty} \underbrace{\int_\Omega |\varepsilon(\bar{v}(t))| |\varepsilon(\bar{u}(t))| dx}_{=:T_1^{(1,2)}} - C\|\mathbf{c}\|_{L^\infty} \|\varepsilon(v^0)\|_{L^2} \|\varepsilon(u^0)\|_{L^2}.
\end{aligned}$$

By using Hölder's and Young's inequalities and the first a priori estimates, we obtain

$$\begin{aligned}
T_1^{(1,1)} &\leq \|\partial_t \chi\|_{L^2(0, \bar{t}_\tau; L^4)} \|\varepsilon(\underline{v})\|_{L^2(0, \bar{t}_\tau; L^4)} \|\varepsilon(\underline{u})\|_{L^\infty(0, \bar{t}_\tau; L^2)} \\
&\quad + \|\varepsilon(\underline{v})\|_{L^2(0, \bar{t}_\tau; L^2)} \|\varepsilon(\bar{v})\|_{L^2(0, \bar{t}_\tau; L^2)} \\
&\leq C_\delta + \delta \|\varepsilon(\underline{v})\|_{L^2(0, \bar{t}_\tau; L^4)}^2
\end{aligned}$$

and

$$\begin{aligned}
T_1^{(1,2)} &\leq C_\delta \|\varepsilon(\bar{u})\|_{L^\infty(0, \bar{t}_\tau; L^2)}^2 + \delta \|\varepsilon(\bar{v}(t))\|_{L^2}^2 \\
&\leq C_\delta + \delta \|\varepsilon(\bar{v}(t))\|_{L^2}^2.
\end{aligned}$$

The term $T_1^{(2)}$ estimates as follows:

$$\begin{aligned}
T_1^{(2)} &= \int_0^{\bar{t}_\tau} \int_\Omega \frac{1}{2\tau} (\mathbf{d}(\bar{\chi})\mathbf{D}\varepsilon(\bar{v}) : \varepsilon(\bar{v}) - \mathbf{d}(\bar{\chi})\mathbf{D}\varepsilon(\underline{v}) : \varepsilon(\underline{v})) dx ds \\
&\quad + \int_0^{\bar{t}_\tau} \int_\Omega \frac{1}{2\tau} \mathbf{d}(\bar{\chi})\mathbf{D}\varepsilon(\bar{v} - \underline{v}) : \varepsilon(\bar{v} - \underline{v}) dx ds \\
&= \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega \frac{1}{2\tau} (\mathbf{d}(\bar{\chi})\mathbf{D}\varepsilon(\bar{v}) : \varepsilon(\bar{v}) - \mathbf{d}(\underline{\chi})\mathbf{D}\varepsilon(\underline{v}) : \varepsilon(\underline{v})) dx ds}_{=:T_1^{(2,1)}} \\
&\quad + \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega \frac{1}{2\tau} (\mathbf{d}(\underline{\chi}) - \mathbf{d}(\bar{\chi})) \mathbf{D}\varepsilon(\underline{v}) : \varepsilon(\underline{v}) dx ds}_{=:T_1^{(2,2)}} \\
&\quad + \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega \frac{1}{2\tau} \mathbf{d}(\bar{\chi})\mathbf{D}\varepsilon(\bar{v} - \underline{v}) : \varepsilon(\bar{v} - \underline{v}) dx ds}_{\geq 0}.
\end{aligned}$$

For further estimations we make use of the *Ladyzhenskaya's inequality* (see [23])

$$\|w\|_{L^4} \leq C \|w\|_{H^1}^{1/2} \|w\|_{L^2}^{1/2} \quad \text{valid for all } w \in H^1(\Omega),$$

which is a special version of *Gagliardo-Nirenberg inequality* in 2D (see [34]). This inequality naturally generalizes to \mathbb{R}^m -valued Sobolev functions because

$$\begin{aligned} \|\mathbf{w}\|_{L^4} &= \left(\sum_{i=1}^m \|w_i\|_{L^4}^4 \right)^{1/4} \leq C \left(\sum_{i=1}^m \|w_i\|_{H^1}^2 \|w_i\|_{L^2}^2 \right)^{1/4} \\ &\leq C \left(\sum_{i=1}^m \|w_i\|_{H^1}^2 \right)^{1/4} \left(\sum_{i=1}^m \|w_i\|_{L^2}^2 \right)^{1/4} \\ &= C \|\mathbf{w}\|_{H^1}^{1/2} \|\mathbf{w}\|_{L^2}^{1/2} \end{aligned} \quad (38)$$

valid for all $\mathbf{w} = (w_1, \dots, w_m) \in H^1(\Omega; \mathbb{R}^m)$. By using (38), the boundedness $\mathbf{d} \geq \eta > 0$ and the Lipschitz continuity of \mathbf{d} (see (A3)), we obtain

$$\begin{aligned} T_1^{(2,1)} &= \sum_{k=1}^{\bar{t}_\tau/\tau} \int_{\Omega} \frac{1}{2} \left(\mathbf{d}(\chi^k) \mathbf{D}\varepsilon(v^k) : \varepsilon(v^k) - \mathbf{d}(\chi^{k-1}) \mathbf{D}\varepsilon(v^{k-1}) : \varepsilon(v^{k-1}) \right) dx \\ &= \int_{\Omega} \frac{1}{2} \left(\mathbf{d}(\bar{\chi}) \mathbf{D}\varepsilon(\bar{v}(t)) : \varepsilon(\bar{v}(t)) - \mathbf{d}(\chi^0) \mathbf{D}\varepsilon(v^0) : \varepsilon(v^0) \right) dx \\ &\geq \int_{\Omega} \left(\frac{\eta}{2} \varepsilon(\bar{v}(t)) : \varepsilon(\bar{v}(t)) - \frac{1}{2} \mathbf{d}(\chi^0) \mathbf{D}\varepsilon(v^0) : \varepsilon(v^0) \right) dx, \\ T_1^{(2,2)} &\geq -C \int_0^{\bar{t}_\tau} \int_{\Omega} |\partial_t \chi| |\varepsilon(\underline{v})|^2 dx ds \\ &\geq -C \int_0^{\bar{t}_\tau} \|\partial_t \chi\|_{L^4} \|\varepsilon(\underline{v})\|_{L^2} \|\varepsilon(\underline{v})\|_{L^4} ds \\ &\geq -\delta_1 \int_0^{\bar{t}_\tau} \|\partial_t \chi\|_{L^4}^2 \|\varepsilon(\underline{v})\|_{L^2}^2 ds - C_{\delta_1} \int_0^{\bar{t}_\tau} \|\varepsilon(\underline{v})\|_{L^4}^2 ds \\ &\geq -\delta_1 \int_0^{\bar{t}_\tau} \|\partial_t \chi\|_{H^1}^2 \|\varepsilon(\underline{v})\|_{L^2}^2 ds - C_{\delta_1} \int_0^{\bar{t}_\tau} \|\varepsilon(\underline{v})\|_{L^2} \|\varepsilon(\underline{v})\|_{H^1} ds \\ &\geq -\delta_1 \int_0^{\bar{t}_\tau} \|\partial_t \chi\|_{H^1}^2 \|\varepsilon(\underline{v})\|_{L^2}^2 ds - C_{\delta_1} C_{\delta_2} \|\varepsilon(\underline{v})\|_{L^2(0, \bar{t}_\tau; L^2)}^2 - C_{\delta_1} \delta_2 \|\varepsilon(\underline{v})\|_{L^2(0, \bar{t}_\tau; H^1)}^2. \end{aligned}$$

Note that by choosing $\delta_1 = \delta$ and $\delta_2 = \delta C_{\delta_1}^{-1}$ and boundedness of $\|\varepsilon(\underline{v})\|_{L^2(0, \bar{t}_\tau; L^2)}^2$ by the first a priori estimates,

$$T_1^{(2,2)} \geq -\delta \int_0^{\bar{t}_\tau} \|\partial_t \chi\|_{H^1}^2 \|\varepsilon(\underline{v})\|_{L^2}^2 ds - \delta \|\varepsilon(\underline{v})\|_{L^2(0, \bar{t}_\tau; H^1)}^2 - C_\delta.$$

The term $T_1^{(3)}$ can be treated by using the Neumann condition (10d) and by applying the discrete integration by parts formula in time

$$T_1^{(3)} = - \int_0^{\bar{t}_\tau} \int_{\Gamma} \partial_t v \cdot \bar{b} dx ds$$

$$\begin{aligned}
&= - \int_0^{\bar{t}_\tau} \int_\Gamma \underline{v} \cdot \partial_t b \, dx \, ds - \int_\Gamma \bar{v}(t) \cdot \bar{b}(t) \, dx + \int_\Gamma v^0 \cdot b^0 \, dx \\
&\geq - \int_0^{\bar{t}_\tau} \|\underline{v}\|_{L^2(\Gamma; \mathbb{R}^n)} \|\partial_t b\|_{L^2(\Gamma; \mathbb{R}^n)} \, ds - \|\bar{v}(t)\|_{L^2(\Gamma; \mathbb{R}^n)} \|\bar{b}(t)\|_{L^2(\Gamma; \mathbb{R}^n)} \\
&\quad - \|v^0\|_{L^2(\Gamma; \mathbb{R}^n)} \|b^0\|_{L^2(\Gamma; \mathbb{R}^n)}
\end{aligned}$$

By using the trace theorem $H^1(\Omega; \mathbb{R}^n) \hookrightarrow L^2(\Gamma; \mathbb{R}^n)$, we obtain

$$\begin{aligned}
T_1^{(3)} &\geq - \frac{1}{2} \|\underline{v}\|_{L^2(0, \bar{t}_\tau; H^1)}^2 - \frac{1}{2} \|\partial_t b\|_{L^2(0, \bar{t}_\tau; L^2(\Gamma; \mathbb{R}^n))}^2 - \delta \|\bar{v}(t)\|_{H^1}^2 - C_\delta \|\bar{b}(t)\|_{L^2(\Gamma; \mathbb{R}^n)}^2 \\
&\quad - \frac{1}{2} \|v^0\|_{H^1}^2 - \frac{1}{2} \|b^0\|_{L^2(\Gamma; \mathbb{R}^n)}^2
\end{aligned}$$

– To (T_2) : We estimate T_2 by

$$\begin{aligned}
T_2 &\geq \eta \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega |\mathbf{d}(\bar{\chi}) \operatorname{div}(\mathbf{D}\varepsilon(\bar{v}))|^2 \, dx \, ds}_{=: T_2^{(1)}} - C_\eta \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega |\mathbf{c}'(\bar{\chi}) \nabla \bar{\chi} \cdot \mathbf{C}\varepsilon(\bar{u})|^2 \, dx \, ds}_{=: T_2^{(2)}} \\
&\quad - C_\eta \left(\underbrace{\int_0^{\bar{t}_\tau} \int_\Omega |\mathbf{c}(\bar{\chi}) \operatorname{div}(\mathbf{C}\varepsilon(\bar{u}))|^2 \, dx \, ds}_{=: T_2^{(3)}} + \underbrace{\int_0^{\bar{t}_\tau} \int_\Omega |\mathbf{d}'(\bar{\chi}) \nabla \bar{\chi} \cdot \mathbf{D}\varepsilon(\bar{v})|^2 \, dx \, ds}_{=: T_2^{(4)}} \right)
\end{aligned}$$

With the help of the following elliptic regularity estimate which follows from [6, Theorem 3.4.1] (remember that Γ is a C^2 -boundary by (A1))

$$\|w\|_{H^2}^2 \leq C \left(\|\operatorname{div}(\mathbf{D}\varepsilon(w))\|_{L^2}^2 + \|w\|_{H^1}^2 + \|\mathbf{D}\varepsilon(w) \cdot \nu\|_{H^{1/2}(\Gamma; \mathbb{R}^n)}^2 \right),$$

valid for all $w \in H^2(\Omega; \mathbb{R}^n)$, we estimate

$$\begin{aligned}
T_2^{(1)} &\geq \eta \int_0^{\bar{t}_\tau} \int_\Omega |\operatorname{div}(\mathbf{D}\varepsilon(\bar{v}))|^2 \, dx \, ds \\
&\geq \eta \|\bar{v}\|_{L^2(0, \bar{t}_\tau; H^2)}^2 - C \|\bar{v}\|_{L^2(0, \bar{t}_\tau; H^1)}^2 - C \|\mathbf{D}\varepsilon(\bar{v}) \cdot \nu\|_{L^2(0, \bar{t}_\tau; H^{1/2}(\Gamma; \mathbb{R}^n))}^2 \\
&= \eta \|\bar{v}\|_{L^2(0, \bar{t}_\tau; H^2)}^2 - C \|\bar{v}\|_{L^2(0, \bar{t}_\tau; H^1)}^2 - C \|\bar{b} - \mathbf{c}(\bar{\chi}) \mathbf{C}\varepsilon(\bar{u}) \cdot \nu\|_{L^2(0, \bar{t}_\tau; H^{1/2}(\Gamma; \mathbb{R}^n))}^2.
\end{aligned}$$

The usage of the continuous embedding $H^1(\Omega; \mathbb{R}^n) \hookrightarrow H^{1/2}(\Gamma; \mathbb{R}^n)$ and the first a priori estimates yields

$$\begin{aligned}
T_2^{(1)} &\geq \eta \|\bar{v}\|_{L^2(0, \bar{t}_\tau; H^2)}^2 - C \left(1 + \|\bar{b}\|_{L^2(0, \bar{t}_\tau; H^{1/2}(\Gamma; \mathbb{R}^n))}^2 + \|\mathbf{c}(\bar{\chi}) \mathbf{C}\varepsilon(\bar{u}) \cdot \nu\|_{L^2(0, \bar{t}_\tau; H^{1/2}(\Gamma; \mathbb{R}^n))}^2 \right) \\
&\geq \eta \|\bar{v}\|_{L^2(0, \bar{t}_\tau; H^2)}^2 - C \left(1 + \|\bar{b}\|_{L^2(0, \bar{t}_\tau; H^{1/2}(\Gamma; \mathbb{R}^n))}^2 \right) \\
&\quad - C \left(\underbrace{\|\varepsilon(\bar{u}) \otimes \nabla \bar{\chi}\|_{L^2(0, \bar{t}_\tau; L^2)}^2}_{=: T_2^{(1,1)}} + \|\nabla \varepsilon(\bar{u})\|_{L^2(0, \bar{t}_\tau; L^2)}^2 \right).
\end{aligned}$$

By using Gagliardo-Nirenberg type inequality (38) and the first a priori estimates, we have

$$T_2^{(1,1)} \leq C \int_0^{\bar{t}_\tau} \|\varepsilon(\bar{u})\|_{L^4}^2 \|\nabla \bar{\chi}\|_{L^4}^2 \, ds$$

$$\begin{aligned}
&\leq C \int_0^{\bar{t}_\tau} \|\varepsilon(\bar{u})\|_{L^2} \|\varepsilon(\bar{u})\|_{H^1} \|\nabla \bar{\chi}\|_{L^2(\Omega)} \|\nabla \bar{\chi}\|_{H^1} \, ds \\
&\leq C \|\varepsilon(\bar{u})\|_{L^\infty(0, \bar{t}_\tau; L^2)} \|\nabla \bar{\chi}\|_{L^\infty(0, \bar{t}_\tau; L^2)} \int_0^{\bar{t}_\tau} \|\varepsilon(\bar{u})\|_{H^1} \|\nabla \bar{\chi}\|_{H^1} \, ds \\
&\leq C \int_0^{\bar{t}_\tau} (\|\varepsilon(\bar{u})\|_{H^1}^2 + \|\nabla \bar{\chi}\|_{H^1}^2) \, ds.
\end{aligned}$$

In the same fashion, we estimate the other terms as

$$\begin{aligned}
T_2^{(2)} &\leq C \int_0^{\bar{t}_\tau} (\|\varepsilon(\bar{u})\|_{H^1}^2 + \|\nabla \bar{\chi}\|_{H^1}^2) \, ds, \\
T_2^{(3)} &\leq C \int_0^{\bar{t}_\tau} \|\bar{u}\|_{H^2}^2 \, ds, \\
T_2^{(4)} &\leq C \int_0^{\bar{t}_\tau} \|\nabla \bar{\chi}\|_{L^4}^2 \|\varepsilon(\bar{v})\|_{L^4}^2 \, ds \\
&\leq C \int_0^{\bar{t}_\tau} \|\nabla \bar{\chi}\|_{L^2} \|\nabla \bar{\chi}\|_{H^1} \|\varepsilon(\bar{v})\|_{L^2} \|\varepsilon(\bar{v})\|_{H^1} \, ds \\
&\leq C_\delta \int_0^{\bar{t}_\tau} \|\varepsilon(\bar{v})\|_{L^2}^2 \|\nabla \bar{\chi}\|_{H^1}^2 \, ds + \delta \|\varepsilon(\bar{v})\|_{L^2(0, \bar{t}_\tau; H^1)}^2.
\end{aligned}$$

– To (T_3) : It can be seen by integration by parts and from the definition of I_β (see Definition 2.3 (iii)-(iv)) that

$$T_3 = \int_0^{\bar{t}_\tau} \int_\Omega \xi'_\beta(\partial_t \chi) |\nabla \partial_t \chi|^2 \, dx \, ds \geq 0.$$

– To (T_4) : The term T_4 can be treated by applying the Gagliardo-Nirenberg type inequality (38). We obtain

$$\begin{aligned}
T_4 &\geq -\delta \|\Delta \partial_t \chi\|_{L^2(0, \bar{t}_\tau; L^2)}^2 - C_\delta \int_0^{\bar{t}_\tau} \|\varepsilon(\underline{u})\|_{L^4}^4 \, ds \\
&\geq -\delta \|\Delta \partial_t \chi\|_{L^2(0, \bar{t}_\tau; L^2)}^2 - C_\delta \int_0^{\bar{t}_\tau} \|\varepsilon(\underline{u})\|_{L^2}^2 \|\varepsilon(\underline{u})\|_{H^1}^2 \, ds \\
&\geq -\delta \|\Delta \partial_t \chi\|_{L^2(0, \bar{t}_\tau; L^2)}^2 - C_\delta \|\varepsilon(\underline{u})\|_{L^\infty(0, \bar{t}_\tau; L^2)}^2 \|\varepsilon(\underline{u})\|_{L^2(0, \bar{t}_\tau; H^1)}^2.
\end{aligned}$$

– To (T_5) : We find by Young's inequality and by boundedness of f' on $[0, 1]$ (see (A4)) and $\bar{\chi} \in [0, 1]$:

$$T_5 \geq -\delta \|\Delta \partial_t \chi\|_{L^2(0, \bar{t}_\tau; L^2)}^2 - C_\delta.$$

In the following we use the estimates (by the fundamental theorem of calculus and Hölder's inequality)

$$\|\bar{u}(s)\|_{H^2}^2 = \left\| u^0 + \int_0^{\bar{s}_\tau} \partial_t u(\iota) \, d\iota \right\|_{H^2}^2 \leq \|u^0\|_{H^2}^2 + \bar{s}_\tau \|\bar{v}\|_{L^2(0, \bar{s}_\tau; H^2)}^2, \quad (39a)$$

$$\|\bar{\chi}\|_{H^2}^2 \leq \|\chi^0\|_{H^2}^2 + \bar{s}_\tau \|\partial_t \chi\|_{L^2(0, \bar{s}_\tau; H^2)}^2. \quad (39b)$$

Now we conclude by taking the above estimates as well as the first a priori estimates into account:

$$\begin{aligned}
T_1 &\geq \eta \|\bar{v}(t)\|_{H^1}^2 - D_\delta - \delta (\|u\|_{H^1(0, \bar{t}_\tau; H^2)}^2 + \|\bar{v}(t)\|_{H^1}^2) - \delta \int_0^{\bar{t}_\tau} \|\partial_t \chi(s + \tau)\|_{H^1}^2 \|\bar{v}(s)\|_{H^1}^2 ds, \\
T_2 &\geq \eta \|u\|_{H^1(0, \bar{t}_\tau; H^2)}^2 - D_\delta - C_\delta \int_0^{\bar{t}_\tau} \left(\|\bar{\chi}\|_{H^2}^2 + \|u\|_{H^1(0, \bar{s}_\tau; H^2)}^2 + \bar{t}_\tau \|\bar{v}\|_{H^1}^2 \|\partial_t \chi\|_{L^2(0, \bar{s}_\tau; H^2)}^2 \right) ds \\
&\quad - \delta \|u\|_{H^1(0, \bar{t}_\tau; H^2)}^2, \\
T_3 &\geq 0, \\
T_4 &\geq -D_\delta - C_\delta \int_0^{\bar{t}_\tau} \|u\|_{H^1(0, \bar{s}_\tau; H^2)}^2 ds - \delta \|\chi\|_{H^1(0, \bar{t}_\tau; H^2)}^2, \\
T_5 &\geq -C_\delta - \delta \|\chi\|_{H^1(0, \bar{t}_\tau; H^2)}^2,
\end{aligned}$$

where the constant $C_\delta > 0$ continuously depends on (besides δ)

$$C_\delta = C_\delta(\|u^0\|_{H^1}, \|v^0\|_{L^2}, \|\chi^0\|_{H^1}, \|b\|_{L^2(0, T; L^2(\Gamma; \mathbb{R}^n))}, \|\ell\|_{L^2(0, T; L^2)})$$

and the constant $D_\delta > 0$ continuously depends on (besides δ)

$$D_\delta = D_\delta(\|u^0\|_{H^2}, \|v^0\|_{H^1}, \|\chi^0\|_{H^2}, \|b\|_{L^2(0, T; H^{1/2}(\Gamma; \mathbb{R}^n)) \cap H^1(0, T; L^2(\Gamma; \mathbb{R}^n))}, \|\ell\|_{L^2(0, T; L^2)}).$$

By adding the identities (36) and (37), using the estimates for T_1, \dots, T_5 developed above, using the H^2 -regularity estimate (see [6, Theorem 3.4.1])

$$\|w\|_{H^2}^2 \leq C(\|\Delta w\|_{L^2}^2 + \|w\|_{H^1}^2) \quad \text{valid for all } w \in H^2(\Omega) \text{ with } \nabla w \cdot \nu = 0 \text{ a.e. on } \Gamma$$

for $\bar{\chi}$ and $\partial_t \chi$ (note the boundary conditions in (12)) and the first a priori estimates, we obtain

$$\begin{aligned}
&\|\bar{v}(t)\|_{H^1}^2 + \|\bar{\chi}(t)\|_{H^2}^2 + \|u\|_{H^1(0, \bar{t}_\tau; H^2)}^2 + \|\chi\|_{H^1(0, \bar{t}_\tau; H^2)}^2 \\
&\quad + \|\operatorname{div}(\mathbf{c}(\bar{\chi})\mathbf{C}\varepsilon(\bar{u}) + \mathbf{d}(\bar{\chi})\mathbf{D}\varepsilon(\bar{v}))\|_{L^2(0, \bar{t}_\tau; L^2)}^2 \\
&\leq D_\delta + C_\delta \int_0^{\bar{t}_\tau} \left(\|\bar{\chi}\|_{H^2}^2 + \|u\|_{H^1(0, \bar{s}_\tau; H^2)}^2 + \bar{t}_\tau \|\bar{v}\|_{H^1}^2 \|\partial_t \chi\|_{L^2(0, \bar{s}_\tau; H^2)}^2 \right) ds \\
&\quad + \delta \left(\|u\|_{H^1(0, \bar{t}_\tau; H^2)}^2 + \|\bar{v}(t)\|_{H^1}^2 + \|\operatorname{div}(\mathbf{c}(\bar{\chi})\mathbf{C}\varepsilon(\bar{u}) + \mathbf{d}(\bar{\chi})\mathbf{D}\varepsilon(\bar{v}))\|_{L^2(0, \bar{t}_\tau; L^2)}^2 \right) \\
&\quad + \delta \int_0^{\bar{t}_\tau} \|\partial_t \chi(s + \tau)\|_{H^1}^2 \|\bar{v}(s)\|_{H^1}^2 ds. \tag{40}
\end{aligned}$$

By choosing $\delta > 0$ small, the first δ -term on the right-hand side of (40) can be absorbed by the left-hand side. Furthermore, for later estimates, δ should also satisfy

$$\delta < \frac{1}{8(\|\partial_t \chi\|_{L^2(0, T; H^1)}^2 + 1)}. \tag{41}$$

Indeed, the right-hand side is bounded from below by the first a priori estimates.

We infer from the estimates (40) and (41)

$$\alpha^k \leq D_\delta + \sum_{j=1}^k \tau \gamma^j \alpha^j \tag{42}$$

with

$$\begin{aligned}\alpha^k &:= \|v^k\|_{H^1}^2 + \|\chi^k\|_{H^2}^2 + \|u\|_{H^1(0,\tau k;H^2)}^2 + \|\chi\|_{H^1(0,\tau k;H^2)}^2, \\ \gamma^k &:= C_\delta + \frac{\|(\chi^{k+1} - \chi^k)/\tau\|_{H^1}^2}{8(\|\partial_t \chi\|_{L^2(0,T;H^1)}^2 + 1)} + C_\delta \tau k \|v^k\|_{H^1}^2.\end{aligned}$$

In the following, we will choose a time $t_0 > 0$ such that for all small $\tau > 0$ and all $k = 1, \dots, (\bar{t}_0)_\tau/\tau$:

$$0 \leq \tau \gamma_\tau^k < \frac{1}{2} \quad (43)$$

Indeed, we know by the first a priori estimate that

$$\sum_{k=1}^M \tau \|v^k\|_{H^1}^2 < \widehat{C} \quad \text{uniformly in } \tau,$$

where $\widehat{C} > 0$ denotes the constant C in (31). Thus

$$\tau \|v^k\|_{H^1}^2 < \widehat{C} \quad \text{uniformly in } \tau \text{ and in } k. \quad (44)$$

Taking (44) into account and choosing

$$t_0 := \frac{1}{4C_\delta \widehat{C}}, \quad (45)$$

we get for all $k = 1, \dots, (\bar{t}_0)_\tau/\tau$:

$$\begin{aligned}\tau \gamma_\tau^k &= \tau C_\delta + \frac{\tau \|(\chi^{k+1} - \chi^k)/\tau\|_{H^1}^2}{8(\sum_{j=0}^{M-1} \tau \|(\chi^{j+1} - \chi^j)/\tau\|_{H^1}^2 + 1)} + \tau C_\delta \times \underbrace{\tau k}_{\leq (\bar{t}_0)_\tau \leq t_0 + \tau} \times \|v^k\|_{H^1}^2 \\ &\leq \tau C_\delta + \frac{1}{8} + \tau C_\delta (t_0 + \tau) \|v^k\|_{H^1}^2 \\ &= \tau C_\delta + \frac{1}{8} + \frac{\tau \omega_\tau^k}{4\widehat{C}} + \tau^2 C_\delta \|v^k\|_{H^1}^2 \\ &\leq \tau C_\delta + \frac{1}{8} + \frac{1}{4} + \tau C_\delta \widehat{C} \\ &\leq \tau C_\delta (1 + \widehat{C}) + \frac{1}{8} + \frac{1}{4}.\end{aligned}$$

Consequently, for small $\tau > 0$, estimate (43) is fulfilled.

Finally, by ensuring (43), (42) rewrites in the desired form

$$\alpha^k \leq \frac{D_\delta}{1 - \tau \gamma^k} + \sum_{j=1}^{k-1} \tau \frac{\gamma^j}{1 - \tau \gamma^k} \alpha^j$$

and, therefore,

$$\alpha^k \leq \frac{D_\delta}{2} + \sum_{j=1}^{k-1} \tau \frac{\gamma^j}{2} \alpha^j.$$

We are now in a position to apply the discrete version of Gronwall's lemma in the sum form (see, e.g., [39, page 26]) and obtain

$$\alpha^k \leq \frac{D_\delta}{2} e^{\sum_{j=1}^{k-1} \tau \frac{\gamma^j}{2}}.$$

We obtain boundedness of α^k uniformly in τ and $k = 1, \dots, (\bar{t}_0)_\tau/\tau$. Therefore, (ii) is shown except the boundedness for $\|\xi_\beta(\partial_t \chi_{\tau,\beta})\|_{L^2(0,(\bar{t}_0)_\tau;L^2)}$. The latter follows by a comparison argument in (10b).

To (ii) – global-in-time result:

The main observation to obtain global-in-time estimates is that the local estimates above can not only be performed on the time interval $[0, (\bar{t}_0)_\tau]$ but also, with minor modifications, to each interval $[\underline{s}_\tau, \bar{t}_\tau] \subseteq [0, T]$ such that $|t - s| \leq t_0$, where $t_0 > 0$ from (45) depends on quantities which can be bounded globally in time by the first a priori estimates. Thus we find a $t_0 > 0$ such that the second a priori estimates can be performed on each interval interval $[\underline{s}_\tau, \bar{t}_\tau] \subseteq [0, T]$ with $|t - s| \leq t_0$.

To conclude the proof, let

$$\begin{aligned} \mathfrak{t}_\tau^k &:= \left(k \frac{t_0}{2}\right)_\tau = \max \left\{ j\tau \mid j \in \mathbb{N} \text{ such that } k \frac{t_0}{2} \geq \tau j \right\}, \\ \mathfrak{l}_\tau &:= (\bar{t}_0)_\tau = \min \left\{ j\tau \mid j \in \mathbb{N} \text{ such that } t_0 \leq \tau j \right\}. \end{aligned}$$

We define the time intervals

$$I_\tau^k := [\mathfrak{t}_\tau^k, \mathfrak{t}_\tau^k + \mathfrak{l}_\tau] \cap [0, T]$$

for all $k = 0, \dots, N$ with $N := \lceil T/(t_0/2) \rceil - 1$ where $\lceil \cdot \rceil$ denotes the ceiling function.

We apply the local-in-time estimates above to each interval I_τ^k and obtain constants $C_0, \dots, C_N > 0$ which continuously depend on

$$\begin{aligned} C_k &= C_k(\|\bar{u}_{\tau,\beta}(\mathfrak{t}_\tau^k)\|_{H^2}, \|\bar{v}_{\tau,\beta}(\mathfrak{t}_\tau^k)\|_{H^1}, \|\bar{\chi}_{\tau,\beta}(\mathfrak{t}_\tau^k)\|_{H^2}, \\ &\quad \|b\|_{L^2(0,T;H^{1/2}(\Gamma;\mathbb{R}^n)) \cap H^1(0,T;L^2(\Gamma;\mathbb{R}^n))}, \|\ell\|_{L^2(0,T;L^2)}), \quad k = 1, \dots, N \end{aligned}$$

such that for all $\tau, \beta > 0$ and all $k = 1, \dots, N$

$$\begin{aligned} \|u_{\tau,\beta}\|_{H^1(I_\tau^k;H^2) \cap W^{1,\infty}(I_\tau^k;H^1)} &\leq C_k, & \|\chi_{\tau,\beta}\|_{H^1(I_\tau^k;H^2)} &\leq C_k, \\ \|\underline{u}_{\tau,\beta}\|_{L^\infty(I_\tau^k;H^2)} &\leq C_k, & \|\underline{\chi}_{\tau,\beta}\|_{L^\infty(I_\tau^k;H^2)} &\leq C_k, \\ \|\bar{u}_{\tau,\beta}\|_{L^\infty(I_\tau^k;H^2)} &\leq C_k, & \|\bar{\chi}_{\tau,\beta}\|_{L^\infty(I_\tau^k;H^2)} &\leq C_k, \\ \|v_{\tau,\beta}\|_{L^2(I_\tau^k;H^2) \cap L^\infty(I_\tau^k;H^1) \cap H^1(I_\tau^k;L^2)} &\leq C_k, & \|\xi_\beta(\partial_t \chi_{\tau,\beta})\|_{L^2(I_\tau^k;L^2)} &\leq C_k. \end{aligned}$$

To obtain a global bound, we can argue by induction. We sketch the argument:

Suppose we have given the a priori bound C_{k-1} for the time interval I_τ^{k-1} . By definition, we find $\mathfrak{t}_\tau^k \in I_\tau^{k-1}$. Thus

$$\|\bar{u}_{\tau,\beta}(\mathfrak{t}_\tau^k)\|_{H^2} \leq \|\bar{u}_{\tau,\beta}\|_{L^\infty(I_\tau^{k-1};H^2)} \leq C_{k-1},$$

$$\begin{aligned}\|\bar{v}_{\tau,\beta}(\mathbf{t}_\tau^k)\|_{H^1} &\leq \|\bar{v}_{\tau,\beta}\|_{L^\infty(I_\tau^{k-1}; H^1)} \leq C_{k-1}, \\ \|\bar{\chi}_{\tau,\beta}(\mathbf{t}_\tau^k)\|_{H^2} &\leq \|\bar{\chi}_{\tau,\beta}\|_{L^\infty(I_\tau^{k-1}; H^2)} \leq C_{k-1}.\end{aligned}$$

Consequently, we find an a priori bound $\tilde{C}_k \geq C_k$ for the solutions on the interval I_τ^k by

$$\tilde{C}_k := \max_{|x|,|y|,|z| \leq C_{k-1}} C_k(x, y, z, \|b\|_{L^2(0,T;H^{1/2}(\Gamma;\mathbb{R}^n)) \cap H^1(0,T;L^2(\Gamma;\mathbb{R}^n))}, \|\ell\|_{L^2(0,T;L^2)}).$$

Note that \tilde{C}_k does only depend on

$$\begin{aligned}\tilde{C}_k &= \tilde{C}_k(\|\bar{u}_{\tau,\beta}(\mathbf{t}_\tau^{k-1})\|_{H^2}, \|\bar{v}_{\tau,\beta}(\mathbf{t}_\tau^{k-1})\|_{H^1}, \|\bar{\chi}_{\tau,\beta}(\mathbf{t}_\tau^{k-1})\|_{H^2}, \\ &\quad \|b\|_{L^2(0,T;H^{1/2}(\Gamma;\mathbb{R}^n)) \cap H^1(0,T;L^2(\Gamma;\mathbb{R}^n))}, \|\ell\|_{L^2(0,T;L^2)}).\end{aligned}$$

□

Remarks to the proof of Theorem 2.11

- (i) The regularizing term $-\Delta\chi_t$ in (10b) is needed in order to obtain an $H^1(H^1)$ -bound for χ in the first estimate. This, in turn, was particularly necessary to estimate δ in (41) and to estimate the term T_4 in the second estimate.
- (ii) In the mathematical literature the elasticity equations (10a) is sometimes tested with the function $-\operatorname{div}(\mathbf{D}\varepsilon(u_t))$ to gain higher-order estimates for u (see [18, 37, 38]). However, due to the nonhomogeneous Neumann boundary condition (3c) in our case, it is more convenient to test with $-\operatorname{div}(c(\chi)\mathbf{C}\varepsilon(u) + d(\chi)\mathbf{D}\varepsilon(u_t))$ since, otherwise, integration by parts in space of the term $-\iint u_{tt} \cdot \operatorname{div}(\mathbf{D}\varepsilon(u_t))$ yields unpleasant terms even after using the boundary condition (3c) (cf. estimates for T_1 in the second estimate).

We perform the limit passage $\tau \downarrow 0$ and $\beta \downarrow 0$ separately in order to show existence of strong solutions for both cases: namely for $\beta > 0$ and $\beta = 0$ in Definition 2.5 (i) and (i). The a priori estimates give rise to the following convergence properties along a suitably chosen subsequence.

Lemma 2.10 (Convergence properties)

There exist limit functions for every $\beta \geq 0$ (we will also write $u := u_0$, $\chi := \chi_0$)

$$\begin{aligned}u_\beta &\in H^1(0, T; H^2(\Omega; \mathbb{R}^n)) \cap W^{1,\infty}(0, T; H^1(\Omega; \mathbb{R}^n)) \cap H^2(0, T; L^2(\Omega; \mathbb{R}^n)), \\ \chi_\beta &\in H^1(0, T; H^2(\Omega))\end{aligned}$$

with

$$\begin{aligned}u_\beta(0) &= u^0 \text{ a.e. in } \Omega, & \partial_t u_\beta(0) &= v^0 \text{ a.e. in } \Omega, & \chi_\beta(0) &= \chi^0 \text{ a.e. in } \Omega, \\ u_\beta &= b \text{ a.e. on } \Sigma, & 0 &\leq \chi_\beta \leq 1 \text{ a.e. in } \Sigma\end{aligned}$$

such that

(i) for fixed $\beta > 0$ and $\tau \downarrow 0$ (along a subsequence):

$$u_{\tau,\beta} \rightarrow u_\beta \quad \text{weakly in } H^1(0, T; H^2(\Omega; \mathbb{R}^n)) \quad (46a)$$

$$\begin{aligned}
& \bar{u}_{\tau,\beta}, \underline{u}_{\tau,\beta} \rightarrow u_\beta && \text{weakly-star in } W^{1,\infty}(0, T; H^1(\Omega; \mathbb{R}^n)), && (46b) \\
& u_{\tau,\beta} \rightarrow u_\beta && \text{weakly-star in } L^\infty(0, T; H^2(\Omega; \mathbb{R}^n)), && (46c) \\
& \bar{u}_{\tau,\beta}, \underline{u}_{\tau,\beta} \rightarrow u_\beta && \text{strongly in } H^1(0, T; H^1(\Omega; \mathbb{R}^n)), && (46d) \\
& u_{\tau,\beta}, \bar{u}_{\tau,\beta}, \underline{u}_{\tau,\beta} \rightarrow u_\beta && \text{strongly in } L^\infty(0, T; H^1(\Omega; \mathbb{R}^n)), && (46e) \\
& v_{\tau,\beta} \rightarrow \partial_t u_\beta && \text{a.e. in } \Omega \times (0, T), && (46f) \\
& \chi_{\tau,\beta} \rightarrow \chi_\beta && \text{weakly in } H^1(0, T; L^2(\Omega; \mathbb{R}^n)), && (46g) \\
& \bar{\chi}_{\tau,\beta}, \underline{\chi}_{\tau,\beta} \rightarrow \chi_\beta && \text{weakly in } H^1(0, T; H^2(\Omega)), && (46h) \\
& \bar{\chi}_{\tau,\beta}, \underline{\chi}_{\tau,\beta} \rightarrow \chi_\beta && \text{weakly-star in } L^\infty(0, T; H^2(\Omega)), && (46i) \\
& \bar{\chi}_{\tau,\beta}, \underline{\chi}_{\tau,\beta} \rightarrow \chi_\beta && \text{strongly in } L^\mu(0, T; H^1(\Omega)) \text{ for all } \mu \geq 1, && (46j) \\
& \bar{\chi}_{\tau,\beta}, \underline{\chi}_{\tau,\beta} \rightarrow \chi_\beta && \text{uniformly on } \bar{\Omega} \times [0, T], && (46k) \\
& \xi_\beta(\partial_t \chi_{\tau,\beta}) \rightarrow \xi_\beta(\partial_t \chi_\beta) && \text{weakly in } L^2(0, T; L^2(\Omega)), && (46l)
\end{aligned}$$

(ii) for $\beta \downarrow 0$ (along a subsequence):

$$\begin{aligned}
u_\beta \rightarrow u && \text{weakly in } H^2(0, T; L^2(\Omega; \mathbb{R}^n)) \cap H^1(0, T; H^2(\Omega; \mathbb{R}^n)) && (47a) \\
&& \text{weakly-star in } W^{1,\infty}(0, T; H^1(\Omega; \mathbb{R}^n)), && (47b) \\
u_\beta \rightarrow u && \text{strongly in } H^1(0, T; H^1(\Omega; \mathbb{R}^n)), && (47c) \\
u_\beta \rightarrow u && \text{a.e. in } \Omega \times (0, T), && (47d) \\
\chi_\beta \rightarrow \chi && \text{weakly in } H^1(0, T; H^2(\Omega)), && (47e) \\
\chi_\beta \rightarrow \chi && \text{strongly in } L^\mu(0, T; H^1(\Omega)) \text{ for all } \mu \geq 1, && (47f) \\
\chi_\beta \rightarrow \chi && \text{uniformly on } \bar{\Omega} \times [0, T], && (47g) \\
\xi_\beta(\partial_t \chi_\beta) \rightarrow \xi && \text{weakly in } L^2(0, T; L^2(\Omega)) \text{ for a } \xi \in {}^2(0, T, L^2(\Omega)) && \\
&& \text{with } \xi \in \partial I_{(-\infty, 0]}(\partial_t \chi) \text{ a.e. in } \Omega \times (0, T). && (47h)
\end{aligned}$$

Proof.

To (i): Properties (46a)-(46c) and (46f)-(46j) can be obtained by standard compact embedding, whereas (46d), (46e) and (46k) can be obtained by an Aubin-Lions type compactness result [40] (please note that $\bar{v}_{\tau,\beta} = \partial_t u_{\tau,\beta}$). It remains to show (46l).

By Lemma 2.9, we find a cluster point $\eta_\beta \in L^2(0, T; L^2(\Omega))$ such that along a subsequence $\tau \downarrow 0$

$$\xi_\beta(\partial_t \chi_{\tau,\beta}) \rightarrow \eta_\beta \text{ weakly in } L^2(0, T; L^2(\Omega)). \quad (48)$$

We have to show $\eta_\beta = \xi_\beta(\partial_t \chi_\beta)$ to finish the proof. To avoid confusion in this proof we will sometimes write “ \times ” for scalar-scalar/vector-scalar multiplication.

Let $\zeta \in L^2(0, T; H^1(\Omega))$ with $\zeta \geq 0$ a.e. in $\Omega \times (0, T)$ be an arbitrary test-function. Testing equation (10b) with $\partial_t \chi_{\tau,\beta} \times \zeta$ and integrating over $\Omega \times (0, T)$ in space and time and passing $\tau \downarrow 0$ by using weak lower-semicontinuity properties for the $\int \int |\partial_t \chi_{\tau,\beta}|^2 \zeta$ -term, we obtain

$$\liminf_{\tau \downarrow 0} \int_0^T \int_\Omega -\xi_\beta(\partial_t \chi_{\tau,\beta}) \times \partial_t \chi_{\tau,\beta} \times \zeta \, dx \, ds$$

$$\begin{aligned}
&= \liminf_{\tau \downarrow 0} \int_0^T \int_{\Omega} |\partial_t \chi_{\tau, \beta}|^2 \zeta \, dx \, ds \tag{49} \\
&\quad + \lim_{\tau \downarrow 0} \int_0^T \int_{\Omega} \left(\nabla \bar{\chi}_{\tau, \beta} + \nabla \partial_t \chi_{\tau, \beta} \right) \cdot \left(\nabla \partial_t \chi_{\tau, \beta} \times \zeta + \nabla \zeta \times \partial_t \chi_{\tau, \beta} \right) \, dx \, ds \\
&\quad + \lim_{\tau \downarrow 0} \int_0^T \int_{\Omega} \left(\frac{1}{2} \mathbf{c}'(\bar{\chi}_{\tau, \beta}) \varepsilon(\underline{u}_{\tau, \beta}) : \varepsilon(\underline{u}_{\tau, \beta}) + f'(\bar{\chi}_{\tau, \beta}) \right) \partial_t \chi_{\tau, \beta} \times \zeta \, dx \, ds \\
&\geq \int_0^T \int_{\Omega} |\partial_t \chi_{\tau}|^2 \zeta \, dx \, ds + \int_0^T \int_{\Omega} \left(\nabla \chi_{\beta} + \nabla \partial_t \chi_{\beta} \right) \cdot \left(\nabla \partial_t \chi_{\beta} \times \zeta + \nabla \zeta \times \partial_t \chi_{\beta} \right) \, dx \, ds \\
&\quad + \int_0^T \int_{\Omega} \left(\frac{1}{2} \mathbf{c}'(\chi_{\beta}) \varepsilon(u_{\beta}) : \varepsilon(u_{\beta}) + f'(\chi_{\beta}) \right) \partial_t \chi_{\beta} \times \zeta \, dx \, ds \\
&= \int_0^T \int_{\Omega} \left(\partial_t \chi_{\beta} - \Delta \chi_{\beta} - \Delta \partial_t \chi_{\beta} + \frac{1}{2} \mathbf{c}'(\chi_{\beta}) \varepsilon(u_{\beta}) : \varepsilon(u_{\beta}) + f'(\chi_{\beta}) \right) \partial_t \chi_{\beta} \times \zeta \, dx \, ds. \tag{50}
\end{aligned}$$

Note that we also get

$$-\eta_{\beta} = \partial_t \chi_{\beta} - \Delta \chi_{\beta} - \Delta \partial_t \chi_{\beta} + \frac{1}{2} \mathbf{c}'(\chi_{\beta}) \varepsilon(u_{\beta}) : \varepsilon(u_{\beta}) + f'(\chi_{\beta}) \quad \text{a.e. in } \Omega \times (0, T)$$

by performing a limit passage $\tau \downarrow 0$ in (10b) after testing with a function, integrating and using (48). In combination with (50), we find

$$\liminf_{\tau \downarrow 0} \int_0^T \int_{\Omega} -\xi_{\beta}(\partial_t \chi_{\tau, \beta}) \times \partial_t \chi_{\tau, \beta} \times \zeta \, dx \, ds \geq \int_0^T \int_{\Omega} -\eta_{\beta} \times \partial_t \chi_{\beta} \times \zeta \, dx \, ds. \tag{51}$$

By the convexity of the potential I_{β} (see Definition 2.3), we obtain a.e. in $\Omega \times (0, T)$:

$$I_{\beta}(\partial_t \chi_{\tau, \beta}) + \xi_{\beta}(\partial_t \chi_{\tau, \beta}) \times (\varphi - \partial_t \chi_{\tau, \beta}) \leq I_{\beta}(\varphi) \quad \text{for all } \varphi \in \mathbb{R}.$$

Multiplying this inequality with a test-function $\zeta(x, t)$ with $\zeta \in L^2(0, T; H^1(\Omega))$ and $\zeta \geq 0$ a.e. in $\Omega \times (0, T)$ and integrating over $\Omega \times (0, T)$, we obtain

$$\begin{aligned}
&\int_0^T \int_{\Omega} I_{\beta}(\partial_t \chi_{\tau, \beta}) \times \zeta \, dx \, ds + \int_0^T \int_{\Omega} \xi_{\beta}(\partial_t \chi_{\tau, \beta}) \times (\varphi - \partial_t \chi_{\tau, \beta}) \times \zeta \, dx \, ds \\
&\leq \int_0^T \int_{\Omega} I_{\beta}(\varphi) \times \zeta \, dx \, ds.
\end{aligned}$$

By using weak lower-semicontinuity of $\int \int I_{\beta}(\cdot) \zeta$ in the limit passage $\tau \downarrow 0$ and the estimate (51), we find

$$\int_0^T \int_{\Omega} I_{\beta}(\partial_t \chi_{\beta}) \times \zeta \, dx \, ds + \int_0^T \int_{\Omega} \eta_{\beta} \times (\varphi - \partial_t \chi_{\beta}) \times \zeta \, dx \, ds \leq \int_0^T \int_{\Omega} I_{\beta}(\varphi) \times \zeta \, dx \, ds. \tag{52}$$

Since the non-negative test-function ζ can be chosen arbitrary, we obtain

$$I_{\beta}(\partial_t \chi_{\beta}) + \eta_{\beta} \times (\varphi - \partial_t \chi_{\beta}) \leq I_{\beta}(\varphi) \quad \text{for all } \zeta \in \mathbb{R}$$

a.e. in $\Omega \times (0, T)$. In other words $\eta_{\beta} = \xi_{\beta}(\partial_t \chi_{\beta})$ a.e. in $\Omega \times (0, T)$.

To (ii): Since the a priori estimates in Lemma 2.9 are also independent of β , we obtain an analogous result for the limit case $\beta \downarrow 0$.

It remains to pass to the limit $\beta \downarrow 0$ in (52). Due to the fact that $I_\beta(\partial_t \chi_\beta) \times \zeta \geq 0$ a.e. in $\Omega \times (0, T)$ and $I_\beta(\varphi) = 0$ for $\varphi \leq 0$, we obtain from (52) for $\varphi \leq 0$

$$\int_0^T \int_\Omega \xi_\beta(\partial_t \chi_\beta) \times (\varphi - \partial_t \chi_\beta) \times \zeta \, dx \, ds \leq 0.$$

Passing to $\beta \downarrow 0$ (for a subsequence) by using (with same reasoning as (51))

$$\liminf_{\tau \downarrow 0} \int_0^T \int_\Omega -\xi_\beta(\partial_t \chi_\beta) \times \partial_t \chi_\beta \times \zeta \, dx \, ds \geq \int_0^T \int_\Omega -\eta \times \partial_t \chi \times \zeta \, dx \, ds,$$

where $\eta \in L^2(0, T; L^2(\Omega))$ denotes a weak cluster point of $\{\xi_\beta(\partial_t \chi_\beta)\}$, and switching to an a.e. formulation (remember that ζ is a freely chosen non-negative test-function) yield

$$\eta \times (\varphi - \partial_t \chi) \leq 0 \quad \text{a.e. in } \Omega \times (0, T).$$

Finally, it remains to show $\partial_t \chi \leq 0$ to conclude $\xi \in \partial I_{(-\infty, 0]}(\partial_t \chi)$ a.e. in $\Omega \times (0, T)$. Indeed, testing (52) with $\zeta \equiv 1$ and $\varphi = 0$ and using the already known a priori bounds yield

$$\int_0^T \int_\Omega I_\beta(\partial_t \chi_\beta) \, dx \, ds \leq C, \tag{53}$$

where $C > 0$ is independent of β . Now, let $\eta \in (0, 1)$ be arbitrary given. By Definition 2.3 (i) we obtain $I_\beta \geq I_\eta$ pointwise in \mathbb{R} for all β with $0 < \beta \leq \eta$. By using this estimate, the lower-semicontinuity of $\int_0^T \int_\Omega I_\eta(\cdot)$ and the weak convergence property (47e), we find

$$\liminf_{\beta \downarrow 0} \int_0^T \int_\Omega I_\beta(\partial_t \chi_\beta) \, dx \, ds \geq \liminf_{\beta \downarrow 0} \int_0^T \int_\Omega I_\eta(\partial_t \chi_\beta) \, dx \, ds \geq \int_0^T \int_\Omega I_\eta(\partial_t \chi) \, dx \, ds.$$

Together with (53), we obtain for every $\eta \in (0, 1)$

$$\int_0^T \int_\Omega I_\eta(\partial_t \chi) \, dx \, ds \leq C.$$

Thus, from property (ii) in Definition 2.3, we infer $\partial_t \chi \leq 0$ a.e. in $\Omega \times (0, T)$. \square

Theorem 2.11 *Let the Assumptions (A1)-(A4) be satisfied and the data $(u^0, v^0, \chi^0, b, \ell)$ from (25) and (26) be given. The following statements are true:*

- (i) *Regularized case ($\beta > 0$): There exists a strong global-in-time solution (u_β, χ_β) in the sense of Definition 2.5 (ii) which satisfies (11).*
- (ii) *Limit case ($\beta = 0$): There exists a strong global-in-time solution (u, χ) in the sense of Definition 2.5 (i). which satisfies (11).*

Proof.

To (i): By multiplying the systems (10a), (10b), (10d) and (10e) with test-functions, integrating over space and time, we may pass to the limit $\tau \downarrow 0$ for fixed $\beta > 0$ by utilizing Lemma 2.8 and Lemma 2.10 (i) and standard convergence arguments. Then, switching back to an (x, t) -a.e. formulation, we obtain a strong solution of system (1)-(3).

To (ii): The transition $\beta \downarrow 0$ can be conducted as in (i) by utilizing Lemma 2.10 (ii). \square

2.4 Continuous dependence on the data

We are going to show continuous dependence on the data of strong solutions of the PDE system given in Definition 2.5 (i) and (ii).

Theorem 2.12 (Continuous dependence) *Let the Assumptions (A1)-(A4) be satisfied. Moreover, assume that $\mathbf{d} \equiv 1$ in (A3) and one of the following condition:*

- *Let (u_1, χ_1) and (u_2, χ_2) be both strong solutions according to Definition 2.5 (ii) (for $\beta > 0$) with data $(u_1^0, v_1^0, \chi_1^0, b_1, \ell_1)$ and $(u_2^0, v_2^0, \chi_2^0, b_2, \ell_2)$.*
- *Let (u_1, χ_1) and (u_2, χ_2) be both strong solutions according to Definition 2.5 (i) (for $\beta = 0$) with data $(u_1^0, v_1^0, \chi_1^0, b_1, \ell_1)$ and $(u_2^0, v_2^0, \chi_2^0, b_2, \ell_2)$.*

Then,

$$\begin{aligned} & \|u_1 - u_2\|_{W^{1,\infty}(0,T;L^2) \cap H^1(0,T;H^1)} + \|\chi_1 - \chi_2\|_{H^1(0,T;H^1)} \\ & \leq C \left(\|u_1^0 - u_2^0\|_{H^1} + \|v_1^0 - v_2^0\|_{L^2} + \|\chi_1^0 - \chi_2^0\|_{H^1} + \|\ell_1 - \ell_2\|_{L^2(0,T;L^2)} + \|b_1 - b_2\|_{L^2(0,T;L^2(\Gamma;\mathbb{R}^n))} \right), \end{aligned} \quad (54)$$

where the constant $C > 0$ continuously depends on

$$C = C \left(\|u_1\|_{\mathcal{U}}, \|u_2\|_{\mathcal{U}}, \|\chi_1\|_{\mathcal{X}}, \|\chi_2\|_{\mathcal{X}} \right).$$

Proof. For notational convenience, define

$$\begin{aligned} \mathbf{u} &:= u_1 - u_2, & \boldsymbol{\chi} &:= \chi_1 - \chi_2, & \mathbf{u}^0 &:= u_1^0 - u_2^0, \\ \mathbf{v}^0 &:= v_1^0 - v_2^0, & \boldsymbol{\chi}^0 &:= \chi_1^0 - \chi_2^0, & \mathbf{b} &:= b_1 - b_2, \\ \boldsymbol{\ell} &:= \ell_1 - \ell_2. \end{aligned}$$

Let $t \in [0, T]$ be arbitrary. Firstly, testing the damage equation (1b) for each solution with $\boldsymbol{\chi}_t$, subtracting the resulting equations and integrating over $\Omega \times (0, t)$, we obtain

$$\int_0^t \int_{\Omega} \left(|\boldsymbol{\chi}_t|^2 + \nabla \boldsymbol{\chi} \cdot \nabla \boldsymbol{\chi}_t + |\nabla \boldsymbol{\chi}_t|^2 + \frac{1}{2} c'(\chi_1) (\mathbf{C}\varepsilon(u_1) : \varepsilon(u_1) - \mathbf{C}\varepsilon(u_2) : \varepsilon(u_2)) \boldsymbol{\chi}_t \right) dx ds$$

$$\begin{aligned}
& + \int_0^t \int_{\Omega} \left(\frac{1}{2} (\mathbf{c}'(\chi_1) - \mathbf{c}'(\chi_2)) \mathbf{C} \varepsilon(u_2) : \varepsilon(u_2) \boldsymbol{\chi}_t + (f'(\chi_1) - f'(\chi_2)) \boldsymbol{\chi}_t + (\xi_1 - \xi_2) \boldsymbol{\chi}_t \right) dx ds \\
& = 0.
\end{aligned} \tag{55}$$

By assumption, we know

$$\begin{aligned}
\xi_i & \in \partial I_{(-\infty, 0]}(\partial_t \chi_i), \quad i = 1, 2 & \text{if } \beta = 0, \\
\xi_i & = I'_{\beta}(\partial_t \chi_i), \quad i = 1, 2 & \text{if } \beta > 0.
\end{aligned}$$

It follows from the monotonicity of $\partial I_{(-\infty, 0]}$ and I'_{β} (see Definition 2.3), respectively, that

$$(\xi_1 - \xi_2) \boldsymbol{\chi}_t = (\xi_1 - \xi_2)(\chi_1 - \chi_2)_t \geq 0.$$

Therefore, by (55),

$$\begin{aligned}
& \|\boldsymbol{\chi}_t\|_{L^2(0,t;H^1)}^2 + \frac{1}{2} \|\nabla \boldsymbol{\chi}(t)\|_{L^2}^2 - \frac{1}{2} \|\nabla \boldsymbol{\chi}^0\|_{L^2}^2 \\
& \leq \underbrace{- \int_0^t \int_{\Omega} \frac{1}{2} \mathbf{c}'(\chi_1) \mathbf{C} \varepsilon(\mathbf{u}) : \varepsilon(u_1 + u_2) \boldsymbol{\chi}_t dx ds}_{=: T_1} \\
& \quad - \underbrace{\int_0^t \int_{\Omega} \frac{1}{2} (\mathbf{c}'(\chi_1) - \mathbf{c}'(\chi_2)) \mathbf{C} \varepsilon(u_2) : \varepsilon(u_2) \boldsymbol{\chi}_t dx ds}_{=: T_2} \\
& \quad - \underbrace{\int_0^t \int_{\Omega} (f'(\chi_1) - f'(\chi_2)) \boldsymbol{\chi}_t dx ds}_{=: T_3}.
\end{aligned} \tag{56}$$

By using Hölder's and Young's inequalities as well as standard Sobolev embeddings, Lipschitz continuity of \mathbf{c}' (see (A2)) and Lipschitz continuity of f' (see (A3)), we find

$$\begin{aligned}
T_1 & \leq C \|\varepsilon(u_1 + u_2)\|_{L^\infty(0,t;L^4)} \int_0^t \|\varepsilon(\mathbf{u})\|_{L^2} \|\boldsymbol{\chi}_t\|_{L^4} ds \\
& \leq \delta \|\boldsymbol{\chi}_t\|_{L^2(0,t;H^1)}^2 + C_{\delta} \|\varepsilon(\mathbf{u})\|_{L^2(0,t;L^2)}^2, \\
T_2 & \leq C \|\mathbf{c}'\|_{Lip} \|\varepsilon(u_2)\|_{L^\infty(0,t;L^4)} \int_0^t \|\boldsymbol{\chi}\|_{L^4} \|\boldsymbol{\chi}_t\|_{L^4} ds \\
& \leq \delta \|\boldsymbol{\chi}_t\|_{L^2(0,t;H^1)}^2 + C_{\delta} \|\boldsymbol{\chi}\|_{L^2(0,t;H^1)}^2, \\
T_3 & \leq \|f'\|_{Lip} \int_0^t \|\boldsymbol{\chi}\|_{L^2} \|\boldsymbol{\chi}_t\|_{L^2} ds \\
& \leq \delta \|\boldsymbol{\chi}_t\|_{L^2(0,t;L^2)}^2 + C_{\delta} \|\boldsymbol{\chi}\|_{L^2(0,t;L^2)}^2.
\end{aligned}$$

Applying the estimates for T_1 , T_2 and T_3 to (56), we obtain

$$\|\boldsymbol{\chi}_t\|_{L^2(0,t;H^1)}^2 + \|\nabla \boldsymbol{\chi}(t)\|_{L^2}^2 \leq C_{\delta} \left(\|\boldsymbol{\chi}^0\|_{H^1}^2 + \|\boldsymbol{\chi}\|_{L^2(0,t;H^1)}^2 + \|\mathbf{u}\|_{L^2(0,t;H^1)}^2 \right) + \delta \|\boldsymbol{\chi}_t\|_{L^2(0,t;H^1)}^2. \tag{57}$$

Secondly, we test each of the corresponding elasticity equations (1a) for u_1 and u_2 with \mathbf{u}_t and obtain by subtraction and integration over $\Omega \times (0, t)$:

$$\int_0^t \int_{\Omega} \left(\mathbf{u}_{tt} \mathbf{u}_t + \mathbf{c}(\chi_1) \mathbf{C} \varepsilon(\mathbf{u}) : \varepsilon(\mathbf{u}_t) + (\mathbf{c}(\chi_1) - \mathbf{c}(\chi_2)) \mathbf{C} \varepsilon(u_2) : \varepsilon(\mathbf{u}_t) + \mathbf{D} \varepsilon(\mathbf{u}_t) : \varepsilon(\mathbf{u}_t) \right) dx ds$$

$$= \int_0^t \int_{\Gamma} \mathbf{b} \cdot \mathbf{u}_t \, dx \, ds + \int_0^t \int_{\Omega} \boldsymbol{\ell} \cdot \mathbf{u}_t \, dx \, ds.$$

This implies

$$\begin{aligned} & \frac{1}{2} \|\mathbf{u}_t(t)\|_{L^2}^2 - \frac{1}{2} \|\mathbf{v}^0\|_{L^2}^2 + \eta \|\varepsilon(\mathbf{u}_t)\|_{L^2(0,t;L^2)}^2 \\ & \leq \underbrace{- \int_0^t \int_{\Omega} \mathbf{c}(\chi_1) \mathbf{C} \varepsilon(\mathbf{u}) : \varepsilon(\mathbf{u}_t) \, dx \, ds}_{=:T_4} - \underbrace{\int_0^t \int_{\Omega} (\mathbf{c}(\chi_1) - \mathbf{c}(\chi_2)) \mathbf{C} \varepsilon(u_2) : \varepsilon(\mathbf{u}_t) \, dx \, ds}_{=:T_5} \\ & \quad + \underbrace{\int_0^t \int_{\Gamma} \mathbf{b} \cdot \mathbf{u}_t \, dx \, ds}_{=:T_6} + \underbrace{\int_0^t \int_{\Omega} \boldsymbol{\ell} \cdot \mathbf{u}_t \, dx \, ds}_{=:T_7} \end{aligned} \quad (58)$$

Standard estimates yield

$$\begin{aligned} T_4 & \leq \delta \|\varepsilon(\mathbf{u}_t)\|_{L^2(0,t;L^2)}^2 + C_{\delta} \|\varepsilon(\mathbf{u})\|_{L^2(0,t;L^2)}^2 \\ & \leq \delta \|\mathbf{u}_t\|_{L^2(0,t;H^1)}^2 + C_{\delta} \|\mathbf{u}\|_{L^2(0,t;H^1)}^2, \\ T_5 & \leq C \|\boldsymbol{\chi}\|_{L^2(0,t;L^3)} \|\varepsilon(u_2)\|_{L^{\infty}(L^6)} \|\varepsilon(\mathbf{u}_t)\|_{L^2(0,t;L^2)} \\ & \leq \delta \|\mathbf{u}_t\|_{L^2(0,t;H^1)}^2 + C_{\delta} \|\boldsymbol{\chi}\|_{L^2(0,t;H^1)}^2, \\ T_6 & \leq \delta \|\mathbf{u}_t\|_{L^2(0,t;L^2(\Gamma;\mathbb{R}^n))}^2 + C_{\delta} \|\mathbf{b}\|_{L^2(0,t;L^2(\Gamma;\mathbb{R}^n))}^2 \\ & \leq \delta \|\mathbf{u}_t\|_{L^2(0,t;H^1)}^2 + C_{\delta} \|\mathbf{b}\|_{L^2(0,t;L^2(\Gamma;\mathbb{R}^n))}^2, \\ T_7 & \leq \frac{1}{2} \|\mathbf{u}_t\|_{L^2(0,t;L^2)}^2 + \frac{1}{2} \|\boldsymbol{\ell}\|_{L^2(0,t;L^2)}^2. \end{aligned}$$

Applying the estimates T_4 , T_5 , T_6 and T_7 to (58) shows

$$\begin{aligned} & \|\mathbf{u}_t(t)\|_{L^2}^2 + \|\varepsilon(\mathbf{u}_t)\|_{L^2(0,t;L^2)}^2 \\ & \leq C_{\delta} \left(\|\mathbf{v}^0\|_{L^2}^2 + \|\mathbf{u}\|_{H^1(0,t;L^2)}^2 + \|\mathbf{u}\|_{L^2(0,t;H^1)}^2 + \|\boldsymbol{\chi}\|_{L^2(0,t;H^1)}^2 + \|\boldsymbol{\ell}\|_{L^2(0,t;L^2)}^2 \right) \\ & \quad + C_{\delta} \|\mathbf{b}\|_{L^2(0,t;L^2(\Gamma;\mathbb{R}^n))}^2 + \delta \|\mathbf{u}\|_{H^1(0,t;H^1)}^2. \end{aligned} \quad (59)$$

Adding (57) and (59), we obtain

$$\begin{aligned} & \|\varepsilon(\mathbf{u}_t)\|_{L^2(0,t;L^2)}^2 + \|\boldsymbol{\chi}_t\|_{L^2(0,t;H^1)}^2 + \|\mathbf{u}_t(t)\|_{L^2}^2 + \|\nabla \boldsymbol{\chi}(t)\|_{L^2}^2 \\ & \leq C_{\delta} \left(\|\mathbf{v}^0\|_{L^2}^2 + \|\boldsymbol{\chi}^0\|_{H^1}^2 + \|\boldsymbol{\chi}\|_{L^2(0,t;H^1)}^2 + \|\mathbf{u}\|_{H^1(0,t;L^2)}^2 + \|\mathbf{u}\|_{L^2(0,t;H^1)}^2 \right) \\ & \quad + C_{\delta} \left(\|\boldsymbol{\ell}\|_{L^2(0,t;L^2)}^2 + \|\mathbf{b}\|_{L^2(0,t;L^2(\Gamma;\mathbb{R}^n))}^2 \right) + \delta \left(\|\mathbf{u}\|_{H^1(0,t;H^1)}^2 + \|\boldsymbol{\chi}_t\|_{L^2(0,t;H^1)}^2 \right). \end{aligned} \quad (60)$$

Now, adding $\|\mathbf{u}\|_{L^2(0,t;H^1)}^2 + \|\mathbf{u}_t\|_{L^2(0,t;L^2)}^2$ on both sides and using

$$\|\boldsymbol{\chi}(t)\|_{L^2}^2 = \|\boldsymbol{\chi}^0\|_{L^2}^2 + \int_0^t \|\boldsymbol{\chi}_t(s)\|_{L^2}^2 \, ds \leq C \left(\|\boldsymbol{\chi}^0\|_{L^2}^2 + \|\boldsymbol{\chi}_t\|_{L^2(0,t;L^2)}^2 \right).$$

and Korn's inequality

$$\|w\|_{H^1} \leq C \left(\|w\|_{L^2}^2 + \|\varepsilon(w)\|_{L^2}^2 \right)$$

holding for all $w \in H^1(\Omega; \mathbb{R}^n)$, the estimate (60) becomes

$$\begin{aligned} & \|\mathbf{u}\|_{H^1(0,t;H^1)}^2 + \|\boldsymbol{\chi}_t\|_{L^2(0,t;H^1)}^2 + \|\mathbf{u}_t(t)\|_{L^2}^2 + \|\boldsymbol{\chi}(t)\|_{H^1}^2 \\ & \leq C_\delta \left(\|\mathbf{v}^0\|_{L^2}^2 + \|\boldsymbol{\chi}^0\|_{H^1}^2 + \|\boldsymbol{\chi}\|_{L^2(0,t;H^1)}^2 + \|\mathbf{u}\|_{H^1(0,t;L^2)}^2 + \|\mathbf{u}\|_{L^2(0,t;H^1)}^2 \right) \\ & \quad + C_\delta \left(\|\boldsymbol{\ell}\|_{L^2(0,t;L^2)}^2 + \|\mathbf{b}\|_{L^2(0,t;L^2(\Gamma;\mathbb{R}^n))}^2 \right) + \delta \left(\|\mathbf{u}\|_{H^1(0,t;H^1)}^2 + \|\boldsymbol{\chi}_t\|_{L^2(0,t;H^1)}^2 \right). \end{aligned}$$

By choosing $\delta > 0$ small and noticing

$$\|\mathbf{u}\|_{L^2(0,t;H^1)}^2 \leq C \left(\|\mathbf{u}^0\|_{H^1}^2 + \int_0^t \|\mathbf{u}_t\|_{L^2(0,s;H^1)}^2 \, ds \right),$$

we get

$$\begin{aligned} & \|\mathbf{u}\|_{H^1(0,t;H^1)}^2 + \|\boldsymbol{\chi}_t\|_{L^2(0,t;H^1)}^2 + \|\mathbf{u}_t(t)\|_{L^2}^2 + \|\boldsymbol{\chi}(t)\|_{H^1}^2 \\ & \leq C \left(\|\mathbf{u}^0\|_{H^1}^2 + \|\mathbf{v}^0\|_{L^2}^2 + \|\boldsymbol{\chi}^0\|_{H^1}^2 + \|\boldsymbol{\ell}\|_{L^2(0,t;L^2)}^2 + \|\mathbf{b}\|_{L^2(0,t;L^2(\Gamma;\mathbb{R}^n))}^2 \right) \\ & \quad + C \int_0^t \left(\|\mathbf{u}_t\|_{L^2}^2 + \|\mathbf{u}_t\|_{L^2(0,s;H^1)}^2 + \|\boldsymbol{\chi}\|_{H^1}^2 \right) \, ds. \end{aligned}$$

The claim follows by Gronwall's lemma. \square

The continuous dependence result in Theorem 2.12 as well as the a priori estimates in Lemma 2.9 yield the following corollaries.

Corollary 2.13 (Uniqueness) *Strong solutions in the sense of Definition 2.5 (i) or (ii) with constant viscosity \mathbb{D} are unique to given initial-boundary data $(\mathbf{u}^0, \mathbf{v}^0, \boldsymbol{\chi}^0, \mathbf{b}, \boldsymbol{\ell})$.*

Corollary 2.14 (A priori estimates) *A strong solution $(\mathbf{u}, \boldsymbol{\chi}) \in \mathcal{U} \times \mathcal{X}$ for the system in Definition 2.5 (i) or (ii) with constant viscosity \mathbb{D} and given data $(\mathbf{u}^0, \mathbf{v}^0, \boldsymbol{\chi}^0, \mathbf{b}, \boldsymbol{\ell})$ satisfies the a priori estimates*

$$\|\mathbf{u}\|_{\mathcal{U}} \leq C, \quad \|\boldsymbol{\chi}\|_{\mathcal{X}} \leq C, \quad \|\boldsymbol{\xi}\|_{L^2(\Omega \times (0,T))} \leq C,$$

where the constant $C > 0$ continuously depends on

$$C = C(\|\mathbf{u}^0\|_{H^2}, \|\mathbf{v}^0\|_{H^1}, \|\boldsymbol{\chi}^0\|_{H^2}, \|\mathbf{b}\|_{L^2(0,T;H^{1/2}(\Gamma;\mathbb{R}^n)) \cap H^1(0,T;L^2(\Gamma;\mathbb{R}^n))}, \|\boldsymbol{\ell}\|_{L^2(0,T;L^2)}).$$

3 Optimal control problem

In this section we establish the announced optimal control problem for the damage-elasticity system (1)-(3). From now on, we assume for the viscosity tensor $\mathbb{D} = \mathbf{D}$, i.e. $\mathbf{d} \equiv 1$, in order to apply the well-posedness result from the last section.

Let \mathcal{U} , \mathcal{X} and \mathcal{B} be given as in Section 2.1. Our aim is to approximate with χ prescribed damage profiles by controlling the Neumann boundary data $b \in \mathcal{B}$ for the stress tensor σ in (3b). The cost functionals measures the deviation from the prescribed profiles at the final time or/and at all time during the evolution in an L^∞ -norm. We make the following assumptions:

(O1) We assume that λ_Q , λ_Ω and λ_Σ are given non-negative constants which do not all vanish.

(O2) The target damage profiles are given by

$$\chi_Q \in L^\infty(Q), \quad \chi_T \in L^\infty(\Omega).$$

(O3) The admissible set of controls $\mathcal{B}_{adm} \subseteq \mathcal{B}$ is assumed to be non-empty, closed and bounded.

Remark 3.1 A typical choice for \mathcal{B}_{adm} would be

$$\mathcal{B}_{adm} = \{b \in \mathcal{B} \mid b_{min} \leq b \leq b_{max} \text{ a.e. in } \Sigma \text{ and } \|b\|_{\mathcal{B}} \leq M\},$$

where $M \in (0, \infty)$ denotes the maximal \mathcal{B} -cost and $b_{min}, b_{max} \in \mathcal{B}$ the minimal and maximal cost functions satisfying $b_{min} \leq b_{max}$ a.e. in Σ .

We define the following tracking type objective functional

$$\mathcal{J}(\chi, b) := \frac{\lambda_Q}{2} \|\chi - \chi_Q\|_{L^\infty(Q)} + \frac{\lambda_\Omega}{2} \|\chi(T) - \chi_T\|_{L^\infty(\Omega)} + \frac{\lambda_\Sigma}{2} \|b\|_{L^2(\Sigma; \mathbb{R}^n)}^2, \quad (61)$$

where our overall optimization problem reads as

$$\left. \begin{array}{l} \text{minimize } \mathcal{J}(\chi, b) \text{ over } \mathcal{X} \times \mathcal{B}_{adm} \\ \text{s.t. the PDE system in Definition 2.5 (i) is satisfied for an } u \in \mathcal{U}. \end{array} \right\} \quad (P_0)$$

Remark 3.2 Let us emphasize that we may also choose $\|\cdot\|_{L^2}$ -terms instead of the $\|\cdot\|_{L^\infty}$ -terms in the cost functional (61). The existence results presented in this section work for both cases.

We recall that the system (1)-(3) is an initial-boundary value problem, which admits by Theorem 2.11 and Corollary 2.13 for every $(u^0, v^0, \chi^0, b, \ell)$ satisfying (25) and (26) a unique solution $(u, \chi) \in \mathcal{U} \times \mathcal{X}$ in the sense of Definition 2.5 (i). Hence, the solution operator

$$\Psi_0 : \mathcal{I} \rightarrow \mathcal{O}, \quad (u^0, v^0, \chi^0, b, \ell) \mapsto (u, \chi)$$

with

$$\begin{aligned} \mathcal{I} &:= \{(u^0, v^0, \chi^0, \ell, b) \mid \text{satisfying (25) and (26)}\}, \\ \mathcal{O} &:= \mathcal{U} \times \mathcal{X} \end{aligned}$$

is well-defined. Moreover, for fixed data (u^0, v^0, χ^0, ℓ) the control-to-state operator

$$S_0 : \mathcal{B}_{adm} \rightarrow \mathcal{O}, \quad b \mapsto (u, \chi)$$

is also well-defined, and the optimal control problem (P_0) is equivalent to minimize the reduced cost functional

$$j(b) := \mathcal{J}(S_{0|2}(b), b)$$

over \mathcal{B}_{adm} , where $S_{0|2}$ denotes the second component of S_0 , i.e. $S_0 = (S_{0|1}, S_{0|2})$.

For $\beta \in (0, 1)$, let us denote by S_β the operator mapping the control $b \in \mathcal{B}_{adm}$ into the unique solution $(u_\beta, \chi_\beta) \in \mathcal{O}$ to the β -regularized problem in Definition 2.5 (ii).

Remark 3.3 In view of the continuous dependence result in Theorem 2.12 the operators Ψ_0 and S_0 are well-posed in a larger target space $(W^{1,\infty}(0, T; L^2(\Omega; \mathbb{R}^n)) \cap H^1(0, T; H^1(\Omega; \mathbb{R}^n))) \times H^1(0, T; H^1(\Omega))$, which contains the space \mathcal{O} . This fact is important for the sensitivity analysis of these operators. But in this section, we are interested only in existence of optimal controls, so this result is not needed. The sensitivity analysis which also establishes the optimality conditions of first-order will be treated in a forthcoming paper.

3.1 Existence of optimal controls to (P_0) via β -regularization

The following lemma is the basis for the main result in this section.

Lemma 3.4 *We have the following continuity properties:*

For a given sequence $\{b_\beta\}_{\beta \in (0,1)} \subseteq \mathcal{B}$ and $\bar{b} \in \mathcal{B}$ with

$$b_\beta \rightharpoonup \bar{b} \quad \text{weakly in } \mathcal{B} \text{ as } \beta \downarrow 0, \quad (62)$$

it holds

$$S_\beta(b_\beta) \rightharpoonup S_0(\bar{b}) \quad \text{weakly-star in } \mathcal{O} \text{ as } \beta \downarrow 0, \quad (63a)$$

$$S_\eta(b_\beta) \rightharpoonup S_\eta(\bar{b}) \quad \text{weakly-star in } \mathcal{O} \text{ as } \beta \downarrow 0 \text{ for every } \eta \in (0, 1), \quad (63b)$$

$$S_0(b_\beta) \rightharpoonup S_0(\bar{b}) \quad \text{weakly-star in } \mathcal{O} \text{ as } \beta \downarrow 0. \quad (63c)$$

Proof. Let $(u_\beta, \chi_\beta) = S_\beta(b_\beta)$. Then, (u_β, χ_β) is a solution to the β -regularized system in the sense of Definition 2.5 (ii) with Neumann data b_β . Since $\{b_\beta\} \subseteq \mathcal{B}$ is bounded by (62), we obtain the a priori estimates from Corollary 2.14. In particular,

$$S_\beta(b_\beta) \rightharpoonup (u, \chi) \quad \text{weakly-star in } \mathcal{O} \text{ as } \beta \downarrow 0 \text{ (for a subsequence).}$$

for some $(u, \chi) \in \mathcal{O}$.

We see from the proof of Lemma 2.10 that the convergence properties in Lemma 2.10 (ii) hold for a subsequence $\beta \downarrow 0$. By using these convergence properties as well as (62), we can pass to the limit for a subsequence in the β -regularized PDE system (9a)-(9e) (cf. proof of Theorem 2.11). We obtain that (u, χ) satisfies the limit system in Definition 2.5 (i) to the Neumann data \bar{b} . In other words, $S_0(\bar{b}) = (u, \chi)$. By uniqueness of solutions shown in Corollary 2.13, we see that $S_\beta(b_\beta)$ converges weakly-star to (u, χ) for the whole sequence $\beta \downarrow 0$. Hence, (63a) is shown and (63b) and (63c) follow with the same reasoning. \square

Corollary 3.5 *We also have the property*

$$\lim_{\beta \downarrow 0} \mathcal{J}(S_{\beta|2}(b), b) = \mathcal{J}(S_{0|2}(b), b)$$

for all $b \in \mathcal{B}_{adm}$.

Proof. Set $b_\beta := b$ and apply Lemma 3.4. In particular, (63a) implies

$$S_{\beta|2}(b) \rightharpoonup S_{0|2}(b) \quad \text{weakly-star in } \mathcal{X} \text{ as } \beta \downarrow 0.$$

Define $\chi_\beta := S_{\beta|2}(b)$ and $\chi := S_{0|2}(b)$. A standard compactness result shows (see [40])

$$\chi_\beta \rightarrow \chi \quad \text{strongly in } C^0(\bar{Q}) \text{ as } \beta \downarrow 0$$

and the claim follows. \square

Theorem 3.6 *Suppose that Assumptions (A1)-(A4) as well as (O1)-(O3) are satisfied. Then the optimal control problem (P_0) admits a solution.*

Before proving Theorem 3.6, we introduce a family of auxilliary optimal control problems (P_β) , which are parametrized by $\beta \in (0, 1)$. For $\beta \in (0, 1)$, let us denote by S_β the operator mapping the control $b \in \mathcal{B}_{adm}$ into the unique solution $(u_\beta, \chi_\beta) \in \mathcal{O}$ to the regularized problem in Definition 2.5 (ii). We define

$$\left. \begin{array}{l} \text{minimize } \mathcal{J}(\chi, b) \text{ over } \mathcal{X} \times \mathcal{B}_{adm} \\ \text{s.t. the } \beta\text{-regularized PDE system in Definition 2.5 (ii) is satisfied.} \end{array} \right\} \quad (P_\beta)$$

The following result guarantees the existence of an optimal control to (P_β) .

Lemma 3.7 *Suppose that the Assumptions (A1)-(A4) as well as (O1)-(O3) are fulfilled. Let $\beta > 0$ be given. Then the optimal control problem (P_β) admits a solution.*

Proof. Let $\{b^n\}_{n \in \mathbb{N}} \subseteq \mathcal{B}_{adm}$ be a minimizing sequence for (P_β) , and let $(u_\beta^n, \chi_\beta^n) = S_\beta(b^n)$, $n \in \mathbb{N}$. By the boundedness and closedness of \mathcal{B}_{adm} (see (O3)), we find a function $\bar{b} \in \mathcal{B}_{adm}$ and a subsequence of $\{b^n\}$ (we omit the subscript) such that

$$b^n \rightarrow \bar{b} \quad \text{weakly in } \mathcal{B} \text{ as } n \uparrow \infty.$$

Lemma 3.4 yields

$$S_\beta(b^n) \rightarrow S_\beta(\bar{b}) \quad \text{weakly-star in } \mathcal{O} \text{ as } n \uparrow \infty.$$

Let $(\bar{u}_\beta, \bar{\chi}_\beta) := S_\beta(\bar{b})$. We particularly find

$$\bar{\chi}_\beta^n = S_{\beta|_2}(b^n) \rightarrow S_{\beta|_2}(\bar{b}) = \bar{\chi}_\beta \quad \text{weakly-star in } \mathcal{X} \text{ as } n \uparrow \infty.$$

A standard compact result reveals

$$\bar{\chi}_\beta^n \rightarrow \bar{\chi}_\beta \quad \text{strongly in } C^0(\bar{Q}) \text{ as } n \uparrow \infty.$$

It follows from the sequentially weak lower semicontinuity of the cost functional \mathcal{J} that \bar{b} is an optimal control for (P_β) , i.e.

$$J(\bar{\chi}_\beta, \bar{b}_\beta) \leq \liminf_{n \uparrow \infty} J(\chi_\beta^n, b_\beta^n).$$

□

Proof of Theorem 3.6. By virtue of Lemma 3.7, for any $\beta \in (0, 1)$, we may pick an optimality pair

$$(\chi_\beta, b_\beta) \in \mathcal{X} \times \mathcal{B}_{adm}$$

for the optimal control problem (P_β) . Obviously, we have $(u_\beta, \chi_\beta) = S_\beta(b_\beta)$, $\beta \in (0, 1)$. By the assumption (O3) and Lemma 3.4, we find functions $(\bar{u}, \bar{\chi}) \in \mathcal{O}$ and $\bar{b} \in \mathcal{B}_{adm}$ with $S_0(\bar{b}) = (\bar{u}, \bar{\chi})$ such that

$$(u_\beta, \chi_\beta) \rightarrow (\bar{u}, \bar{\chi}) \quad \text{weakly-star in } \mathcal{O}, \quad (64a)$$

$$b_\beta \rightarrow \bar{b} \quad \text{weakly in } \mathcal{B} \quad (64b)$$

as $\beta \downarrow 0$ (for a subsequence).

It remains to show that $(\bar{\chi}, \bar{b})$ is in fact an optimality pair of (P_0) . To this end, let $b \in \mathcal{B}_{adm}$ be arbitrary. In view of the convergence properties (64) and the sequentially weak lower semicontinuity of the cost functional, we have

$$\mathcal{J}(\bar{\chi}, \bar{b}) \leq \liminf_{\beta \downarrow 0} \mathcal{J}(\chi_\beta, b_\beta) = \liminf_{\beta \downarrow 0} \mathcal{J}(S_{\beta|2}(b_\beta), b_\beta).$$

By using the optimality property of (P_β) , we obtain

$$\liminf_{\beta \downarrow 0} \mathcal{J}(S_{\beta|2}(b_\beta), b_\beta) \leq \liminf_{\beta \downarrow 0} \mathcal{J}(S_{\beta|2}(b), b).$$

Finally, the convergence property in Corollary 3.5 shows

$$\liminf_{\beta \downarrow 0} \mathcal{J}(S_{\beta|2}(b), b) = \mathcal{J}(S_{0|2}(b), b).$$

In conclusion, we have proven $\mathcal{J}(S_{0|2}(\bar{b}), \bar{b}) \leq \mathcal{J}(S_{0|2}(b), b)$. \square

Remark 3.8 *Theorem 3.6 can also be shown in the spirit of Lemma 3.7. However, the proof presented via convergence of β -approximations might be of interest in view of the implementation of optimality systems.*

3.2 An adapted optimal control problem to (P_0)

Theorem 3.6 does not yield any information on whether every solution to the optimal control problem (P_0) can be approximated by a sequence of solutions to the problem (P_β) . Such a result is also called *global* result. However, we can give a *local* answer for every individual optimizer of (P_0) . For this purpose, we employ a trick due to [2].

To this end, let $((\bar{u}, \bar{\chi}), \bar{b}) \in \mathcal{O} \times \mathcal{B}_{adm}$, where $(\bar{u}, \bar{\chi}) = S_0(\bar{b})$, be an arbitrary but fixed solution to (P_0) . We associate with this solution the *adapted cost functional*

$$\tilde{\mathcal{J}}(\chi, b) := \mathcal{J}(\chi, b) + \frac{1}{2} \|b - \bar{b}\|_{L^2(\Sigma; \mathbb{R}^n)}^2$$

and the corresponding *adapted optimal control problem*

$$\left. \begin{array}{l} \text{minimize } \tilde{\mathcal{J}}(\chi, b) \text{ over } \mathcal{X} \times \mathcal{B}_{adm} \\ \text{s.t. the } \beta\text{-regularized PDE system in Definition 2.5 (ii) is satisfied.} \end{array} \right\} \quad (\tilde{P}_\beta)$$

With a proof that resembles that of Lemma 3.7 and needs no repetition here, we can show the following result:

Lemma 3.9 *Suppose that the Assumptions (A1)-(A4) as well as (O1)-(O3) are fulfilled. Let $\beta \in (0, 1)$ be given. Then, the optimal control problem (\tilde{P}_β) admits a solution.*

We are now in the position to give a partial answer to the question raised above. More precisely, we show the following theorem:

Theorem 3.10 *Let the Assumptions (A1)-(A4) and (O1)-(O3) be satisfied. Suppose that $(\bar{\chi}, \bar{b}) \in \mathcal{X} \times \mathcal{B}_{adm}$ is any fixed solution to the optimal control problem (P_0) . Then, there exists a pair $(\bar{\chi}_\beta, \bar{b}_\beta) \in \mathcal{X} \times \mathcal{B}_{adm}$ solving the adapted problem (\tilde{P}_β) such that $\tilde{\mathcal{J}}(\bar{\chi}_\beta, \bar{b}_\beta) \rightarrow \mathcal{J}(\bar{\chi}, \bar{b})$ as $\beta \downarrow 0$.*

Proof. For every $\beta \in (0, 1)$ we pick an optimal pair $(\bar{\chi}_\beta, \bar{b}_\beta) \in \mathcal{X} \times \mathcal{B}_{adm}$ for the adapted problem (\tilde{P}_β) . By the boundedness and closedness of \mathcal{B}_{adm} (see (O3)), there exists a $b \in \mathcal{B}_{adm}$ satisfying

$$\bar{b}_\beta \rightarrow b \quad \text{weakly in } \mathcal{B} \text{ as } \beta \downarrow 0. \quad (65)$$

Owing to Lemma 3.4 we find

$$(\bar{u}_\beta, \bar{\chi}_\beta) = S_\beta(\bar{b}_\beta) \rightarrow S_0(b) =: (u, \chi) \quad \text{weakly-star in } \mathcal{O} \text{ as } \beta \downarrow 0.$$

and, particularly,

$$\bar{\chi}_\beta \rightarrow \chi \quad \text{strongly in } C^0(\bar{Q}) \text{ as } \beta \downarrow 0. \quad (66)$$

We now aim to prove that $b = \bar{b}$. Once this is shown, we can infer from the unique solvability of the state system (see Theorem 2.12) that also $(u, \chi) = (\bar{u}, \bar{\chi})$.

Indeed, we have, owing to (65), (66), the sequentially weak lower semicontinuity of $\tilde{\mathcal{J}}$, and the optimality property of $(\bar{\chi}, \bar{b})$ for problem (P_0) ,

$$\begin{aligned} \liminf_{\beta \downarrow 0} \tilde{\mathcal{J}}(\bar{\chi}_\beta, \bar{b}_\beta) &\geq \mathcal{J}(\chi, b) + \frac{1}{2} \|b - \bar{b}\|_{L^2(\Sigma; \mathbb{R}^n)}^2 \\ &\geq \mathcal{J}(\bar{\chi}, \bar{b}) + \frac{1}{2} \|b - \bar{b}\|_{L^2(\Sigma; \mathbb{R}^n)}^2. \end{aligned} \quad (67)$$

On the other hand, the optimality property of $(\bar{\chi}_\beta, \bar{b}_\beta)$ for problem (\tilde{P}_β) yields that

$$\tilde{\mathcal{J}}(\bar{\chi}_\beta, \bar{b}_\beta) = \tilde{\mathcal{J}}(S_{\beta|2}(\bar{b}_\beta), \bar{b}_\beta) \leq \tilde{\mathcal{J}}(S_{\beta|2}(\bar{b}), \bar{b}).$$

Whence, taking the limes superior as $\beta \downarrow 0$ on both sides and invoking Corollary 3.5, we find

$$\limsup_{\beta \downarrow 0} \tilde{\mathcal{J}}(\bar{\chi}_\beta, \bar{b}_\beta) \leq \tilde{\mathcal{J}}(S_{0|2}(\bar{b}), \bar{b}) = \tilde{\mathcal{J}}(\bar{\chi}, \bar{b}) = \mathcal{J}(\bar{\chi}, \bar{b}). \quad (68)$$

We obtain by combining (67) and (68)

$$\frac{1}{2} \|b - \bar{b}\|_{L^2(\Sigma; \mathbb{R}^n)}^2 = 0.$$

Thus $b = \bar{b}$ and, consequently, $(u, \chi) = (\bar{u}, \bar{\chi})$ by Theorem 2.12.

Finally, by using $b = \bar{b}$ in (67) and (68), we end up with $\lim_{\beta \downarrow 0} \tilde{\mathcal{J}}(\bar{\chi}_\beta, \bar{b}_\beta) = \mathcal{J}(\bar{\chi}, \bar{b})$. \square

References

- [1] L. Ambrosio and V.M. Tortorelli. Approximation of functionals depending on jumps by elliptic functionals via Γ -convergence. *Commun. Pure Appl. Math.*, 43(8):999–1036, 1990.

- [2] V. Barbu. Necessary conditions for nonconvex distributed control problems governed by elliptic variational inequalities. *J. Math. Anal. Appl.*, 80:566–597, 1981.
- [3] E. Bonetti and G. Schimperna. Local existence for Frémond’s model of damage in elastic materials. *Contin. Mech. Thermodyn.*, 16(4):319–335, 2004.
- [4] E. Bonetti, G. Schimperna, and A. Segatti. On a doubly nonlinear model for the evolution of damaging in viscoelastic materials. *J. of Diff. Equations*, 218(1):91–116, 2005.
- [5] S. Burke, C. Ortner, and E. Süli. An adaptive finite element approximation of a generalized Ambrosio-Tortorelli functional. *Math. Models Methods Appl. Sci.*, 23(9):1663–1697, 2013.
- [6] M. Costabel, M. Dauge, and S. Nicaise. *Corner Singularities and Analytic Regularity for Linear Elliptic Systems. Part I: Smooth domains*. HAL Archives, 2010.
- [7] P. Destuynder. Remarques sur le contrôle de la propagation des fissures en régime stationnaire. (Remarks on a crack propagation control for stationary loaded structures). *C. R. Acad. Sci., Paris, Sér. II*, 308(8):697–701, 1989.
- [8] G.A. Francfort and J.-J. Marigo. Revisiting brittle fracture as an energy minimization problem. *J. Mech. Phys. Solids*, 46(8):1319–1342, 1998.
- [9] M. Frémond. *Non-smooth thermomechanics*. Berlin: Springer, 2002.
- [10] M. Frémond. *Phase Change in Mechanics*. Lecture Notes of the Unione Matematica Italiana. Springer, 2012.
- [11] M. Frémond, K.L. Kuttler, B. Nedjar, and M. Shillor. One-dimensional models of damage. *Adv. Math. Sci. Appl.*, 8(2):541–570, 1998.
- [12] M. Frémond, K.L. Kuttler, and M. Shillor. Existence and uniqueness of solutions for a dynamic one-dimensional damage model. *J. Math. Anal. Appl.*, 229(1):271–294, 1999.
- [13] M. Frémond and B. Nedjar. Damage, gradient of damage and principle of virtual power. *Int. J. Solids Structures*, 33(8):1083–1103, 1996.
- [14] A. Giacomini. Ambrosio–Tortorelli approximation of quasi–static evolution of brittle fractures. *Calc. Var. Partial Differ. Equ.*, 22(2):129–172, 2005.
- [15] P. Grisvard. *Elliptic Problems in Nonsmooth Domains*. Pitman Advanced Publishing Program, 1985.
- [16] C. Heinemann and C. Kraus. Existence of weak solutions for Cahn-Hilliard systems coupled with elasticity and damage. *Adv. Math. Sci. Appl.*, 21(2):321–359, 2011.
- [17] C. Heinemann and C. Kraus. Existence of weak solutions for a pde system describing phase separation and damage processes including inertial effects. *Discrete Contin. Dyn. Syst. Ser. A (to appear)*, 2014.
- [18] C. Heinemann and E. Rocca. Damage processes in thermoviscoelastic materials with damage-dependent thermal expansion coefficients. *Math. Methods Appl. Sci. (to appear)*, 2014.
- [19] P. Hild, A. Münch, and Y. Ousset. On the control of crack growth in elastic media. *C. R., Méc., Acad. Sci. Paris*, 336(5):422–427, 2008.
- [20] A. Khludnev, G. Leugering, and M. Specovius-Neugebauer. Optimal control of inclusion and crack shapes in elastic bodies. *J. Optim. Theory Appl.*, 155(1):54–78, 2012.
- [21] D. Knees, R. Rossi, and C. Zanini. A vanishing viscosity approach to a rate-independent damage model. *Math. Models Methods Appl. Sci.*, 23(4):565–616, 2013.
- [22] D. Knees, R. Rossi, and C. Zanini. A quasilinear differential inclusion for viscous and rate-independent damage systems in non-smooth domains. *WIAS preprint 1867*, 2013.
- [23] O.A. Ladyzhenskaya. Solution ”in the large” to the boundary value problem for the Navier- Stokes equations in two space variables. *Sov. Phys., Dokl.*, 3:1128–1131, 1958.
- [24] G. Lazzaroni, R. Rossi, M. Thomas, and R. Toader. Rate-independent damage in thermo-viscoelastic materials with inertia. *WIAS preprint 2025*, 2014.
- [25] J. Lemaitre and R. Desmorat. *Engineering Damage Mechanics: Ductile, Creep, Fatigue and Brittle Failures*. Springer-Verlag, Berlin, 2005.
- [26] G. Leugering, J. Sokołowski, and A. Żochowski. Control of crack propagation by shape-topological optimization. *Discrete Contin. Dyn. Syst. Ser. A*, 35(6):2625–2657, 2015.
- [27] G. Dal Maso and R. Toader. A model for the quasi-static growth of brittle fractures: Existence and approximation results. *Arch. Ration. Mech. Anal.*, 162(2):101–135, 2002.
- [28] A. Mielke. Complete-damage evolution based on energies and stresses. *Discrete Contin. Dyn. Syst., Ser. S*, 4(2):423–439, 2011.
- [29] A. Mielke and T. Roubíček. Rate-independent damage processes in nonlinear elasticity. *Mathematical Models and Methods in Applied Sciences*, 16:177–209, 2006.

- [30] A. Mielke, T. Roubíček, and J. Zeman. Complete Damage in elastic and viscoelastic media. *Comput. Methods Appl. Mech. Engrg.*, 199:1242–1253, 2010.
- [31] A. Mielke and M. Thomas. Damage of nonlinearly elastic materials at small strain — Existence and regularity results. *ZAMM Z. Angew. Math. Mech.*, 90:88–112, 2010.
- [32] P. Neittaanmäki and D. Tiba. *Optimal control of nonlinear parabolic systems: theory, algorithms, and applications*. New York: Marcel Dekker, Inc., 1994.
- [33] E. Di Nezza, G. Palatucci, and E. Valdinoci. Hitchhiker’s guide to the fractional Sobolev spaces. *Preprint arXiv:1104.4345*, April 2011.
- [34] L. Nirenberg. On elliptic differential equations. *Ann. Scuola Norm. Pisa (III)*, 13:1–48, 1959.
- [35] K. Pham, H. Amor, J.-J. Marigo, and C. Maurini. Gradient Damage Models and Their Use to Approximate Brittle Fracture. *International Journal of Damage Mechanics*, 20(4):618–652, 2011.
- [36] M. Prechtel, G. Leugering, P. Steinmann, and M. Stingl. Towards optimization of crack resistance of composite materials by adjustment of fiber shapes. *Engineering Fracture Mechanics*, 78:944–960, 2011.
- [37] E. Rocca and R. Rossi. A degenerating PDE system for phase transitions and damage. *Math. Models Methods Appl. Sci.*, 24:1265–1341, 2014.
- [38] E. Rocca and R. Rossi. “Entropic” solutions to a thermodynamically consistent PDE system for phase transitions and damage. *Preprint arXiv:1403.2577*, 2014.
- [39] T. Roubíček. *Nonlinear Partial Differential Equations with Applications*. International Series of Numerical Mathematics. Birkhäuser-Verlag, second edition, 2013.
- [40] J. Simon. Compact sets in the space $L^p(0, T; B)$. *Annali di Matematica Pura ed Applicata*, 146:65–96, 1986.
- [41] M. Thomas. Quasistatic damage evolution with spatial BV-regularization. *Discrete and Continuous Dynamical Systems - Series S*, 6(1):235–255, 2013.