

The geometry of the space of branched rough paths

Nikolas Tapia¹, Lorenzo Zambotti²

submitted: November 21, 2019

¹ Weierstrass Institute
Mohrenstr. 39
10117 Berlin
Germany
E-Mail: nikolasesteban.tapiamunoz@wias-berlin.de

² Laboratoire de Probabilités, Statistiques et Modélisation
4 place Jussieu
75005 Paris
France
E-Mail: zambotti@lpsm.paris

No. 2645
Berlin 2019



2010 *Mathematics Subject Classification.* 60H10, 16T05.

Key words and phrases. rough paths, Hopf algebras, renormalization.

The authors thank Jean-David Jacques for pointing out a mistake in a previous version. N.T. acknowledges support by the CONICYT/Doctorado Nacional doctoral scholarship grant number 2013-21130733, *Núcleo Milenio Modelos Estocásticos de Sistemas Complejos y Desordenados* and the Berlin Mathematical School MATH+ EF1-5 project “On robustness of deep neural networks”. L.Z. gratefully acknowledges support by the project of the Agence Nationale de la Recherche ANR-15-CE40-0020-01 grant LSD.

Edited by
Weierstraß-Institut für Angewandte Analysis und Stochastik (WIAS)
Leibniz-Institut im Forschungsverbund Berlin e. V.
Mohrenstraße 39
10117 Berlin
Germany

Fax: +49 30 20372-303
E-Mail: preprint@wias-berlin.de
World Wide Web: <http://www.wias-berlin.de/>

The geometry of the space of branched rough paths

Nikolas Tapia, Lorenzo Zambotti

Abstract

We construct an explicit transitive free action of a Banach space of Hölder functions on the space of branched rough paths, which yields in particular a bijection between these two spaces. This endows the space of branched rough paths with the structure of a principal homogeneous space over a Banach space and allows to characterize its automorphisms. The construction is based on the Baker-Campbell-Hausdorff formula, on a constructive version of the Lyons-Victoir extension theorem and on the Hairer-Kelly map, which allows to describe branched rough paths in terms of anisotropic geometric rough paths.

1 Introduction

The theory of Rough Paths has been introduced by Terry Lyons in the '90s with the aim of giving an alternative construction of stochastic integration and stochastic differential equations. More recently, it has been expanded by Martin Hairer to cover stochastic partial differential equations, with the invention of regularity structures.

A rough path and a *model* of a regularity structure are mathematical objects which must satisfy some algebraic and analytical constraints. For instance, a rough path can be described as a Hölder function defined on an interval and taking values in a non-linear finite-dimensional Lie group; models of regularity structures are a generalization of this idea. A crucial ingredient of regularity structures is the *renormalisation* procedure: given a family of regularized models, which fail to converge in an appropriate topology as the regularization is removed, one wants to modify it in a such a way that the algebraic and analytical constraints are still satisfied and the modified version converges. This procedure has been obtained in [6, 9] for a general class of models with a *stationary* character.

The same question about rough paths has been asked recently in [3, 4, 5], and indeed it could have been asked much earlier. Maybe this has not happened because the motivation was less compelling; although one can construct examples of rough paths depending on a positive parameter which do not converge as the parameter tends to 0, this phenomenon is the exception rather than the rule. However the problem of characterizing the automorphisms of the space of rough paths is clearly of interest; one example is the transformation from Itô to Stratonovich integration, see e.g. [1, 16, 17]. However our aim is to put this particular example in a much larger context.

We recall that there are several possible notions of rough paths; in particular we have *geometric* RPs and *branched* RPs, two notions defined respectively by Terry Lyons [29] and Massimiliano Gubinelli [25], see Sections 3 and 4 below. These two notions are intimately related to each other, as shown by Hairer and Kelly [28], see Section 4 below. We note that regularity structures [27] are a natural and far-reaching generalization of branched RPs.

In this paper we concentrate on the automorphisms of the space of branched RPs, see below for a discussion of the geometric case. Let \mathcal{F} be the collection of all non-planar rooted forests with nodes

decorated by $\{1, \dots, d\}$, see Section 4 below. For instance the following forest



is an element of \mathcal{F} . We call $\mathcal{T} \subset \mathcal{F}$ the set of *rooted trees*, namely of *non-empty forests* with a single connected component. Grading elements $\tau \in \mathcal{F}$ by the number $|\tau|$ of their nodes we set

$$\mathcal{T}_n := \{\tau \in \mathcal{T} : |\tau| \leq n\}, \quad n \in \mathbb{N}.$$

Let now \mathcal{H} be the linear span of \mathcal{F} . It is possible to endow \mathcal{H} with a product and a coproduct $\Delta : \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{H}$ which make it a Hopf algebra, also known as the Butcher-Connes-Kreimer Hopf algebra, see Section 4.2 below. We let G denote the set of all *characters* over \mathcal{H} , that is, elements of G are functionals $X \in \mathcal{H}^*$ that are also multiplicative in the sense that

$$\langle X, \tau\sigma \rangle = \langle X, \tau \rangle \langle X, \sigma \rangle$$

for all forests (and in particular trees) $\tau, \sigma \in \mathcal{F}$. Furthermore, the set G can be endowed with a product \star , dual to the coproduct, defined pointwise by $\langle X \star Y, \tau \rangle = \langle X \otimes Y, \Delta\tau \rangle$. We work on the compact interval $[0, 1]$ for simplicity, and all results can be proved without difficulty on $[0, T]$ for any $T \geq 0$.

Definition 1.1 (Gubinelli [25]). *Given $\gamma \in]0, 1[$, a *branched γ -rough path* is a path $X : [0, 1]^2 \rightarrow G$ which satisfies Chen's rule*

$$X_{su} \star X_{ut} = X_{st}, \quad s, u, t \in [0, 1],$$

and the analytical condition

$$|\langle X_{st}, \tau \rangle| \lesssim |t - s|^{\gamma|\tau|}, \quad \tau \in \mathcal{F}.$$

Setting $x_t^i := \langle X_{0t, \bullet i} \rangle$, $t \in [0, 1]$, we say that X is a *branched γ -rough path over the path $x = (x^1, \dots, x^d)$* . We denote by \mathbf{BRP}^γ the set of all *branched γ -rough paths* (for a fixed finite alphabet $\{1, \dots, d\}$).

By introducing the *reduced coproduct* $\Delta' : \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{H}$

$$\Delta'\tau := \Delta\tau - \tau \otimes 1 - 1 \otimes \tau,$$

where 1 denotes the empty forest, Chen's rule can be rewritten as follows

$$\delta \langle X, \tau \rangle_{sut} = \langle X_{su} \otimes X_{ut}, \Delta'\tau \rangle, \quad s, u, t \in [0, 1], \tag{1.1}$$

where for $F : [0, 1]^2 \rightarrow \mathbb{R}$ we set $\delta F : [0, 1]^3 \rightarrow \mathbb{R}$,

$$\delta F_{sut} := F_{st} - F_{su} - F_{ut}, \tag{1.2}$$

which is the second order finite increment considered by Gubinelli [24]. Note that the right-hand side of (1.1) depends on the values of X on trees with strictly fewer nodes than τ ; if we can invert the operator δ , then the right-hand side of (1.1) determines the left-hand side. This is however not a trivial result. In fact, a simple (but crucial for us) remark is the following: if $\gamma|\tau| \leq 1$, then for any $g^\tau : [0, 1] \rightarrow \mathbb{R}$ such that $g^\tau \in C^{\gamma|\tau|}([0, 1])$, the classical homogeneous Hölder space on $[0, 1]$ with Hölder exponent $\gamma|\tau|$, the function

$$[0, 1]^2 \ni (s, t) \mapsto F_{st} := \langle X_{st}, \tau \rangle + g_t^\tau - g_s^\tau \tag{1.3}$$

also satisfies

$$\delta F_{sut} = \langle X_{su} \otimes X_{ut}, \Delta' \tau \rangle, \quad |F_{st}| \lesssim |t - s|^{\gamma|\tau|}, \quad s, u, t \in [0, 1]. \quad (1.4)$$

Inversely, if $F: [0, 1]^2 \rightarrow \mathbb{R}$ satisfies (1.4), then F must satisfy (1.3) with $g^\tau \in C^{\gamma|\tau|}([0, 1])$.

If $\gamma|\tau| > 1$, then Gubinelli's *Sewing Lemma* [24] yields that the function $(s, t) \mapsto \langle X_{st}, \tau \rangle$ is uniquely determined by (1.4) i.e. by the values of X on trees with at most $|\tau| - 1$ nodes, and therefore, applying a recursion, on trees with at most $N := \lfloor \gamma^{-1} \rfloor$ nodes. More explicitly, the Sewing Lemma is an *existence and uniqueness result* for $[0, 1]^2 \ni (s, t) \mapsto \langle X_{st}, \tau \rangle$ with $\gamma|\tau| > 1$, once the right-hand side of (1.1) is known. However, for $\gamma|\tau| \leq 1$ we have no uniqueness, as we have already seen, and existence is not trivial.

As we have seen in (1.3), the value of $\langle X, \tau \rangle$ can be modified by adding the increment of a function in $C^{\gamma|\tau|}([0, 1])$, as long as $\gamma|\tau| \leq 1$. It seems reasonable to think that it is therefore possible to construct an action on the set of branched γ -rough paths of the abelian group (under pointwise addition)

$$\mathcal{C}^\gamma := \{(g^\tau)_{\tau \in \mathcal{T}_N} : g_0^\tau = 0, g^\tau \in C^{\gamma|\tau|}([0, 1]), \forall \tau \in \mathcal{T}, |\tau| \leq N\},$$

namely the set of all collections of functions $(g^\tau \in C^{\gamma|\tau|}([0, 1]) : \tau \in \mathcal{T}, |\tau| \leq N)$ indexed by rooted trees with fewer than $N := \lfloor \gamma^{-1} \rfloor$ nodes, such that $g_0^\tau = 0$ and $g^\tau \in C^{\gamma|\tau|}([0, 1])$. This is indeed the content of the following

Theorem 1.2. *Let $\gamma \in]0, 1[$ such that $\gamma^{-1} \notin \mathbb{N}$. There is a transitive free action of \mathcal{C}^γ on \mathbf{BRP}^γ , namely a map $(g, X) \mapsto gX$ such that*

- 1 for each $g, g' \in \mathcal{C}^\gamma$ and $X \in \mathbf{BRP}^\gamma$ the identity $g'(gX) = (g + g')X$ holds.
- 2 if $(g^\tau)_{\tau \in \mathcal{T}_N} \in \mathcal{C}^\gamma$ is such that there exists a unique $\tau \in \mathcal{T}_N$ with $g^\tau \neq 0$, then

$$\langle (gX)_{st}, \tau \rangle = \langle X_{st}, \tau \rangle + g_t^\tau - g_s^\tau$$

and $\langle gX, \sigma \rangle = \langle X, \sigma \rangle$ for all $\sigma \in \mathcal{T}$ not containing τ .

- 3 For every pair $X, X' \in \mathbf{BRP}^\gamma$ there exists a unique $g \in \mathcal{C}^\gamma$ such that $gX = X'$.

We say that a tree $\sigma \in \mathcal{T}$ contains a tree $\tau \in \mathcal{T}$ if there exists a subtree τ' of σ , not necessarily containing the root of σ , such that τ and τ' are isomorphic as rooted trees, where the root of τ' is its node which is closest to the root of σ . Note that every $(g^\tau)_{\tau \in \mathcal{T}_N} \in \mathcal{C}^\gamma$ is the sum of finitely many elements of \mathcal{C}^γ having satisfying the property required in point (2) of Theorem 1.2.

If $\gamma > 1/2$ then the result of Theorem 1.2 is trivial. Indeed, in this case $N = 1$, $\mathcal{T}_N = \{\bullet_i : i = 1, \dots, d\}$, and $\mathcal{C}^\gamma = \{g^{\bullet_i} \in C^\gamma([0, 1]) : g_0^{\bullet_i} = 0, i = 1, \dots, d\}$. Then the action is

$$(g, X) \mapsto gX, \quad \langle (gX)_{st}, \bullet_i \rangle := \langle X_{st}, \bullet_i \rangle + g_t^{\bullet_i} - g_s^{\bullet_i}, \quad (1.5)$$

while the value of $\langle gX, \tau \rangle$ for $|\tau| \geq 2$ is uniquely determined by (1.1) via the Sewing Lemma. For example

$$\langle (gX)_{st}, \bullet_i^j \rangle := \int_s^t (x_u^j - x_s^j + g_u^{\bullet_i} - g_s^{\bullet_i}) d(x_u^j + g_u^{\bullet_i}), \quad (1.6)$$

where $x_u^j := \langle X_{0u}, \bullet_i \rangle$ and the integral is well-defined in the Young sense, see [24, section 3].

If $1/3 < \gamma \leq 1/2$ then $N = 2$ and $\mathcal{T}_2 = \mathcal{T}_1 \sqcup \{\mathfrak{t}_i^j : i, j = 1, \dots, d\}$. Then the action at level $|\tau| = 1$ is still given by (1.5), while at level $|\tau| = 2$ we must have by (1.1)

$$\delta \langle gX, \mathfrak{t}_i^j \rangle_{sut} = \langle (gX)_{su} \otimes (gX)_{ut}, \Delta' \tau \rangle = (x_u^j - x_s^j + g_u^{\bullet j} - g_s^{\bullet j})(x_t^i - x_u^i + g_t^{\bullet i} - g_u^{\bullet i}). \quad (1.7)$$

Although the right-hand side of (1.7) is explicit and simple, in this case there is no canonical choice for $\langle gX, \mathfrak{t}_i^j \rangle$. An expression like (1.6) is ill-defined in the Young sense, and the same is true if we try the formulation

$$\langle (gX)_{st}, \mathfrak{t}_i^j \rangle = \langle X_{st}, \mathfrak{t}_i^j \rangle + \int_s^t ((x_u^j - x_s^j + g_u^{\bullet j} - g_s^{\bullet j}) dg_u^{\bullet i} + (g_u^{\bullet j} - g_s^{\bullet j}) dx_u^i), \quad (1.8)$$

which satisfies formally (1.7), but the Young integrals are ill defined since $2\gamma \leq 1$. The construction of $\langle gX, \mathfrak{t}_i^j \rangle$ is therefore not trivial in this case.

The same argument applies for any $\gamma \leq 1/2$ and any tree τ such that $2 \leq |\tau| \leq N = \lfloor \gamma^{-1} \rfloor$, and the fact that the above Young integrals are not well defined shows why existence of the map $X \rightarrow gX$ is not trivial.

Since Theorem 1.2 yields an action of \mathcal{C}^γ on \mathbf{BRP}^γ which is *regular*, i.e. free and transitive, then \mathbf{BRP}^γ is a *principal \mathcal{C}^γ -homogeneous space* or *\mathcal{C}^γ -torsor*. In particular, \mathbf{BRP}^γ is a copy of \mathcal{C}^γ , but there is no canonical choice of an origin in \mathbf{BRP}^γ .

Therefore, Theorem 1.2 also yields the following

Corollary 1.3. *Given a branched γ -rough path X , the map $g \rightarrow gX$ yields a bijection between \mathcal{C}^γ and the set of branched γ -rough paths.*

Therefore Corollary 1.3 yields a complete parametrization of the space of branched rough paths. This result is somewhat surprising, since rough paths form a non-linear space, in particular because of the Chen relation; however Corollary 1.3 yields a natural bijection between the space of branched γ -rough paths and the linear space \mathcal{C}^γ .

Corollary 1.3 also gives a complete answer to the question of existence and characterization of branched γ -rough paths over a γ -Hölder path x . Unsurprisingly, for our construction we start from a result of T. Lyons and N. B. Victoir's [30] of 2007, which was the first general theorem of existence of a geometric γ -rough path over a γ -Hölder path x , see our discussion of Theorem 1.4 below.

An important point to stress is that the action constructed in Theorem 1.2 is neither unique nor canonical. In the proof of Theorem 3.4 below, some parameters have to be fixed arbitrarily, and the final outcome depends on them, see Remark 3.6. In this respect, the situation is similar to what happens in regularity structures with the reconstruction operator on spaces \mathcal{D}^γ with a negative exponent $\gamma < 0$, see [27, Theorem 3.10].

1.1 Outline of our approach

A key point in Theorem 1.2 is the construction of branched γ -rough paths. In the case of *geometric* rough paths, see Definition 4.1, the *signature* [11, 29] of a smooth path $x: [0, 1] \rightarrow \mathbb{R}^d$ yields a canonical construction. Other cases where geometric rough paths over non-smooth paths have been constructed are Brownian motion and fractional Brownian motion (see [13] for the case $H > \frac{1}{4}$ and [33] for the general case) among others. However, until T. Lyons and N. B. Victoir's paper [30] in 2007, this question remained largely open in the general case. The precise result is as follows

Theorem 1.4 (Lyons–Victoir extension). *If $p \in [1, \infty) \setminus \mathbb{N}$ and $\gamma := 1/p$, a γ -Hölder path $x: [0, 1] \rightarrow \mathbb{R}^d$ can be lifted to a geometric γ -rough path. For any $p \geq 1$ and $\varepsilon \in]0, \gamma[$, a γ -Hölder path can be lifted to a geometric $(\gamma - \varepsilon)$ -rough path.*

Our first result is a version of this theorem which holds for rough paths in a more general algebraic context, see Theorem 3.4 below. We use the Lyons–Victoir approach and an explicit form of the Baker–Campbell–Hausdorff formula by Reutenauer [34], see formula (2.11) below. Whereas Lyons and Victoir used in one passage the axiom of choice, our method is completely constructive.

Using the same idea we extend this construction to the case where the collection (x^1, \dots, x^d) is allowed to have different regularities in each component, which we call *anisotropic (geometric) rough paths (aGRP)*, see Definition 4.8.

Theorem 1.5. *To each collection $(x^i)_{i=1, \dots, d}$, with $x^i \in C^{\gamma_i}([0, 1])$, we can associate an anisotropic rough path \bar{X} over $(x^i)_{i=1, \dots, d}$. For every collection $(g^i)_{i=1, \dots, d}$, with $g^i \in C^{\gamma_i}([0, 1])$, denoting by $g\bar{X}$ the anisotropic geometric rough path over $(x^i + g^i)_{i=1, \dots, d}$, we have*

$$g'(g\bar{X}) = (g + g')\bar{X}.$$

This kind of extension to rough paths has already been explored in the papers [2, 26] in the context of isomorphisms between geometric and branched rough paths. It turns out that the additional property obtained by our method enables us to explicitly describe the propagation of suitable modifications from lower to higher degrees.

We then go on to describe the interpretation of the above results in the context of branched rough paths. The main tool is the Hairer–Kelly map [28], that we introduce and describe in Lemma 5.1 and then use to encode branched rough paths via anisotropic geometric rough paths, along the same lines as in [2, Theorem 4.3].

Theorem 1.6. *Let X be a branched γ -rough path. There exists an anisotropic geometric rough path \bar{X} indexed by words on the alphabet \mathcal{T}_N , with exponents $(\gamma_\tau = \gamma|\tau|, \tau \in \mathcal{T}_N)$, and such that $\langle X, \tau \rangle = \langle \bar{X}, \psi(\tau) \rangle$, where ψ is the Hairer–Kelly map.*

The main difference of this result with [28, Theorem 1.9] is that we obtain an anisotropic geometric rough path instead of a classical geometric rough path. This means that we do not construct unneeded components, i.e. components with regularity larger than 1, and we also obtain the right Hölder estimates in terms of the size of the indexing tree. This addresses two problems mentioned in Hairer and Kelly’s work, namely Remarks 4.14 and 5.9 in [28].

We then use Theorem 1.5 and Theorem 1.6 to construct our action on branched rough paths. Given $(g, X) \in \mathcal{C}^\gamma \times \mathbf{BRP}^\gamma$, we construct the anisotropic geometric rough paths \bar{X} and $g\bar{X}$ and then define the branched rough path $gX \in \mathbf{BRP}^\gamma$ as $\langle gX, \tau \rangle = \langle g\bar{X}, \psi(\tau) \rangle$, where ψ is the Hairer–Kelly map.

Our approach also does not make use of Foissy–Chapoton’s Hopf-algebra isomorphism [10, 20] between the Butcher–Connes–Kreimer Hopf algebra and the shuffle algebra over a complicated set I of trees as is done in [2]. This allows us to construct an action of a *larger* group on the set of branched rough paths; indeed, using the above isomorphism one would obtain a transformation group parametrized by $(g^\tau)_{\tau \in I}$ where I is the aforementioned set of trees of Foissy–Chapoton’s results and

$g^\tau \in C^{|\tau|}$; on the other hand our approach yields a transformation group parametrized by $(g^\tau)_{\tau \in \mathcal{T}_N}$. With the smaller set $I \cap \mathcal{T}_N$, transitivity of the action $g \mapsto gX$ would be lost.

Finally we note that we use a special property of the Butcher-Connes-Kreimer Hopf algebra: the fact that it is freely generated as an algebra by the set of trees, so defining characters over it is significantly easier than in the geometric case. To define an element $X \in G$ it suffices to give the values $\langle X, \tau \rangle$ for all trees $\tau \in \mathcal{T}$; by freeness there is a unique multiplicative extension to all of \mathcal{H} . This is not at all the case for geometric rough paths: the shuffle algebra $T(A)$ over an alphabet A is not free over the linear span of words so if one is willing to define a character X over $T(A)$ there are additional algebraic constraints that the values of X on words must satisfy.

Outline. We start by reviewing all the theoretical concepts needed to make the exposition in this section formal. In Section 3 we state and prove the main result of this chapter. We extend the notion of rough path and we give an explicit construction of such a generalized rough path above any given path $x \in C^\gamma$. Next, in Section 4.3 we extend this result to the class of anisotropic geometric rough paths. Finally, in Section 4 we connect our construction with M. Gubinelli's branched rough paths, and we extend M. Hairer and D. Kelly's work in Section 5.1. We also explore possible connections with renormalisation in Section 6 by studying how our construction behaves under modification of the underlying paths. Then, we connect this approach with a recent work by Bruned, Chevyrev, Friz and Preiß [4] in Section 6.1, who borrowed ideas from the theory of Regularity Structures [6, 27] and proposed a renormalisation procedure for geometric and branched rough paths [4] based on pre-Lie morphisms.

The main difference between our result and the BCFP procedure is that they consider translation only by time-independent factors, whereas –under reasonable hypotheses– we are also able to handle general translations depending on the time parameter. We also mention that some further algebraic aspects of renormalisation in rough paths have been recently developed in [5].

2 Preliminaries

A Hopf algebra \mathcal{H} is a vector space endowed with an *associative product* $m: \mathcal{H} \otimes \mathcal{H} \rightarrow \mathcal{H}$:

$$m(m \otimes \text{id}) = m(\text{id} \otimes m),$$

and a *coassociative coproduct* $\Delta: \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{H}$:

$$(\text{id} \otimes \Delta)\Delta = (\Delta \otimes \text{id})\Delta,$$

satisfying moreover certain compatibility assumptions; \mathcal{H} is also supposed to have a *unit* $1 \in \mathcal{H}$, a *counit* $\varepsilon \in \mathcal{H}^*$ and an *antipode* $S: \mathcal{H} \rightarrow \mathcal{H}$ such that

$$m(\text{id} \otimes S)\Delta x = \varepsilon(x)1 = m(S \otimes \text{id})\Delta x$$

for all $x \in \mathcal{H}$. As usual we will use the more compact notation $m(x \otimes y) = xy$. The reader is referred to the papers [8, 31] for further details.

Definition 2.1. We say that the Hopf algebra \mathcal{H} is graded if it can be decomposed as a direct sum

$$\mathcal{H} = \bigoplus_{n=0}^{\infty} \mathcal{H}_{(n)} \tag{2.1}$$

with

$$m: \mathcal{H}_{(n)} \otimes \mathcal{H}_{(m)} \rightarrow \mathcal{H}_{(n+m)}, \quad \Delta: \mathcal{H}_{(n)} \rightarrow \bigoplus_{p+q=n} \mathcal{H}_{(p)} \otimes \mathcal{H}_{(q)}. \quad (2.2)$$

In a graded Hopf algebra, each element $x \in \mathcal{H}$ can be decomposed as a sum

$$x = \sum_{n=0}^{\infty} x_n, \quad x_n \in \mathcal{H}_{(n)}, \quad (2.3)$$

where only a finite number of the summands are non-zero. We call each x_n the *homogeneous part of degree n* of x , and elements of $\mathcal{H}_{(n)}$ are said to be homogeneous of degree n . In this case we write $|x_n| = n$.

Definition 2.2. The graded Hopf algebra \mathcal{H} is connected if the degree 0 part is one-dimensional. It is locally finite if $\dim \mathcal{H}_{(n)} < \infty$ for all $n \geq 0$.

From now on we consider a graded connected locally finite Hopf algebra \mathcal{H} . Then, for any homogeneous element $x \in \mathcal{H}_{(n)}$ the coproduct can be written as

$$\Delta x = x \otimes 1 + 1 \otimes x + \Delta' x, \quad \text{where } \Delta' x \in \bigoplus_{\substack{p+q=n \\ p, q \geq 1}} \mathcal{H}_{(p)} \otimes \mathcal{H}_{(q)}$$

and $\Delta': \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{H}$ is known as the *reduced coproduct*. Furthermore, the coassociativity of Δ and of Δ' , i.e. the identity $(\Delta' \otimes \text{id})\Delta' = (\text{id} \otimes \Delta')\Delta'$, allows to unambiguously define their iterates $\Delta_n, \Delta'_n: \mathcal{H} \rightarrow \mathcal{H}^{\otimes(n+1)}$ by setting for $n \geq 2$

$$\Delta_n = (\text{id} \otimes \Delta_{n-1})\Delta, \quad \Delta'_n = (\text{id} \otimes \Delta'_{n-1})\Delta'.$$

Then we have, for a homogeneous element $x \in \mathcal{H}_{(k)}$ of degree k ,

$$\Delta'_n x \in \bigoplus_{\substack{p_1 + \dots + p_{n+1} = k \\ p_j \geq 1}} \mathcal{H}_{(p_1)} \otimes \dots \otimes \mathcal{H}_{(p_{n+1})}.$$

Remark 2.3. These properties of the iterated coproduct imply that the bialgebra (\mathcal{H}, Δ) is *conilpotent*, that is, for each homogeneous $x \in \mathcal{H}_{(k)}$ there is an integer $n \leq k$ such that $\Delta'_n x = 0$. We obtain also the inclusion

$$\Delta'_n \mathcal{H}_{(n+1)} \subset \mathcal{H}_{(1)}^{\otimes(n+1)},$$

that is, the n -fold reduced coproduct of a homogeneous element of degree $n+1$ is a sum of $(n+1)$ -fold tensor products of homogeneous elements of degree 1.

We recall that in general the dual space \mathcal{H}^* carries an algebra structure given by the convolution product \star , dual to the coproduct Δ , defined by

$$\langle f \star g, x \rangle := \langle f \otimes g, \Delta x \rangle.$$

For a collection of maps $f_1, \dots, f_k \in \mathcal{H}^*$ we have the formula

$$f_1 \star \dots \star f_k = (f_1 \otimes \dots \otimes f_k) \circ \Delta_{k-1}. \quad (2.4)$$

Definition 2.4. A character on \mathcal{H} is a non-zero linear map $X: \mathcal{H} \rightarrow \mathbb{R}$

$$\langle X, xy \rangle = \langle X, x \rangle \langle X, y \rangle, \quad \forall x, y \in \mathcal{H}.$$

for all $x, y \in \mathcal{H}$. We call G the set of all characters on \mathcal{H} . An infinitesimal character (or derivation) on \mathcal{H} is a linear map $\alpha: \mathcal{H} \rightarrow \mathbb{R}$ such that

$$\langle \alpha, xy \rangle = \langle \alpha, x \rangle \langle \varepsilon, y \rangle + \langle \varepsilon, x \rangle \langle \alpha, y \rangle, \quad \forall x, y \in \mathcal{H}.$$

We call \mathfrak{g} the set of all infinitesimal characters on \mathcal{H} .

We observe that necessarily $\langle X, 1 \rangle = 1$ and $\langle \alpha, 1 \rangle = 0$ for all $X \in G$ and $\alpha \in \mathfrak{g}$. It is well known that the (G, \star, ε) is a group with product \star , unit ε and inverse $X^{-1} = X \circ S$ where S is the antipode defined above. Moreover $(\mathfrak{g}, [\cdot, \cdot])$ is a Lie algebra with bracket $[\alpha, \beta] := \alpha \star \beta - \beta \star \alpha$. See e.g. [31].

2.1 Nilpotent Lie algebras

From (2.2) we have

Lemma 2.5. For any $N \in \mathbb{N}$ the subspace

$$\mathcal{H}_N := \bigoplus_{k=0}^N \mathcal{H}_{(k)} \subset \mathcal{H}$$

is a counital subcoalgebra of $(\mathcal{H}, \Delta, \varepsilon)$.

By Lemma 2.5 we can consider the dual algebra $(\mathcal{H}_N^*, \star, \varepsilon)$. This algebra is also graded and connected, since we have the natural grading

$$\mathcal{H}_N^* = \bigoplus_{k=0}^N \mathcal{H}_{(k)}^*, \quad \mathcal{H}_N^* \ni \alpha = \sum_{k=0}^N \alpha_{(k)}, \quad (2.5)$$

where $\alpha_{(k)}: \mathcal{H}_N \rightarrow \mathbb{R}$ is defined by $\alpha_{(k)}(x) := \alpha(x_k)$ with the notation (2.3).

Since \mathcal{H}_N is not a subalgebra of \mathcal{H} , the notions of character and infinitesimal character on \mathcal{H}_N^* are not well-defined. We can however introduce their *truncated* versions.

Definition 2.6. We say that $X \in \mathcal{H}_N^* \setminus \{0\}$ is a truncated character on \mathcal{H}_N if

$$\langle X, xy \rangle = \langle X, x \rangle \langle X, y \rangle$$

holds for all $x \in \mathcal{H}_{(n)}, y \in \mathcal{H}_{(m)}$ with $n + m \leq N$. We call G^N the space of truncated characters on \mathcal{H}_N .

Likewise, we say that $\alpha \in \mathcal{H}_N^*$ is a truncated infinitesimal character if

$$\langle \alpha, xy \rangle = \langle \alpha, x \rangle \langle \varepsilon, y \rangle + \langle \varepsilon, x \rangle \langle \alpha, y \rangle$$

holds for all $x \in \mathcal{H}_{(n)}, y \in \mathcal{H}_{(m)}$ with $n + m \leq N$. We call \mathfrak{g}^N the space of truncated infinitesimal characters on \mathcal{H}_N .

Lemma 2.7. *There are a canonical inclusions $\mathcal{H}_N^* \hookrightarrow \mathcal{H}_{N+1}^* \hookrightarrow \mathcal{H}^*$, which induce canonical inclusion $\mathfrak{g}^N \hookrightarrow \mathfrak{g}^{N+1} \hookrightarrow \mathfrak{g}$. Moreover such canonical inclusions are right-inverse for the corresponding restriction maps $\mathcal{H}^* \rightarrow \mathcal{H}_{N+1}^* \rightarrow \mathcal{H}_N^*$.*

Proof. Using the notation (2.5), we can extend $\alpha \in \mathcal{H}_N^*$ to $\alpha \in \mathcal{H}_{N+1}^*$ (respectively \mathcal{H}^*) by setting $\alpha_{(N+1)} \equiv 0$ (respectively $\alpha_{(k)} \equiv 0$ for all $k \geq N+1$). Trivially this extension takes \mathcal{H}_N^* to \mathcal{H}_{N+1}^* . If $\alpha \in \mathfrak{g}^N$ and $x, y \in \mathcal{H}_N$ are such that $|x| + |y| \leq N+1$ then

$$\begin{aligned} \langle \alpha, xy \rangle &= \left\langle \alpha, \sum_{j=0}^{N+1} (xy)_j \right\rangle = \sum_{j=0}^N \langle \alpha, (xy)_j \rangle = \sum_{j=0}^N \sum_{k=0}^j \langle \alpha, x_k y_{k-j} \rangle \\ &= \sum_{j=0}^N (\langle \alpha, x_j \rangle \langle \varepsilon, y \rangle + \langle \varepsilon, x \rangle \langle \alpha, y_j \rangle) = \langle \alpha, x \rangle \langle \varepsilon, y \rangle + \langle \varepsilon, x \rangle \langle \alpha, y \rangle. \end{aligned}$$

so that the extension of α is in \mathfrak{g}^{N+1} . The same argument yields the inclusion $\mathfrak{g}^N \hookrightarrow \mathfrak{g}$. \square

There are also the truncated exponential $\exp_N: \mathcal{H}_N^* \rightarrow \mathcal{H}_N^*$ and logarithm $\log_N: \mathcal{H}_N^* \rightarrow \mathcal{H}_N^*$, defined by the sums

$$\exp_N(\alpha) := \sum_{k=0}^N \frac{1}{k!} \alpha^{*k} \Big|_{\mathcal{H}_N}, \quad \log_N(X) := \sum_{k=1}^N \frac{(-1)^{k+1}}{k} (X - \varepsilon)^{*k} \Big|_{\mathcal{H}_N}. \quad (2.6)$$

The proof of the next result can be found for instance in [19, Thm 77].

Lemma 2.8. *$(G^N, \star, \varepsilon)$ is a group and $(\mathfrak{g}^N, [\cdot, \cdot])$ is a Lie algebra. Moreover, $\exp_N: \mathfrak{g}^N \rightarrow G^N$ is a bijection with inverse $\log_N: G^N \rightarrow \mathfrak{g}^N$.*

For every $k \geq 0$ we define now, using the notation (2.5),

$$W_k := \{ \alpha \in \mathfrak{g} : \alpha = \alpha_{(k)} \}.$$

Lemma 2.9. *For all $n, m \geq 0$ we have $[W_n, W_m] \subset W_{n+m}$.*

Proof. Let $x \in \mathcal{H}$. With the notation (2.3) we have for $\alpha \in W_n$ and $\beta \in W_m$

$$(\alpha \star \beta - \beta \star \alpha)(x) = (\alpha \otimes \beta - \beta \otimes \alpha) \Delta x = (\alpha \otimes \beta - \beta \otimes \alpha) \Delta x_{n+m}$$

by (2.2). \square

By the canonical inclusion of Lemma 2.7, we observe that

$$\mathfrak{g}^N = \bigoplus_{k=1}^N W_k, \quad \mathfrak{g}^N \ni \alpha = \sum_{k=0}^N \alpha_{(k)} \quad (2.7)$$

in the notation (2.5). With this decomposition \mathfrak{g}^N becomes by Lemma 2.9 a graded Lie algebra. We recall that the *center* of \mathfrak{g}^N is the subspace of all $w \in \mathfrak{g}^N$ such that $[\alpha, w] = 0$ for all $\alpha \in \mathfrak{g}^N$, while the *center* of G^N is the set of all $X \in G^N$ such that $X \star Y = Y \star X$ for all $Y \in G^N$.

Proposition 2.10. *W_N is contained in the center of \mathfrak{g}^N and $\exp_N(W_N)$ is a subgroup contained in the center of G^N .*

Proof. Let $\alpha \in \mathfrak{g}^N$ and $w \in W_N$. Clearly, $\langle [\alpha, w], x \rangle$ is zero unless $|x| = N$. In this case

$$\langle [\alpha, w], x \rangle = \langle \alpha \otimes w - w \otimes \alpha, \Delta x \rangle = \langle \alpha, 1 \rangle \langle w, x \rangle - \langle w, x \rangle \langle \alpha, 1 \rangle = 0$$

since $\langle w, y \rangle = \langle w, y_N \rangle$, in the notation (2.3). The second assertion follows easily: it is enough to write $X = \exp_N(w)$ and $Y = \exp_N(\alpha)$ with $\alpha \in \mathfrak{g}^N$ and $w \in W_N$ and use the explicit representation (2.6) of \exp_N and the fact that $\alpha \star w = w \star \alpha$. \square

The next (famous) result describes the group law on G^N in terms of an operation on \mathfrak{g}^N via the exponential/logarithmic map.

Theorem 2.11 (Baker–Campbell–Hausdorff). *For all $\alpha, \beta \in \mathfrak{g}^N$, we have*

$$\log_N(\exp_N(\alpha) \star \exp_N(\beta)) \in \mathfrak{g}^N.$$

We define the map $\text{BCH}_N: \mathfrak{g}^N \times \mathfrak{g}^N \rightarrow \mathfrak{g}^N$ by

$$\text{BCH}_N(\alpha, \beta) := \log_N(\exp_N(\alpha) \star \exp_N(\beta)). \quad (2.8)$$

Another way to interpret this theorem is to say that there exists an element $\gamma = \text{BCH}_N(\alpha, \beta) \in \mathfrak{g}^N$ such that $\exp_N(\alpha) \star \exp_N(\beta) = \exp_N(\gamma)$.

It is a classical result that the map BCH_N is formed by a sum of iterated Lie brackets of α and β , where the first terms are

$$\text{BCH}_N(\alpha, \beta) = \alpha + \beta + \frac{1}{2}[\alpha, \beta] + \frac{1}{12}[\alpha, [\alpha, \beta]] - \frac{1}{12}[\beta, [\alpha, \beta]] + \cdots, \quad (2.9)$$

and the following ones are explicit but difficult to compute. Nevertheless, fully explicit formulas have been known since 1947 by Dynkin [15].

For our purposes, however, Dynkin's formula is too complicated (for example, the regularity argument in step 2 of the proof of Theorem 3.4 would not be as evident) so we rely on a different expression first shown by Reutenauer [34]. In order to describe it, let $\varphi_k: (\mathcal{H}^*)^{\otimes k} \rightarrow \mathcal{H}^*$ be the linear map

$$\varphi_k(\alpha_1 \otimes \cdots \otimes \alpha_k) = \sum_{\sigma \in S_k} a_\sigma \alpha_{\sigma(1)} \star \cdots \star \alpha_{\sigma(k)} \quad (2.10)$$

where S_k denotes the symmetric group of order k , and $a_\sigma := \frac{(-1)^{d(\sigma)}}{k} \binom{k-1}{d(\sigma)}^{-1}$ is a constant depending only on the *descent number* $d(\sigma)$ of the permutation $\sigma \in S_k$, namely the number of $i \in \{1, \dots, k-1\}$ such that $\sigma(i) > \sigma(i+1)$.

Lemma 2.12 (Reutenauer's formula). *For all $\alpha, \beta \in \mathfrak{g}^N$*

$$\text{BCH}_N(\alpha, \beta) = \sum_{k=1}^N \sum_{i+j=k} \frac{1}{i!j!} \varphi_k(\alpha^{\otimes i} \otimes \beta^{\otimes j}). \quad (2.11)$$

Moreover, for all $i \in \{0, \dots, N\}$, we have $\varphi_N(\alpha^{\otimes i} \otimes \beta^{\otimes(N-i)}) \in W_N$.

Proof. Let us suppose first that $T(V)$ is the (completed) tensor algebra over a two-dimensional vector space V , with V linearly generated by $\{e_1, e_2\}$. Then the result is contained in Reutenauer's paper [34] where the free step- N nilpotent Lie algebra \mathfrak{L}^N plays the rôle of \mathfrak{g}^N . We want now to show how this implies the same result in our more general setting.

Let $\alpha, \beta \in \mathfrak{g}^N$ and let $\Phi: (T(V)_N, \otimes) \rightarrow (\mathcal{H}_N^*, \star)$ be the unique algebra morphism such that $\Phi(e_1) = \alpha$, $\Phi(e_2) = \beta$. Then Φ restricts to a Lie-algebra morphism $\Phi: \mathfrak{L}^N \rightarrow \mathfrak{g}^N$ such that $\text{BCH}_N(\alpha, \beta) = \Phi(\text{BCH}_N(e_1, e_2))$ and therefore (2.11) follows.

In order to prove the first formula, we first note that Φ is not a graded morphism, since the generators e_1 and e_2 are homogeneous of degree 1 in $T(V)_N$, but α and β are in general not homogeneous in \mathcal{H}_N^* . However, from the bilinearity of the Lie bracket and Lemma 2.9 we obtain

$$[W_n \oplus \cdots \oplus W_N, W_m \oplus \cdots \oplus W_N] \subset W_{n+m} \oplus W_{n+m+1} \oplus \cdots \oplus W_N.$$

Then, if $\alpha_1, \dots, \alpha_k \in \mathfrak{g}^N$ then $\varphi_k(\alpha_1 \otimes \cdots \otimes \alpha_k) \in W_k \oplus \cdots \oplus W_N$. \square

From all these considerations we obtain the following result on the map

$$\text{BCH}_{(n+1)}: \mathfrak{g}^{n+1} \times \mathfrak{g}^{n+1} \rightarrow W_{n+1}, \quad \text{BCH}_{(n+1)} := \text{BCH}_{n+1} - \text{BCH}_n, \quad n \geq 0. \quad (2.12)$$

Note that $\text{BCH}_{(n+1)}$ takes indeed values in W_{n+1} rather than in \mathfrak{g}^{n+1} by (both assertions of) Lemma 2.12.

Lemma 2.13. *Let $x \in \mathcal{H}_{(n+1)}$ and $\alpha, \beta \in \mathfrak{g}^{n+1}$. Then*

$$\langle \text{BCH}_{(n+1)}(\alpha, \beta), x \rangle = \sum_{i+j=n+1} \frac{1}{i!j!} \sum_{(x)} \sum_{\sigma \in S_{n+1}} a_\sigma \prod_{p=1}^i \langle \alpha, x_{(\sigma^{-1}(p))} \rangle \prod_{q=i+1}^{n+1} \langle \beta, x_{(\sigma^{-1}(q))} \rangle, \quad (2.13)$$

where

$$\Delta'_n x = \sum_{(x)} x_{(1)} \otimes \cdots \otimes x_{(n+1)} \in \mathcal{H}_{(1)}^{\otimes(n+1)}.$$

Proof. Set $\alpha_1 = \cdots = \alpha_i := \alpha$, $\alpha_{i+1} = \cdots = \alpha_{n+1} := \beta$. Then the result follows directly from the definition of φ_k in (2.10) together with (2.4) and the fact that since $\langle \alpha_j, 1 \rangle = 0$ we can write

$$\alpha_1 \star \cdots \star \alpha_{n+1} = (\alpha_1 \otimes \cdots \otimes \alpha_{n+1}) \Delta'_n \quad (2.14)$$

instead (note the reduced coproduct in place of the full coproduct). \square

2.2 A distance on the group of truncated characters

Now we introduce a distance on G^N which is well adapted to the notion of rough paths, to be introduced in Definition 3.1 below. We fix a basis B of \mathcal{H}_N and define a norm $\|\cdot\|$ on this space by requiring that B is orthonormal. There is a unique function $c: B \times B \times B \rightarrow \mathbb{R}$ such that

$$\Delta v = \sum_{v_1, v_2 \in B} c(v, v_1, v_2) v_1 \otimes v_2, \quad \forall v \in B.$$

Then we define

$$K := \max_{v \in B} \sum_{v_1, v_2 \in B} |c(v, v_1, v_2)| < \infty, \quad |||f||| := K \sup_{v \in B} |\langle f, v \rangle|, \quad f \in \mathcal{H}_N^*.$$

Then, if $f, g \in \mathcal{H}_N^*$, for any $v \in B$

$$|\langle f \star g, v \rangle| \leq \sum_{v_1, v_2 \in B} |c(v, v_1, v_2)| |\langle f, v_1 \rangle| |\langle g, v_2 \rangle| \leq \frac{1}{K} |||f||| |||g|||,$$

thus $|||f \star g||| \leq |||f||| |||g|||$. We set now for all $X \in G^N$

$$|X| := \max_{k=1, \dots, N} (k! |||X_{(k)}|||)^{1/k} + \max_{k=1, \dots, N} (k! |||(X^{-1})_{(k)}|||)^{1/k}, \quad (2.15)$$

where for $X \in G^N \subset \mathcal{H}_N^*$ we use the notation (2.5). We define $G^N \times G^N \ni (X, Y) \mapsto \rho^N(X, Y) := |X^{-1} \star Y| \in \mathbb{R}_+$, i.e. by (2.15)

$$\rho_N(X, Y) = \max_{k=1, \dots, N} (k! |||(Y^{-1} \star X)_{(k)}|||)^{1/k} + \max_{k=1, \dots, N} (k! |||(X^{-1} \star Y)_{(k)}|||)^{1/k} \quad (2.16)$$

Proposition 2.14. *The map ρ_N defines a left-invariant distance on the group G^N such that the metric space (G^N, ρ_N) is complete.*

Proof. We only need to prove that the function $|\cdot|$ defined in (2.15) is sub-additive, the other properties being clear. Note that for $X, Y \in G^N$, with the notation (2.5) we have

$$X \star Y = \left(\sum_{k=0}^N X_{(k)} \right) \star \left(\sum_{k=0}^N Y_{(k)} \right) = \sum_{k=0}^N \sum_{j=0}^k X_{(j)} \star Y_{(k-j)}. \quad (2.17)$$

Therefore

$$|||(X \star Y)_{(k)}||| \leq \sum_{j=0}^k |||X_{(j)}||| |||Y_{(k-j)}||| \leq \frac{1}{k!} \sum_{j=0}^k \binom{k}{j} |X|^j |Y|^{k-j} = \frac{1}{k!} (|X| + |Y|)^k$$

whence the result. \square

The next result is the analog of [30, Prop. 7].

Lemma 2.15. *If $X = \exp_N(w_1 + \dots + w_N)$ with $w_i \in W_i$, then*

$$C_N \max_{k=1, \dots, N} |||w_k|||^{1/k} \leq |X| \leq C_N \max_{k=1, \dots, N} |||w_k|||^{1/k}.$$

Proof. Using the notation (2.5), we have

$$X_{(k)} = \sum_{i=1}^k \frac{1}{i!} \sum_{j_1 + \dots + j_i = k} w_{j_1} \star \dots \star w_{j_i}$$

so that for all $k = 1, \dots, N$

$$\begin{aligned} (k! \|\|X_{(k)}\|\|)^{1/k} &\leq \left(\sum_{i=1}^k \frac{k!}{i!} \sum_{j_1+\dots+j_i=k} \left(\|\|w_{j_1}\|\|^{1/j_1} \right)^{j_1} \dots \left(\|\|w_{j_i}\|\|^{1/j_i} \right)^{j_i} \right)^{1/k} \\ &\leq \left(\sum_{i=1}^k \frac{k!}{i!} \sum_{j_1+\dots+j_i=k} \left(\max_{\ell=1,\dots,k} \|\|w_\ell\|\|^{1/\ell} \right)^{j_1+\dots+j_i} \right)^{1/k}. \end{aligned}$$

There are exactly $\binom{k-1}{i-1} \leq \frac{(k-1)^{i-1}}{(i-1)!}$ solutions to $j_1 + \dots + j_i = k$ so that

$$(k! \|\|X_{(k)}\|\|)^{1/k} \leq (k!(e^k - 1))^{1/k} \max_{\ell=1,\dots,k} \|\|w_\ell\|\|^{1/\ell}.$$

Since $X^{-1} = \exp_N(-w_1 - \dots - w_N)$, the bound for X^{-1} follows in the same way and we have therefore proved the desired upper bound for $|X|$. For the lower bound, we use the truncated logarithm

$$w_k = \sum_{i=1}^k \frac{(-1)^{i-1}}{i} \sum_{j_1+\dots+j_i=k} X_{(j_1)} \star \dots \star X_{(j_i)}.$$

Then we can estimate

$$\begin{aligned} \|\|w_k\|\|^{1/k} &\leq \left(\sum_{i=1}^k \frac{1}{i} \sum_{j_1+\dots+j_i=k} \left(\|\|X_{(j_1)}\|\|^{1/j_1} \right)^{j_1} \dots \left(\|\|X_{(j_i)}\|\|^{1/j_i} \right)^{j_i} \right)^{1/k} \\ &\leq \left(\sum_{i=1}^k \frac{1}{i} \binom{k-1}{i-1} \left(\max_{\ell=1,\dots,k} \|\|X_\ell\|\|^{1/\ell} \right)^{j_1+\dots+j_i} \right)^{1/k} \leq \frac{1}{c_N} |X| \end{aligned}$$

and the proof is complete. □

We now note that the function $|\cdot|$ and the distance ρ^N make G^N a *homogeneous group*, see [22] for an extensive treatment of this subject, and [30] for the case of tensor algebras and the relation with geometric rough paths.

To put it briefly, for all $r > 0$ we can define the following linear operator $\Omega_r: \mathcal{H}^* \rightarrow \mathcal{H}^*$

$$\Omega_r \alpha := \sum_{k \geq 0} r^k \alpha_{(k)}.$$

This family satisfies $\Omega_r \circ \Omega_s = \Omega_{rs}$, $r, s > 0$. Moreover $\Omega_r: \mathfrak{g}^N \rightarrow \mathfrak{g}^N$ is a Lie-algebra automorphism of \mathfrak{g}^N for all $r > 0$. Then they induce group automorphisms $\Lambda_r := \exp_N \circ \Omega_r \circ \log_N: G^N \rightarrow G^N$, $r > 0$. In the terminology of [22], $(\Omega_r)_{r>0}$ is a family of *dilations* on the finite-dimensional Lie algebra \mathfrak{g}^N and G^N is a *homogeneous group*.

Note that the function $|\cdot|: G^N \rightarrow \mathbb{R}_+$ is continuous, satisfies $|\Lambda_r X| = r|X|$ for all $r > 0$ and $X \in G^N$, and $|X| = 0$ for $X \in G^N$ if and only if $X = 1$. These three properties make $|\cdot|$ a *homogeneous norm* on G^N , see [22]. The homogeneity property plays an important role in the proof of Theorem 3.4 below.

3 Construction of Rough paths

As in the previous section, we fix a locally-finite graded connected Hopf algebra \mathcal{H} . We also fix a number $\gamma \in]0, 1[$ and let $N := \lfloor \gamma^{-1} \rfloor$ be the biggest integer such that $N\gamma \leq 1$. Without loss of generality we can fix a basis B of \mathcal{H}_N consisting only of homogeneous elements and in particular we let $\{e_1, \dots, e_d\} = B \cap \mathcal{H}_{(1)}$ where $d := \dim \mathcal{H}_{(1)}$.

Definition 3.1. A (\mathcal{H}, γ) -rough path is a function $X : [0, 1]^2 \rightarrow G^N$, with $N = \lfloor \gamma^{-1} \rfloor$, which satisfies Chen's rule

$$X_{su} \star X_{ut} = X_{st}, \quad s, u, t \in [0, 1], \quad (3.1)$$

and such that for all $v \in B$

$$|\langle X_{st}, v \rangle| \lesssim |t - s|^{\gamma|v|}. \quad (3.2)$$

If $x^i : [0, 1] \rightarrow \mathbb{R}$, $i = 1, \dots, d$, is such that $x_t^i - x_s^i = \langle X_{st}, e_i \rangle$, $s, t \in [0, 1]$, we say that X is a γ -rough path over (x^1, \dots, x^d) .

Remark 3.2. By specializing this definition to different choices of \mathcal{H} we recover both *geometric rough paths* [29] where \mathcal{H} is the shuffle Hopf algebra over an alphabet, *branched rough paths* [25] where \mathcal{H} is the Butcher–Connes–Kreimer Hopf algebra on decorated non-planar rooted trees, and also *planarly branched rough paths* [14].

We remark that there is a bijection between

- 1 functions $X : [0, 1]^2 \rightarrow G^N$ such that $X_{su} \star X_{ut} = X_{st}$, for all $s, u, t \in [0, 1]$,
- 2 functions $\mathbb{X} : [0, 1] \rightarrow G^N$ such that $\mathbb{X}_0 = 1$,

given by

$$X \mapsto \mathbb{X}, \quad \mathbb{X}_t := X_{0t}, \quad \mathbb{X} \mapsto X, \quad X_{st} := \mathbb{X}_s^{-1} \star \mathbb{X}_t, \quad s, t \in [0, 1]. \quad (3.3)$$

Proposition 3.3. Let $\mathbb{X} : [0, 1] \rightarrow G^N$ and $X : [0, 1]^2 \rightarrow G^N$ as in (3.3). Then X is a (\mathcal{H}, γ) -rough path as in Definition 3.1 if and only if \mathbb{X} is γ -Hölder with respect to the metric ρ_N defined in (2.15).

Proof. First note that the distance in (2.16) is defined with respect to a fixed (but arbitrary) basis so we use the basis B fixed at the beginning of this section. Also, due to the above remark we only have to verify that \mathbb{X} is γ -Hölder with respect to ρ_N if and only if X satisfies (3.2) using the same basis. In one direction, if \mathbb{X} is γ -Hölder then, by definition

$$|X_{st}| = \rho_N(\mathbb{X}_s, \mathbb{X}_t) \lesssim |t - s|^\gamma$$

and so, for a basis element $v \in B$ we have

$$|\langle X_{st}, v \rangle| \lesssim |t - s|^{\gamma|v|}.$$

Conversely, if (3.2) holds then $|X_{st}| \lesssim |t - s|^\gamma$ and so by definition also $\rho_N(\mathbb{X}_s, \mathbb{X}_t) \lesssim |t - s|^\gamma$, i.e. \mathbb{X} is γ -Hölder with respect to ρ_N . \square

We now come to the problem of existence. Our construction of a rough path in the sense of Definition 3.1 over an arbitrary collection of γ -Hölder paths (x^1, \dots, x^d) relies in the following extension theorem. We note that the proof is a reinterpretation of the approach of Lyons-Victoir [30, Theorem 1] in the context of a more general graded Hopf-algebra \mathcal{H} .

Theorem 3.4 (Rough path extension). *Let $1 \leq n \leq N - 1$ and $\gamma \in]0, 1[$ such that $\gamma^{-1} \notin \mathbb{N}$. Suppose we have a γ -Hölder path $\mathbb{X}^n : [0, 1] \rightarrow (G^n, \rho_n)$. There is a γ -Hölder path $\mathbb{X}^{n+1} : [0, 1] \rightarrow (G^{n+1}, \rho_{n+1})$ extending \mathbb{X}^n , i.e. such that $\mathbb{X}^{n+1}|_{\mathcal{H}_n} = \mathbb{X}^n$.*

A key tool is the following technical lemma whose proof can be found in [30, Lemma 2].

Lemma 3.5. *Let (E, ρ) be a complete metric space and set*

$$D = \{t_k^m := k2^{-m} : m \geq 0, k = 0, \dots, 2^m - 1\}.$$

Suppose $y : D \rightarrow E$ is a path satisfying the bound $\rho(y_{t_k^m}, y_{t_{k+1}^m}) \lesssim 2^{-\gamma m}$ for some $\gamma \in (0, 1)$. Then, there exists a γ -Hölder path $x : [0, 1] \rightarrow E$ such that $x|_D = y$.

Proof of Theorem 3.4. The construction of \mathbb{X}^{n+1} is made in two steps.

Step 1. For $m \geq 0$ and $k \in \{0, \dots, 2^m\}$ we define $t_k^m := k2^{-m} \in [0, 1]$. Then we define the following sets of dyadics in $[0, 1]$

$$D_{(m)} := \{t_k^m \mid k = 0, \dots, 2^m\}, \quad D_m := \bigcup_{n=0, \dots, m} D_{(n)}, \quad D := \bigcup_{m \geq 0} D_{(m)}.$$

Set $X_{st} = (\mathbb{X}_s^n)^{-1} \star \mathbb{X}_t^n \in G^n$ and $L_{st} = \log_n(X_{st}) \in \mathfrak{g}^n$ where \log_n was defined in (2.6). Then, the Baker–Campbell–Hausdorff formula (2.8) and Chen’s rule (3.1) imply that

$$L_{st} = \text{BCH}_n(L_{su}, L_{ut}). \quad (3.4)$$

We look for $Y : [0, 1]^2 \rightarrow G^{n+1}$ such that Y satisfies Chen’s rule (3.1) and $Y|_{\mathcal{H}_N} = X$. We use throughout the proof that $\mathfrak{g}^n \subset \mathfrak{g}^{n+1}$, see Lemma 2.7.

In a first step, we define $Y : D \times D \rightarrow G^{n+1}$. In the second step we show that Y has suitable uniform continuity properties and can thus be extended to $[0, 1]^2$ using Lemma 3.5.

The construction of $Y : D \times D \rightarrow G^{n+1}$ goes through a construction of $Y^m : D_m \times D_m \rightarrow G^{n+1}$ by recursion on $m \geq 0$. We claim that for all $m \geq 0$ we can find Y^m such that

- 1 Y^m satisfies Chen’s relation on D_m , namely $Y_{ab}^m \star Y_{bc}^m = Y_{ac}^m$ for all $a, b, c \in D_m$
- 2 for any $n \in \{0, \dots, m\}$ and $k, \ell \in \{0, \dots, 2^{m-n}\}$, we have the compatibility relation

$$Y_{t_{k2^n}^m t_{\ell 2^n}^m}^m = Y_{t_k^{m-n} t_\ell^{m-n}}^{m-n}.$$

- 3 Y^m restricted to \mathcal{H}_n is equal to $X : D_m \times D_m \rightarrow G^n$, in the sense that

$$Y_{ab}^m|_{\mathcal{H}_n} = X_{ab}, \quad \forall a, b \in D_m.$$

- 4 for all $k = 0, \dots, 2^m - 1$, setting

$$Z_{t_k^m t_{k+1}^m}^m := \log_{n+1} \left(Y_{t_k^m t_{k+1}^m}^m \star \exp_{n+1} \left(-L_{t_k^m t_{k+1}^m} \right) \right),$$

we have $Z_{t_k^m t_{k+1}^m}^m \in W_{n+1}$.

For $m = 0$, we set $Y_{01}^0 = \exp_{n+1}(L_{01})$, $Y_{00}^0 = Y_{11}^0 := \varepsilon$, and $Z_{01}^0 := 0 \in W_{n+1}$. For $x \in \mathcal{H}_n$, we have $\langle \exp_{n+1}(L_{01}), x \rangle = \langle \exp_n(L_{01}), x \rangle$, so that Y^0 restricted to \mathcal{H}_n is equal to $X: D_0 \times D_0 \rightarrow G^n$.

Let now $m \geq 1$, and suppose that $Y^{m-1}: D_{m-1} \times D_{m-1} \rightarrow G^{n+1}$ has been constructed with the above properties. We start by defining $Y_{tt}^m = \varepsilon$ for all $t \in D_{(m)}$. Let us consider three consecutive points in $D_{(m)}$ of the form

$$s = t_{2k}^m, \quad u = t_{2k+1}^m, \quad t = t_{2k+2}^m$$

for some $k = 0, \dots, 2^{m-1} - 1$. Note that $s = t_k^{m-1}$ and $t = t_{k+1}^{m-1}$, so that $Z_{st}^m := Z_{st}^{m-1} \in W_{n+1}$ is already defined by the recurrence hypothesis. We define Z_{su}^m and Z_{ut}^m as follows

$$Z_{su}^m = Z_{ut}^m := \frac{1}{2} (Z_{st}^{m-1} - \text{BCH}_{(n+1)}(L_{su}, L_{ut})), \quad (3.5)$$

where $\text{BCH}_{(n+1)} = \text{BCH}_{n+1} - \text{BCH}_n: \mathfrak{g}^{n+1} \times \mathfrak{g}^{n+1} \rightarrow W_{n+1}$, see (2.12). Since by recurrence $Z_{st}^{m-1} \in W_{n+1}$, we obtain that $Z_{su}^m, Z_{ut}^m \in W_{n+1}$ and

$$Z_{su}^m + Z_{ut}^m = Z_{st}^{m-1} - \text{BCH}_{(n+1)}(L_{su}, L_{ut}) = L_{st} + Z_{st}^m - \text{BCH}_{n+1}(L_{su}, L_{ut}) \quad (3.6)$$

where in the last equality we have applied (3.4). Then we set

$$Y_{su}^m := \exp_{n+1}(L_{su} + Z_{su}^m), \quad Y_{ut}^m := \exp_{n+1}(L_{ut} + Z_{ut}^m).$$

Since $\exp_{n+1}(W_{n+1})$ is in the center of G^{n+1} by Proposition 2.10, we obtain that

$$Y_{su}^m = \exp_{n+1}(L_{su}) \star \exp_{n+1}(Z_{su}^m), \quad Y_{ut}^m = \exp_{n+1}(L_{ut}) \star \exp_{n+1}(Z_{ut}^m).$$

By (2.8) and (3.6) the product is equal to

$$Y_{su}^m \star Y_{ut}^m = \exp_{n+1}(\text{BCH}_{n+1}(L_{su}, L_{ut}) + Z_{su}^m + Z_{ut}^m) = \exp_{n+1}(L_{st} + Z_{st}^m) = Y_{st}^m.$$

Let now $t_j^m, t_k^m \in D_{(m)}$ with $0 \leq j < k \leq 2^m$. We set

$$Y_{t_j^m t_k^m}^m := Y_{t_j^m t_{j+1}^m}^m \star \dots \star Y_{t_{k-1}^m t_k^m}^m, \quad Y_{t_k^m t_j^m}^m := \left(Y_{t_j^m t_k^m}^m \right)^{-1}$$

so that the identity $Y_{ab}^m \star Y_{bc}^m = Y_{ac}^m$ is valid for any $a, b, c \in D_{(m)}$.

We need now to check that this definition is compatible with the values already constructed on $D_{m-1} \times D_{m-1}$. By the recursion assumption, it is enough to show that for all $k, \ell \in \{0, \dots, 2^{m-1}\}$

$$Y_{t_{2k}^m t_{2\ell}^m}^m = Y_{t_k^{m-1} t_\ell^{m-1}}^{m-1}.$$

If $k = \ell$ or $|k - \ell| = 1$, then this is true by construction. Otherwise, if for example $k + 1 < \ell$ then

$$Y_{t_{2k}^m t_{2\ell}^m}^m = Y_{t_{2k}^m t_{2k+2}^m}^m \star \dots \star Y_{t_{2\ell-2}^m t_{2\ell}^m}^m = Y_{t_k^{m-1} t_{k+1}^{m-1}}^{m-1} \star \dots \star Y_{t_{\ell-1}^{m-1} t_\ell^{m-1}}^{m-1} = Y_{t_k^{m-1} t_\ell^{m-1}}^{m-1}$$

by the recursion property and the Chen relation satisfied by Y^m (respectively Y^{m-1}) on $D_{(m)}$ (resp. $D_{(m-1)}$).

We also have to check the extension property: for $x \in \mathcal{H}_n$ we have

$$\langle Y_{t_j^m t_{j+1}^m}^m, x \rangle = \langle \exp_{n+1}(L_{t_j^m t_{j+1}^m}) \star \exp_{n+1}(Z_{t_j^m t_{j+1}^m}^m), x \rangle = \langle \exp_n(L_{t_j^m t_{j+1}^m}), x \rangle = \langle X_{t_j^m t_{j+1}^m}, x \rangle.$$

By recurrence, we have proved that $Y^m: D_m \times D_m \rightarrow G^{n+1}$ is well defined for all $m \geq 0$, with the above properties. Therefore, we can unambiguously define $Y: D \times D \rightarrow G^{n+1}$,

$$Y_{st} := Y_{st}^m, \quad s, t \in D_m,$$

and Y indeed satisfies the Chen relation on D , namely $Y_{ab} \star Y_{bc} = Y_{ac}$ for all $a, b, c \in D$, and the restriction property

$$\langle Y_{ab}, x \rangle = \langle X_{ab}, x \rangle, \quad \forall a, b \in D, x \in \mathcal{H}_n.$$

Step 2. In order to have a $(\mathcal{H}_{n+1}, \gamma)$ -Hölder path, Definition 3.1 requires us to construct a γ -Hölder path with values in G^{n+1} , and for this we will use Lemma 3.5. Set

$$a_m := 2^{m(n+1)\gamma} \max_{k=0, \dots, 2^m-1} \left\| Z_{t_k^m, t_{k+1}^m}^m \right\|_{n+1}.$$

Then, if v is a basis element in $\mathcal{H}_{(n+1)}$ we have by (2.13), for $s = t_k^m$, $u = t_{k+1}^m$ and $t = t_{k+2}^m$

$$|\langle \text{BCH}_{(n+1)}(L_{su}, L_{ut}), v \rangle| \leq \sum_{(v)} \sum_{i+j=n+1} \frac{1}{i!j!} \sum_{\sigma \in S_{n+1}} |a_\sigma| \prod_{p=1}^i |\langle L_{su}, v_{(\sigma^{-1}(p))} \rangle| \prod_{q=i+1}^{n+1} |\langle L_{ut}, v_{(\sigma^{-1}(q))} \rangle|.$$

Now, since $v_{(j)} \in \mathcal{H}_{(1)}$ for all $j = 1, \dots, n+1$ we actually have that

$$|\langle L_{su}, v_{(j)} \rangle| \leq \sum_{k=1}^d |x_u^k - x_s^k| |v_{(j)}^k| \leq 2^{-m\gamma} \sum_{k=1}^d |v_{(j)}^k|$$

for some coefficients $v_{(j)}^k \in \mathbb{R}$ such that $v_{(j)} = \sum_{k=1}^d v_{(j)}^k e_k$, and we have a similar estimate for L_{ut} instead of L_{su} . Therefore we obtain that

$$\left\| \text{BCH}_{(n+1)}(L_{su}, L_{ut}) \right\|_{n+1} \leq C 2^{-m(n+1)\gamma},$$

where

$$C = K \max_v \sum_{(v)} \sum_{i+j=n+1} \frac{1}{i!j!} \sum_{\sigma \in S_{n+1}} |a_\sigma| \sum_{k_1, \dots, k_{n+1}=1}^{n+1} \prod_{\ell=1}^{n+1} |v_{(\ell)}^{k_\ell}|.$$

Therefore, from (3.5) we get

$$\max_{k=0, \dots, 2^m-1} \left\| Z_{t_k^m, t_{k+1}^m}^m \right\|_{n+1} \leq \frac{1}{2} \max_{k=0, \dots, 2^{m-1}-1} \left\| Z_{t_k^{m-1}, t_{k+1}^{m-1}}^{m-1} \right\|_{n+1} + \frac{1}{2} C 2^{-m(n+1)\gamma}$$

hence

$$a_m \leq 2^{(n+1)\gamma-1} a_{m-1} + \frac{C}{2}, \quad m \geq 1.$$

Since $a_0 = 0$ we can show by recurrence on $m \geq 0$

$$a_m \leq \frac{C}{2} \sum_{j=0}^{m-1} 2^{-j(1-(n+1)\gamma)}.$$

Since we are in the regime where $(n+1)\gamma < 1$ (here we use that $\gamma^{-1} \notin \mathbb{N}$) we obtain that

$$\sup_{m \geq 0} a_m \leq \frac{C}{2 - 2^{(n+1)\gamma}}.$$

Therefore

$$\left\| Z_{t_k^m, t_{k+1}^m}^m \right\|_{n+1} \lesssim 2^{-m(n+1)\gamma}, \quad \forall m \geq 0, k = 0, \dots, 2^m - 1. \quad (3.7)$$

Let now fix $m \geq 0$, $i \in \{0, \dots, 2^m - 1\}$, and set $s := t_j^m$, $t := t_{j+1}^m$. Then we want to prove that $|Y_{st}| \lesssim 2^{-m\gamma}$, see (2.15) for the definition of $|\cdot|$. By subadditivity of $|\cdot|$ w.r.t. the convolution product \star we have

$$|Y_{st}| \leq |\exp_{n+1}(L_{st})| + |\exp_{n+1}(Z_{st}^m)|.$$

By Lemma 2.15 and (3.7)

$$|\exp_{n+1}(Z_{st}^m)| \lesssim \| \| Z_{t_k^m, t_{k+1}^m}^m \| \|_{n+1}^{\frac{1}{n+1}} \lesssim 2^{-m\gamma}.$$

Moreover, using Lemma 2.15 again (first the upper bound, then the lower bound) and the fact that $\mathbb{X}^n: [0, 1] \rightarrow G^n$ is γ -Hölder by assumption,

$$\begin{aligned} |\exp_{n+1}(L_{st})| &\leq C_{n+1} \sup_{k=1, \dots, n+1} \| \| (L_{st})_k \| \|^{1/k} = C_{n+1} \sup_{k=1, \dots, n} \| \| (L_{st})_k \| \|^{1/k} \leq \\ &\leq \frac{C_{n+1}}{c_n} |\exp_n(L_{st})| = \frac{C_{n+1}}{c_n} |X_{st}| = \frac{C_{n+1}}{c_n} \rho_n(\mathbb{X}_{t_j^m}^n, \mathbb{X}_{t_{j+1}^m}^n) \lesssim 2^{-m\gamma}. \end{aligned}$$

Therefore, the path $\mathbb{X}^{n+1}: D \rightarrow G^{n+1}$ defined by $\mathbb{X}_{t_j^m}^{n+1} := Y_{0, t_j^m}$ satisfies

$$\rho_{n+1}(\mathbb{X}_{t_j^m}^{n+1}, \mathbb{X}_{t_{j+1}^m}^{n+1}) \lesssim 2^{-m\gamma},$$

thus by Lemma 3.5 we obtain a γ -Hölder path $\mathbb{X}^{n+1}: [0, 1] \rightarrow G^{n+1}$ extending \mathbb{X}^n . □

Remark 3.6. Our construction depends on a finite number of choices, namely we set $Z_{01} = 0$ to start the recursion in (3.6), and this for each level; moreover in (3.6) we make the choice $Z_{t_{2k}^m, t_{2k+1}^m}^m = Z_{t_{2k+1}^m, t_{2k+2}^m}^m$. These choices are the same as in [30, Proof of Theorem 1] and are indeed the most natural ones, but one could change them and the final outcome would be different.

Remark 3.7. While in [30, Proof of Proposition 6] Lyons and Victoir use the axiom of choice, our proof is completely constructive. In particular, we use the explicit map $\exp_{k+1} \circ \log_k: G^k(\mathcal{T}_n) \rightarrow G^{k+1}(\mathcal{T}_n)$ which plays the role of the injection $i_{G/K, G}: G/K \rightarrow G$ in [30, Proposition 6]. The fact that this map has good continuity estimates is based on Lemma 2.15.

Corollary 3.8. *Given $\gamma \in]0, 1[$ with $\gamma^{-1} \notin \mathbb{N}$ and a collection of γ -Hölder paths $x^i: [0, 1] \rightarrow \mathbb{R}$, $i = 1, \dots, d$, there exists a γ -Hölder path $\mathbb{X}: [0, 1] \rightarrow G^N$ such that $\langle \mathbb{X}, e_i \rangle = x^i - x_0^i$, $i = 1, \dots, d$. Then $X_{st} := \mathbb{X}_s^{-1} \star \mathbb{X}_t$ defines a (\mathcal{H}, γ) -rough path over (x^1, \dots, x^d) .*

Proof. We start with the following observation: for $n = 1$, the group $G^1 \subset \mathcal{H}_{(1)}^*$ is abelian, and isomorphic to the additive group $\mathcal{H}_{(1)}^*$. Indeed, let $X, Y \in G^1$ and $x \in \mathcal{H}_{(1)}$. Then, as $\Delta x = x \otimes 1 + 1 \otimes x$ by the grading, we have that

$$\langle X \star Y, x \rangle = \langle X, x \rangle + \langle Y, x \rangle,$$

that is, $X \star Y = X + Y$. Moreover, in \mathcal{H}_1 the product $xy = 0$. Therefore, we may set $\langle \mathbb{X}_t^1, e_i \rangle := x_t^i - x_0^i$ where $\{e_1, \dots, e_d\}$ is a basis of $\mathcal{H}_{(1)}$ and this path is γ -Hölder with respect to ρ_1 .

By Theorem 3.4 there is a γ -Hölder path $\mathbb{X}^2: [0, 1] \rightarrow (G^2, \rho_2)$ extending \mathbb{X}^1 so in particular $\langle \mathbb{X}_t^2, e_i \rangle = x_t^i - x_0^i$ also. Continuing in this way we obtain successive γ -Hölder extensions $\mathbb{X}^3, \dots, \mathbb{X}^N$ and we set $\mathbb{X} := \mathbb{X}^N$. □

The following result has already been proved in the case where the underlying Hopf algebra \mathcal{H} is *combinatorial* by Curry, Ebrahimi-Fard, Manchon and Munthe-Kaas in [14, Theorem 4.3]. We remark that their proof works without modifications in our context so we have

Theorem 3.9. *Let $\mathbb{X}: [0, 1] \rightarrow G^N$ be a γ -Hölder path with $\mathbb{X}_0 = 1$ and suppose that $\gamma^{-1} \notin \mathbb{N}$. There exists a path $\hat{\mathbb{X}}: [0, 1] \rightarrow G$ such that $|\langle \hat{\mathbb{X}}_s^{-1} \star \hat{\mathbb{X}}_t, v \rangle| \lesssim |t - s|^{\gamma|v|}$ for all homogeneous $v \in \mathcal{H}$ and extending \mathbb{X} , in the sense that $\hat{\mathbb{X}}|_{\mathcal{H}_N} = \mathbb{X}$.*

Remark 3.10. In view of Theorem 3.9 we can replace the truncated group in Definition 3.1 by the full group of characters G . What this means is that γ -rough paths are uniquely defined once we fix the first N levels and since \mathcal{H} is locally finite, this amounts to a finite number of choices. This is of course a generalization of the extension theorem of [29], see also [25, Theorem 7.3] for the branched case.

4 Applications

We now apply Theorem 3.4 to various kinds of Hopf algebras in order to link this result with the contexts already existing in the literature.

4.1 Geometric rough paths

In this setting we fix a finite alphabet $A := \{1, \dots, d\}$. As a vector space $\mathcal{H} := T(A)$ is the linear span of the *free monoid* $M(A)$ generated by A . The product on \mathcal{H} is the shuffle product $\sqcup: \mathcal{H} \otimes \mathcal{H} \rightarrow \mathcal{H}$ defined recursively by $1 \sqcup v = v \sqcup 1 = v$ for all $v \in \mathcal{H}$, where $1 \in M(A)$ is the unit for the monoid operation, and

$$(au \sqcup bv) = a(u \sqcup bv) + b(au \sqcup v)$$

for all $u, v \in \mathcal{H}$ and $a, b \in A$, where au and bv denote the product of the letters a, b with the words u, v in $M(A)$.

The coproduct $\bar{\Delta}: \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{H}$ is obtained by *deconcatenation* of words,

$$\bar{\Delta}(a_1 \cdots a_n) = a_1 \cdots a_n \otimes 1 + 1 \otimes a_1 \cdots a_n + \sum_{k=1}^{n-1} a_1 \cdots a_k \otimes a_{k+1} \cdots a_n.$$

It turns out that $(\mathcal{H}, \sqcup, \bar{\Delta})$ is a commutative unital Hopf algebra, and $(\mathcal{H}, \bar{\Delta})$ is the cofree coalgebra over the linear span of A . The antipode is the linear map $S: \mathcal{H} \rightarrow \mathcal{H}$ given by

$$S(a_1 \cdots a_n) = (-1)^n a_n \cdots a_1.$$

Finally, we recall that \mathcal{H} is graded by the length $\ell(a_1 \cdots a_n) = n$ and it is also connected. The homogeneous components $\mathcal{H}_{(n)}$ are spanned by the sets $\{a_1 \cdots a_n : a_i \in A\}$.

Definition 3.1 specializes in this case to *geometric rough paths* (GRP) as defined in [28] (see just below for the precise definition) and Theorem 3.4 coincides with [30, Theorem 6].

Definition 4.1. Let $\gamma \in]0, 1[$ and set $N := \lfloor \gamma^{-1} \rfloor$. A geometric γ -rough path is a map $X: [0, 1]^2 \rightarrow G^N$ which satisfies Chen's rule

$$X_{st} = X_{su} \star X_{ut}$$

for all $s, u, t \in [0, 1]$ and the analytic bound $|\langle X_{st}, v \rangle| \lesssim |t - s|^{\gamma \ell(v)}$ for all $v \in \mathcal{H}_N$.

Then Proposition 3.3 and the existence results Theorem 3.4-Corollary 3.8 are the content of the paper [30] by Lyons and Victoir.

4.2 Branched rough paths

Let \mathcal{T} be the collection of all non-planar non-empty rooted trees with nodes decorated by $\{1, \dots, d\}$. Elements of \mathcal{T} are written as 2-tuples $\tau = (T, c)$ where T is a non-planar tree with node set N_τ and edge set E_τ , and $c: N_\tau \rightarrow \{1, \dots, d\}$ is a function. Edges in E_τ are oriented away from the root, but this is not reflected in our graphical representation. Examples of elements of \mathcal{T} include the following

$$\bullet_i, \quad \begin{array}{c} \bullet_j \\ | \\ \bullet_i \end{array}, \quad \begin{array}{c} \bullet_j^k \\ / \quad \backslash \\ \bullet_i \end{array}, \quad \begin{array}{c} \bullet_j^k \quad \bullet_l^m \\ / \quad \backslash \\ \bullet_i \end{array}.$$

For $\tau \in \mathcal{T}$ write $|\tau| = \#N_\tau$ for its number of nodes. Also, given an edge $e = (x, y) \in E_\tau$ we set $s(e) = x$ and $t(e) = y$. There is a natural partial order relation on N_τ where $x \leq y$ if and only if there is a path in T from the root to y containing x .

We denote by \mathcal{F} the collection of decorated rooted forests and we let $\mathcal{H} := \mathcal{H}_{\text{BCK}}$ denote the vector space spanned by \mathcal{F} . There is a natural commutative and associative product on \mathcal{F} , denoted by \cdot and given by the disjoint union of forests, where the empty forest 1 acts as the unit. Then, \mathcal{H} is the free commutative algebra over \mathcal{T} , with grading $|\tau_1 \cdots \tau_k| = |\tau_1| + \cdots + |\tau_k|$. Given $i \in \{1, \dots, d\}$ and a forest $\tau = \tau_1 \cdots \tau_k$ we denote by $[\tau_1 \cdots \tau_k]_i$ the tree obtained by grafting each of the trees τ_1, \dots, τ_k to a new root decorated by i , e.g.

$$[\bullet]_i = \mathfrak{!}_i^j, \quad [\bullet \bullet \bullet]_i = \mathfrak{!}_i^{j^k}.$$

The decorated Butcher–Connes–Kreimer coproduct [12, 25] is the unique algebra morphism $\Delta : \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{H}$ such that

$$\Delta[\tau]_i = [\tau]_i \otimes 1 + (\text{id} \otimes [\cdot]_i)\Delta\tau.$$

This coproduct admits a representation in terms of *cuts*. An *admissible cut* C of a tree T is a non-empty subset of E_T such that any path from any vertex of the tree to the root contains at most one edge from C ; we denote by $\mathfrak{A}(T)$ the set of all admissible cuts of the tree T . Any admissible cut C containing k edges maps a tree T to a forest $C(T) = T_1 \cdots T_{k+1}$ obtained by removing each of the edges in C . Observe that only one of the remaining trees T_1, \dots, T_{k+1} contains the root of T , which we denote by $R^C(T)$; the forest formed by the other k factors is denoted by $P^C(T)$. This naturally induces a map on decorated trees by considering cuts of the underlying tree, and restriction of the decoration map to each of the rooted subtrees T_1, \dots, T_{k+1} . Then,

$$\Delta\tau = \tau \otimes 1 + 1 \otimes \tau + \sum_{C \in \mathfrak{A}(\tau)} P^C(\tau) \otimes R^C(\tau). \tag{4.1}$$

This, together with the counit map $\varepsilon : \mathcal{F} \rightarrow \mathbb{R}$ such that $\varepsilon(\tau) = 1$ if and only if $\tau = 1$ endows \mathcal{F} with a connected graded commutative non-cocommutative bialgebra structure, hence a Hopf algebra structure [31].

As before we denote by \mathcal{H}^* the linear dual of \mathcal{H} which is an algebra via the convolution product $\langle X \star Y, \tau \rangle = \langle X \otimes Y, \Delta\tau \rangle$ and we denote by G the set of characters on \mathcal{H} , that is, linear functionals $X \in \mathcal{H}^*$ such that $\langle X, \sigma \cdot \tau \rangle = \langle X, \sigma \rangle \langle X, \tau \rangle$. For each $n \in \mathbb{N}$ the finite-dimensional vector space \mathcal{H}_n spanned by the set \mathcal{F}_n of forests with at most n nodes is a subcoalgebra of \mathcal{H} , hence its dual is an algebra under the convolution product, and we let G_n be the set of characters on \mathcal{H}_n .

We have already defined branched rough paths in Definition 1.1. Proposition 3.3 yields the following characterization

Proposition 4.2. *A path $X : [0, 1]^2 \rightarrow G^N$ is a branched rough path if and only if $\mathbb{X}_t := X_{0t}$ is γ -Hölder path with respect to the distance ρ_N defined in (2.16).*

Directly applying Theorem 3.4 to the Butcher–Connes–Kreimer Hopf algebra \mathcal{H} we obtain

Corollary 4.3. *Given $\gamma \in]0, 1[$ with $\gamma^{-1} \notin \mathbb{N}$ and a family of γ -Hölder paths $(x^i : i = 1, \dots, d)$, there exists a branched rough path X above $(x^i : i = 1, \dots, d)$, i.e. $X : [0, 1]^2 \rightarrow G^N$ is such that $\langle X_{st}, \bullet^i \rangle = x_t^i - x_s^i$ for all $i = 1, \dots, d$.*

Remark 4.4. Given the level of generality in which Theorem 3.4 is developed, our results also apply to the case when \mathcal{H} is a combinatorial Hopf algebra as defined in [14]. In particular, we also have a construction theorem for *planarly branched rough paths* [14] which are characters over Munthe-Kaas and Wright’s Hopf algebra of Lie group integrators [32].

4.3 Anisotropic geometric rough paths

We now apply our results to another class of rough paths which we call *anisotropic geometric rough paths* (aGRPs for short). L. Gyurkó introduced a similar concept in [26], which he called Π -rough paths; unlike us, he uses a “primal” presentation, i.e. paths taking values in the tensor algebra $T(\mathbb{R}^d)$, and p -variation norms rather than Hölder norms. Geometric rough paths over an inhomogeneous (or anisotropic) set of paths can be traced back to Lyons’ original paper [29].

As in the geometric case, see Section 4.1, fix a finite alphabet $A = \{1, \dots, d\}$ and denote by $M(A)$ the free monoid generated by A . We denote again by $\mathcal{H} := T(A)$ the shuffle Hopf algebra over the alphabet A .

Let $(\gamma_a : a \in A)$ be a sequence of real numbers such that $0 < \gamma_a < 1$ for all a , and let $\hat{\gamma} = \min_{a \in A} \gamma_a$. For a word $v = a_1 \cdots a_k \in M(A)$ of length k define

$$\omega(v) := \gamma_{a_1} + \dots + \gamma_{a_k}$$

and observe that ω is additive in the sense that $\omega(uv) = \omega(u) + \omega(v)$ for each pair of words $u, v \in M(A)$. The set

$$\mathfrak{L} := \{v \in M(A) : \omega(v) \leq 1\}$$

is finite; if $\hat{N} := \lfloor \hat{\gamma}^{-1} \rfloor$ then $\mathfrak{L} \subset \mathcal{H}_{\hat{N}}$. In analogy with Lemma 2.5, the additivity of ω implies

Lemma 4.5. *The subspace $\mathcal{H}_a \subset \mathcal{H}_{\hat{N}}$ spanned by \mathfrak{L} is a subcoalgebra of $(\mathcal{H}, \bar{\Delta}, \varepsilon)$.*

Consequently, we will consider the dual algebra $(\mathcal{H}_a^*, \star, \varepsilon)$. In this case, we define \mathfrak{g}_a to be the space of truncated infinitesimal characters on \mathcal{H}_a , namely the linear functionals $\alpha \in \mathcal{H}_a^*$ such that

$$\langle \alpha, x \sqcup y \rangle = \langle \alpha, x \rangle \langle \varepsilon, y \rangle + \langle \varepsilon, x \rangle \langle \alpha, y \rangle$$

for all $x, y \in \mathcal{H}_a$ such that $x \sqcup y \in \mathcal{H}_a$, and let $G_a := \{X = \exp_{\hat{N}}(\alpha)|_{\mathcal{H}_a} : \alpha \in \mathfrak{g}_a\}$. As before, there is a canonical injection $\mathcal{H}_a^* \hookrightarrow \mathcal{H}^*$ so we suppose that $\langle X, v \rangle = 0$ for all $X \in \mathcal{H}^*$ and $v \notin \mathfrak{L}$.

For each $\lambda > 0$ there is a unique coalgebra automorphism $\Omega_\lambda : \mathcal{H} \rightarrow \mathcal{H}$ such that $\Omega_\lambda a = \lambda^{\gamma_a/\hat{\gamma}} a$ for all $a \in A$. We also define $\|\cdot\| : G_a \rightarrow \mathbb{R}$,

$$\|X\| := \max_{v \in \mathfrak{L}} |\langle X, v \rangle|^{\hat{\gamma}/\omega(v)}. \tag{4.2}$$

As at the end of Section 2, $(\Omega_\lambda)_{\lambda>0}$ is a one-parameter family of Lie-algebra automorphisms of \mathfrak{g}_a and $\|\Omega_\lambda X\| = \lambda \|X\|$ for all $\lambda > 0$ and $X \in G_a$, namely $\|\cdot\|$ is a homogeneous norm on G_a . However, unlike $\|\cdot\|$ this norm is not subadditive and it therefore does not define a distance on G_a .

4.3.1 Signatures

In order to construct an appropriate metric on G_a we consider *signatures* of smooth paths. We observe that $A \subset \mathfrak{L}$. Let $x = (x^a : a \in A)$ be a collection of (piecewise) smooth paths, and define a map $S(x) : [0, 1]^2 \rightarrow \mathcal{H}^*$ by

$$\langle S(x)_{st}, v \rangle := \int_s^t dx_{s_k}^{v_k} \int_s^{s_k} dx_{s_{k-1}}^{v_{k-1}} \cdots \int_s^{s_2} dx_{s_1}^{v_1}.$$

In his seminal work [11], K. T. Chen showed that $S(x)$ is a character of $(T(A), \sqcup)$; in particular, $S(x)_{st}|_{\mathcal{H}_a} \in G_a$.

Consider the metric $d_a(X, Y) = \sum_{a \in A} |\langle X - Y, a \rangle|^{\hat{\gamma}/\gamma_a}$ on $\mathcal{H}_{(1)}^*$, where we recall that $\mathcal{H}_{(1)}$ is the vector space spanned by A . The *anisotropic length* of a smooth curve $\theta: [0, 1] \rightarrow \mathcal{H}_{(1)}^*$ is defined to be its length with respect to this metric and will be denoted by $L_a(\theta)$. Observe that since $d_a(\Omega_\lambda X, \Omega_\lambda Y) = \lambda d_a(X, Y)$ we have that $L_a(\Omega_\lambda \theta) = \lambda L_a(\theta)$.

We now define a homogeneous norm (see the end of Section 2) $|\cdot|_{CC}: G_a \rightarrow \mathbb{R}_+$, called the *anisotropic Carnot–Carathéodory norm*, by setting

$$|X|_{CC} := \inf\{L_a(x) : x^a \in C^\infty, S(x)_{01} = X\}.$$

Since curve length is invariant under reparametrization in any metric space we obtain, as in [23, Section 7.5.4]:

Proposition 4.6. *The infimum defining the anisotropic Carnot–Carathéodory norm is finite and attained at some minimizing path \hat{x} .*

Proposition 4.7. *The anisotropic Carnot–Carathéodory norm is homogeneous, that is, $|\Omega_\lambda X|_{CC} = \lambda |X|_{CC}$.*

Proof. Let \hat{x} be the curve such that $|X|_{CC} = L_a(\hat{x})$. For any $\lambda > 0$ and word $v \in \mathfrak{L}$ we have

$$\langle S(\Omega_\lambda \hat{x})_{01}, v \rangle = \lambda^{\omega(v)/\hat{\gamma}} \langle S(\hat{x})_{01}, v \rangle = \langle \Omega_\lambda S(\hat{x})_{01}, v \rangle = \langle \Omega_\lambda X, v \rangle,$$

thus $|\Omega_\lambda X|_{CC} \leq L_a(\Omega_\lambda \hat{x}) = \lambda L_a(\hat{x}) = \lambda |X|_{CC}$. The reverse inequality is obtained by noting that $X = (\Omega_{\lambda^{-1}} \circ \Omega_\lambda) X$. \square

The anisotropic Carnot–Carathéodory norm can also be seen to satisfy $|X|_{CC} = |X^{-1}|_{CC}$ and $|X \star Y|_{CC} \leq |X|_{CC} + |Y|_{CC}$ for all $X, Y \in G_a$, see e.g. the proof of [23, Proposition 7.40]; hence it induces a left-invariant metric $\rho_a(X, Y) := |X^{-1} \star Y|_{CC}$ on G_a . Moreover, arguing as in the proof of [23, Theorem 7.44] we see that there exist positive constants c, C such that

$$c|X|_{CC} \leq \|X\| \leq C|X|_{CC}, \quad \forall X \in G_a. \quad (4.3)$$

Definition 4.8. *An anisotropic geometric γ -rough path, with $\gamma = (\gamma_a, a \in A)$, is a map $X: [0, 1]^2 \rightarrow G_a$ which satisfies*

- 1 the Chen rule $X_{su} \star X_{ut} = X_{st}$ for all $(s, u, t) \in [0, 1]^3$,
- 2 the bound $|\langle X_{st}, v \rangle| \lesssim |t - s|^{\omega(v)}$ for all $v \in \mathfrak{L}$.

Proposition 4.9. *Anisotropic geometric γ -rough paths are in one-to-one correspondence with $\hat{\gamma}$ -Hölder paths $\mathbb{X}: [0, 1] \rightarrow (G_a, \rho_a)$ with $\mathbb{X}_0 = 1$.*

Proof. Let X be an anisotropic geometric γ -rough path and v a word. By definition we have that $|\langle X_{st}, v \rangle| \lesssim |t - s|^{\omega(v)}$, hence $\|X_{st}\| \lesssim |t - s|^{\hat{\gamma}}$. The equivalence between $\|\cdot\|$ and $|\cdot|_{CC}$ of (4.3) implies that $\rho_a(\mathbb{X}_s, \mathbb{X}_t) = |X_{st}|_{CC} \lesssim |t - s|^{\hat{\gamma}}$, hence $t \mapsto \mathbb{X}_t$ is $\hat{\gamma}$ -Hölder with respect to ρ_a . The other direction follows in a similar manner. \square

Theorem 3.4 also applies to this situation, and we obtain the following

Corollary 4.10. *Let $(\gamma_a : a \in A)$ be real numbers in $]0, 1[$ such that $1 \notin \sum_{a \in A} \gamma_a \mathbb{N}$. Let $(x^a : a \in A)$ be a collection of real-valued paths such that x^a is γ_a -Hölder. Then there exists an anisotropic geometric γ -rough path X such that $\langle X_{st}, a \rangle = x_t^a - x_s^a$ for all $a \in A$.*

Proof. We start by constructing the *homogeneous* geometric rough path X given by the $\hat{\gamma}$ -Hölder path $\mathbb{X}: [0, 1] \rightarrow G^{\hat{N}}$ of Corollary 3.8. Then we restrict \mathbb{X} to $\mathcal{H}_a \subset \mathcal{H}_{\hat{N}}$ and we show that on this space it satisfies the stronger bound $|\langle X_{st}, v \rangle| \lesssim |t - s|^{\omega(v)}$ for all $v \in \mathcal{L}$.

Recalling the proof of Theorem 3.4, we consider $v \in \mathcal{H}_n \cap \mathcal{H}_a$, and we proceed by recurrence on n . For $n = 0$ there is nothing to prove. Suppose we have proved the result for n and let $v \in \mathcal{H}_{n+1} \cap \mathcal{H}_a$. In this case

$$\begin{aligned} \langle X_{st}^{n+1}, v \rangle &= \langle \exp_{n+1}(L_{st} + Z_{st}), v \rangle = \sum_{i=0}^{n+1} \frac{1}{i!} \langle (L_{st})^{i*} \star (\varepsilon + Z_{st}), v \rangle \\ &= \sum_{i=0}^{n+1} \frac{1}{i!} \langle (L_{st})^{i*}, v \rangle + \langle Z_{st}, v \rangle = \langle X_{st}^n, v \rangle + \frac{1}{(n+1)!} \langle (L_{st})^{(n+1)*}, v \rangle + \langle Z_{st}, v \rangle. \end{aligned}$$

We want to prove now that

$$\left| \langle X_{t_k^m t_{k+1}^m}^{n+1}, v \rangle \right| \lesssim 2^{-m\omega(v)}, \quad \forall m \geq 0, k = 0, \dots, 2^m - 1. \quad (4.4)$$

For $m \geq 0$ set

$$b_m := 2^{m\omega(v)} \max_{k=0, \dots, 2^m - 1} \left| \langle Z_{t_k^m t_{k+1}^m}^m, v \rangle \right|.$$

Then, for $s = t_k^m$, $u = t_{k+1}^m$ and $t = t_{k+2}^m$ and $v = v_1 \cdots v_{n+1}$

$$\left| \langle \text{BCH}_{(n+1)}(L_{su}, L_{ut}), v \rangle \right| \leq \sum_{i+j=n+1} \frac{1}{i!j!} \sum_{\sigma \in \mathcal{S}_{n+1}} |a_\sigma| \prod_{p=1}^i |\langle L_{su}, v_{\sigma^{-1}(p)} \rangle| \prod_{q=i+1}^{n+1} |\langle L_{ut}, v_{\sigma^{-1}(q)} \rangle|.$$

Now, since $v_j \in \mathcal{H}_{(1)}$ for all $j = 1, \dots, n+1$ we actually have that by the assumption $x^a \in C^{\gamma_a}$

$$|\langle L_{su}, a \rangle| = |x_u^a - x_s^a| \lesssim 2^{-m\gamma_a}$$

and we have a similar estimate for L_{ut} instead of L_{su} . Therefore we obtain that

$$\left| \langle \text{BCH}_{(n+1)}(L_{su}, L_{ut}), v \rangle \right| \lesssim 2^{-m\omega(v)}.$$

Therefore, from (3.5) we get

$$b_m \leq 2^{m(\omega(v)-1)} b_{m-1} + C, \quad m \geq 1,$$

hence since $b_0 = 0$ we can show by recurrence on $m \geq 0$

$$b_m \leq C \sum_{j=0}^{m-1} 2^{-j(1-\omega(v))}.$$

Since we are in the regime where $\omega(v) < 1$ (here we use that $1 \notin \sum_{a \in A} \gamma_a \mathbb{N}$) we obtain that

$$\sup_{m \geq 0} b_m \leq \frac{C}{1 - 2^{\omega(v)-1}}.$$

Therefore

$$\left| \langle Z_{t_k^m t_{k+1}^m}^m, v \rangle \right| \lesssim 2^{-m\omega(v)}, \quad m \geq 0, k = 0, \dots, 2^m - 1.$$

Analogously, since $L_{st} \in \mathfrak{g}$, arguing as in (2.14) we have

$$\langle (L_{st})^{(n+1)*}, v \rangle = \prod_{i=1}^{n+1} \langle L_{st}, v_i \rangle = \prod_{i=1}^{n+1} (x_t^{v_i} - x_s^{v_i}) \implies |\langle (L_{st})^{(n+1)*}, v \rangle| \lesssim 2^{-m\omega(v)}$$

and (4.4) is proved. This implies that $\|X_{t_k^m t_{k+1}^m}^{n+1}\| \lesssim 2^{-m\hat{\gamma}}$ and by equivalence of homogeneous norms (4.3) we obtain

$$\rho_a \left(\mathbb{X}_{t_k^m}^{n+1}, \mathbb{X}_{t_{k+1}^m}^{n+1} \right) \lesssim 2^{-m\hat{\gamma}}.$$

Then we can use Lemma 3.5 and obtain that the path \mathbb{X}^{n+1} constructed in the proof of Theorem 3.4 is in fact $\hat{\gamma}$ -Hölder path with values in G_a . □

5 The Hairer-Kelly construction

In this section we develop further results specifically for branched rough paths as introduced in Section 4.2 by using our general results from Section 3. We analyze in detail the Hairer-Kelly map introduced in [28], which plays a very important role in our construction, and we use it to prove Theorem 1.2 and Corollary 1.3.

5.1 The Hairer–Kelly map

Recall that \mathcal{T} denotes the set of all decorated rooted trees, \mathcal{F} denotes the collection of all decorated rooted forests, and \mathcal{H}_{BCK} is the Butcher–Connes–Kreimer Hopf algebra. As in Section 4.2, Δ denotes the Connes–Kreimer coproduct on \mathcal{H}_{BCK} . For each $n \in \mathbb{N}$, $n \geq 1$, we denote by \mathcal{T}_n the set of (non-empty) trees with at most n vertices.

Recall also from Section 4.1 that given an alphabet A we denote by $T(A)$ the shuffle Hopf algebra generated by A , and that $\bar{\Delta}$ denotes the deconcatenation coproduct on it. We fix $N \in \mathbb{N}$ and we consider the shuffle Hopf algebras $T(\mathcal{T})$ and $T(\mathcal{T}_N)$, namely we choose as letters of our alphabet the (non-empty) decorated rooted trees (respectively rooted trees with at most N vertices). Note that we can identify every non-empty tree $\tau \in \mathcal{T}$ with the word in $T(\mathcal{T})$ composed by the single letter τ . We also remark that, in order to avoid confusion with the forest product on \mathcal{H}_{BCK} we denote the concatenation of letters in $T(\mathcal{T})$ by a tensor symbol.

We note that $T(\mathcal{T})$ and $T(\mathcal{T}_N)$ admit two different natural gradings, both of which make them locally finite graded Hopf-algebras. One grading, as in Section 4.1, is given by the number of letters (trees) of each word, namely the degree of $v = \tau_1 \otimes \cdots \otimes \tau_k$ is k . The other grading is given by the sum of the number of nodes of each letter (tree), namely the degree of $v = \tau_1 \otimes \cdots \otimes \tau_k$ is $|\tau_1| + \cdots + |\tau_k|$, where we recall that forests and trees are graded in \mathcal{H}_{BCK} by the number of nodes, with the notation $|\tau| = \#N_\tau$. We remark the latter grading is always greater or equal to the former. As an example, take $v = \bullet_i \otimes \uparrow_j^k$; then, as a word v has length 2 but the total number of nodes is 3.

We recall the following result from [28, Lemma 4.9].

Lemma 5.1. *We grade $T(\mathcal{T})$ according to the number of nodes. Then there exists a graded morphism of Hopf algebras $\psi: \mathcal{H}_{\text{BCK}} \rightarrow T(\mathcal{T})$ satisfying $\psi(\tau) = \tau + \psi_{n-1}(\tau)$ for all $\tau \in \mathcal{T}_n$, where ψ_{n-1} denotes the projection of ψ onto $T(\mathcal{T}_{n-1})$.*

We call ψ the Hairer-Kelly map. Since ψ is graded, for any forest $\tau \in \mathcal{F}$ the image $\psi(\tau)$ is a sum of words of the form $\tau_1 \otimes \cdots \otimes \tau_k$ where all terms satisfy $|\tau_1| + \cdots + |\tau_k| = |\tau|$. Observe that since ψ is a Hopf algebra morphism, in particular a coalgebra morphism, then

$$(\psi \otimes \psi)\Delta'\tau = \bar{\Delta}'\psi(\tau) = \bar{\Delta}'\psi_{n-1}(\tau), \quad \tau \in \mathcal{T}_n,$$

since trees are primitive elements in $T(\mathcal{T})$, being single-letter words. From the proof of [28, Lemma 4.9] we are able to see that in fact ψ_{n-1} is given by the recursion $\psi_{n-1} = m_{\otimes}(\psi \otimes \text{id})\Delta'$ on the linear span of \mathcal{T}_n , see also [3, Definition 1, section 6].

Example 5.2. Here are some examples of the action of ψ on some trees:

$$\begin{aligned} \psi(\bullet_i) &= \bullet_i, & \psi(\bullet_{aa}) &= \psi(\bullet_a) \sqcup \psi(\bullet_b) = \bullet_a \otimes \bullet_b + \bullet_b \otimes \bullet_a, & \psi(\mathfrak{V}_a^b) &= \mathfrak{V}_a^b + \bullet_b \otimes \bullet_a \\ \psi\left(\mathfrak{V}_a^d\left(\mathfrak{V}_a^c\left(\mathfrak{V}_a^b\right)\right)\right) &= \mathfrak{V}_a^d\left(\mathfrak{V}_a^c\left(\mathfrak{V}_a^b\right)\right) + \bullet_b \otimes \mathfrak{V}_a^d\left(\mathfrak{V}_a^c\right) + \bullet_c \otimes \mathfrak{V}_a^d\left(\mathfrak{V}_a^b\right) + \mathfrak{V}_a^d\left(\mathfrak{V}_a^c\right) \otimes \mathfrak{V}_a^b + \bullet_c \otimes \bullet_b \otimes \mathfrak{V}_a^d + \bullet_c \otimes \bullet_b \otimes \mathfrak{V}_a^c + \bullet_b \otimes \bullet_c \otimes \mathfrak{V}_a^d \\ &+ \mathfrak{V}_a^d \otimes \bullet_b \otimes \bullet_a + \bullet_b \otimes \mathfrak{V}_a^d \otimes \bullet_a + \bullet_c \otimes \bullet_c \otimes \bullet_b \otimes \bullet_a + \bullet_c \otimes \bullet_b \otimes \bullet_c \otimes \bullet_d \\ &+ \bullet_b \otimes \bullet_d \otimes \bullet_c \otimes \bullet_a. \end{aligned}$$

5.2 A special class of anisotropic geometric rough paths

We have already discussed anisotropic geometric rough paths (aGRPs) in Section 4.3. For the Hairer-Kelly construction we need a very particular subclass of aGRPs, where the base paths $(x^a)_{a \in A}$ are such that each x^a is γ_a -Hölder and there exists $\gamma \in]0, 1[$ and $(k_a)_{a \in A} \subset \mathbb{N}$ such that $\gamma_a = k_a \gamma$; therefore the Hölder exponents are all integer multiples of a fixed exponent γ .

We may of course apply the extension result of Corollary 4.10, but it turns out that in this setting we can avoid using the Carnot-Carathéodory distance and rather use a more explicit metric, which is a simple generalization of the homogeneous case (2.16).

We have already seen that the space $\mathcal{H} := T(\mathcal{T}_N)$ can be graded in two ways. We can even define a *bigrading* on this space: for $1 \leq n \leq N$ and $n \leq j \leq nN$, we define the space $\mathcal{H}_{(n,j)}$ as the linear span of the words $\tau_1 \otimes \cdots \otimes \tau_n \in T(\mathcal{T}_N)$ such that $|\tau_1| + \cdots + |\tau_n| = j$. Then, in analogy with (2.2), we have

$$\sqcup: \mathcal{H}_{(n,j)} \otimes \mathcal{H}_{(m,h)} \rightarrow \mathcal{H}_{(n+m,j+h)}, \quad \bar{\Delta}: \mathcal{H}_{(n,j)} \rightarrow \bigoplus_{p=0, \dots, n, q=1, \dots, j-1} \mathcal{H}_{(p,q)} \otimes \mathcal{H}_{(n-p,j-q)}.$$

Then, recalling that $\mathcal{H}_0 = \mathbb{R}1$, we set

$$\mathcal{H}_{N,N} := \mathcal{H}_0 \oplus \bigoplus_{n=1}^N \bigoplus_{j=n}^N \mathcal{H}_{(n,j)}.$$

In other words, $\mathcal{H}_{N,N}$ is the linear span of all words $\tau_1 \otimes \cdots \otimes \tau_n$ with $n \leq N$ and $|\tau_1| + \cdots + |\tau_n| \leq N$. Therefore, analogously to (2.3) and (2.5), we have decompositions

$$\mathcal{H}_{N,N} \ni x = x_0 + \sum_{n=1}^N \sum_{j=n}^N x_{n,j}, \quad \mathcal{H}_{N,N}^* \ni \alpha = \alpha_{(0)} + \sum_{n=1}^N \sum_{j=n}^N \alpha_{(n,j)}, \quad \alpha_{(n,j)}(x) = \alpha(x_{n,j}).$$

We define now $\mathfrak{g}^{N,N}$ as the space of truncated characters on $\mathcal{H}_{N,N}$, namely of all linear $\alpha : \mathcal{H}_{N,N} \rightarrow \mathbb{R}$ such that

$$\langle \alpha, x \sqcup y \rangle = \langle \alpha, x \rangle \langle \varepsilon, y \rangle + \langle \varepsilon, x \rangle \langle \alpha, y \rangle$$

for all $x, y \in \mathcal{H}_{N,N}$ such that $x \sqcup y \in \mathcal{H}_{N,N}$. Moreover we define $G^{N,N} := \exp_N(\mathfrak{g}^{N,N}) \subset \mathcal{H}_N^*$. Then we set in analogy with (2.17) for $X \in G^{N,N}$

$$|X| := N \max_{n=1,\dots,N} \left(\max_{j=n,\dots,N} (j! \|X_{(n,j)}\|)^{1/j} + \max_{j=n,\dots,N} (j! \|(X^{-1})_{(n,j)}\|)^{1/j} \right), \quad (5.1)$$

and we can see that

Lemma 5.3. *The map $G^{N,N} \times G^{N,N} \ni (X, Y) \mapsto \rho^{N,N}(X, Y) := |X^{-1} \star Y| \in \mathbb{R}$ defines a distance on $G^{N,N}$.*

Proof. We only have to check the triangular inequality, which is equivalent to the sub-additivity property $|X \star Y| \leq |X| + |Y|$ for all $X, Y \in G^{N,N}$. Arguing as in the proof of Proposition 2.14

$$\begin{aligned} \| (X \star Y)_{(n,j)} \| &\leq \sum_{m=0}^n \sum_{i=1}^{j-1} \| X_{(m,j)} \| \| Y_{(n-m,j-i)} \| \\ &\leq N \frac{1}{j!} \frac{1}{N^2} \sum_{i=0}^j \binom{j}{i} |X|^i |Y|^{j-i} = \frac{1}{N j!} (|X| + |Y|)^j \end{aligned}$$

whence the result. □

Let $\gamma \in]0, 1[$ and $N := \lfloor \gamma^{-1} \rfloor$. In accordance with Definition 4.8, an *anisotropic geometric γ -rough path* in this setting is a map $X : [0, 1]^2 \rightarrow G^{N,N}$ which satisfies

- 1 the Chen rule $X_{su} \star X_{ut} = X_{st}$ for all $(s, u, t) \in [0, 1]^3$,
- 2 $|\langle X_{st}, v \rangle| \lesssim |t - s|^{j\gamma}$ for all $v \in \mathcal{H}_{(n,j)}$ with $1 \leq n \leq N$ and $j \leq N$.

Then, arguing as in Proposition 3.3, it is easy to show that $\mathbb{X} : [0, 1] \rightarrow G^{N,N}$ is γ -Hölder with respect to the metric $\rho^{N,N}$ if and only if $X : [0, 1]^2 \rightarrow G^{N,N}$, defined as $X_{st} := \mathbb{X}_s^{-1} \star \mathbb{X}_t$, is an anisotropic geometric γ -rough path with $\gamma_v = j\gamma$ for $v = \tau_1 \otimes \dots \otimes \tau_n$ with $n \leq N$ and $|\tau_1| + \dots + |\tau_n| = j \leq N$.

The next result is the analog of Corollary 4.10 in this setting. The proof is the same, with one exception: we can use the explicit norm (5.1) rather than the Carnot-Carathéodory norm $|\cdot|_{CC}$ and we do not need the equivalence of norms result (4.3).

Proposition 5.4. *Given $\gamma \in]0, 1[$ with $\gamma^{-1} \notin \mathbb{N}$ and a collection of paths $x^\tau : [0, 1] \rightarrow \mathbb{R}$, $\tau \in \mathcal{T}_N$, such that $x^\tau \in C^{|\tau|}$, there exists a γ -Hölder path $\mathbb{X} : [0, 1] \rightarrow G^{N,N}$ such that $\langle \mathbb{X}, \tau \rangle = x^\tau$ for all $\tau \in \mathcal{T}_N$.*

Corollary 5.5. *In the setting of Proposition 5.4, let $(g^\tau : \tau \in \mathcal{T}_N)$ be a collection of functions with $g^\tau \in C^{|\tau|}$. Set $\bar{x}_t^\tau = x_t^\tau + g_t^\tau$ and denote by gX the anisotropic geometric γ -rough path constructed in Proposition 5.4 above the path*

$$\bar{x}_t = \sum_{\tau \in \mathcal{T}_N} \bar{x}_t^\tau \tau \in \mathcal{H}_{(1)}, \quad t \in [0, 1].$$

Then, for any two such functions g and g' we have that $g'(gX) = (g + g')X$.

Proof. Let g, g' be two collections of functions as in the statement of the theorem. We have the identity

$$\langle [g'(gX)]_t, \tau \rangle = \langle (gX)_t, \tau \rangle + (g')_t^\tau = x_t^\tau + g_t^\tau + (g')_t^\tau = \langle [(g' + g)X]_t, \tau \rangle.$$

Since both $g'(gX)$ and $(g' + g)X$ are constructed iteratively by adding at each step a function Z satisfying (3.6) on the dyadics, if we let L^n and \bar{L}^n denote the logarithms corresponding to $g'(gX)$ and $(g' + g)X$, Lemma 2.13 and the previous identity imply that

$$\text{BCH}_{n+1}(L_{su}^n, L_{ut}^n) = \text{BCH}_{n+1}(\bar{L}_{su}^n, \bar{L}_{ut}^n)$$

and so $g'(gX) = (g' + g)X$. \square

5.3 Branched rough paths are anisotropic geometric rough paths

The next theorem is almost the same statement as Theorem 4.10 in [28], the only difference being that we construct an *anisotropic* geometric rough path \bar{X} while Hairer-Kelly need only that \bar{X} is geometric in the usual sense (see also [28, Remark 4.14]).

Theorem 5.6. *Let $\gamma \in]0, 1[$ with $\gamma^{-1} \notin \mathbb{N}$, and let X be a branched γ -rough path. There exists an anisotropic geometric rough path $\bar{X}: [0, 1]^2 \rightarrow G^{N,N}$ with exponents $\gamma = (\gamma_\tau = \gamma|\tau|, \tau \in \mathcal{T}_N)$, and such that*

$$\langle X, \tau \rangle = \langle \bar{X}, \psi(\tau) \rangle, \quad \forall \tau \in \mathcal{F}_N.$$

Proof. We construct \bar{X} iteratively as follows. Let $\bar{X}^{(1)}$ be the anisotropic geometric rough path indexed by $\mathcal{T}_1 = \{\bullet_1, \dots, \bullet_d\}$ over the paths $(x_t^i := \langle X_t, \bullet_i \rangle : i = 1, \dots, d)$ with exponents $(\gamma_{\bullet_i} = \gamma)$ given by Proposition 5.4 (alternatively we could have used Theorem 3.4 since all the exponents are equal). This will give us an anisotropic rough path $X: [0, 1]^2 \rightarrow G_a(\mathcal{T}_1)$ with exponents $(\gamma_\tau = \gamma, \tau \in \mathcal{T}_1)$.

Suppose we have constructed anisotropic geometric rough paths $\bar{X}^{(k)}: [0, 1]^2 \rightarrow G_a(\mathcal{T}_k)$ over the paths $(x^\tau : \tau \in \mathcal{T}_k)$ such that $x_t^\tau - x_s^\tau = \langle X_{st}, \tau \rangle - \langle \bar{X}_{st}^{(k-1)}, \psi_{k-1}(\tau) \rangle$ for $k = 1, \dots, n$. This is true for $n = 1$ by the previous paragraph, since $\psi(\bullet_i) = \bullet_i$ for all $i = 1, \dots, d$.

If we let $F_{st}^\tau = \langle X_{st}, \tau \rangle$ and $G_{st}^\tau = \langle \bar{X}_{st}^{(n)}, \psi_n(\tau) \rangle$ for $\tau \in \mathcal{T}_{n+1}$ we have, by Chen's rule, that

$$\delta F_{sut}^\tau = \langle X_{su} \otimes X_{ut}, \Delta' \tau \rangle = \langle \bar{X}_{su}^{(n)} \circ \psi \otimes \bar{X}_{ut}^{(n)} \circ \psi, \Delta' \tau \rangle.$$

Since ψ is in particular a coalgebra morphism between (\mathcal{H}, Δ) and $(T(\mathcal{T}_N), \bar{\Delta})$ we obtain the identity $\delta F_{sut}^\tau = \langle \bar{X}_{su}^{(n)} \otimes \bar{X}_{ut}^{(n)}, \bar{\Delta}' \psi(\tau) \rangle$, which then, by Lemma 5.1 becomes

$$\delta F_{sut}^\tau = \langle \bar{X}_{su}^{(n)} \otimes \bar{X}_{ut}^{(n)}, \bar{\Delta}' \psi_n(\tau) \rangle = \delta G_{sut}^\tau. \quad (5.2)$$

since every $\tau \in \mathcal{T}$ is primitive in $(T(\mathcal{T}_N), \bar{\Delta})$ being a single-letter word.

The finite increment operator δ has the following property: if $J: [0, 1]^2 \rightarrow \mathbb{R}$ is such that $\delta J = 0$ then there exists $f: [0, 1] \rightarrow \mathbb{R}$ such that $J_{st} = f_t - f_s$, and the function f is unique up to an additive constant shift, see also [25, formula (5)]. Thus, by this fundamental property, for each $\tau \in \mathcal{T}_{n+1}$ there exists a function $x^\tau: [0, 1] \rightarrow \mathbb{R}$ such that $x_t^\tau - x_s^\tau = F_{st}^\tau - G_{st}^\tau$ and then

$$|x_t^\tau - x_s^\tau| \leq |\langle X_{st}, \tau \rangle| + |\langle \bar{X}_{st}^{(n)}, \psi_n(\tau) \rangle| \lesssim |t - s|^{\gamma|\tau|}$$

since $\psi_n(\tau)$ preserves the number of nodes by Lemma 5.1.

Repeatedly using Proposition 5.4 we obtain an anisotropic geometric rough path $\bar{X}^{(n+1)}: [0, 1] \rightarrow G_a(\mathcal{T}_{n+1})$ over $(x^\tau : \tau \in \mathcal{T}_{n+1})$ whose restriction to $T(\mathcal{T}_n)$ coincides with $\bar{X}^{(n)}$.

Finally notice that if $\tau \in \mathcal{T}_{n+1}$ is a tree then

$$\begin{aligned} \langle \bar{X}_{st}^{(n+1)}, \psi(\tau) \rangle &= \langle \bar{X}_{st}^{(|\tau|)}, \tau \rangle + \langle \bar{X}_{st}^{(|\tau|)}, \psi_{|\tau|-1}(\tau) \rangle \\ &= x_t^\tau - x_s^\tau + \langle X_{st}, \tau \rangle - (x_t^\tau - x_s^\tau) = \langle X_{st}, \tau \rangle \end{aligned}$$

and the corresponding identity for arbitrary forests follows by multiplicativity. The anisotropic geometric rough path sought for is $\bar{X} = \bar{X}^{(N)}$. □

We note that our proof is shorter and simpler than that of [28, Theorem 4.10], so we will now dedicate a few paragraphs to highlight the differences between our approach and that of Hairer and Kelly. They define first

$$\hat{X}_t^1 = \exp_N \left(\sum_{a \in A} x_t^a \bullet_a \right) \in G^N(\mathcal{T}_1)$$

then they note that this is not γ -Hölder with values in $G^N(\mathcal{T}_1)$, but it is γ -Hölder with values in $G^N(\mathcal{T}_1)/K_1$, where $K_1 := \exp_N(W_2 + \dots + W_N)$, see (2.7). By the Lyons-Victoir extension theorem there exists a γ -Hölder path $\bar{X}_t^1 \rightarrow G^N(\mathcal{T}_1)$ such that $\pi_{G^N(\mathcal{T}_1) \rightarrow G^N(\mathcal{T}_1)/K_1}(\bar{X}^1) = \hat{X}^1$. Then, in order to add a new tree τ with $|\tau| = 2$, they define

$$(\delta \bar{X}^\tau)_{st} = \langle X_{st}, \tau \rangle - \langle \bar{X}_{st}^{(1)}, \psi_1(\tau) \rangle$$

and this defines the new function $t \mapsto \langle \hat{X}_t, \tau \rangle$. Then they define

$$\hat{X}_t^2 = \exp_N \left(\sum_{a \in A} x_t^a \bullet_a + \sum_{|\tau|=2} \langle \hat{X}_t, \tau \rangle \tau \right) \in G^N(\mathcal{T}_2)$$

and again they note that this path is not γ -Hölder with values in $G^N(\mathcal{T}_2)$, but it is with values in $G^N(\mathcal{T}_2)/K_2$, where $K_2 := \exp_N(W_3 + \dots + W_N)$, and again the Lyons-Victoir extension theorem yields a γ -Hölder path $\bar{X}_t^2 \rightarrow G^N(\mathcal{T}_2)$ such that $\pi_{G^N(\mathcal{T}_2) \rightarrow G^N(\mathcal{T}_2)/K_2}(\bar{X}^2) = \hat{X}^2$. Finally they construct recursively in this way \hat{X}^k and \bar{X}^k for all $k \leq N$.

At this point we see the difference with our approach. We do not define \hat{X}_t^2 nor \hat{X}^k but rather we construct \bar{X} step by step, namely on all $G^k(\mathcal{T}_n)$ with $1 \leq k, n \leq N$, first by recursion on k for fixed n and then by recursion on n ; at each step we enforce the Hölder continuity on $G^k(\mathcal{T}_n)$ and the compatibility with the previous levels. This is done using the Lyons-Victoir technique, but in a very explicit and constructive way, in particular without ever using the axiom of choice, since we have the explicit map $\exp_{k+1} \circ \log_k: G^k(\mathcal{T}_n) \rightarrow G^{k+1}(\mathcal{T}_n)$ which plays the role of the injection $i_{G/K,G}: G/K \rightarrow G$ in [30, Proposition 6].

6 An action on branched rough paths

In this section we prove Theorem 1.2.

Given $\gamma \in]0, 1[$, let $N = \lfloor \gamma^{-1} \rfloor$ and denote by \mathcal{C}^γ the set of collections of functions $(g^\tau)_{\tau \in \mathcal{T}_N}$ such that $g^\tau \in C^{\gamma|\tau|}$ and $g_0^\tau = 0$ for all $\tau \in \mathcal{T}_N$. It is easy to see that \mathcal{C}^γ is a group under pointwise addition in t , that is,

$$(g + h)^\tau := g^\tau + h^\tau.$$

As a consequence of Proposition 5.4, $(g, \bar{X}) \mapsto g\bar{X}$ is an action of \mathcal{C}^γ on the space of anisotropic geometric rough paths.

We use the Hairer-Kelly map ψ of Lemma 5.1 to induce an action of \mathcal{C}^γ on branched rough paths. Given a branched rough path X and $g \in \mathcal{C}^\gamma$ we let gX be the branched rough path defined by

$$\langle gX_{st}, \tau \rangle = \langle g\bar{X}_{st}, \psi(\tau) \rangle,$$

where \bar{X} is the anisotropic geometric rough path given by Theorem 5.6. As a simple consequence of Proposition 5.4 we obtain

Proposition 6.1. *Let $X \in BRP^\gamma$.*

- 1 We have $g'(gX) = (g' + g)X$ for all $g, g' \in \mathcal{C}^\gamma$.
- 2 If $(g^\tau)_{\tau \in \mathcal{T}_N} \in \mathcal{C}^\gamma$ is such that there exists a unique $\tau \in \mathcal{T}_N$ with $g^\tau \neq 0$, then

$$\langle (gX)_{st}, \tau \rangle = \langle X_{st}, \tau \rangle + g_t^\tau - g_s^\tau$$

and $\langle gX, \sigma \rangle = \langle X, \sigma \rangle$ for all $\sigma \in \mathcal{T}$ not containing τ as a subtree.

Proof. The first claim follows from point (1) in Proposition 5.4. In order to prove the second claim, let $g = (g^\tau)_{\tau \in \mathcal{T}_N} \in \mathcal{C}^\gamma$ be such that there exists a unique $\tau \in \mathcal{T}_N$ with $g^\tau \neq 0$. Then by the property of g we have

$$\begin{aligned} \langle gX, \tau \rangle &= \langle g\bar{X}, \psi(\tau) \rangle = \langle g\bar{X}, \tau + \psi_{|\tau|-1}(\tau) \rangle \\ &= \langle \bar{X}, \tau \rangle + \delta g^\tau + \langle g\bar{X}, \psi_{|\tau|-1}(\tau) \rangle \end{aligned}$$

where $\delta g_{st}^\tau := g_t^\tau - g_s^\tau$. By Lemma 5.1 the tree τ does not appear as a factor in any of the tensor products appearing in $\psi_{|\tau|-1}(\tau)$, hence one can recursively show that $\langle g\bar{X}, \psi_{|\tau|-1}(\tau) \rangle = \langle \bar{X}, \psi_{|\tau|-1}(\tau) \rangle$ so that the above expression becomes

$$\begin{aligned} \langle gX, \tau \rangle &= \langle \bar{X}, \tau + \psi_{|\tau|-1}(\tau) \rangle + \delta g^\tau \\ &= \langle X, \tau \rangle + \delta g^\tau. \end{aligned}$$

For the last assertion, it is enough to note that $\sigma \in \mathcal{T}$ contains $\tau \in \mathcal{T}$ if and only if τ appears in the expression for $\psi(\sigma)$; this can be expressed more precisely by saying that $\sigma \notin T(\mathcal{T}_N \setminus \{\tau\})$. But if $\sigma \in T(\mathcal{T}_N \setminus \{\tau\})$, then $\langle g\bar{X}, \psi(\sigma) \rangle = \langle \bar{X}, \psi(\sigma) \rangle$. \square

Proposition 6.2. *The action of \mathcal{C}^γ on branched γ -rough paths is transitive: for every pair of branched γ -rough paths X and X' there exists $g \in \mathcal{C}^\gamma$ such that $gX = X'$.*

Proof. We define $g \in \mathcal{C}^\gamma$ inductively by imposing the desired identity. For trees $\tau \in \mathcal{T}_1 = \{\bullet_1, \dots, \bullet_d\}$ we set $g_t^\tau = \langle X'_{0t}, \tau \rangle - \langle X_{0t}, \tau \rangle \in C^\gamma$ so that

$$\langle gX, \tau \rangle = \langle g\bar{X}, \psi(\tau) \rangle = \langle g\bar{X}, \tau \rangle = \langle \bar{X}, \tau \rangle + \delta g^\tau = \langle X', \tau \rangle$$

where $\delta g_{st}^\tau := g_t^\tau - g_s^\tau$. Suppose we have already defined g^τ for all $\tau \in \mathcal{T}_n$ for some $n \geq 1$, satisfying the constraints in the definition of \mathcal{C}^γ . For a tree τ with $|\tau| = n + 1$ we define

$$F_{st}^\tau = \langle X'_{st}, \tau \rangle - \langle \bar{X}_{st}, \tau \rangle - \langle g\bar{X}_{st}, \psi_n(\tau) \rangle.$$

Then

$$\begin{aligned} \delta F_{sut}^\tau &= \langle X'_{su} \otimes X'_{ut}, \Delta' \tau \rangle - \langle g\bar{X}_{su} \otimes g\bar{X}_{ut}, \bar{\Delta}' \psi_n(\tau) \rangle \\ &= \langle X'_{su} \otimes X'_{ut}, \Delta' \tau \rangle - \langle g\bar{X}_{su} \otimes g\bar{X}_{ut}, \bar{\Delta}' \psi(\tau) \rangle \\ &= \langle X'_{su} \otimes X'_{ut}, \Delta' \tau \rangle - \langle g\bar{X}_{su} \circ \psi \otimes g\bar{X}_{ut} \circ \psi, \Delta' \tau \rangle \\ &= \langle X'_{su} \otimes X'_{ut}, \Delta' \tau \rangle - \langle gX_{su} \otimes gX_{ut}, \Delta' \tau \rangle = 0 \end{aligned}$$

by the induction hypothesis. Hence there is $g^\tau : [0, 1] \rightarrow \mathbb{R}$ such that $g_0^\tau = 0$ and

$$g_t^\tau - g_s^\tau = \langle X'_{st}, \tau \rangle - \langle \bar{X}_{st}, \tau \rangle - \langle g\bar{X}_{st}, \psi_n(\tau) \rangle \tag{6.1}$$

whence $g \in C^{\gamma|\tau|}$; by construction

$$\begin{aligned} \langle gX, \tau \rangle &= \langle g\bar{X}, \psi(\tau) \rangle = \langle g\bar{X}, \tau \rangle + \langle g\bar{X}, \psi_n(\tau) \rangle \\ &= \langle \bar{X}, \tau \rangle + \delta g^\tau + \langle g\bar{X}, \psi_n(\tau) \rangle = \langle X', \tau \rangle, \end{aligned}$$

where $\delta g_{st}^\tau = g_t^\tau - g_s^\tau$. This concludes the proof. □

Proposition 6.3. *The action of \mathcal{C}^γ on branched γ -rough paths is free, namely if $gX = g'X$ then $g = g'$.*

Proof. This follows from the fact that by (6.1) the function g^τ is defined up to a constant shift. Therefore, the condition $g_0^\tau = 0$ determines g^τ uniquely. □

Together, Proposition 6.1, Proposition 6.2 and Proposition 6.3 imply Theorem 1.2.

6.1 The BCFP renormalisation

In [4] a different kind of modification is proposed. There, a new decoration 0 is considered so rough paths –branched and geometric– are over paths taking values in \mathbb{R}^{d+1} . Recall that since branched rough paths are seen as Hölder paths taking values in the character group of the Butcher-Connes-Kreimer Hopf algebra, we may think of them as an infinite forest series of the form

$$X_{st} = \sum_{\tau \in \mathcal{F}} \langle X_{st}, \tau \rangle \tau \tag{6.2}$$

where we regard τ as a linear functional on \mathcal{H} , such that $\langle \tau, \sigma \rangle = 1$ if $\sigma = \tau$ and zero else. The aforementioned modification procedure then acts as a translation of the series (6.2). Specifically, for each collection $v = (v_0, \dots, v_d) : \mathcal{T} \rightarrow \mathbb{R}^{d+1}$ an operator $M_v : \mathcal{H}^* \rightarrow \mathcal{H}^*$ is defined, such that for a γ -branched rough path, $(M_v X)_{st} := M_v(X_{st})$ is a γ/N -branched rough path.

In the particular case where $v_j = 0$ except for v_0 , the action of this operator can be described in terms of an extraction/contraction map¹ $\Psi : \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{H}$. This map acts on a tree τ by extracting

¹In [4] this map is named δ but we choose to call it Ψ in order to avoid confusion with the operator defined here.

subforests and placing them in the left factor; the right factor is obtained by contracting the extracted forest and decorating the resulting node with 0. As an example, consider

$$\begin{aligned} \Psi(\mathbb{V}_i^{j,k}) &= 1 \otimes \mathbb{V}_i^{j,k} + \bullet_i \otimes \mathbb{V}_0^{j,k} + \bullet_j \otimes \mathbb{V}_i^{0,k} + \bullet_k \otimes \mathbb{V}_i^{j,0} + \mathbb{I}_i^j \otimes \mathbb{I}_0^k + \mathbb{I}_i^k \otimes \mathbb{I}_0^j \\ &+ \bullet_i \bullet_j \otimes \mathbb{V}_0^{0,k} + \bullet_i \bullet_k \otimes \mathbb{V}_0^{j,0} + \bullet_j \bullet_k \otimes \mathbb{V}_i^{0,0} + \mathbb{I}_i^{j \bullet k} \otimes \mathbb{I}_0^0 + \mathbb{I}_i^k \bullet_j \otimes \mathbb{I}_0^0 + \mathbb{V}_i^{j \bullet k} \otimes \bullet_0. \end{aligned}$$

Extending $\nu = \nu_0: \mathcal{T} \rightarrow \mathbb{R}$ to all of \mathcal{H}^* as an algebra morphism it is shown that

$$\langle (M_\nu X)_{st}, \tau \rangle = \langle X_{st}, (\nu \otimes \text{id})\Psi(\tau) \rangle. \tag{6.3}$$

Furthermore, in this case $M_\nu X$ is a γ -branched rough path if coefficients corresponding to trees with decoration zero are required to satisfy the stronger analytical condition

$$\sup_{0 \leq s, t \leq 1} \frac{|\langle X_{st}, \tau \rangle|}{|t - s|^{(1-\gamma)|\tau|_0 + \gamma|\tau|}} < \infty, \tag{6.4}$$

where $|\tau|_0$ counts the times the decoration 0 appears in τ . Essentially, this condition imposes that the components corresponding to the zero decoration be Lipschitz on the diagonal $s = t$.

We now show how this setting can be recovered from the results of Section 6. Let X be a γ -branched rough path on \mathbb{R}^{d+1} satisfying (6.4). Since $M_\nu X$ is again a γ -branched rough path, by Proposition 6.2 there exists a collection of functions $g \in \mathcal{C}^\gamma$ such that $gX = M_\nu X$. Moreover, this collection is the unique one satisfying

$$g_t^\tau - g_s^\tau = \langle X_{st}, (\nu \otimes \text{id})\Psi(\tau) \rangle - \langle \bar{X}_{st}, \tau \rangle - \langle g \bar{X}_{st}, \psi_{|\tau|-1}(\tau) \rangle \tag{6.5}$$

for all $\tau \in \mathcal{T}(\mathbb{R}^{d+1})$ where we have used (6.3) in order to express $M_\nu X$ in terms of Ψ . Theorem 28 in [4] ensures that the first term on the right-hand side is in $C_2^{\gamma|\tau|}$ hence g is actually in $C^{\gamma|\tau|}$ as required.

The approach of [4] is based on pre-Lie morphisms and crucially on a *cointeraction property*, which has been explored by [7], see in particular [4, Lemma 18]. The cointeraction property can be used for *time-independent* modifications, indeed note that the functional ν in [4] is always constant.

Let us see why this is the case. The approach of [4] is based on a cointeraction property studied by [6, 7, 21] between the Butcher-Connes-Kreimer coproduct and another *extraction-contraction* coproduct $\delta: \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{H}$. The formula is the following

$$(\text{id} \otimes \Delta)\delta = \mathcal{M}_{1,3}(\delta \otimes \delta)\Delta.$$

Let us consider now a character $\nu \in \mathcal{H}^*$. If we multiply both sides by $(\nu \otimes \text{id} \otimes \text{id})$ and set $M_\nu^* = (\nu \otimes \text{id})\delta: \mathcal{H} \rightarrow \mathcal{H}$ as in [4, Proposition 17], then we obtain

$$\Delta M_\nu^* = (M_\nu^* \otimes M_\nu^*)\Delta,$$

namely M_ν^* is a coalgebra morphism on \mathcal{H} . Then one can define a modified rough path as $\nu X := M_\nu X = X \circ M_\nu^*$. The crucial Chen property is still satisfied since

$$\begin{aligned} (\nu X)_{st} &= (\nu \otimes X_{st})\delta = (\nu \otimes X_{su} \otimes X_{ut})(\text{id} \otimes \Delta)\delta \\ &= (\nu \otimes X_{su} \otimes X_{ut})\mathcal{M}_{1,3}(\delta \otimes \delta)\Delta \\ &= ((\nu \otimes X_{su}) \otimes (\nu \otimes X_{ut}))(\delta \otimes \delta)\Delta \\ &= ((\nu X)_{su} \otimes (\nu X)_{ut})\Delta \end{aligned}$$

However this does not work if $v: [0, 1]^2 \rightarrow \mathcal{H}^*$ is a time-dependent character. Indeed in this case we set $(vX)_{st} := (v_{st} \otimes \text{id})\delta$ and we obtain

$$\begin{aligned} (vX)_{st} &= (v_{st} \otimes X_{st})\delta = (v_{st} \otimes X_{su} \otimes X_{ut})(\text{id} \otimes \Delta)\delta \\ &= (v_{st} \otimes X_{su} \otimes X_{ut})\mathcal{M}_{1,3}(\delta \otimes \delta)\Delta \\ &= ((v_{st} \otimes X_{su}) \otimes (v_{st} \otimes X_{ut}))(\delta \otimes \delta)\Delta \end{aligned}$$

but we can not conclude that this is equal to $((vX)_{su} \otimes (vX)_{ut})\Delta$. Our construction, as explained after formula (1.8), is not purely algebraic but is based on a (non-canonical) choice of generalized Young integrals with respect to the rough path X . Moreover our transformation group, infinite-dimensional, is much larger than that finite-dimensional group studied in [4].

7 Perspectives

In this paper we have shown that the space of branched γ -rough paths is a principal homogeneous space with respect to the linear group \mathcal{C}^γ . This is related to the analytical properties of the operator δ defined in (1.2), which is invertible under the conditions of Gubinelli's Sewing Lemma, but not in general, and in particular not in the context of the Chen relation on trees with low degree.

It would be now interesting to see how this action can be translated on the level of controlled paths [24]. The space of paths controlled by a rough path $X \in \mathbf{BRP}^\gamma$ should be interpreted as the tangent space to \mathbf{BRP}^γ at X , and the action on rough paths should induce an action on controlled paths. In particular it should be possible to write an action on solutions to rough differential equations.

The proof of Proposition 6.2, and in particular (6.1), gives a recursive way of computing the unique $g \in \mathcal{C}^\gamma$ translating a given branched γ -rough path into another. An interesting feature of the BCFP scheme is that is given in terms of a coaction so explicit calculations are somewhat easier in this more restricted case as one can compute g^τ for each tree $\tau \in \mathcal{T}_N$ directly by extracting and contracting subforests of τ without doing any recursions (see (6.5).) However, we do not have a computational rule for an important case: suppose that \mathbb{X} is branched rough path lift of a stochastic process with a.s. C^γ -trajectories; it would be nice to have a way of finding $g \in \mathcal{C}^\gamma$ such that $g\mathbb{X}$ is centered with respect to the underlying distribution of the process, provided this is possible. Even this last problem, namely giving precise conditions under which this centering is possible is interesting in itself. This should be related to the notion of Wick polynomials and deformations of products as considered in [18].

More generally, in the physics literature there are various renormalisation procedures which allow to obtain convergent iterated integrals from divergent ones by subtracting suitable "counterterms". In the context of rough paths, implementing one of the most accepted such procedures due to Bogoliubov–Parasiuk–Hepp–Zimmerman (BPHZ) has been carried out by J. Unterberger in [35, 36] by means of the Fourier normal ordering algorithm and using a technique relating the trees in the Butcher–Connes–Kreimer Hopf algebra to certain Feynman diagrams. In our context, this could provide a canonical choice for $g \in \mathcal{C}^\gamma$ implementing the BPHZ renormalization procedure in a way analogous to what is done in [6] for Regularity Structures.

References

- [1] G. Ben Arous, "Flots et séries de Taylor stochastiques," *Probab. Theory Related Fields* **81** no. 1, (1989) 29–77.

- [2] H. Boedihardjo and I. Chevyrev, “An isomorphism between branched and geometric rough paths,” [arXiv:1712.01965](https://arxiv.org/abs/1712.01965) [math.PR].
- [3] Y. Bruned, I. Chevyrev, and P. K. Friz, “Examples of renormalized SDEs,” in *Stochastic partial differential equations and related fields*, vol. 229 of *Springer Proc. Math. Stat.*, pp. 303–317. Springer, Cham, 2018.
- [4] Y. Bruned, I. Chevyrev, P. K. Friz, and R. Preiß, “A Rough Path Perspective on Renormalization,” to appear in *Journal of Functional Analysis* (2019), [arXiv:1701.01152](https://arxiv.org/abs/1701.01152) [math.PR].
- [5] Y. Bruned, C. Curry, and K. Ebrahimi-Fard, “Quasi-shuffle algebras and renormalisation of rough differential equations,” [arXiv:1801.02964](https://arxiv.org/abs/1801.02964) [math.PR].
- [6] Y. Bruned, M. Hairer, and L. Zambotti, “Algebraic renormalisation of regularity structures,” *Invent. Math.* **215** no. 3, (2019) 1039–1156.
<https://doi.org/10.1007/s00222-018-0841-x>.
- [7] D. Calaque, K. Ebrahimi-Fard, and D. Manchon, “Two interacting Hopf algebras of trees: A Hopf-algebraic approach to composition and substitution of B-series,” *Adv. Appl. Math.* **47** no. 2, (2011) 282–308, [arXiv:0806.2238](https://arxiv.org/abs/0806.2238) [math.CO].
- [8] P. Cartier, *A Primer of Hopf Algebras*, pp. 537–615. Springer-Verlag Berlin Heidelberg, 2007.
- [9] A. Chandra and M. Hairer, “An analytic BPHZ theorem for regularity structures,” [arXiv:1612.08138](https://arxiv.org/abs/1612.08138) [math.PR].
- [10] F. Chapoton, “Free Pre-Lie Algebras are Free as Lie Algebras,” *Canad. Math. Bull.* **53** no. 3, (2010) 425–437, [arXiv:0704.2153](https://arxiv.org/abs/0704.2153) [math.RA].
- [11] K.-T. Chen, “Iterated path integrals,” *Bull. Amer. Math. Soc.* **83** no. 5, (1977) 831–880.
- [12] A. Connes and D. Kreimer, “Hopf Algebras, Renormalization and Noncommutative Geometry,” *Comm. Math. Phys.* **199** no. 1, (1998) 203–242, [arXiv:hep-th/9808042](https://arxiv.org/abs/hep-th/9808042) [hep-th].
- [13] L. Coutin and Z. Qian, “Stochastic analysis, rough path analysis and fractional Brownian motions,” *Probab. Theory Related Fields* **122** no. 1, (2002) 108–140.
- [14] C. Curry, K. Ebrahimi-Fard, D. Manchon, and H. Z. Munthe-Kaas, “Planarly branched rough paths and rough differential equations on homogeneous spaces,” [arXiv:1804.08515](https://arxiv.org/abs/1804.08515) [math.CA].
- [15] E. B. Dynkin, *Calculation of the coefficients in the Campbell–Hausdorff formula*, pp. 31–35. Collected Works. American Mathematical Society and International Press, 2000.
- [16] K. Ebrahimi-Fard, S. J. A. Malham, F. Patras, and A. Wiese, “The exponential Lie series for continuous semimartingales,” *Proc. A.* **471** no. 2184, (2015) 20150429, 19.
- [17] K. Ebrahimi-Fard, S. J. A. Malham, F. Patras, and A. Wiese, “Flows and stochastic Taylor series in Itô calculus,” *J. Phys. A* **48** no. 49, (2015) 495202, 17.
- [18] K. Ebrahimi-Fard, F. Patras, N. Tapia, and L. Zambotti, “Hopf-algebraic deformations of products and Wick polynomials,” to appear in *Int. Math. Res. Not.* (2018), [arXiv:1710.00735](https://arxiv.org/abs/1710.00735) [math.PR].

- [19] L. Foissy, “Algèbres de Hopf combinatoires,”
<http://loic.foissy.free.fr/pageperso/Hopf.pdf>. Master 2 lecture notes.
- [20] L. Foissy, “Finite dimensional comodules over the Hopf algebra of rooted trees,” *J. Algebra* **255** no. 1, (2002) 89–120, arXiv:math/0105210 [math.QA].
- [21] L. Foissy, “Commutative and non-commutative bialgebras of quasi-posets and applications to Ehrhart polynomials,” arXiv:1605.08310 [math.RA].
- [22] G. B. Folland and E. M. Stein, *Hardy Spaces on Homogeneous Groups*. No. 28 in Princeton mathematical notes. Princeton University Press, 1 ed., 1982.
- [23] P. K. Friz and N. B. Victoir, *Multidimensional Stochastic Processes as Rough Paths*. Cambridge University Press, Cambridge, 2010.
- [24] M. Gubinelli, “Controlling Rough Paths,” *J. Funct. Anal.* **216** no. 1, (2004) 86–140, arXiv:math/0306433 [math.PR].
- [25] M. Gubinelli, “Ramification of rough paths,” *J. Differential Equations* **248** no. 4, (2010) 693–721, arXiv:math/0610300 [math.CA].
- [26] L. G. Gyurkó, “Differential Equations Driven by Π -Rough Paths,” *Proc. Edinb. Math. Soc.* **59** no. 3, (2016) 741–758, arXiv:1205.1832 [math.CA].
- [27] M. Hairer, “A theory of regularity structures,” *Invent. Math.* **198** no. 2, (2014) 269–504, arXiv:1303.5113 [math.AP].
- [28] M. Hairer and D. Kelly, “Geometric versus non-geometric rough paths,” *Annales de l’Institut Henri Poincaré Probabilités et Statistiques* **198** no. 2, (2014) 269–504, arXiv:1210.6294 [math.PR].
- [29] T. Lyons, “Differential equations driven by rough signals,” *Rev. Mat. Iberoam.* **14** (1998) 215–310.
- [30] T. Lyons and N. B. Victoir, “An extension theorem to rough paths,” *Ann. Inst. H. Poincaré Anal. Non Linéaire* **24** no. 5, (2007) 835–847.
- [31] D. Manchon, “Hopf Algebras in Renormalisation,” in *Handbook of Algebra*, M. Hazewinkel, ed., vol. 5, pp. 365–427. North-Holland, 2008.
- [32] H. Munthe-Kaas and W. Wright, “On the Hopf Algebraic Structure of Lie Group Integrators,” *Foundations of Computational Mathematics* **8** no. 2, (2008) 227–257.
- [33] D. Nualart and S. Tindel, “A construction of the rough path above fractional Brownian motion using Volterra’s representation,” *Ann. Probab.* **39** no. 3, (2011) 1061–1096.
- [34] C. Reutenauer, “Theorem of Poincare-Birkhoff-Witt, logarithm and symmetric group representations of degrees equal to stirling numbers,” in *Combinatoire énumérative*, G. Labelle and P. Leroux, eds., vol. 1234 of *Lecture Notes in Mathematics*. Springer, Berlin, Heidelberg, 1986.
- [35] J. Unterberger, “Hölder-continuous rough paths by Fourier normal ordering,” *Comm. Math. Phys.* **298** no. 1, (2010) 1–36.
- [36] J. Unterberger, “A renormalized rough path over fractional Brownian motion,” *Comm. Math. Phys.* **320** no. 3, (2013) 603–636.