ORGANIZATION OF AFRICAN UNITY SCIENTIFIC, TECHNICAL AND RESEARCH COMMISSION (D A U / S T R C)

WEST AND CENTRAL AFRICA COWPEA NETWORK

"Réseau Niébé de l'Afrique Centrale et Occidentale"

(R E N A C O)



REGIONAL APPROACH TO COWPEA RESEARCH IN WEST AND CENTRAL AFRICA

Dr. N. Muleba





RENACO-IITA/SAFGRAD
O1, B.P. 1495,
OUAGADOUGOU O1, BURKINA FASO



TABLE OF CONTENTS **********

| I. BACKGROUNI | D |
|----------------------------|---|
| 1) Agro- | -ecologies |
| | atic constraints |
| 3) Biolo | ogical constraints |
| 4) Soil | constraints |
| 5) Socio | o-economic constraints |
| 6) Finar | ncial constraints |
| 7) Insuf techr | fficient number of skilled scientists, nicians and extension personnel |
| II. NETWORK O | DBJECTIVES |
| III. NETWORK | PROGRAMME AND IMPLEMENTATION |
| IV. SUMMARY C | OF ACHIEVEMENTS OF THE COWPEA NETWORK |
| 2) Regio | ngthening national research system |
| V. FUTURE THE FINANCIAL | RUST OF NETWORK PROGRAM AND AREAS OF WHICH SUPPORT IS REQUIRED |
| APPENDIX 1 : | List of national cowpea scientists - RENACO |
| APPENDIX 2 : | Varieties reseased by different RENACO national cowpea programmes since 1987. |
| APPENDIX 3 : | Varieties in pre-extension stage in various RENACO national programmmes since 1987. |
| APPENDIX 4 : | List of national and international scientists who attended the 1987 workshop at Ouagadougou, Burkina Faso in March, 1987. |
| APPENDIX 5 : | RENACO Steering Committee Meetings |

Name of Project: West and Central Africa Cowpea Network (RENACO)

Date Commenced: March 1987.

I. - BACKGROUND

The SAFGRAD Phase I Cowpea Research mandate in Burkina Faso was successfully completed in 1978-1986.

From the laudable scientific breakthrough of the Phase I research activities, it was unanimously agreed at two Workshops held at Ouagadougou, Burkina Faso from 23-27 February 1987 and from 23-27 March 1987 by national directors of agricultural research and their cowpea scientists of the 18 SAFGRAD member countries as well as Regional and International Research Centers that the SAFGRAD Research Project should be extended into a second phase.

The second phase of the SAFGRAD cowpea research project has a primary objective of boosting the capacity of national scientists to direct cowpea research activities themselves in the subregion in the long run.

In order to prepare the foundation for the eventual take over of cowpea research activities by national scientists, a collective venture on cowpea research was established by SAFGRAD-IITA involving 17 SAFGRAD member countries known as the West and Central Africa Cowpea Collaborative Research Network (RENACO). Cowpea production constraints, research personnel, infrastructure as well as the research strengths and weaknesses of each national programme were presented. The needs, researchable topics and the state of art on cowpea research in Central and West Africa were also enumerated and discussed.

The national directors of research and cowpea scientists were sincere in appraising their individual country's research capacities and they fully endorsed the laudable idea of networking.

They believed that the network exercise was the most feasible solution to tackling cowpea production constraints by sharing scientific information and technologies so developed from the network exercise or by other regional and international agricultural centers.

A Steering Committee comprising of six national cowpeascientists was elected during the workshop. The committee immediately met with responsible authorities of the SAFGRAD Coordination Office (SCO), IITA-GLIP, USAID and the IITA-GLIP seconded Coordinator of the network.



A review of cowpea research and production programmes as well as research infrastructure and personnel of each country was carried out. Common constraints were identified and relatively strong national programmes were given the role of Lead Centers. Researchable topics were assigned to Lead Centers according to their strengths and ecological zones.

The resolutions and decisions taken by the Steering Committee during the March 1987 workshop are summarized as follows:

1) Agro-ecologies

Three agro-ecological zones from north to south were recognized:

- 1) The Sahel: 200-600 mm rainfall from mid-June to mid-September;
- 2) Sudan savanna: 600-900 mm rainfal from June to September;
- 3) Northern Guinea savanna: 900-1200 mm rainfall from June to mid-October.

2) Climatic Constraints

Drought (inadequate, poor distribution and erratic rainfall) and heat (high air and soil temperatures) stresses, and sandblasts due to high wind velocity are major climatic constraints. They are gradually increasing from south to northwards.

3) Biological Constraints

Diseases (scab, brown blotch, <u>Septoria</u> leaf spot, viral diseases, bacterial blight, ashy-stem rot), insect pests (thrips, aphids, bruchids, pod sucking bugs and <u>Maruca</u> pod borers), parasitic weeds (<u>Striga</u> and <u>Alectra</u>).

4) Soil Constraints

Low water retention capacity, low fertility and high soil temperatures.

5) Socio-economic constraints

Poor on farm-testing, inadequate seed production and distribution system and continued cultivation without use of appropriate inputs.

6) Financial Constraints

7) Insufficient number of skilled scientists, technicians and extension personnel

In conclusion, after prioritizing cowpea production constraints and evaluating the capacity of national programmes, the Steering Committee allocated research responsibilities to strong national programmes according to their geographical locations. The Steering Committee also realized that peasant farmers adopt new technologies not necessarily because of better adaptation to the physical environment and high yield, but also consumers preference and requirements (grain type, colour, texture, etc). Therefore, national scientists were urged to pay more attention to these spectifics more than ever before.

II. - NETWORK OBJECTIVES

The primary objective of RENACO is to develop the capacity and initiative of the national cowpea scientists to direct the network themselves by (i) properly identifying cowpea production constraints and (ii) generating through networking in collaboration with IITA-GLIP, the appropriate technologies overcoming the constraints. The purpose of networking is to enable national cowpea programmes of West and Central Africa to pool together their resources to tackle common cowpea production problems in the subregion and to find appropriate solutions for the benefit of their inhabitants. The rationale of networking is based on this very simple but effective adage "United, we stand, Divided, we fall". This becomes more relevant when needs are numerous, and resources are limited.

III. - NETWORK PROGRAMME AND IMPLEMENTATION

Based on the commonality of the constraints and the existence of strong and weak national programmes within the subregion, in order to ensure the cost effectiveness and sustainability of networking, the cowpea network Steering Committee adopted the strategy of assigning technology-development research responsibilities to strong national programmes (Lead Centers) depending on the ecological zone; the technology adaptive research responsibilities being handled by all national programmes, especially the weak ones, of the participating countries, while IITA-GLIP at Ibadan, Nigeria, and its outreach sub-stations in Nigeria and Niger, continue to backstop the activities (research, training, etc) of the network. Varieties developed by IITA are channelled, through Lead Centers, directly or indirectly within the network; directly if varieties were recommended for regional testing after being identified as promising by Lead Centers; and indirectly if they were used in cowpea genetic improvement by Lead Centers.

The cowpea Steering Committee assigned research responsibilities to 6 national programmes, which accepted the role as Lead Centers as follows:

1) Burkina Faso

- Breeding for drought, <u>Striga</u>, insect pests and disease resistance;
- Entomology and pathology (including viral diseases) for the three ecological zones of semi-arid West Africa.

2) Cameroun

- Entomology with emphasis on cowpea storage pest problems.

3) Niger

- Breeding for drought, <u>Striga</u> and <u>Macrophomina</u> disease resistance;
- Agronomic studies (millet-cowpea intercropping) and cowpea pathology (Macrophomina spp) for the Sahelian zones.

4) Nigeria

- Breeding for drought, Striga, Alectra, insect pests and disease resistance;
- Cowpea agronomy, pathology (including scab, brown blotch, Septoria leaf spot, Striga and Alectra) and entomology for the three ecological zones of West Africa with emphasis on mode of inheritance of diseases, Striga and Alectra resistance in cowpea.

5) Senegal

- Breeding for drought, insect pests and disease resistance;
- Cowpea entomology for the Sahelo-Sudanian zones.

6) Ghana

- Breeding for adaptation to transition zones;
- Cowpea entomology for transition zones.

Owing to variations in <u>Striga</u> strains, two countries (Benin and Mali) were assigned the responsibilities of selecting and testing for <u>Striga</u> resistance in 1990.

IV. - SUMMARY OF ACHIEVEMENTS OF THE COWPEA NETWORK

1) Strengthening national research system

As at October 30, 1990, RENACO Lead Centers were conducting activities in all aspects of cowpea research. While capitalizing on multiple insect pests and disease resistance developed by IITA-GLIP, national scientists of Burkina Faso attempting to incorporate them into agronomic backgrounds, acceptable to peasant farmers. Their ultimate goal is to have drought, multiple disease, <u>Striga</u>, aphids and other insect pest resistant cowpea varieties. Nigerian scientists are attempting to identify new sources of Striga and Alectra tolerance and mode of inheritance which is a commendable step towards an initiation of a breed ing programme for resistance to these parasitic weeds. Scientists from Niger are looking for new and stable sources of Striga resistance. Whereas, scientists in Mali are studying the virulence of Striga gesnerioides strains including the ones parasitizing weeds such as Ipomeas sp.. Scientists in Cameroon are working on cowpea storage methods acceptable to peasant farmers.

2) Regional trials

In 1987, the biennial regional testing (1987-88) consisted of a total of 7 trials:

- 1) Drought resistance
- 2) Striga resistance
- 3) Sorghum-cowpea intercropping
- 4) Millet-cowpea intercropping
- 5) Maize-cowpea relay cropping
- 6) Observation nursery
- 7) Minimum insecticide

in 81 sets. The inputs of the trials were obtained from IITA-SAFGRAD resident research in Ouagadougou and from IITA headquarters, Ibadan, Nigeria. They were dispatched to member countries as shown in Table 1; a total of 78 feedbacks was received from participating countries by the end of 1988 (Table 2).

In 1989, the biennial regional testing (1989-90) consisted again of a total of 7 trials:

- 1) Resistance to aphids
- 2) Resistance to bruchids
- 3) Resistance to virus
- 4) Resistance to Striga
- 5) Adaptation to transition zones
- 6) Adaptation to Sudano-Sahelian zones
- 7) Adaptation to Northern-Guinea zones

in 53 sets. Lines included in the 1989 trials were developed by Burkina Faso, Niger, Nigeria, Ghana and IITA-GLIP. Trials were dispatched to member countries as shown on Table 3; and a total of 35 feedbacks was received from participating countries (Table 4).

TABLE 1. COWPEA REGIONAL TRIALS DISPATCHED TO MEMBER COUNTRIES IN 1987

| | | | NUMBE | R OF TH | RIALS RE | QUESTED | | |
|---------------|--|-------------|---------|-------------------------------------|------------------|--------------------------------|----|-------|
| Country | Drought <u>Striga</u> resis- resis- | | Intercr | opping | Maize/ Cowpea | Maize/ Obser- Cowpea vation | | Total |
| en a v | tance trial | tance tance | | sorghum/ millet/ cowpea cowpea c | | Relay nursery cropping | | Total |
| Benin | 1 | | 2 | _ | _ | 1 | 2 | 6 |
| Burkina Faso | 1 | 1 | _ | _ | | 1 | | 3 |
| Cameroon | _ | | 1 | _ | 1 | _ | 1 | 3 |
| Cape Verde | _ | | _ | _ | _ | 1 | _ | 1 |
| Central Afric | an | | | | | | | |
| Republic | - | - | _ | _ | _ | (2)-1 | _ | 1 |
| Tchad | 2 | | _ | 2 | 1 | 2 | | 7 |
| The Gambia | 1 | _ | 2 | 2 | | 1 | | 6 |
| Ghana | 1 | 1 | 1 | _ | | This - | 1 | 4 |
| Guinea Bissau | 1 1 | _ | 1 | - | _15.00 | 1 | | 3 |
| Guinea Conakr | | - | 1 | _ | 2 | (3)-2 | 1 | 5 |
| Côte d'Ivoire | , – | - | - | _ | | 1 | | 1 |
| Mali | 2 | 2 | - | 3 | - 1984 | 1 | _ | 8 |
| Mauritania | _ | - | - 1 7 7 | 7 1 | - | | _ | , 0 |
| Niger | 3 | 3 | - | - 100 | | (3)-1 | 2 | 10 |
| Nigeria | 2 | 3 | 1 | 3 | 1 | 1 | 2 | 13 |
| Senegal | 2 | - | 1 | 1 | 12 857 | (3)-2 | 2 | 7 |
| Sierra Leone | - 200 | W-Made | 2-1 | September 15 Child | TO THE STATE OF | | _ | 0 |
| Togo | | | 2 | | 1 | | | 3 |
| Total | 16 | 10 | 12 | 11 | 6 | 15 | 11 | 81 |

TABLE 2. FEEDBACKS RECEIVED FROM MEMBER COUNTRIES FOR THE 1987-88 REGIONAL TRIALS

| NAME OF TRIALS | | | | | | | | | | |
|-----------------|-------------------|------------------|-----------------|----------|-------------------|------------------|--------------------|------------------|-------------------|-------|
| Country | Drought resis- | Striga resis- | Virus resis- | Intercro | pping | Maize/ Cowpea | Bruchids resis- | Aphids resis- | Minimum insec- | Total |
| | tance trial | tance trial | tance trial | | millet/ cowpea | Relay crop | tance trial | tance trial | ticide trial | |
| Benin | 2 | 1 | 1 | 1 | _ | _ | 1 | 1 | _ | 6 |
| Burkina Faso | 3 | 3 | 1 | 2 | _ | 2 | 1 | 1 | 3 | 16 |
| Cameroon | - | 1 | - | 1 | _ | _ | _ | _ | 1 | 3 |
| Central African | | | | | | | | | | |
| Republic | | - | - | - | - | _ | - | | _ | 0 |
| Cape Verde | - | - | _ | - | _ | _ | _ | 1 | - | 1 |
| Tchad | 2 | - | - | _ | - | 1 | | - 11 | | 3 |
| The Gambia | 1 | - | - | - | - | - | - | - | 1 | 2 |
| Ghana | 2 | 1 | - | 2 | - | - | 1 | 1 | 1 | 8 |
| Guinea Bissau | 1 | - | - | - | - | - | - | - | - | 1 |
| Guinea Conakry | - | - | - | 1 | | 2 | _ | - | 1 | 4 |
| Côte d'Ivoire | - | - | - | - | - | _ | - | - | 1.10- 7 | 0 |
| Mali | 2 | 2 | Fair ca | - | - | | - | - | - | 4 |
| Mauritania | 3 | - | - | - | W | - | 2 | 2 | - | 7 |
| Niger | 3 | 3 | - | 7.5 | | - | - | - | 1 | 7 |
| Nigeria | 2 | 3 | - | - | | 1 | | - | 1 | 7 |
| Senegal | 2 | - | _ | | - | - | - | - | 1- | 2 |
| Sierra Leone | - | - | - | _ | T | - | 7 | - | | 0 |
| Togo | - | - | 2 | 2 | - | 1 | 1 | 1 | - | 7 |
| Total | 23 | 13 | 4 | 9 | 0 | 7 | 6 | 7 | 9 | 78 |

-

TABLE 3. COWPEA REGIONAL TRIALS DISPATCHED TO MEMBER COUNTRIES IN 1989

| | | | | NAME OF | TRIAL | | | |
|-----------------|--------|----------|--------|---------|-------------------------|-----------------------------|----------------------------|-------|
| Country | | Resistar | ice to | | | Adaptatio | on to | Total |
| 2 | Aphids | Bruchids | Virus | Striga | Tran- sition zone | Sudano- Sahelian zone | Northern Guinea zone | |
| Benin | _ | | | 1 | | | 1 | 2 |
| Burkina Faso | 1 | 1 | 1 | 1 | _ | | 1 | 5 |
| Cameroon | _ | 1 | 1 | _ | _ | 1 | 1 | 4 |
| Cape Verde | 1 | - S | _ | | - | | | 1 |
| Central African | | | | | | | | |
| Republic | - | - | - | - | - | _ | _ | 0 |
| Côte d'Ivoire | _ | - | - | - | - Ded | _ | 1 | 1 |
| The Gambia | - | - | - | - | - 1 | | 1 | 1 |
| Ghana | | -0 | - | | - 1 3 | | 1 | 1 |
| Guinea Bissau | 1 | 1 | _ | - | 1 - | 1 | | 3 |
| Guinea Conakry | 2 | 4 | 1 | - | 4 | | - | 11 |
| Niger | 1 | - | 1 | 1 | - | 1 | - | 4 |
| Nigeria | 1 | 1 | 1 | 1 | - 3 | 1 | 1 | 6 |
| Mali | - | 1 | - | 1 | - | - 4 | - | 2 |
| Mauritania | - | 1 | - | - | - | 1 | al -0 | 2 |
| Senegal | - | - | - | 1 | - 4 | | | 1 |
| Tchad | 1 | 1 | 1 | 1 | - | 1 | 160 - mile | 5 |
| Northern Togo | - | 12 | 1 | - | - | | 1 | 2 |
| Southern Togo | - | 1 '. | - | _ | 1 | - 45 50 | - | 2 |
| Total | 8 | 12 | 7 | 7 | 5 | 6 | 8 | 53 |

TABLE 4. FEEDBACKS RECEIVED FROM MEMBER COUNTRIES FOR THE 1989 REGIONAL COWPEA TRIALS

| | - 31 | | DE LO | NAME | OF TR | RIAL | | | |
|------------------------------|--------|----------|--------|---------|--------------|-----------|-----------------------------|----------------------------|-------|
| Country | | Re | sistan | ce to | A | daptation | to | Total | |
| | Aphids | Bruchids | Virus | Drought | Striga | | Sudano- Sahelian zone | Northern Guinea zone | Total |
| Benin | _ | | _ | - | | _ | _ | 1 | 1 |
| Burkina Faso | 1 | 1 | - | _ | 7 - | _ | 1 | 1 | 4 |
| Cameroon | _ | _ | - | | _ | _ | 1 | 1 | 2 |
| Cape Verde Central Africa | - | | _ | | - | - | | | 0 |
| Republic | 11 | | | | | | | | 0 |
| Côte d'Ivoire | | Py L | | | | 6.8 | | 1 | 1 |
| The Gambia | | | | | Near Control | | | 1 | 1 |
| Ghana | | 73.7 | | | 7 90 - | | | 1 | 1 |
| Guinea Bissau | _ | | _ | | | | | | Ô |
| Guinea Conakry | 1 | 1 | _ | | | 2 | _ | | 4 |
| Mali | _ | 1 | _ | | 1 | - | _ | di.d | 2 |
| Mauritania | | 1 | _ | 1 | 1 | _ | _ | _ | 3 |
| Niger | - | _ | _ | | | _ | 1 | | 1 |
| Nigeria | 1 | 1 | 1 | | 1 | - | | 1 | 4 |
| Senegal | _ | _ | _ | _ | - | - | _ | _ | 0 |
| Tchad | 1 | 1 | 1 | _ | 1 | _ | 1 | _ | 5 |
| Togo | 2 | 1 | 1 | - | - | 1 | - | 1 | 6 |
| Total | 6 | 7 | 2 | 1 | 4 | 3 | 4 | 8 | 35 |

3) On-farm testing

Although the network is not directly involved in multilocational trials and on-farm testings; it is significant that through the network efforts (regional trials included), there have been renewed interests in cowpea research activities in all participating countries. The following cultivars obtained through RENACO activities have been released or are about to be released in the underlisted member countries (Table 5).

4) Training of National Scientists

With the ultimate goal of boosting the capacity and initiative of national cowpea scientists to identify cowpea production constraints and develop or identify appropriate technologies to overcome such constraints, following training activities are being carried out.

Monitoring tour:

The objective of the monitoring tour is to enable scientists from Technology Adopting Centers, Lead Centers and IITA-GLIP to interact on the field with regard to production constraints, research methodologies and appropriate new technologies. A monitoring tour was organised in 1988 to IITA-Ibadan, northern Nigeria, Niger and Burkina Faso. Six national scientists from Niger, Burkina Faso, Senegal, Cape Verde and Guinea Bissau participated (Table 6). In 1990, the same afore-mentioned countries were toured by eight scientists from Benin, Burkina Faso, Cameroon, The Gambia, Ghana, Niger and Nigeria (Table 7).

Short term in-service training for scientists:

The objective is the same as the monitoring tour, but discussion are held in classrooms and laboratories with lectures given by national as well international scientists.

A seminar was organised in November 1988 at IITA, Ibadan for 12 scientists from Lead Centers and Ghana (Table 8). The scientists included breeders, agronomists, pathologists and entomologists; the subject discussed centered mainly on appropriate research methodologies.

A group training course was organized in 1989 at Kamboinse/Ouagadougou in cooperation with the national cowpea programme of Burkina Faso. Ten scientists and technicians from Côte d'Ivoire, Niger, Guinea Conakry, Mali, Benin, Guinea Bissau and Tchad participated (Table 9). The subject matters centered mainly on technology development and transfer.

TABLE 5. CULTIVARS RELEASED OR ABOUT TO BE RELEASED FROM THE NETWORK EFFORTS.

| Country | Cultiv | Cultivars | | | | | | |
|--------------------------------|---|---|---|--|--|--|--|--|
| | Released | To be released | Area of adaptation | | | | | |
| Benin | Vita-5 | IT82E-32 IT81D-1137 TVx 1850-01F | Coastal zone Coastal zone Transition zone | | | | | |
| Burkina Faso | Gorom L. (Suvita-2) KN-1 | KVx61-1 KVx396-4-4 KVx396-4-2 | Sahel Sahel Sud. zone Sudano-Guinean zone | | | | | |
| Cameroon | Br1 (IT81D-985) | IT81D-994 | Sudano-Guinean zone | | | | | |
| Ghana | Asonteme (IT82E-32) Valenga (IT82E-16) | | Transition zone Guinea savanna zone | | | | | |
| Guinea Bissau | IT82E-9 | | Guinea savanna zone, | | | | | |
| Mali | Gorom L. (Suvita-2) | KVx61-1 | Sahel | | | | | |
| | TN88-63 KN-1 | KVx61-74 | Sahelo-Sudanian Sudano-Guinea | | | | | |
| Gambia | IT81D-994 | - | Sudano-Guinean | | | | | |
| Niger | | KVx100-2 KVx30-309-6G KVx61-74 TN27-80 | Sudano-Sahelian zone | | | | | |
| Nigeria | Sampea-7 (IAR-48) (IAR-339-1) | | Sudano-Guinean savanna zone | | | | | |
| | (INV-000-1) | TVx3236 IT81D-994 | Sudano-Guinea savanna zone | | | | | |
| Senegal | | IS86-275 B 89 | Sahelo-Sudanian zone | | | | | |
| Tchad | IT81D-994 KN-1 TVx3236 | TN88-63 | Sudano-Sahelian zone | | | | | |
| Togo | Vitoco (IT81D-985) (Vita-5) | IT81D-1137 | Coastal, transition and Guinea savanna zones | | | | | |
| Central African Republic | KN-1 TVx 1948-01F | | Transition and Guinea savanna zones | | | | | |

Bibliothèque UA/SAFGRAD 01 BP. 1783 Ouagadougeu 01 Tél. 30 - 60 - 71/31 - 15 - 98 Burkina Faso

TABLE 6. LIST OF THE 1988 COWPEA MONITORING TOUR PARTICIPANTS.

| Country | Name of Scientist | Address |
|----------------|-------------------------|---|
| BURKINA FASO | 1. Dr. Sérémé Paco | Cowpea Pathologist CRAF, O1 B.P. 476 Ouagadougou O1 |
| CAPE VERDE | 2. Mr. Carlos Silva | Cowpea Agronomist INIA, B.P. 50, Praia |
| GUINEA BISSAU | 3. Mr. Malam Sadjo | Cowpea Agronomist MDR/DEPA, C.P. 71 Bissau |
| GUINEA CONAKRY | 4. Dr. F. L. Guilavogui | Cowpea Entomologist IRAG-MEN, B.P. 1003 Conakry |
| NIGER | 5. Mr. Adamou Moutari | Cowpea Breeder INRAN, B.P. 429 Niamey |
| SENEGAL | 6. Mr. Cissé Ndiaga | Cowpea Breeder ISRA/CNRA, B.P. 53 Bambey |

TABLE 7. LIST OF THE 1990 COWPEA MONITORING TOUR PARTICIPANTS

| Country | Name of Scientist | Address |
|--------------|------------------------|--|
| BENIN | 1. Dr. J. Detongnon, | Cowpea Breeder SRCV-Niaouli B.P. 3, ATTOGON |
| BURKINA FASO | 2. Dr. C. Dabire (Mrs) | Cowpea Entomologist CRAF, 01 B.P. 476 OUAGADOUGOU 01 |
| | 3. Mr. J. Ouedraogo | Cowpea Breeder INERA, 01 B.P. 7192 OUAGADOUGOU 01 |
| CAMEROON | 4. Mr. G. N'Toukam | Cowpea Entomologist IRA, B.P. 33, MAROUA |
| GAMBIA | 5. Mr. M. Bojang | Cowpea Agronomist Yundum Agric. Research Station P.O. Box 739, Yundum |
| GHANA | 6. Dr. M.O. Akyaw | Cowpea Entomologist Crops Research Institute P.O. Box 3785 Kumasi |
| NIGER | 7. Dr. S.D. Maiga | Cowpea Entomologist INRAN, B.P. 429, Niamey |
| NIGERIA | 8. Mr. A.A. Zaria | Cowpea Breeder IAR/ABU, PMB 1044, Zaria |
| | 9. Dr. 0.0. Olufajo | Cowpea Agronomist IAR/ABU, PMB 1044, Zaria |

TABLE 8. LIST OF THE PARTICIPANTS TO THE RENACO'S SEMINAR HELD IN NOVEMBER 1988 AT IITA, IBADAN, NIGERIA

| Country | Name of Scientist | Address |
|--------------|-----------------------|--|
| BURKINA FASO | Dr. C. DABIRE (Mrs) | Cowpea Entomologist, CRAF, 01 B.P. 476, Ouagadougou 01, |
| | Mr. OUEDRAOGO J. | Cowpea Breeder, INERA, 01 B.P. 7192, Ouagadougou 01 |
| CAMEROON | Mr. NTOUKAM, G. | Cowpea Entomologist B.P. 33, Maroua, |
| GHANA | Dr. OWUSU-AKYAW, M. | Cowpea Entomologist Crops Research Institute P.O.Box 3785, Kumasi, |
| NIGER | Dr. ADAM Toudou | Cowpea Pathologist, INRAN, B.P 429, Niamey |
| | Mr. HAMMA Hassane | Cowpea Pathologist, INRAN, B.P. 429, Niamey |
| NIGERIA | Prof. LELEJI, O.I. | Cowpea Breeder, IAR/ABU, PMB 1044, Zaria |
| | Dr. AMATOBI, A. M. | Cowpea Breeder, IAR/ABU, Kano |
| | Prof. EMECHEBE, A. M. | Cowpea Pathologist IAR/ABU, PMB 1044, Zaria |
| | Mr. ODION, C. E. | Cowpea Agronomist, IAR/ABU, Kano |
| SENEGAL | Dr. BAL, A.B. | Cowpea Entomologist, CNRA, B.P. 53, Bambey |
| | Mr. NDIAGA C. | Cowpea Breeder ISRA/CNRA, B.P. 55, Bambey |

TABLE 9. LIST OF THE PARTICIPANTS TO THE RENACO'S GROUP TRAINING COURSE AT THE INERA RESEARCH STATION, KAMBOINSE/OUAGADOUGOU IN 1989.

| Country | Name of Scientist | Address |
|----------------|------------------------|--|
| BENIN . | 1. Dr. J. Detongnon | Cowpea Breeder Station RCV-Niaouli, B.P. 3 ATTOGON |
| COTE D'IVOIRE | 2. Mr. Adou Amalaman | Cowpea Agronomist IDESSA-DCV, BP 635, Bouake 01 |
| GUINEA BISSAU | 3. Mr. Abu Biai | Cowpea Agronomist M.D.R.E Agricultura C.P. 71, Bissau- DEPA/CENEMAC, Contuboel |
| GUINEA CONAKRY | 4. Dr. F.L. Guilavogui | Cowpea Entomologist IRAG-MEN B.P. 1003, Conakry |
| MALI | 5. Mr. Kodio Ondié | Cowpea Breeder IER/DRA/SRCVO, B.P. 438, Sotuba |
| | 6. Mme D. N. Yaro | Cowpea Entomologist, IER/DRA/SRCVO, B.P. 438, Sotuba |
| | 7. Mr. D. Sogodogo | Cowpea Agronomist IER/DRA/SRCVO B.P. 438, Sotuba |
| | 8. Mr. S.O. Katilé | Cowpea Pathologist, IER/DRA/SRCVO B.P. 438, Sotuba |
| NIGER | 9. Mr. A. Moutari | Cowpea Breeder INRAN, B.P. 429, Niamey, |
| TCHAD | 10. Mr. Ouéitar Gam | Cowpea Agronomist Projet CHD82/003/PNUD/FAO B.P. 101, Gassi |

Workshop: Scientific information and technology exchange:

During the 1989 joint maize-cowpea workshop held in Lome, Togo, in March 1989, a day and a half was dedicated to scientific communication. Thirty original scientific papers by maize and cowpea national and international scientists were presented and discussed during the workshop. All aspects of maize and cowpea research activities were covered. Also scientists had the opportunity to interact with one another for five days. Forty-three cowpea scientists attended the Lome workshop (Table 10).

The proceedings of the workshop were published in two volumes - technical papers and country reports.

National and IITA-GLIP cowpea research activities were presented and discussed by all participants.

Visits to national programmes:

Seasonal visits to national programmes by either the Steering Committee members, Lead Centers Staff, IITA-GLIP scientists or the Cowpea Network Coordinator offered an informal on-the-spot training opportunity to national scientists and support staff by enabling them to discuss cowpea production technologies.

The following countries were visited by either the Network Coordinator or RENACO national scientists or IITA-GLIP scientists:

- In 1987: Burkina Faso, Guinea Conakry, Mali, Mauritania, Niger, Nigeria, Senegal and Togo;
- In 1988: Burkina Faso, Cameroon, Cape Verde, Niger, Nigeria, Senegal and Togo;
- In 1989: Benin, Burkina Faso, Côte d'Ivoire, Ghana, Guinea Bissau, Mali, Niger, Nigeria and Togo;
- In 1990: Burkina Faso, Cape Verde, Central African Republic, The Gambia, Mali, Niger, Nigeria, Senegal, Tchad.

TABLE 10. LIST OF NATIONAL AND INTERNATIONAL COWPEA SCIENTISTS WHO ATTENDED THE 1989 WORKSHOP AT LOME, TOGO, IN MARCH 1989.

| Country | Name of Scientist | Address |
|----------------|----------------------|---|
| BENIN | 1. Jean DETONGNON | Cowpea Breeder, SRCV-Niaouli, B.P.3 ATTOGON |
| BURKINA FASO | 2. C. DABIRE (Mrs) | Cowpea Entomologist, CRAF, 01 B.P. 476, Ouagadougou 01 |
| | 3. Jeremy OUEDRAOGO | Cowpea Breeder, INERA, 01 BP 7192, Ouagadougou 01, |
| | 4. Michel SEDOGO | Cowpea Agronomist, INERA, 01 B.P.7192, Ouagadougou 01, |
| CAMEROON | 5. Moffi TA'AMA | Cowpea Entomologist IRA/USAID/CRSP, B.P. 33, Maroua |
| CAPE VERDE | 6. C. E. P. SILVA | Cowpea Agronomist, MDR-DEPA B.P. 50, Praia |
| COTE D'IVOIRE | 7. Adou AMALAMAN | Cowpea Agronomist IDESSA, B.P 635, Bouake 01, |
| GAMBIA | 8. Musa BOJANG | Cowpea Agronomist Dept. of Agric. Research Yundum Research Station P.O. Box 739, Yundum |
| GHANA | 9. Asafu AGYEI | Cowpea Agronomist GGDP/CRI, Box 3785, Kumasi |
| | 10. G. A. AMANKWA | Cowpea Breeder, GGDP/CRI, Box 3785, Kumasi |
| | 11. Thimoty KIPO | Crops Research Institute P.O Box 3785, Kumasi |
| | 12. A. A. MAHAMA | Cowpea Breeder, CRI/NAES Box 52 or 483, Tamale |
| | 13. M. O. AKYAW | Cowpea Entomologist CRI, Box 3785, Kumasi |
| GUINEA BISSAU | 14. I. MIRANDA (Mrs) | Cowpea Agronomist MDR/DEPA, C.P. 71, Bissau |
| GUINEA CONAKRY | 15. F.L. GUILAVOGUI | Cowpea Entomologist IRAG. B.P. 1003, Conakry |
| MALI | 16. D. SOGODOGO | Cowpea Agronomist, ICRISAT B.P. 34, Bamako |
| | 17. B.A. KANTE (Mrs) | Seed Technologist IER, BP 438, Bamako |
| | 18. Ondie KODIO | Cowpea Breeder IER, B.P. 438, Bamako |

TABLE 10 (CONT'D-1). LIST OF NATIONAL AND INTERNATIONAL COWPEA SCIENTISTS WHO ATTENDED THE 1989 WORKSHOP AT LOME, TOGO, IN MARCH 1989.

| Country | Name of Scientist | Address |
|--------------|-------------------------|---|
| MAURITANIA | 19. R'Chid SIDI | Agronomist CNRADA, BP 22, Kaedi, |
| NIGER | 20. Toudou ADAM | Cowpea Pathologist Faculté d'Agronomie B.P. 10960, Niamey |
| | 21. Hassane HAMMA | Cowpea Pathologist INRAN, BP 240, Maradi, |
| | 22. Alzouma INEZDANE | Cowpea Entomologist Université de Niamey F.S. Département de Biologie B.P. 10662, Niamey |
| | 23. Adamou MOUTARI | Cowpea Breeder INRAN, B.P. 429, Niamey, |
| NIGERIA | 24. J.K. ADU | Microbiologist, IAR/ABU, PMB 1044, Zaria |
| | 25. C.I. AMATOBI | Cowpea Entomologist Agricultural Research Station, IAR, P.O. Box 1062, Kano, |
| | 26. K.A. ELEMO | Agronomist IAR/ABU, PMB 1044, Zaria |
| | 27. Prof. A.M. Emechebe | Cowpea Pathologist IAR/ABU, PMB 1044, Zaria |
| | 28. O.O. OLUFAJO | Cowpea Agronomist, IAR/ABU, PMB 1044, Zaria |
| SENEGAL | 29. N. CISSE | Cowpea Breeder ISRA, B.P. 55, Bambey |
| TCHAD | 30. C. D. BICHARA | Cowpea Agronomist, Station de Gassi, BP 441, N'Djamena |
| TOGO | 31. C. A. AGBOBLI | D.R.A., B.P. 2318, Lome, |
| | 32. A. DUYIBOE (Mrs) | Cowpea Agronomist D.R.A. B.P. 2318, Lomé |
| | 33. Mr. Toky PAYARO | Cowpea Agronomist RPAA, B.P. 218, Kara, |
| IITA, IBADAN | 34. E.F. DEGANUS | Administrator, ICP, IITA, PMB, 5320 Ibadan, |
| | 35. L.E.N. JACKAI | Cowpea Entomologist IITA, PMB 5320, Ibadan |
| | 36. G.O. MYERS | Cowpea Breeder, GLIP IITA, PMB 5320, Ibadan |

TABLE 10 (CONT'D-2). LIST OF NATIONAL AND INTERNATIONAL COWPEA SCIENTISTS WHO ATTENDED THE 1989 WORKSHOP AT LOME, TOGO, IN MARCH 1989.

| Country | Name | of Scientist | Address |
|---------|------|------------------|---|
| | | | |
| | 37. | B. R. NTARE | Cowpea Breeder, IITA/ICRISAT Sahelian Center, BP 12404, Niamey |
| | 38. | S.R. SINGH | Cowpea Entomologist Director, GLIP, PMB 5320, Ibadan |
| | 39. | Joseph Benah SUH | Cowpea Entomologist IITA, PMB 5320, Ibadan |
| USAID | 40. | Gerbrand KINGMA | Breeder, C/O O1 BP 1783, USAID/SAFGRAD, Ouagadougou O1 |
| | 41. | James C. SENTZ | Breeder USAID/IITA PMB 5320, Ibadan |
| SAFGRAD | 42. | Taye BEZUNEH | Physiologist, Director of Research, OAU/STRC, O1 B.P. 1783, Ouagadougou 01, |
| | 43. | Nyanguila MULEBA | Cowpea Agronomist RENACO Coordinator IITA/SAFGRAD,01 BP 1495, Ouagadougou 01 |

5) Network Impact

The impact of the network can be viewed on several grounds as follows:

- Management of research activities: A strong link has been established between SAFGRAD Coordination Office (SCO) and the Directors of Research of participating countries. The Council of Directors meet to review network activities and to establish guidelines to be followed or implemented by the networks. Through an Oversight Committee (emanating from the Council of Directors), the Council monitors and oversees the activities of networks. The Oversight Committee meets once a year. Thus, the Directors have been very active and responsive to all network activities (Steering Committee meetings, monitoring tours, workshops, training and regional trials) by either encouraging the contribution and the participation of their scientists and/or hosting meetings. In many countries, steps are underway towards specializing some scientists in cowpea research work (as opposed to a scientist or group of scientists working on several crops). It should be noted that without the full cooperation of the Directors of research, the success of the network in any form would not have been possible.
- Cowpea research: The greatest impact of the cowpea network is the renewed interests and total commitment of national programs to cowpea research activities. Sixty national scientists throughout West and Central Africa are not only enthusiastic in carrying out their respective responsibilities, but are also very keen in collaborating with one another within the network area and IITA in developing appropriate technologies meeting farmers needs and requirements. Thus, the linguistic barrier that has always separated anglophone and francophone countries from learning from one another has been broken!

Within each country, an unprecedented strong link has been established between cowpea scientists and peasant farmers through the farming system research scientists and extension workers. This has resulted in the conduct of multilocational trials and on-farm testings and release of new cultivars (Table 5 and Appendix 2) with several others in the pipe line for release (Appendix 3).

New varieties have been developed with the following attributes:

. Striga resistant varieties: The varieties shown on Table 11 were identified to be resistant to Striga gesnerioides and are being incorporated in good agronomic background.

TABLE 11. COWPEA VARIETIES RESISTANT TO STRIGA IN WEST AND CENTRAL AFRICA

| Name of variety | Origin | Pedigree | Country for which it is resistant to Striga | National programs incorporating it in good agronomic background |
|-----------------------------|-----------------|-----------------------------------|--|--|
| - Gorom Local (Suvita-2) | Burkina Faso | A selection from a landrace | Burkina Faso, Mali Senegal | Burkina Faso, Mali |
| - B301 | Botswana | _ | Burkina Faso, Mali Senegal, Niger, Nigeria, Benin | Burkina Fasl, Mali Niger, Nigeria |
| - IT82D-849 | IITA-Ibadan | | Burkina Faso, Mali, Senegal, Niger, Nigeria, Benin | Burkina Faso |
| - TN93-80 | Niger | Landrace | Burkina Faso, Mali, Senegal, Niger, Nigeria | |
| - TN121-80 | Niger | Landrace | Burkina Faso, Mali, Senegal, Niger, Nigeria | |
| - KVx61-1 | Burkina Faso | - | Burkina Faso, Mali | Burkina Faso |
| - KVx61-74 | Burkina Faso | - | Burkina Faso, Mali | Burkina Faso |
| - IT81D-994 | IITA-Ibadan | - | Burkina Faso, Nigeria | - |

. Drought resistant varieties:

- Gorm Local (SUVITA-2) (Burkina Faso)
- 58-57 (Senegal)

- TN88-63 (Niger)

- KVx 30-309-6G (Burkina Faso) - KVx 396-4 (Burkina Faso) - IS86-275 (Senegal)

. Varieties adapted to drought and excess moisture

- KVx 398-18 and KVx 396-4 (Burkina Faso)

. Aphids resistant varieties

- IT82E-25, IT83S-742-2, IT85D-3577 (IITA, Ibadan)

. Bruchid resistant varieties

- IT84S-275-9, KVx 30-6467-5-10K, IT84S-2246 (IITA, Ibadan and Burkina Faso)

In conclusion, the impact of the Network on agricultural production and development will largely depend on the extent to which technologies developed by RENACO Lead and International Agricultural Research Centers are transferred to local farmers.

Technology transfer does not simply mean moving technologies, say, from Point A (Experiment Station) to Point B (Farmers' fields). It also includes the adoption of the technologies by farmers. Therefore, such technologies must be proven to be more profitable and meeting the needs and requirements of the targeted farmers. To achieve this, scientists must, therefore, familiarize themselves with the problems and contrainsts confronting farmers, so that they can design the appropriate research methodologies for the development of the most efficient and beneficial technologies for farmers immediate use.

Due to the longtime neglect by policy makers, agricultural research in West and Central Africa has not been geared specifically to meeting the needs and requirements of peasant farmers. In order to remedy this situation, the SAFGRAD West and Central African Cowpea Network underlined this point as one of its top priority research effort since its inception in March, 1987. Consequently, Workshops, monitoring tours, Seminars and other training activities have been regularly carried out with the view of identifying the basic constraints limiting cowpea production and the best ways to go about solving them.

To this effect, five RENACO Lead Centres were identified in 1987 and became fully and actively operational in 1988. A sixth Lead Center was added in 1989. IITA core activities, which have been redeployed at two sub-stations located at strategic points: Niamey, Niger in the Sahel (in collaboration with ICRISAT) and Kano, Nigeria in the Sudan savanna (in collaboration with the Institute for Agricultural Research (IAR)), offer technical

backstopping for the conduct of relevant research and technology development, etc., for the interest of the semi-arid West and Central African region. It is gratifying to note that such technologies were put out for regional testing in 1989 and have been reported in the 1989-90 regionl trial preliminary results to be of outstanding performance.

With further efforts and investment on training activities, greater and relevant research achievements shall be obtained and transferrable technologies shall be developed for the enhancement of increased cowpea productivity and production in the not-toodistant future.

V. - FUTURE THRUST OF NETWORK PROGRAM AND AREAS OF WHICH FINANCIAL SUPPORT IS REQUIRED

Besides drought, heat, <u>Striga</u>, <u>Alectra</u> and disease resistance or tolerance, cowpea research in West and Central Africa has a long way to go. The use of chemical poisons in controlling insect pests in order to incerease cowpea yield from 200-300 kg/ha to 1000-1500 kg/ha is not a viable option for the generally poor African farmer.

The African economy is not sound enough to accept the massive use of chemical products in its agricultural production system. African scientists are therefore, confronted with the chanllenge of finding relatively cheap ways requiring minimum or no input. With this challenge, the most feasible approach is to embark on a mass breeding program, incorporating all genes and conferring resistance or tolerance to all major physical, chemical and biological cowpea production constraints in good agronomic backgrounds. The success of this minimum input strategy will no doubt boost cowpea yield of 600-1000 kg/ha in the near future.

With the ever increasing African population and given the importance of cowpea diet (supplies about 50% of the needed proteins) in low income African families, every effort must be made to up-grade cowpea production in the next 5-10 years. This dream can only come true if each national program is able to put in place a multidisciplinary team of cowpea scientists comprising a breeder, an agronomist, a pathologist, an entomologist and a social scientist. Since most national programs, perhaps with the exception of Nigeria and Ghana, do not have this facility, training effort cannot be neglected in the network program of activities.

APPENDIX-1

LIST OF NATIONAL COWPEA SCIENTISTS - RENACO

| | ntry/ e of Scientist | Qualification | Crops Research Time Areas on co (% | wpea |
|----------------------|--|---|---|-----------------------|
| BEN | IN | | | |
| 1. 2. 3. 4. | Jean Detongnon Moustapha Adamou Kouessi Aihou David Arodokoun | Cowpea Breeder (Ph.D) Soil Scientist (Ing. Agr) Agro-chemist (Ing. Agr) Entomologist (Ing. Agr) | Cowpea breeding Cowpea agronomy Cowpea agronomy Cowpea entomology | 100 40 30 30 |
| BUR | KINA FASO: | | | |
| 5. 6. | Issa Drabo Clementine Dabire | Cowpea Breeder (M.Sc) Cowpea Entomologist (Dr. 3eme C.) | (On-Ph.D study leave) Cowpea entomology | 100 |
| 7. 8. 9. | Jeremy Ouedraogo Paco Sereme Gnissa Konate | Cowpea Breeder (Ing. Agr) Phytopathologist (Dr./Ing) Virologist (Dr. D'Etat) | Cowpea breeding Cowpea pathology Cowpea virology | 10 3 4 |
| CAM | EROON | | | |
| | Georges Ntoukam Chevalier Endondo | Entomologist (M.Sc) Agronomist (Ing. agr) | Cowpea entomology (On-M.Sc study leave) | 10 10 |
| CAP | E VERDE: | | | |
| 12. | Carlos Silva | Agronomist/Breeder (B.Sc) | Cowpea agronomy/ | 4 |
| COT | E D'IVOIRE: | | | |
| 13. | Adou Amalaman | Agronomist (Diploma) | Cowpea agronomy | 10 |
| THE | GAMBIA: | | | |
| 14. | Musa Bojang | Agronomist (B.Sc) | Cowpea agronomy | 10 |
| TIA | NTA . | | | |
| GHA | | | | |
| | NYANKPALA STATION | | | |
| | K.O. Marfo M.A. Assibi P.B. Tanzubil | Legume Breeder Legume Breeder (B.Sc) Entomologist (M.Sc) | Ph.D candidate (On-M.Sc study leave) Cowpea entomology | 10 |

APPENDIX. - 1 (CONT'D-1):

LISTE OF NATIONAL COWPEA SCIENTIST - RENACO

| | ntry/ e of Scientist | | Crops Research Time spen on cowpea (%) | |
|---------------------------------|--|--|---|------------------------------|
| | KWADASSO/KUMASI ST | PATION | | |
| 19. 20. 21. 22. 23. | B. Asafu Agyei G.A. Amankwa Stella Ennin M.O. Akyaw J.K. Twumasi V.J. Affun J.N.A. Agyei | Legume Breeder Legume Breeder (M.Sc) Agronomist Entomologist (Ph.D) Pathologist (Ph.D) Entomologist (M.Sc) Agronomist (M.Sc) | (On study leave) Cowpea breeding (On-M.Sc study leave) Cowpea entomology Cowpea pathology Cowpea entomology Cowpea agronomy | 100 |
| GUI | NEE BISSAU | | | |
| 25. | Abu Biai | Agronomist (Diploma) | Cowpea agronomy | 100 |
| GUI | NEA CONAKRY | | | |
| 26. | F.L. Guilavogui | Entomologist (Ph.D) | Cowpea entomology | 100 |
| MAL | I | | | |
| 28. 29. | Ondie Kodio Mamadou Toure Aliou Traore D.N. Yaro (Mrs) | Cowpea breeder (Ing. Agr) Cowpea Breeder (Ing. Agr) Cowpea breeder (Ing. Agr) Cowpea entomologist (M.Sc) | (On-M.Sc study leave) (On-Ph.D study leave) Cowpea breeding Cowpea entomology | 100 100 100 80 |
| 31. | Seriba O. Katile | Pathologist (Ing. Agr) | Cowpea pathology | 40 |
| 32. | Diakalia Sogodogo | Cereal Agronomist (Ing. Ag | gr) Cowpea agronomy | 40 |
| MAU | RITANIA | | | |
| | Sidi Fall Sidi R'Chid | Plant Breeder (Ing. Agr) Agronomist (Diploma) | (On-M.Sc study leave) Cowpea agronomy | 40 |
| NIG | ER | | | |
| 36. 37. 38. | Issaka Maga Adamou Moutari Maman Nouri Ahamadou N'Diaye Adam Toudou | Cowpea Breeder (Ing. Agr.) Cowpea Breeder (Ing. Agr) Agronomist (Ing. Agr) Entomologist (Ing. Agr) Phytopathologist (Ph.D) | (On-Ph.D study leave) Cowpea breeding Cowpea agronomy Cowpea entomology Cowpea pathology | 100 100 40 40 75 |
| | Hassane Hamma Alzouma Indesdane | (University) Phytopathologist (Ph.D) Entomologist (Ph.D) (University) | Cowpea pathology Cowpea entomology | 50 |
| | Oumarou Moussa Seyni D. Maīga | Seed Technologist (Ing.Agr Entomologist (Ph.D) | Cowpea seed technology Cowpea entomolgy | 40 |

Varieties released by different RENACO national cowpea programs since 1987.

| Country/Name of variety | Origin | Areas of Adaptation | Year relea- sed | Quantity of seeds released (kg/ha) | Areas cultiva- ted in 1990 | Yield poten- tial (kg/ha) | Yield under farmers' conditions (kg/ha) | r Remarks |
|---|-------------|----------------------------------|-----------------------|---|-------------------------------------|------------------------------------|--|--|
| BURKINA FASO: TVx3236 | IITA | 300-1000 mm | 1987 | 2.500 | 250 | 1.500 | 800-1000 400-600 | Pure crop mixed cropping Amount of seed requested less than what is produced by extension services. |
| CAPE VERDE: KN-1 | | | | | | | | |
| Local Santiago | _ | | _ | - | _ | | - | |
| (IT82E-16) | | Northern Ghana Southern Ghana | 1987 1987 | 1100 100 | 23,000 29,000 | 1,700 | 800-1200 1000 | Good yield potential but low price paid in market Good yield potential but low price paid in market |
| GUINEA BISSAU IT82E-9) Bambey-21) | - | | - | _ | _ | - | | - |
| GUINEA CONAKRY | | | | | | | | |
| | ITA/SAFGRAD | Lower Guinea Medium Guinea | 1990 1990 | 600 700 | 40 46 | 1.000 | | High rainfall zone High Altitude |
| IT83D-338-1 IT84S-2246-4 | | Upper Guinea Upper Guinea | 1989 1990 | 500 800 | 33 53 | 650 800 | 350 500 | Low temperature Southern Sudan Climate |

APPENDIX. - 1 (CONT'D-2)

LIST OF NATIONAL COWPEA SCIENTISTS - RENACO

| | ntry/ me of Scientist | Qualification | Crops Research Areas | Time spent on cowpea (%) |
|--|---|--|---|---|
| NIG | ERIA | | | |
| 45. 46. 47. 48. 49. 50. | A.A. Zaria A.M. Emechebe E.C. Odion C. Amatobi O.O. Olufajo J.K. Adu J.A.Y. Sheybayan S.T.O. Lagoke | Cowpea Breeder (M.Sc) Cowpea Pathologist (Ph.D) Cowpea Agronomist (M.Sc) Cowpea Entomolgist (Ph.D)) Cowpea Agronomist (Ph.D) Microbiologist (Ph.D) Weed Scientist (M.Sc) Weed Scientist (Ph.D) | Cowpea breeding Cowpea pathology Cowpea agronomy Cowpea entomology Cowpea agronomy Microbiology Weed science Weed science | 100 100 100 100 100 40 40 20 |
| SEN | EGAL | | | |
| 53. | Ndiaga Cisse Samba Thiaw Mamadou Gaye | Cowpea Breeder (M.Sc) Agronomist (M.Sc) Microbiologist (Dr./Ing) | Cowpea breeding Cowpea agronomy Microbiology | 100 100 40 |
| TCH | AD | | | |
| 55. | Mr. Daniel Valenghi | | | - |
| TOG | Q | | | |
| 57. 58. 59. | Poda Assiongbou K. Adri Daou Ekou-Edi Yawo A. Akpaloo Akossiwa Duyiboe | (Ing. Agr) Cowpea Agronomist (Ing.Agr) Entomologist (Ing.Agr) Entomologist (Ing.Agr) Agronomist | Seed multiplication Cowpea agronomy Cowpea entomology Cowpea entomology Cowpea agronomy | 50 50 50 50 |

APPENDIX. - 2 (CONT'D)

Varieties released by different RENACO national cowpea programs since 1987.

| Country/Name of variety | Origin | Areas of Adaptation | Year relea- sed | Quantity of seeds released (kg/ha) | culti- | Yield potential (kg/ha) | Yield under farmers' conditions (kg/ha) | Remarks |
|---|------------------|---------------------------|-----------------------|---|-------------------------|-------------------------------|--|--|
| MAURITANIA: IT835-343-5-5 Suvita-2 KVx256-K17-11 | SAFGRA SAFGRA | AD Attabi | 1987/88 | 25 <mark>000</mark> | 1000-2000 500 500 | 1.500 1.000 1.000 | 500-700 300-400 | Drought resistant. Acceptability difficult because of seed color. |
| NIGERIA: Sampea-7 (IAR-48)x Ife brown(local | Air Nigeria | savanna & forest zones | 1987 | 10000 | 75000 | 1500-2500 | 600 | Area cultivated is an estimate. It may actually be more than 75.000 ha. |
| SENEGAL: IS86-275 | ISRA Senegal | Sahelian | u | - : | 20000-30000 | 2200-2500 | 600-1100 | |
| rogo: rt81D-985 | IITA | Savannas | 1987-88 | - | ND | 1000-2000 | 900 | Pre-extension stage, also for sowing date |
| 58-146 | | e whole ountry | 1987-88 | - | ND. | 1100-1600 | 400-1000 | Still in pre-release stage in certains zones |

Varieties in a pre-extension stage in various RENACO national programs since 1987.

| Country/Name of variety | Origin | Area of adaptation | Potential areas of cultivation (ha) | potential | Yield in farmers' conditions (kg/ha) | Remarks |
|--|--------------------|-----------------------------------|--|---------------|---|-----------------------------------|
| BURKINA FASO: KVx30-309-6G | Burkina Faso | 300-900 mm | 110 | 1000 en pure | 800 | These areas of cultiva- |
| NVX30-303-00 | burkina raso | 300-900 mm | 110 | 450 en assoc. | 400 | tion are those covered b |
| KVx61-1 | -do- | -do- | 350 | 1500 | 900 | the 1990 on-farm trials |
| KVx396-4-4 | | 300-1200 mm | 350 | 1500 | 900 | and farmers field which |
| | | | | 450 | 400 | received seeds from our |
| KVx396-4-5 | -do- | -do- | 250 | 1500 | 900 | stocks. |
| | 7.9 | , | | 400 | 400 | |
| KVx396-18-10 | -do- | -do- | -do- | 1500 | 400 | |
| CAMEDOOM | | | | 500 | 400 | |
| CAMEROON IT81D-994 | IITA | Sudan & Norther Guinea Savanna | rn - | 1200 | 400 | Extension stage |
| CAPE VERDE | | dariica bavaiiia | | | | |
| IT83D-442 | | _ | - | - | _ | - |
| Mississipi Silver | | - | - | _ | | |
| GHANA | | | | | | |
| IT81D-1137 | IITA/Ibadan | Savanna areas | The whole of Ghana | f 1700 | 900 | Highly acceptable seed coat color |
| IT83S-818 | IITA/Ibadan | -do- | -do- | 1000 | 650 | -do- |
| 11000 010 | IIIn/Ibadaii | -40- | -40- | 1000 | 050 | -40- |
| GUINEA BISSAU IT83-219 IT85D-3516-2 IT86D-498 IT87S-1390 IT85-3577 IT83D-889 TVx309-66 IS86-275N IS87-416N | nt sa - | - | | | | |

APPENDIX. - 4

LIST OF NATIONAL AND INTERNATIONAL COWPEA SCIENTISTS WHO ATTENDED THE 1987 WORKSHOP AT OUAGADOUGOU, BURKINA FASO, IN MARCH, 1987.

| Country | Name of Scientist | Address |
|----------------|---------------------|--|
| BENIN: | J. Detongnon | Cowpea Breeder, SRCV-Niaouli B.P. 3, ATTOGON |
| BURKINA FASO: | Issa Drabo | Cowpea Breeder, CRAF, 01 B.P. 476, Ouagadougou 01 |
| CAMEROUN | Georges Ntoukam | Cowpea Entomologist IRA, B.P. 33, Maroua |
| COTE D'IVOIRE | Adou Amalaman | Agronomist, IDESSA, B.P. 635, Bouaké 01 |
| GAMBIE | Tijan Jallow | Yundum Agric. Research Station P.O. Box 786, Banjul |
| GHANA | Antoni Assibi | Legume Breeder Nyankpala Agric. Research Station, Box 52, Tamale |
| | A. Atuahene-Amankwa | Crops Research Insitute P.O. Box 3785, Kumasi |
| GUINEE BISSAU | Malam Sadjo | MDR/DEPA, C.P. 71, Bissau |
| GUINEE CONAKRY | Saikou S. Bah | IRAG, B.P. 1003, Conakry |
| MALI | Kodio Ondié | Cowpea Breeder, IER/DRA/SRCVO, B.P. 438, Bamako |
| MAURITANIA | Sidi R'chid | Agronomist, CNRADA, B.P. 22, Kaedi |
| NIGER | Tijan Jallow | INRAN, B.P. 429, Niamey |
| | Alzouma Inesdane | Entomologist, INRAN, B.P. 429, Niamey |
| NIGERIA | Ono Leleji | Agronomist, IAR/ABU, PMB 1044 Zaria |
| | G.O. Aballu | IAR/ABU, PMB 104, Zaria |
| | A.M. Emechebe | Pathologist, IAR/ABU, PMB 1044, Zaria |
| SENEGAL | Mamadou Ndiaye | ISRA, B.P. 3120, Dakar |

Varieties in a pre-extension stage in various RENACO national programs since 1987.

| Country/Name of variety | Origin | Area of adaptation | Potential areas of cultivation (ha) | potential | Yield in farmers' conditions (kg/ha) | Remarks |
|-------------------------|--------------------------|--------------------------|--|-------------|---|---|
| GUINEA CONAKR | Y | | | | | |
| IT84S-2246-4 | IITA/SAFGRA | D Lower Guinea | 20 | 1000 | 500 | Insecticide Protection |
| IT82E-32 | -do- | -do- | 5 | 590 | 400 | - |
| IT86D-1048 | -do- | -do- | 5 | 675 | 400 | |
| IT86D-1056 | -do- | -do- | 5 | 600 | 350 | |
| IT85F-867-5 | | Upper Guinea | 5 | 800 | 500 | 스 교육도 있는 생각을 보면 있다. 전체를 받으는 <u>.</u> |
| MAURITANIA | | | | | | |
| IT86V-472 | SAFGRAD | Valley/Senegal | _ | 1600-2000 | 400-600 | All these varieties have been |
| IT82D-544-4 | -do- | -do- | _ | -do- | -do- | accepted for their bruchid |
| IT81D-897 | -do- | -do- | _ | -do- | -do- | tolerant characteristics |
| IT82D-716 | -do- | -do- | _ | -do- | -do- | volotano onaraovoribolos |
| IT82D-927 | -do- | River valley | _ | 10000-20000 | 10000 | Interesting for forage production |
| TVx1948-MF | -do- | and dams | - | 1500 | -do- | and supply of green leaves for |
| ISRA | - | - | - · | 1000 | 5.800 | human consumption. |
| NIGERIA: | | | | | | |
| TVx3236 | - | Sudano-Guinea savanna | Sudano-Guine & savanna & forest zone | | 600k kg/ha | |
| IT81D-994 (| IVu−1190 x | | TOTOBO ZONO | | | |
| TVu | 16 x TVu2027) TVu625) | -do- | -do- | -do- | -do- | |
| TOGO | 144020) | | | | | |
| IVx 1850-01E | IITA The | whole country | _ | 1000-1300 | 600-1000 | Yield of all varieties are |
| T81D-985 | | whole country | _ | 1000-2000 | 900 en | highly variable, depending |
| | | ept savanna zone | | 1000 1000 | milieu | on the region and crop |
| | 0110 | 1 - 20.000 | | | humide | season as well as cultural |
| 58-146 | ISRA The | whole country | - | 1100-1600 | 400-1000 | |
| IT83S-818 | IITA Rég | ion des plateaux | _ | 1000-1300 | | Potential cultivated areas come |
| IT82E-16 | | Maritime | | 1400-1700 | | under the extension services unit. Non available yet. |

APPENDIX. - 4 (CONT'D)

LIST OF NATIONAL AND INTERNATIONAL COWPEA SCIENTISTS WHO ATTENDED THE 1987 WORKSHOP AT OUAGADOUGOU, BURKINA FASO, IN MARCH, 1987.

| Country | Name of Scientist | Address |
|--------------|---------------------|--|
| TCHAD | Yaouga Djekounkosse | Ministère du Développement Rural, Direction Générale de L'Agriculture, Bureau de la Recherche Agronomique, B.P. 441, N'Djamena |
| TOGO: | Akossiwa Duyiboe | Agronomist, Direction de la Recherche Agronomique, B.P. 2318, Lome |
| SAFGRAD | Kassu Yilala | Farming System Research P.O. Box 476, Kamboinse Ouagadougou |
| | Tadesse Kibreab | Farming System Research P.O. Box 476, Kamboinse Ouagadougou |
| | Toky Payaro | SAFGRAD/RPAA, B.P. 218, Kara |
| | J.B. Suh | Entomologist, IITA/SAFGRAD 01 B.P. 1495, Ouagadougou 01 |
| | T. Bezuneh | Director of Research, OAU-SAFGRAD, O1 B.P. 1783, Ouagadougou O1 |
| IITA/SAFGRAD | V.D. Aggarwal | Cowpe Breeder, IITA/SAFGRAD 01 B.P. 1495, Ouagadougou 01 |
| | N. Muleba | Cowpea Agronomist and Cowpea Network Coordinator IITA/SAFGRAD, O1 B.P. 1495 Ouagadougou O1 |
| JSAID/BF | A. Fleming | USAID/BF |
| | M. Sullivan | USAID/BF |
| IITA/ICRISAT | B.R. N'Tare | Cowpea Breeder, ICRISAT Sahelian Center, B.P. 12404 Niamey |
| ITA/IBADAN | B.B. Singh | Cowpea Breeder, IITA-Kano sub station, PMB 3112, Kano |

APPENDIX. - 5

RENACO STEERING COMMITTEE MEETINGS

| No. Order | Date | Venue p | Number of articipants |
|-----------------------------------|---------------------|-----------------------------|-----------------------|
| 1st Steering Committee meeting | 23-27 March, 1987 | Ouagadougou Burkina Faso | 7 |
| 2nd Steering Committee meeting | 9-12 November, 1987 | Ouagadougou Burkina Faso | 14 |
| 3rd Steering Committee meeting | 28-31 March, 1988 | Ouagadougou Burkina Faso | 15 |
| 4rd Steering Committee meeting | 7-11 November, 1988 | Zaria, Nigeria | 13 |
| 5th Steering Committee meeting | 23-24 March, 1989 | Lome, Togo | 12 |
| 6th Steering Committee meeting | 6-10 November, 1989 | Ouagadougou Burkina Faso | 13 |
| 7th Steering Committee meeting | 26-30 March, 1990 | Ouagadougou Burkina Faso | 9 |
| 8th Steering Committee meeting | 5-9 November, 1990 | Cotonou, Benin | 14 |

AFRICAN UNION UNION AFRICAINE

African Union Common Repository

http://archives.au.int

Department of Rural Economy and Agriculture (DREA)

African Union Specialized Technical Office on Research and Development

1987-03

REGIONAL APPROACH TO COWPEA RESEARCH IN WEST AND CENTRAL AFRICA (RENACO)

MULEBA, N.

AU-SAFGRAD

https://archives.au.int/handle/123456789/8837

Downloaded from African Union Common Repository