

Neutron Shell Strengths at $N = 152$ and towards $N = 162$

F.P. Heßberger^{1,2} and D. Ackermann¹

¹GSI, Darmstadt, Germany; ²HIM, Mainz, Germany

Determination of the shell strengths is an important feature to characterize the properties of superheavy elements and to test the power of theoretical predictions. Recently ground-state masses of several nobelium ($Z=102$) and lawrencium ($Z=103$) isotopes have been measured directly with high precision at SHIPTRAP [1, 2]. The results have been used to determine the strength of the $N = 152$ subshell for $Z = 102$ and $Z = 103$ [2]. The experimental data from [2] are compared in fig. 1 with the results from a microscopic - macroscopic approach [3] and a self-consistent calculation using the SLy4 - force [4]. The latter has been used recently to calculate properties of neutron stars, e.g. the relation between mass and radius (see e.g. [7]). This circumstance demonstrates that properties of the nuclear force (or strong force, respectively) derived from the structure and stability of nuclei can be used to describe astrophysical phenomena or in general, phenomena where the strong force plays an essential role (e.g. quark-gluon plasma). Vice versa, information on the strong force obtained from 'such other' studies will have a feedback on the description of nuclei. Superheavy nuclei are a specific, but of course not the only one, laboratory for such studies, as they exist only due to a delicate balance between the nuclear force and the Coulomb force.

Another source of information on shell strength in the transactinide region are the α - decay chains passing through ^{252,254}No. Based on the measured masses of these isotopes [1] the $2n$ -binding energies of the $N-Z = 50$ nuclei could be determined up to ²⁶⁶Hs [5]. The results are compared with predicted values in fig. 2. It is seen that the $2n$ -binding energies obtained from the microscopic - macroscopic models [3, 6] describe the experimental trend, which shows a maximum value at $N = 152$, at least qualitatively, while the agreement between experimental and calculated values is somewhat better for [3] than for [6]. The self-consistent calculation using the SkP force, however, does not show a local maximum at $N = 152$ and gives too low values for $N \leq 152$. Towards higher neutron numbers all calculations show a steeper increase of the $2n$ - binding energies than measured. This may indicate that the $N = 162$ neutron shell might be weaker or stronger localized. To check this possibility it is necessary to estimate the $2n$ - binding energy of ²⁷⁰Ds, which requires the identification of the so far unknown nuclide ²⁶⁸Ds. Based on production cross-sections for ^{269,270,271}Ds measured previously at SHIP it can be expected to be produced with $\sigma \approx 1-2$ pb in the reaction ²⁰⁷Pb(⁶²Ni,n)²⁶⁸Ds.

This finding is in-line with the trend which is indicated by the spontaneous fission half-lives, which show a slower in-

crease towards $N = 162$ than the values predicted in [8], which are based masses and shell effects from [3].




Figure 1: Experimental shell strength parameters of No isotopes in comparison with results from calculations.




Figure 2: Experimental $2n$ - separation energies in comparison with results from calculations.

References

- [1] M. Dworschak et al., PRC 81, 064312 (2010)
- [2] E. Minaya Ramirez et al., Science 337, 1207 (2012)
- [3] R. Smolanczuk, A. Sobczewski, Proc. EPS Conf. 'Low Energy Nucl. Dyn.', St. Petersburg (Russia), 1995
- [4] J. Dobaczewski et al., arXiv:nucl-th/0404077v1 (2004) (<http://fuw.edu.pl/~dobaczew/thodri/thodri.htm>)
- [5] D. Ackermann et al. GSI-Scientific Report 2011, PHN-NUSTAR-SHE04
- [6] P. Möller et al. ADNDT 59, 185 (1995)
- [7] M. Hempel, Physik in unserer Zeit 01/2014(45), 12 (2014)
- [8] R. Smolanczuk et al., PRC 52, 1871 (1995)