

## Application of Calorimetric Low Temperature Detectors (CLTD's) for Precise Stopping Power Measurements of Heavy Ions in Matter

A. Echler<sup>1,2,3</sup>, P. Egelhof<sup>1,3</sup>, P. Grabitz<sup>1,3</sup>, H. Kettunen<sup>4</sup>, S. Kraft-Bermuth<sup>2</sup>, M. Laitinen<sup>4</sup>, K. Müller<sup>2</sup>, M. Rossi<sup>4</sup>, W.H. Trzaska<sup>4</sup>, and A. Virtanen<sup>4</sup>

<sup>1</sup>GSI, Darmstadt, Germany; <sup>2</sup>University of Gießen, Germany; <sup>3</sup>University of Mainz, Germany; <sup>4</sup>University of Jyväskylä, Finland

Precise data on electronic stopping powers, i.e. specific energy loss  $dE/dx$ , for heavy ions are of high interest in various fields of research [1]. Unfortunately the accuracy of theoretical ab-initio  $dE/dx$ -calculations is nowadays still unsatisfactory. Therefore, predictions of stopping powers are usually calculated by semi-empirical computer codes, which are based on best fits to available experimental data. To improve the reliability of these calculations, precise  $dE/dx$  data are needed for various projectile-target combinations and over a wide range of projectile energies, in particular for heavy projectiles at low energies, as measured data are scarce in that region [2]. However, for high ion masses and low energies ionization based energy detectors, that are commonly used in experiments for stopping power determination [3, 4], suffer from incomplete energy detection, resulting in pulse height defect and a relatively poor energy resolution. As CLTD's provide substantially better energy resolution and linearity for heavy ion detection, with the absence of any pulse height defect [5, 6], this type of energy detectors has the potential to increase sensitivity and accuracy for  $dE/dx$  measurements and to extend the accessible energy range towards lower energies.

For that purpose a CLTD array, that has been developed at GSI in the past years [5, 6, 7], has been used to replace the Si-detector in an established setup for  $dE/dx$  measurements at the K-130 cyclotron at the University of Jyväskylä [3, 8], and to perform measurements with 0.05–1 MeV/u <sup>131</sup>Xe-ions in different absorber materials. As an example the measured electronic stopping power of Xe ions in carbon is displayed in fig. 1. A summary of all results for carbon as well as nickel and gold absorbers can be found in [9], where also a more detailed description of the experiment is given. The obtained  $dE/dx$  values show in general a good agreement with previously measured data in the higher energy range for all studied absorber materials, what provides a consistency check of the new experimental setup. Moreover the application of CLTD's for the present investigations allowed to considerably reduce calibration errors as compared to previous measurements with Si-detectors [10], and to extend the datasets to lower energies, not accessible before.

### References

- [1] ICRU Report 73, *Journal of the ICRU* 5, No 1 (2005).
- [2] H. Paul, web site: [www.exphys.jku.at/stopping/](http://www.exphys.jku.at/stopping/)
- [3] W.H. Trzaska et al., *Nucl. Instr. Meth. B* **195**, 147, (2002).




Figure 1: Electronic stopping power of Xe ions in carbon obtained from the present experiment [9] together with literature values that have been taken from the electronic library for stopping powers of H. Paul [2] (Trz09 [10], Gs98 [11], Pap78 [12]). The lines represent theoretical calculations with the semi-empirical computer code SRIM [13] and the ab-initio models CasP [14] and PASS [15].

- [4] J. Zhang, *Nucl. Instr. Meth. B* **196**, 1, (2002).
- [5] P. Egelhof and S. Kraft-Bermuth, *Topics in Applied Physics* **99**, 469, (2005).
- [6] S. Kraft-Bermuth et al., *Rev. Sci. Instrum.* **80**, 103304, (2009).
- [7] A. Echler et al., *J. Low Temp. Phys.* **167**, 5-6, 949, (2012).
- [8] A. Javanainen et al., *IEEE Trans. Nuc. Sci.* **54**, 4, 1158, (2007).
- [9] A. Echler et al., *J. Low Temp. Phys.* DOI 10.1007/s10909-013-1043-y (2014).
- [10] W.H. Trzaska et al., *Nucl. Instr. Meth. B* **267**, 3403, (2009).
- [11] H. Geissel et al., *various data summarized in electronic library of H. Paul* [2].
- [12] H. Pape et al., *Zeitschr. f. Phys. A* **286**, 159, (1978).
- [13] J.F. Ziegler, *Nucl. Instr. Meth. B* **268**, 11-12, 1818, (2010), [www.srim.org](http://www.srim.org).
- [14] G. Schiwietz and P.L. Grande, *Nucl. Instr. Meth. B* **273**, 1, (2012), CasP download: <http://www.casp-program.org/>
- [15] PASS-code based on *Binary theory of electronic stopping*, P. Sigmund and A. Schinner, *Nucl. Instr. Meth. B* **195**, 64, (2002)