

In-Medium strangeness dynamics at $\bar{\text{PANDA}}^*$

T. Gaitanos¹ and H. Lenseke^{1,2}

¹Inst. für Theor. Physik, Universität Giessen, Germany; ²GSI Darmstadt, Germany

Hypernuclear physics opens a unique opportunity to explore the hyperon-nucleon (YN) and the hyperon-hyperon (YY) in-medium interactions at terrestrial laboratories. Such studies are important for a better understanding of the in-medium interactions with strangeness degree of freedom, which is still uncertain, but of relevance for nuclear astrophysics [1]. The experimental knowledge on multi-strangeness hypernuclei has been so far scarce. However, recent experiments at GSI are very promising. In the HyPHI [2] experiment precise production rates of single- Λ ($S = -1$) hypernuclei were measured. An abundant production of double- Λ ($S = -2$) clusters is also expected in the $\bar{\text{PANDA}}$ experiment [3] at FAIR.

We investigate theoretically the multi-strangeness dynamics within the Giessen-BUU (GiBUU) approach [4]. In particular, we have studied antiproton-induced reactions allowing for reactions of the secondary Ξ -beam on a second target. The formation of double- Λ hypernuclei occurs in the Ξ -interaction with the second target [5]. Two issues are of importance for the production of double-strangeness hypermatter, see Fig. 1. At first, the absorption of the Ξ -beam inside the target matter decreases with increasing energy. This is due to the strong decrease of the elementary $\Xi N \rightarrow \Lambda\Lambda$ channel [5]. Secondly, the abundance of bound Λ -hyperons also strongly decreases with rising incident Ξ -energy. This is mainly due to the repulsive vector field which becomes more pronounced as the particle energy increases. These effects lead to a rather strong energy dependent rise of the double- Λ hypernuclear production cross section. Thus low-energetic cascade beams should be used at $\bar{\text{PANDA}}$ in order to obtain high production rates of double-strangeness hyperfragments [5].

So far bare interactions for elementary YN, YY -channels have been used in transport approaches. However, at $\bar{\text{PANDA}}$ the strangeness dynamics takes place at densities closely up to saturation. We have studied in-medium effects on various YN -processes by solving the Lippmann-Schwinger equation with a Pauli-exclusion operator for intermediate states as the leading-order in-medium effect [6, 7]. The elementary YN cross sections are indeed influenced by in-medium effects, in particular, at low energies, as shown in Fig. 2 (similar effects occur for quasi-elastic channels with strangeness exchange).

In the $S=-2$ sector the situation is still very controversial theoretically [8]. Presently we are investigating the influence of various calculations on $S=-2$ -channels [9] on the strangeness production in \bar{p} -induced reactions and extending our studies to the $S=-3$ sector by accounting for the




Figure 1: GiBUU results for the time dependence of all and free Λ s and of Ξ s for central Ξ -induced reactions on Cu-target at beam energies as indicated [5].

formation of Ω -baryons [10]. The preliminary results are very promising concerning a possible formation of multi-strangeness bound systems at $\bar{\text{PANDA}}$. We emphasize the relevance of our theoretical results for the future activities at FAIR.




Figure 2: Energy dependence of elastic in-medium cross sections for ΛN and ΣN scattering at various densities as indicated [7].

References

- [1] J. Schaffner, I. Mishustin, Phys. Rev. C53 (1996) 1416.
- [2] C. Rappold, *et al.*, Nucl. Phys. A622 (2010) 231.
- [3] A.S. Lorente, *et al.*, Phys. Lett. B 697 (2011) 222.
- [4] O. Buss, *et al.*, Phys. Rep. 512 (2012) 1.
- [5] T. Gaitanos, *et al.*, Nucl. Phys. A881 (2012) 240.
- [6] T. Gaitanos, *et al.*, Nucl. Phys. A914 (2013) 405.
- [7] A. Obermann, Masters Thesis, 2011, Univ. Giessen, unpublished.
- [8] J.K. Ahn, *et al.*, Phys. Lett. B 633 (2006) 214.
- [9] T.A. Rijken, Y. Yamamoto, nucl-th/0608074; Y. Fujiwara, *et al.*, Phys. Rev. C64 (2001) 054001.
- [10] T. Gaitanos, *et al.*, in preparation.

*Work supported by BMBF, DFG, HIC for FAIR, and GSI-JLU Giessen collaboration agreement