

Manuscript received April 24, 2022; Revised April 30, 2022; accepted May 1, 2022; date of publication June 30, 2022

Digital Object Identifier (DOI): <https://doi.org/10.35882/teknokes.v15i2.199>

This work is an open-access article and licensed under a Creative Commons Attribution-ShareAlike 4.0 International License ([CC BY-SA 4.0](https://creativecommons.org/licenses/by-sa/4.0/))



Spectro-analytical Research of Selected Heavy Metals (Cu, Cd, Cr, and Pb) in Four Different Single-use Plastics Commonly in Contact with Food from Sokoto, Nigeria

A.I Umar¹; Sarkingobir Y², and Dikko, M³

¹ Department of Biochemistry, Sokoto State University, Sokoto, Nigeria

² Department of Biology, Shehu Shagari College of Education, Sokoto, Nigeria

³ Sultan Abdulrahman College of Health Technology Gwadabawa, Sokoto State Nigeria

#Corresponding author: A.I Umar (email: superoxidisedismutase594@gmail.com)

ABSTRACT Single use plastics are types of plastics discarded after one usage. They are everywhere you go because of their wide range of applications in life. In plastics, additives are incorporated to supplant properties of plastics. One of the common additives is the heavy metals, which could leach out along the plastics lifecycle, thereby creating a problem. This study was stimulated to determine the levels of heavy metals in selected single-use-plastics commonly in contact with food (as containers or wrappings or relations) in Sokoto. The Atomic Absorption Spectroscopy methodology was utilized to determine the levels of copper, chromium, lead, zinc and cadmium. The results for determination of selected heavy metals in selected single use plastics from Sokoto State University, Nigeria were revealed using atomic absorption spectroscopy. Therein, the concentrations of copper (Cu) ranges from 0.335+0.002 to 0.540+0.002(ug/kg), levels of cadmium (Cd) ranges from 0.0551+0.003 to 0.0041+0.0003(ug/kg), levels of chromium (Cr) ranges from 0.0959 +0.004 to 0.0265+0.001(ug/kg), levels of lead (Pb) ranges from -1.791+0.02 to -0.0706+0.008(ug/kg). Thus, the results revealed disparity in the levels of the analyzed heavy metals in the selected single use plastics. However, Cu concentration was the most elevated in plastic bottle (PB) and lowest in Takeaway (T); Cd was highest in Straw (S), and lowest in PB; Cr was highest in Ice cream cup (IC) and lowest in S; Pb was highest in PB, and lowest in S. The concentrations of Copper, Chromium, Lead, and Cadmium in plastic bottle, Takeaway, Straw and Ice cream cup, have shown disparity. However, Cu concentration was the most elevated in plastic bottle (PB) and lowest in Takeaway; Cd was highest in Straw, and lowest in PB; Cr was highest in Ice cream cup and lowest in S; Pb was highest in PB, and lowest in S. They levels determined are within the permissible limits.

INDEX TERMS Heavy metals, single-use plastics, straw, lead, cadmium, copper, food

I. INTRODUCTION

Plastic are polymers with large molecular weight. The plastic was originated from Greek words “plasticos” meaning moulded. The term plastics refer to the property of a material having ductility during the process of manufacturing [1] [2]. The nature of plastic renders them the ability to be converted into several products that are diversely applicable in human endeavour [3] [4]. Plastics are diverse, like their functions.

Some plastics are thermosets, thermoplastics, micro plastics, macro plastics, nanoplastics, elastomers, bioplastics and single use plastics [1]. Commonly available plastics include: polyethene Terephthalate, found in soft drinks, salad trays, mouth wash bottle; high density polyethylene, found in food packaging, rest mats, cosmetics, raincoats; polyvinyl chloride, found in toys, dolls, beach balls, rubber medicine bottles; polypropylene, found in yoghurt bottles, medicine

bottle; polystyrene, found in cassette cases, coffee cups, cutlery, computers; and polypropylenes, found in yoghurt bottles, medicine bottles [1][2][4]. From 1950 to date, plastic industry has increased to a very significant extent and serves as an important economic sector of the world [2] [5]. Plastic Nobel properties such as malleability, durability, low-cost, multiple application and knowledge gap has led to a drastic growth in sector of plastic production over the years. Globally, in 2015 around 322 million tons of plastics were produced, leading to 6,300 million tons of plastic waste, 9% of it was recycled, 12% incinerated and 79% was in landfills or leaked into the environment [6][7][8]. If nothing tangible was implemented, the trend will skyrocket [2]. However, one of the leading types of plastics of concern is the single use-plastics. Single use-plastic are plastic products that design to be use and dispose once. Examples include: light-weight plastic bags, disposable utensils, beverage containers, coffee capsules, straws, food wrappers, takeaways, plastic bag, and drink bottles [9]. They are the leading plastics produced and used every day and are the most prevalent in the waste produced and discarded in our environment [10][11]. Similarly, every plastic is manufactured as a mixture of specific monomer resins and additives. In some of the additives added to these plastic materials are heavy metals [2]. Forsooth, there are revelations that shows these heavy metals in the plastics can leach to the environment or other contacts (such as food, water, human body) and might elicit negative effects [2][12]. All elements on earth can either be metals or nonmetals based on sets of criteria. Metals are large group of substances, which are opaque, form alloys, conduct electricity and heat. They are usually malleable [13]. Metals occur naturally in the environment, and in the crust. They vary significantly between different regions in spatial variations of background concentrations. Out of the 92 naturally occurring elements, about 30 metals and metalloids are potentially harmful to humans, Be, B, Li, Al, Ti, V, Cr, Mn, Co, Cu, Ni, As, Se, Sr, Mo, Pd, Li, Cd, Sn, Te, Sb, Cs, Ba, Lu, Pt, Au, Hg, Pb and Bi [14][15][16]. Heavy metals are the family of metals having an atomic weight above 40.04. Heavy metals enter the environment via natural or anthropogenic processes, such as in natural weathering, mining, soil erosion, industrial discharge, pest control methods, sewage effluents, air pollution fallout e.t.c. [13]. Heavy metals explored from the earth crust or industrial processes are released to the environment. Therein, humans contacted them through ingestion (drinking or eating) or inhalation (breathing) or absorption by the skin/other body openings [8][17].

Heavy metals cause a wide array of harmful effects to the body. Therein, main general effects of heavy metals resulted when they elicit neurotoxicity, oxidative stress, loss of cellular functions, carcinogenesis and cell damage [13]. That is why this work tried to determine the levels of heavy metals

in some selected single use plastics in Sokoto, Nigeria. Plastic materials are remarkably resourceful and are utilized in variety of applications. More than half of the plastics produced are disposed when they are used once [18]. These single use plastics are ubiquitous in our environment and everywhere you go in Sokoto state and many other places in the country [2][8]. Therefore, they make majority of our waste which find it ways into the environment and exert negative effects to living organisms including humans, because of their component monomers and additives [1]. One of the additives found in plastics are the heavy metals, which are added deliberately and through impurities during production [2][19][20].

However, heavy metals have been shown to cause negative effects, such as cancer effect, neurotoxicity, oxidative stress, cell damage, apoptosis etc [13]. Parable, cancerous breast biopsies show higher level of nickel, chromium, cadmium, mercury and lead. Lead is proven to cause neurotoxicity, learning and land defects, behavioural effects, infertility, miscarriage, reduced fertility, etc. mercury is linked to nervous system toxicity, immune system toxicity, etc [13]. Nickel is a common cause of allergy [21]. Mercury induced nephrotoxicity, neurotoxicity, etc [22]. Arsenic is associated gastro intestinal disturbance, abnormal heart beat, respiration defects, enzyme defects, etc. Equally, other heavy metals exert their harmful effects on the body [21]. Several reports from certain organizations and reports availed that plastics contain heavy metals as additives or as impurities [2][3]. However, across the globe few empirical studies have reported levels of heavy metals in an array of plastics types. Most of the researches are dealing with levels of heavy metals in electronics and single use plastics in contacts with food or plastics in waste [20][23]. For instance, A study on plastic toys imported from China to Northwestern Nigeria determined lead, cadmium, nickel, copper, zinc, chromium, cobalt and manganese with 17% of them having high levels more than USFDA recommendations; which poses threat to children exposed to these toys, and public health [19]. [20] performed an evaluation of lead and copper of selected components of waste computers, the results therein, exceeded the Toxicity Threshold Limit Concentration (TTLC) used in characterizing wastes as toxic, which in turn could be toxic to humans and environment in the inappropriate water disposal fashion. [24] performed a comparative assessment of heavy metals in plastic bottles and sachets. Therein, manganese, iron, copper, cobalt, cadmium, and nickel were detected. Virgin plastics contain very low heavy metals compared to used plastics. The metals are either added intentionally, or through impurities. [23] Reported the levels of heavy metals in selected plastic bags from China. Heavy metals content of polyethylene, High Density Polyethylene, Low Density Polyethylene and Polyvinyl Chloride bags revealed the presence of lead, chromium, cadmium, arsenic, copper, and

TABLE 1
Results of determination of heavy metals (Cu, Cd, Cr, and Pb) in four different single use plastics collected from Sokoto State Nigeria.

Heavy Metal(Ug/Kg) Plastic Type	Copper	Cadmium	Chromium	Lead
Plastic Bottle	0.1335+0.002b	0.0041+0.0003b	0.0544+0.003a	-1.791+0.02b
Straw	0.0540+0.002a	0.0551+0.003a	0.0265+0.001a	-0.0706+0.008c
Take Away	0.0612+0.003a	0.0097+0.003c	0.0687+0.002a	-0.4095+0.2 B
Ice Cream Cup	0.0978+0.001a	0.0050+0.0002c	0.0959+0.004a	-0.8347+0.09b

diarrhea, stomach pain, vomiting, damage central nervous system, immune damage, psychological disorders etc [5][18] 21,]. The finding is similar to result of Khan and Khan (2015)[18]; but lower that of Omolaoye et al., (2010)[19] from Nigeria and [23]Alam et al., (2019).

Cu is an essential element to humans, but critical doses lead to anemia, acne, cancer, diabetes, allergies, hair loss, depression, panic attack, tooth decay, kidney problems, and vitamin deficiencies. In this study, the levels of Cu found were below the limit of 1 ppm. The Cu levels (0.335+0.002 to 0.540+0.002 (ug/kg) found in this study are smaller than the levels found by Olubanjo et al., (2015)[20] in plastic computers in Nigeria. Also Omolaoye et al (2010)[19] found lower levels of Cu from soft plastic toys imported from China to Nigeria.

Cr levels determined were below the permissible limits. Cr is a micronutrient required for certain purposes in the body. It helps in glucose mobilization from the blood to cells, fats metabolism. On the other hand, chromium compounds are toxic at low concentration to plants and animals. They behave as carcinogens, causes breathing problems, asthma, cough etc. Long-term exposure to Cr may lead to liver, kidney damage [18]. The results (0.0959 +0.004 to 0.0265+0.001 (ug/kg), of Cr was lower than the results of other studies reported by Omolaoye et al (2010)[19] and Olubanjo et al (2015)[20].

Pb too had shown very lower levels in the analyzed single use plastics of this study. The levels are below the permissible limit, albeit there is no known safe lead level to humans. It has ability to affects all organs in the body [21][24]. The finding is similar to Khan and Khan (2015)[18]. The lead levels in this study (-1.791+0.02 to -0.0706+0.008 (ug/kg) are lower than the findings by Olubanjo et al (2015)[20] in plastic computers in Nigeria; and Omolaoye et al (2010)[19] who found lower levels of Pb from soft plastic toys imported from China to Nigeria.

Noteworthy, heavy metals including Pb, Cd, Cr, and Cu are natural elements that have a relatively high atomic mass and high density compared to water. Even through heavy metals are naturally present in our environment, but their environmental pollution and exposure to humans is speed up by anthropogenic processes of humans. One of the uses of heavy metals is the application in plastics. They are used as additives in plastics. Parable, Cr, Pb, Cd and Cu are used as

colorants in plastics to improve the properties of the polymers. Other source of heavy metals to plastics is the ability plastics to absorb heavy metals from the environment whenever there is contact. The concerns about the presence of heavy metals in plastics are many. The metals can easily leach into the contacts (food, plant, and animal) when there is heat or rise in temperature because they are already loosely attached to the polymer chain. Moreover, the metals can be release when the plastics are heated, burned or incinerated. The waste of plastics has ability to dispose heavy metals in soils and water where they are discarded. Therefore there is need to carefully monitor heavy metals present in plastics especially the single use ones [8][23][24,].

Implications and disadvantages of finding heavy metals in plastics, especially those that are used once and discarded, are diverse. Single use plastics are indiscriminately discarded in the study area, in turn they act as major sources of pollution and contamination with heavy metals affecting humans, plants and environment [28]-[30]. Presence of heavy metals in single use plastics migrate to the humans in contacts causing devastating health effects, due to their toxicity at low concentration, persistence and bioaccumulation potential [28]. Humans and animals can be affected through consumption (feeding), burning of plastics (especially the volatile metals like lead, cadmium, zinc, chromium, zinc etc), intake of plants grown in polluted soils, drinking water, and many others[23][25]. Invariably, quantification of environmental load of toxicants is an initial store in monitoring and management of pollutants. It will serve scientists, public health experts, policy makers, and regulatory measures or phase-out some heavy metals additives from plastics [28]-[32].

V. CONCLUSION

The main objective of this work was to conduct Spectro-analytical Research of selected heavy metals (Cu, Cd, Cr, and Pb) in four different single-use plastics commonly in contact with food from Sokoto, Nigeria. The concentrations of Copper, Chromium, Lead, and Cadmium in plastic bottle, Takeaway, Straw and ice cream cup, have shown disparity. However, Cu concentration was the most elevated in plastic bottle (PB) and lowest in Takeaway; Cd was highest in Straw, and lowest in PB; Cr was highest in Ice cream cup and lowest in S; Pb was highest in PB, and lowest in S. All they

levels determined are within the permissible limits. Based on this study the following recommendations are made: a) Further studies are needed to ascertain the levels of heavy metals in single use plastics, b) There is need to seek for legislation to screen the levels of heavy metals in Nigeria as there are no safe levels for most of the heavy metals such as Lead, Cadmium, c) there is need for bodies that should screen the levels of heavy metals in plastics to guard public health, d) If possible, alternative less harmful chemicals should be used as plasticizers or colorants in single use plastics, e) people should adhere to the norms of reuse, improvise, recycle, refuse, to limits the plastics pollution.

REFERENCE

- [1] O.A. Alabi, K.I. Ologbunjaye, O., Awosolu, O.E., Alalade, "Public and Environmental health effects of plastic wastes disposal: A review," *J. Toxicol. Risk Assess.* vol.5, no.2, pp.1-13, mon. 2019, doi.
- [2] Y. Sarkingobir, M. Dikko, S. Aliyu, U. Tambari, A. Adamu, I.A. Salau and M.A. "The dangers of plastics to public health: A review". *NIPES J. Sci. Technol. Res.*, vol.2, no.2, pp.195-200, 2020.
- [3] R.U. Halden, "Plastics and health risks. *Ann. Rev. Pub. Health*, 31: 179-194, 2010.
- [4] C.M., Rochman, M.A., Browne, B.S., Halpern, B.T., Hentschel, E., Hoh, et al. "Policy: Classify plastic waste as hazardous," *Nature*, vol. 494, pp.169-171, 2013.
- [5] European Union "Commission Regulation no 1935 and articles intended to come into contact with food," 2011.
- [6] J. Kaiser, "The dirt on ocean garbage patches," *Sci.* vol. 328, pp. 1506, 2010.
- [7] R. Geyer, J.R. Jambeck, and K.L. Law, "Production, use, and fate of all plastics ever made," *Sci. Advancem.*, vol.3, no.e1700782, 2017.
- [8] Y. Sarkingobir, A.M. Bello, and H.M. Yabo. "Harmful effects of plastics on air quality," *Acad. Lett.*, no.2967. <https://doi.org/10.20935/AL296.2021>.
- [9] S. Muhammad, I., Ashiri, D.K. Ibrahim, D. Salawu, I.N. Muhammad and A. Muhammad "Determination of some heavy metals in wastewater sediment of artisanal gold in local mining site of Abare Area Nigeria," *J. Environ. Treat. Techniq.*, vol.1, no.3, pp.174-182, 2014.
- [10] M.S. Ibrahim, A.J. Jafar, S.A.S. Abubakar, U. Saleh, L. Mahmud and A. Ibrahim. "Bioplastics as better alternative to petoplastics and their role in national sustainability: A review". *Advan. Biosci. Bioeng.*, vol.6, no.4, pp.63-70, 2017.
- [11] R.C. Thompson, C.J. Moore, F.S. vom Saal, and S.H. Swan. "Plastics, the environment and human health: Current consensus and future trends, philosophical transactions of the Royal Society," *Biol. Sci.*, vol.364, no.1526, pp.2153-2166, 2009.
- [12] R. Li, X. Lee, Z. Zhan, and R. Clifford, "Evaluation of heavy metal migration from different types of plastic food packaging materials into aqueous simulants using ICP-MS," *Math. Probl. Eng.*, vol.2, no.3, pp.1-5, 2014.
- [13] A.A. Adepoju-Bello, O.O. Oguntibeju, R.A. Adebisi, N. Okpala, and H.A.B. Coker, "Evaluation of the concentration of toxic metals in cosmetics products in Nigeria," *Afri. J. Biotechnol.*, vol.11, no.97, pp.16360-16364, 2012.
- [14] M. Bala, R.A. Shehu, and M. Lawal, "Determination of the level of some heavy metals in water collected from two pollution-prone irrigation areas around Kano Metropolis," *Bayero J. Pure Appl. Sci.*, vol.1, no.1, pp.36-38, 2008.
- [15] S., Khan, Q. Cao, Y.M. Zheng, Y.Z. Huang, and Y.G. Zhu, "Health risks of heavy metals in contaminated soils and food crops irrigated with wastewater in Beijing, China," *Environ. Pollu.*, vol.152, pp.686-692, 2008.
- [16] A. Turner. "Heavy metals, metalloids and other hazardous elements in marine plastic litter," *Mar. Pollu. Bull.*, vol.111, no.1-2, pp.136-142, 2016.
- [17] S.D. Guala, F. A. Vega and E.F. Covelo, "The dynamics of heavy metals in plant-soil interactions," *Ecol. Modell.*, vol.221, pp. 1148-1152, 2010.
- [18] S. Khan, and A. Khan, "Toxic heavy metal concentration in locally made plastic food container," *Int. J. Scien. Eng. Res.*, vol.6, no.6, pp.45-47, 2015.
- [19] J.A. Omolayo, A. Uzairu, and C.E. Gonna. "Heavy metal assessment of some sort plastic toys imported into Nigeria from China," *J. Environ. Chem. Ecotoxicol.*, vol.2, no.8, pp.26-130, 2010.
- [20] K. Olubanjo, O.M. Osibanjo, and I.C Nmorom, "Evaluation of Pb and Cu contents of selected component parts of waste personal computers". *J. Appli. Sci. and Environ. Manage.*, vol.19, no.3, pp. 470-477, 2015.
- [21] WHO. "Adverse health effects of heavy metals in children," 2011 www.who.int
- [22] Plastics Europe "Plastics-The Facts. An analysis of European plastics production, Demand and waste data," Brussels, Belg: Plast Eur <http://www.wikipedia.com/>, 2016.
- [23] O. Alam, L. Yang, and X. Yanchun, "Determination of the selected heavy metal and metalloid contents in various types of plastic toys," *J. Environ. Health Sci. Eng.* vol.17, pp. 161-170, 2019.
- [24] Aliyu A. Ochigbo, S.S. and Akoyi J.N. "Comparative assessment of the levels of some heavy metals in virgin and used water plastic bottles and sachets in Nigeria. *Afri. J. Chemi. Educ.*, vol.7, no.2, pp.93-104, 2017.
- [25] Okenwa-Ani, J.A. and Gimba, C.E., "Heavy metal levels in plastics housing of televisions: Is there a changing trend across the three decades (1980-2000)," *Iranian J. Toxicol.*, vol.13, no. 3, pp.27-32, 2019.
- [26] Seyaboh, E.I. Angaye T.C.N. Seiyaboh, and Z. "Assessment of heavy metal contaminants associated with locally processed beverages in reused plastic container". *Direct Research J. Biol. Biotechnol.*, vol.6, no.4, pp.34-36, 2020.
- [27] A.I. Umar, Y. Sarkingobir, Z. Shehu, and M. Dikko "Determination of micronutrients (zinc, iodine, selenium) in three different egg varieties from Sokoto, Nigeria" *Adamawa State Uni. J. Scien. Res.*, vol.10, no.1, pp.1-6, 2022.
- [28] Olagunju, T.E. Olagunju, A.O. Akawu, I.H. Ugokwe, C.U. "Quantification and risk assessment of heavy metals in groundwater and soil of residential areas around Awotan landfill, Ibadan, Southwest-Nigeria" *J. Toxicol. Risk Assess.*, vol.6, no.1, pp.1-12, 2020.
- [29] Jallo, I.U. Kodiya M.A. Molu M.A. "Waste management practices and its environmental implications in Hadejia Metropolis, Jigawa state, Nigeria," *Dutse J. Pure Appli. Sci.* vol.7, no.2b, pp.228-236.
- [30] Aliyu, S. "Stakeholders views on ways to reduce use of plastic bags and its environmental and public health impacts in Sokoto Metropolis, Sokoto state, Nigeria" *Int. J. Res. Scient. Innov.* vol.vii, no.iv, pp.196-200, 2020.
- [31] Sarkingobir, Y. Sahabi, M. Bello, S.Y. "Effects of plastic pollution: An overview," *Subsaha. Afri. Acad. Res. Publi.: J. Agri. Agricul. Tech.*, vol.22, no.1, pp.163-177, 2021.
- [32] C. Campanale, C. Massarelli, I. Savino, V. Locaputo, and V. F. Uricchio, "A detailed review study on potential effects of microplastics and concern on human health," *Int. J. Environ. Res. Public Health*, vol.17, no.1212, pp.1-26, 2020.

BIOGRAPHY



Umar A.I. The author is an indigene of Sokoto State, Nigeria. He has obtained Bachelor's Degree in Biochemistry as his background, and has obtained Msc later. He holds PhD in Biochemistry from Usmanu Danfodiyo University Sokoto, Nigeria. Dr Aminu Umar Imam/ Dr Umar AI is currently the Head of Department of Biochemistry Sokoto State University Sokoto, Nigeria. His interest is mostly in nutritional biochemistry.



Yusuf Sarkingobir was born in Gwadabawa, Sokoto state Nigeria. He holds BSc. Biochemistry from Usmanu danfodiyo University Sokoto, Nigeria; and MSc Public Health from Maryam Abacha American University of Niger Maradi, Niger Republic. Presently, he taught at Shehu Shagari Colege of Education Sokoto, Nigeria which is now upgraded by the Executive Governor to serve as Shehu Shagari University of Education Sokoto. He is also currently pursuing PhD in Public health from Crown University International Chartered Incorporated. He study Almajiri and Public health, determinants of health, plastics and public health, drugs/ substances abuse and public health, and open defecation.



Malami Dikko. The author, Dr malami Dikko Tambuwal was born in Tambuwal local government, Sokoto State, Nigeria. He holds BSc Microbiology, MSc and Phd Pharmacology from Usmanu Danfodiyo University Sokoto, Nigeria. Presently, he is the College Examination Officer, Sultan Abdurrahman College of Health Technology, Gwadabawa, Sokoto, Nigeria.