

Northumbria Research Link

Citation: Kolodziejczak, Anna, Guerrini-Rousseau, Lea, Planchon, Julien Masliah, Ecker, Jonas, Selt, Florian, Mynarek, Martin, Obrecht, Denise, Sill, Martin, Hirsch, Steffen, Sturm, Dominik, Waszak, Sebastian M, Ramaswamy, Vijay, Pentikainen, Virve, Demir, Haci Ahmet, Clifford, Steven C, Schwalbe, Ed, Massimi, Luca, Snuderl, Matija, Galbraith, Kristyn, Karajannis, Matthias A, Hill, Katie, Li, Bryan, White, Christine L, Redmond, Shelagh, Loizos, Loizou, Jakob, Marcus, Kordes, Uwe, Schmid, Irene, Hauer, Julia, Blattmann, Claudia, Filippidou, Maria, Scheurlen, Wolfram, Kontny, Udo, Grund, Kerstin, Sutter, Christian, Pietsch, Torsten, van Tilburg, Cornelis M, Frank, Stephan, Schewe, Denis M, Malkin, David, Taylor, Michael D, Tabori, Uri, Bouffet, Eric, Kool, Marcel, Sahm, Felix, von Deimling, Andreas, Korshunov, Andrey, Von Hoff, Katja, Kratz, Christian, Jones, David T W, Rutkowski, Stefan, Witt, Olaf, Bougeard, Gaelle, Pajtler, Kristian W, Pfister, Stefan M, Bourdeaut, Franck and Milde, Till (2022) MEDB-14. Clinical outcome of pediatric medulloblastoma patients with Li-Fraumeni syndrome. *Neuro-Oncology*, 24 (Supp_1). i107-i107. ISSN 1522-8517

Published by: Oxford University Press

URL: <https://doi.org/10.1093/neuonc/noac079.389>
<<https://doi.org/10.1093/neuonc/noac079.389>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/49500/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Northumbria
University**
NEWCASTLE



University**Library**

the course of LE, and to investigate its impact on long-term neurocognitive and behavioural outcome. **METHODS:** A French retrospective, multicenter study including 35 children under 5 years of age, treated between 2009 and 2017, with a median follow up of 72 months. All follow-up MRIs including assessment of the severity of the LE (Fazekas and CTCAE grading) and all NP evaluations were centrally reviewed. **RESULTS:** 25/34 evaluable patients presented a LE during follow up, in a median delay of 2 months (1 - 17 months) after the start of chemotherapy. Grade 2 and 3 abnormalities were correlated with higher cumulative dose of ITV -MTX ($p=0,01$). Full Scale IQ (FSIQ) and Wechsler indexes were in the average or low average of the reference population. FSIQ was deficient in 7/20 evaluable patients. Processing speed (PSI) was the most frequently impaired neurocognitive domain: 9/20 patients with borderline or very low score, all having received a significantly higher cumulative dose of ITV-MTX ($p=0,04$). A decrease in overall NP scores was observed in patients for whom grade 2 or 3 LE persisted at the end of follow-up with an average FSIQ estimated at 82.1 (SD 16.9) versus 94.2 (SD 20.6). This decrease was significant for PSI ($p=0,049$). LE and neurocognitive impairments were not correlated with a younger age at diagnosis. **CONCLUSION:** This study confirmed the responsibility of MTX, and in particular ITV-MTX therapy in the onset and, most often, persistence of LE and its association with neurocognitive disorders.

MEDB-14. CLINICAL OUTCOME OF PEDIATRIC MEDULLOBLASTOMA PATIENTS WITH LI-FRAUMENI SYNDROME

Anna Kolodziejczak^{1,2}, **Lea Guerrini-Rousseau**³, **Julien Masliash Planchon**⁴, **Jonas Ecker**^{1,5}, **Florian Selt**^{1,5}, **Martin Mynarek**^{6,7}, **Denise Obrecht**⁸, **Martin Sill**^{1,8}, **Steffen Hirsch**^{1,9}, **Dominik Sturm**^{1,5}, **Sebastian M. Waszak**^{10,11}, **Vijay Ramaswamy**^{12,13}, **Virve Pentikainen**¹⁴, **Haci Ahmet Demir**¹⁵, **Steven C. Clifford**¹⁶, **Ed Schwab**^{16,17}, **Luca Massimi**¹⁸, **Matiija Snuderl**¹⁹, **Kristyn Galbraith**¹⁹, **Matthias A. Karajannis**²⁰, **Katie Hill**²⁰, **Bryan Li**²⁰, **Christine L. White**²¹, **Shelagh Redmond**²², **Loizou Loizos**²³, **Marcus Jakob**²⁴, **Uwe Kordes**⁶, **Irene Schmid**²⁵, **Julia Hauer**²⁶, **Claudia Blattmann**²⁷, **Maria Filippidou**²⁸, **Wolfram Scheurlen**²⁹, **Udo Kontny**³⁰, **Kerstin Grund**⁹, **Christian Sutter**⁹, **Torsten Pietsch**³¹, **Cornelis M. van Tilburg**^{1,32}, **Stephan Frank**³³, **Denis M. Schewe**³⁴, **David Malkin**³⁵, **Michael D. Taylor**¹², **Uri Tabori**¹³, **Eric Bouffet**³⁶, **Marcel Kool**^{1,8}, **Felix Sahn**^{1,37}, **Andreas von Deimling**^{2,37}, **Andrey Korshunov**^{2,37}, **Katja Von Hoff**³⁸, **Christian Kratz**³⁹, **David T.W. Jones**^{1,40}, **Stefan Rutkowski**⁶, **Olaf Witt**^{1,5}, **Gaelle Bougeard**⁴¹, **Kristian W. Pajtler**^{1,5}, **Stefan M. Pfister**^{1,5}, **Franck Bourdeaut**⁴, **Till Milde**^{1,5}; ¹Hopp Children's Cancer Center (KiTZ), Heidelberg, Germany. ²CCU Pediatric Oncology, German Cancer Research Center (DKFZ) and German Cancer Consortium (DKTK), Heidelberg, Germany. ³Gustave Roussy Institute, Paris, France. ⁴SIREDO Pediatric Oncology Center, Curie Institute, Paris-Science Lettres University, Paris, France. ⁵Pediatric Oncology, Hematology, and Immunology, Center for Child and Adolescent Medicine, Heidelberg University Hospital, Heidelberg, Germany. ⁶Pediatric Hematology and Oncology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany. ⁷Mildred Scheel Cancer Career Center HaTriCS⁴, University Medical Center Hamburg-Eppendorf, Hamburg, Germany. ⁸Pediatric Neurooncology, German Cancer Research Center (DKFZ) and German Cancer Consortium (DKTK), Heidelberg, Germany. ⁹Institute of Human Genetics, University Hospital Heidelberg, Heidelberg, Germany. ¹⁰Centre for Molecular Medicine Norway (NCMM), Nordic EMBL Partnership, University of Oslo and Oslo University Hospital, Oslo, Norway. ¹¹Department of Neurology, University of California, San Francisco, CA, USA. ¹²Division of Neurosurgery, Program in Developmental and Stem Cell Biology, Arthur and Sonia Labatt Brain Tumour Research Centre, Hospital for Sick Children, Toronto, Ontario, Canada. ¹³Division of Hematology and Oncology, Hospital for Sick Children, Toronto, Ontario, Canada. ¹⁴Division of Hematology-Oncology and Stem Cell Transplantation, Children's Hospital, Helsinki University Hospital, Helsinki, Finland. ¹⁵Department of Pediatric Hematology-Oncology, Private Memorial Ankara Hospital, Ankara, Turkey. ¹⁶Wolfson Childhood Cancer Research Centre, Newcastle University Centre for Cancer, Newcastle, United Kingdom. ¹⁷Applied Sciences, Northumbria University, Newcastle, United Kingdom. ¹⁸Department of Pediatric Neurosurgery, Catholic University Medical School, Rome, Italy. ¹⁹Department of Pathology, New York University Langone Health, New York City, NY, USA. ²⁰Department of Pediatrics, Memorial Sloan Kettering Cancer Center, New York City, NY, USA. ²¹Victorian Clinical Genetics Services, Parkville, Australia. ²²Swiss Childhood Cancer Registry, Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland. ²³Pediatric Oncology/Hematology/Immunology at the Medical School of the University of Nicosia, Nicosia, Cyprus. ²⁴Department of Paediatric Haematology, Oncology and Stem-Cell Transplantation, University Hospital Regensburg, Regensburg, Germany. ²⁵Paediatric Haematology and Oncology, Dr. von Hauner Children's Hospital, Ludwig Maximilian University, Munich, Germany. ²⁶Pediatric Haematology and Oncology, University Hospital Carl Gustav Carus, Dresden, Germany. ²⁷Paediatric

Haematology, Oncology and Immunology, Olgahospital, Stuttgart, Germany. ²⁸Division of Pediatric Hematology-Oncology, First Department of Pediatrics, National and Kapodistrian University of Athens, "Aghia Sophia" Children's Hospital, Athens, Greece. ²⁹Paediatric Haematology and Oncology, Cnopfsche Paediatric Clinic, Nurnberg, Germany. ³⁰Department of Paediatric Haematology, Oncology and Stem-Cell Transplantation, Paediatric Clinic, University Hospital Aachen, Aachen, Germany. ³¹Neuropathology Institute, DGNN Brain Tumor Reference Center, University of Bonn Medical Center, Bonn, Germany. ³²Department of Pediatric Oncology, Hematology, Immunology and Pulmonology, Heidelberg University Hospital, Heidelberg, Germany. ³³Division of Neuropathology, Institute of Pathology, Basel University Hospital, Basel, Switzerland. ³⁴Department of Pediatrics, Otto-von-Guericke University Magdeburg, Magdeburg, Germany. ³⁵Genetics and Genome Biology Program, The Hospital for Sick Children, Toronto, Ontario, Canada. ³⁶The Arthur and Sonia Labatt Brain Tumour Research Centre, The Hospital for Sick Children, Toronto, ON M⁵G 1X⁸, Canada; Division of Haematology / Oncology, The Hospital for Sick Children, Toronto, Ontario, Canada. ³⁷Neuropathology, Institute of Pathology, Heidelberg University Hospital, Heidelberg, Germany. ³⁸Department of Pediatric Oncology and Hematology, Charité – Universitätsmedizin Berlin, Berlin, Germany. ³⁹Pediatric Haematology and Oncology, Hannover Medical School, Hannover, Germany. ⁴⁰Pediatric Glioma Research, German Cancer Research Center (DKFZ) and German Cancer Consortium (DKTK), Heidelberg, Germany. ⁴¹Inserm U1245 Cancer and Brain Genomics, UNIROUEN, Normandie Univ, and Department of Genetics, Rouen University Hospital, Rouen, France

PURPOSE: The prognosis for SHH-medulloblastoma (MB) patients with Li-Fraumeni syndrome (LFS) is poor. Due to lack of comprehensive data for these patients, it is challenging to establish effective therapeutic recommendations. We here describe the largest retrospective cohort of pediatric LFS SHH-MB patients to date and their clinical outcomes. **PATIENTS AND METHODS:** N=31 patients with LFS SHH-MB were included in this retrospective multicenter study. TP53 variant type, clinical parameters including treatment modalities, event-free survival (EFS) and overall survival (OS), as well as recurrence patterns and incidence of secondary neoplasms, were evaluated. **RESULTS:** All LFS-MBs were classified as SHH subgroup, in 30/31 cases based on DNA methylation analysis. The majority of constitutional TP53 variants (72%) represented missense variants, and all except two truncating variants were located within the DNA-binding domain. 54% were large cell anaplastic, 69% gross totally resected and 81% had M0 status. The 2-(y)ear and 5-(y)ear EFS were 26% and 8,8%, respectively, and 2y- and 5y-OS 40% and 12%. Patients who received post-operative radiotherapy (RT) followed by chemotherapy (CT) showed significantly better outcomes (2y-EFS:43%) compared to patients who received CT before RT (30%) ($p<0.05$). The 2y-EFS and 2y-OS were similar when treated with protocols including high-dose chemotherapy (EFS:22%, OS:44%) compared to patients treated with maintenance-type chemotherapy (EFS:31%, OS:45%). Recurrence occurred in 73.3% of cases independent of resection or M-status, typically within the radiation field (75% of RT-treated patients). Secondary malignancies developed in 12.5% and were cause of death in all affected patients. **CONCLUSIONS:** Patients with LFS-MBs have a dismal prognosis. This retrospective study suggests that upfront RT may increase EFS, while intensive therapeutic approaches including high-dose chemotherapy did not translate into increased survival of this patient group. To improve outcomes of LFS-MB patients, prospective collection of clinical data and development of treatment guidelines are required.

MEDB-15. DYNAMIC CHROMATIN ALTERATION INDUCES ONCOGENIC HIJACKING BY ESSENTIAL TRANSCRIPTIONAL FACTORS DURING SHH MEDULLOBLASTOMA TUMORIGENESIS

Ryo Shiraishi¹, **Gabriele Cancila**², **Kohei Kumegawa**³, **Patricia da Silva**⁴, **Owen Chapman**⁵, **Wang Wanchen**¹, **Maki Jami**¹, **Mikio Hoshino**¹, **Stefan Pfister**⁴, **Lukas Chavez**⁵, **Reo Maruyama**³, **Olivier Ayrault**², **Daisuke Kawauchi**¹; ¹National Center of Neurology and Psychiatry, Tokyo, Japan. ²Institut Curie, Paris, France. ³Japanese Foundation for Cancer Research, Tokyo, Japan. ⁴Hopp Children's Cancer Center Heidelberg (KiTZ), German Cancer Research Center (DKFZ) and Heidelberg University Hospital, Heidelberg, Germany. ⁵University of California San Diego, San Diego, USA

Medulloblastoma is a malignant brain tumor that occurs in the cerebellum, most frequently in children. Medulloblastoma is molecularly classified into four major groups, and therapies are now being developed according to the nature of these groups and subgroups. However, there are currently no effective molecularly targeted drugs for most of these groups. In recent years, we have been analyzing the genomes of medulloblastomas to identify genetic mutations involved in tumorigen-