



Seniority in quantum many-body systems

P. Van Isacker

► **To cite this version:**

P. Van Isacker. Seniority in quantum many-body systems. Symmetries in Nature : Symposium in Memoriam Marcos Moshinsky, Aug 2010, Cuernavaca, Mexico. 1323, pp.141-152, 2010, <10.1063/1.3537842>. <in2p3-00525728>

HAL Id: in2p3-00525728

<http://hal.in2p3.fr/in2p3-00525728>

Submitted on 12 Oct 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Seniority in quantum many-body systems

P. Van Isacker

Grand Accélérateur National d'Ions Lourds, CEA/DSM-CNRS/IN2P3,

B.P. 55027, F-14076 Caen Cedex 5, France

Abstract

The use of the seniority quantum number in many-body systems is reviewed. A brief summary is given of its introduction by Racah in the context of atomic spectroscopy. Several extensions of Racah's original idea are discussed: seniority for identical nucleons in a single- j shell, its extension to the case of many, non-degenerate j shells and to systems with neutrons and protons. To illustrate its usefulness to this day, a recent application of seniority is presented in Bose-Einstein condensates of atoms with spin.

PACS numbers: 03.65.Fd, 21.60.Fw, 21.60.Cs, 03.75.Fi

I. RACAH'S SENIORITY NUMBER

The seniority quantum number was introduced by Racah for the classification of electrons in an atomic ℓ^n configuration [1]. He assumed a spin-independent interaction \hat{V} between the electrons with the property

$$\langle \ell^2; LM_L | \hat{V} | \ell^2; LM_L \rangle = g(2\ell + 1)\delta_{L0}, \quad (1)$$

that is, there is no interaction unless the two electrons' orbital angular momenta ℓ are coupled to a combined angular momentum of $L = 0$. Racah was able to derive a closed formula for the interaction energy among n electrons and to prove that any eigenstate of the interaction (1) is characterized by a 'seniority number' v , a quantum number additional to the total orbital angular momentum L , the total spin S and the number of electrons n . He also showed that v corresponds to the number of electrons that are not in pairs coupled to $L = 0$ [2]. Racah's original definition of seniority made use of coefficients of fractional parentage. He later noted that simplifications arose through the use of group theory [3]. Seniority turned out to be a label associated with the orthogonal algebra $\text{SO}(2\ell + 1)$ in the classification

$$\text{U}(4\ell + 2) \supset (\text{U}(2\ell + 1) \supset \text{SO}(2\ell + 1) \supset \cdots \supset \text{SO}(3)) \otimes \text{SU}_S(2), \quad (2)$$

where the dots indicate intermediate algebras, if any exist. The number of states available to a single electron in an ℓ orbit is $4\ell + 2$. All states of the ℓ^n configuration therefore belong to the totally antisymmetric irreducible representation (IR) $[1^n]$ of $\text{U}(4\ell + 2)$. Furthermore, the natural scheme for electrons in an atom is LS coupling which corresponds to the reduction $\text{U}(4\ell + 2) \supset \text{U}(2\ell + 1) \otimes \text{SU}_S(2)$, where the orbital degrees of freedom are contained in $\text{U}(2\ell + 1)$ and the spin degrees of freedom in $\text{SU}_S(2)$. For any value of ℓ the unitary algebra $\text{U}(2\ell + 1)$ contains the orthogonal subalgebra $\text{SO}(2\ell + 1)$ which in turn contains $\text{SO}(3)$, associated with the total orbital angular momentum L .

The group-theoretical classification (2) allowed Racah to derive a number of important results in the theory of complex atomic spectra. The pairing force (1), however, is a poor approximation to the Coulomb interaction between electrons and for a more physically relevant application of seniority we have to turn to nuclei.

II. SENIORITY IN A SINGLE j SHELL

The discussion of seniority in atoms and in nuclei differs in two aspects: (i) LS coupling is a good first-order approximation in atoms while in nuclei it is rather jj coupling and (ii) electrons are identical particles while nucleons come in two kinds, neutrons and protons. Let us postpone the discussion of the second complication until Sect. IV and concentrate in this section on the case of identical nucleons (either all neutrons or all protons). We impose the additional restriction that the identical nucleons are confined to a single- j shell, deferring the discussion of the many- j case to Sect. III.

It turns out that a pairing force of the type

$$\langle j^2; JM_J | \hat{V} | j^2; JM_J \rangle = -g(2j+1)\delta_{J0}, \quad (3)$$

is a reasonable first-order approximation to the strong interaction between identical nucleons. In Eq. (3) j is the total (orbital+spin) angular momentum of a single nucleon and J results from the coupling of two of them. Since the pairing property now refers to the total j of the nucleons, there is no need for a separate treatment of orbital and spin degrees of freedom as in Eq. (2), and the classification becomes in fact simpler:

$$\begin{array}{ccccc} \text{U}(2j+1) & \supset & \text{Sp}(2j+1) & \supset & \cdots & \supset & \text{SO}(3) \\ \downarrow & & \downarrow & & & & \downarrow \\ [1^n] & & [1^v] & & & & J \end{array} \quad (4)$$

Seniority is associated with the (unitary) symplectic algebra $\text{Sp}(2j+1)$ which replaces the orthogonal algebra $\text{SO}(2\ell+1)$ of the atomic case. Since the nucleons are identical, all states of the j^n configuration belong to the totally antisymmetric IR $[1^n]$ of $\text{U}(2j+1)$. The IRs of $\text{Sp}(2j+1)$ therefore must be totally antisymmetric of the type $[1^v]$. The allowed values of seniority are $v = n, n-2, \dots, 1$ or 0 . The angular momentum content for a given seniority v can also be worked out [4] but no simple general rule is available.

An alternative, simpler definition of seniority can be given which relies on the existence of an $\text{SU}(2)$ symmetry of the pairing hamiltonian [5, 6]. In second quantization the pairing interaction (3) is written as

$$\hat{V} = -g\hat{S}_+^j\hat{S}_-^j, \quad (5)$$

with

$$\hat{S}_+^j = \frac{1}{2}\sqrt{2j+1}(a_j^\dagger \times a_j^\dagger)_0^{(0)}, \quad \hat{S}_-^j = (\hat{S}_+^j)^\dagger, \quad (6)$$

where $a_{jm_j}^\dagger$ creates a nucleon in the shell j with projection m_j . The commutator of \hat{S}_+^j and \hat{S}_-^j leads to the operator $[\hat{S}_+^j, \hat{S}_-^j] = (2\hat{n}_j - 2j - 1)/2 \equiv 2\hat{S}_z^j$, which thus equals, up to a constant, the number operator \hat{n}_j . Since the three operators $\{\hat{S}_z^j, \hat{S}_\pm^j\}$ close under commutation, $[\hat{S}_z^j, \hat{S}_\pm^j] = \pm\hat{S}_\pm^j$ and $[\hat{S}_+^j, \hat{S}_-^j] = 2\hat{S}_z^j$, they form an SU(2) algebra, referred to as the quasi-spin algebra.

This algebraic structure allows an analytical solution of the pairing hamiltonian. From the commutation relations it follows that $\hat{S}_+^j \hat{S}_-^j = (\hat{S}^j)^2 - (\hat{S}_z^j)^2 + \hat{S}_z^j$, which shows that the pairing hamiltonian can be written as a combination of Casimir operators belonging to SU(2) and SO(2) $\equiv \{\hat{S}_z^j\}$. The associated eigenvalue problem can be solved instantly, yielding the energy expression $-g[S(S+1) - M_S(M_S - 1)]$. The quantum numbers S and M_S can be put in relation to the seniority ν and the nucleon number n , $S = (2j - 2\nu + 1)/4$ and $M_S = (2n - 2j - 1)/4$, leading to the energy expression $-g(n - \nu)(2j - n - \nu + 3)/4$. This coincides with the original expression given by Racah, Eq. (50) of Ref. [1], after the replacement of the degeneracy in LS coupling, $4\ell + 2$, by the degeneracy in jj coupling, $2j + 1$.

While this analysis shows that the eigenstates of a pairing interaction carry good seniority, it does not answer the question what are the necessary and sufficient conditions for a general interaction to conserve seniority. Let us specify a rotationally invariant two-body interaction \hat{V} by the matrix elements $\nu_J \equiv \langle j^2; JM_J | \hat{V} | j^2; JM_J \rangle$ with $J = 0, 2, \dots, 2j - 1$. The necessary and sufficient conditions for the conservation of seniority can then be written as

$$\sum_{J=2}^{2j-1} \sqrt{2J+1} \left(\delta_{JI} + 2\sqrt{(2J+1)(2I+1)} \left\{ \begin{matrix} j & j & J \\ j & j & I \end{matrix} \right\} - \frac{4\sqrt{(2J+1)(2I+1)}}{(2j-1)(2j+1)} \right) \nu_J = 0, \quad (7)$$

with $I = 2, 4, \dots, 2j - 1$, and where the symbol between curly brackets is a Racah coefficient. These conditions have been derived previously in a variety of ways [7–9]. Although (7) determines all constraints on the matrix elements ν_J by varying $I = 2, 4, \dots, 2j - 1$, it does not tell us how many of those are independent. This number turns out to be $\lfloor (2j - 3)/6 \rfloor$, the number of independent seniority $\nu = 3$ states [10]. No condition on the matrix elements ν_J is obtained for $j = 3/2, 5/2$ and $7/2$, one condition for $j = 9/2, 11/2$ and $13/2$, and so on. As a result, identical nucleons in a single shell with $j \leq 7/2$ conserve seniority for *any* interaction [7].

Clearly, the conditions (7) are much weaker than the requirement that the interaction be of pairing character but still many of the results of the quasi-spin formalism remain valid.

For instance, the ground state of an even–even nucleus still can be written in the form (8). The main restriction of the concept of seniority as defined so far, concerns the fact that the nucleons are confined to a single- j shell. To lift this restriction, we turn to the generalization presented in the next section.

III. SENIORITY IN SEVERAL j SHELLS

The quasi-spin algebra can be generalized to the case of several degenerate shells (which we assume to be s in number) by making the substitutions $\hat{S}_+^j \mapsto \hat{S}_+ \equiv \sum_j \hat{S}_+^j$ and $2j + 1 \mapsto \sum_j (2j + 1)$. Therefore, if a semi-magic nucleus can be approximated as a system of identical nucleons interacting through a pairing force and distributed over several degenerate shells, the formulas of the quasi-spin formalism should apply. In particular, the ground states of even–even semi-magic nuclei will have a ‘superfluid’ structure of the form

$$\left(\hat{S}_+\right)^{n/2} |o\rangle, \quad (8)$$

where $|o\rangle$ represents the vacuum (*i.e.*, the doubly-magic core nucleus). The SU(2) quasi-spin solution of the pairing hamiltonian (5) leads to several characteristic predictions: a constant excitation energy (independent of n) of the first-excited 2^+ state in even–even isotopes, the linear variation of two-nucleon separation energies as a function of n , the odd–even staggering in nuclear binding energies, the enhancement of two-nucleon transfer.

A more generally valid model is obtained if one imposes the following condition on the hamiltonian:

$$[[\hat{H}, \hat{S}_+], \hat{S}_+] = \Delta \left(\hat{S}_+\right)^2, \quad (9)$$

where \hat{S}_+ creates the lowest two-nucleon eigenstate of \hat{H} and Δ is a constant. This condition of generalized seniority, which was proposed by Talmi [11], is much weaker than the assumption of a pairing interaction and, in particular, it does not require the commutator $[\hat{S}_+, \hat{S}_-]$ to yield (up to a constant) the number operator—a property which is central to the quasi-spin formalism. In spite of the absence of a closed algebraic structure, it is still possible to compute the exact ground-state eigenvalue but hamiltonians satisfying (9) are no longer necessarily completely solvable.

An exact method to solve the problem of identical nucleons distributed over non-degenerate levels interacting through a pairing force was proposed a long time ago by

Richardson [12] based on the Bethe *ansatz* [13]. As an illustration of Richardson's approach, we supplement the pairing interaction with a one-body term, to obtain the following hamiltonian:

$$\hat{H} = \sum_j \epsilon_j \hat{n}_j - g \hat{S}_+ \hat{S}_- = \sum_j \epsilon_j \hat{n}_j - g \sum_j \hat{S}_+^j \sum_{j'} \hat{S}_-^{j'}, \quad (10)$$

where ϵ_j are single-particle energies. The solvability of the hamiltonian (10) arises as a result of the symmetry $SU(2) \otimes SU(2) \otimes \dots$ where each $SU(2)$ algebra pertains to a specific j . Whether the solution of (10) can be called superfluid depends on the differences $\epsilon_j - \epsilon_{j'}$ in relation to the strength g . In all cases the solution is known in closed form for all possible choices of ϵ_j . It is instructive to analyze first the case of $n = 2$ nucleons because it gives insight into the structure of the general problem. The two-nucleon, $J = 0$ eigenstates can be written as $\hat{S}_+ |o\rangle = \sum_j x_j \hat{S}_+^j |o\rangle$ with x_j coefficients that are determined from the eigenequation $\hat{H} \hat{S}_+ |o\rangle = E \hat{S}_+ |o\rangle$ where E is the unknown eigenenergy. With some elementary manipulations this can be converted into the secular equation $2\epsilon_j x_j - g \sum_{j'} \Omega_{j'} x_{j'} = E x_j$, with $\Omega_j = j + 1/2$, from where x_j can be obtained up to a normalization constant, $x_j \propto g/(2\epsilon_j - E)$. The eigenenergy E can be found by substituting the solution for x_j into the secular equation, leading to

$$\sum_j \frac{\Omega_j}{2\epsilon_j - E} = \frac{1}{g}. \quad (11)$$

This equation can be solved graphically which is done in Fig. 1 for a particular choice of single-particle energies ϵ_j and degeneracies Ω_j , appropriate for the tin isotopes with $Z = 50$ protons and neutrons distributed over the 50–82 shell. In the limit $g \rightarrow 0$ of weak pairing, the solutions $E \rightarrow 2\epsilon_j$ are obtained, as should be. Of more interest is the limit of strong pairing, $g \rightarrow +\infty$. From the graphical solution we see that in this limit there is one eigenstate of the pairing hamiltonian which lies well below the other eigenstates with approximately constant amplitudes x_j since for that eigenstate $|E| \gg 2|\epsilon_j|$. Hence, in the limit of strong pairing one finds a $J = 0$ ground state which can be approximated as

$$\hat{S}_+^c |o\rangle \approx \sqrt{\frac{1}{\Omega}} \sum_j \hat{S}_+^j |o\rangle, \quad (12)$$

where $\Omega = \sum_j \Omega_j$. Because of this property this state is often referred to as the collective S state, in the sense that all single-particle orbits contribute to its structure.

This result can be generalized to n particles, albeit that the general solution is more complex. On the basis of the two-particle problem one may propose, for an even number




FIG. 1: Graphical solution of the Richardson equation for $n = 2$ fermions distributed over $s = 5$ single-particle orbits. The sum $\sum_j \Omega_j / (2\epsilon_j - E) \equiv y(E)$ is plotted as a function of E ; the intersections of this curve with the line $y = 1/g$ (dots) then correspond to the solutions of the Richardson equation.

of particles n , a ground state of the hamiltonian (10) of the form (up to a normalization constant)

$$\prod_{\alpha=1}^{n/2} \left(\sum_j \frac{1}{2\epsilon_j - E_\alpha} \hat{S}_+^j \right) |o\rangle, \quad (13)$$

which is known as the Bethe *ansatz* [13]. Each pair in the product is defined through coefficients $x_j = (2\epsilon_j - E_\alpha)^{-1}$ in terms of an energy E_α depending on α which labels the $n/2$ pairs. This product indeed turns out to be the ground state provided the E_α are solutions of $n/2$ coupled, non-linear equations

$$\sum_j \frac{\Omega_j}{2\epsilon_j - E_\alpha} - \sum_{\beta(\neq\alpha)}^{n/2} \frac{2}{E_\beta - E_\alpha} = \frac{1}{g}, \quad \alpha = 1, \dots, n/2, \quad (14)$$

known as the Richardson equations [12]. Note the presence of a second term on the left-hand side with differences of the unknowns $E_\beta - E_\alpha$ in the denominator, which is absent in the two-particle case. In addition, the energy of the state (13) is given by $\sum_\alpha E_\alpha$. A characteristic feature of the Bethe *ansatz* is that it no longer consists of a superposition of *identical* pairs since the coefficients $(2\epsilon_j - E_\alpha)^{-1}$ vary as α runs from 1 to $n/2$. Richardson's model thus provides a solution that covers all possible hamiltonians (10), ranging from those with superfluid character to those with little or no pairing correlations [14].

An important remaining restriction on the form of the pairing hamiltonian (10) is that it contains a single strength parameter g whereas, in general, the interaction might depend

on j and j' , leading to $s(s+1)/2$ strengths $g_{jj'} = g_{j'j}$. In nuclei, often the assumption of a separable interaction is made which, in the case of pairing, leads to strengths $g_{jj'} = gc_jc_{j'}$ in terms of s parameters c_j . This restriction leads to the following pairing hamiltonian:

$$\hat{H} = \sum_j \epsilon_j \hat{n}_j - g \sum_{jj'} c_j c_{j'} \hat{S}_+^j \hat{S}_-^{j'}. \quad (15)$$

As yet, no closed solution of the general hamiltonian (15) is known but three solvable cases have been worked out:

1. The strengths c_j are constant (independent of j). This case was discussed above.
2. The single-particle energies ϵ_j are constant (independent of j). The solution was given by Pan *et al.* [15]
3. There are two levels. The solution was given by Balantekin and Pehlivan [16].

IV. SENIORITY WITH NEUTRONS AND PROTONS

About ten years after its introduction by Racah, seniority was adopted in nuclear physics for the jj -coupling classification of nucleons in a single- j shell [17, 18]. The main additional difficulty in nuclei is that one deals with a system of neutrons and protons, and hence the isospin T of the nucleons should be taken into account. The generalization of the classification (4) for identical nucleons toward neutrons and protons reads as follows:

$$\begin{array}{ccccccccc} \text{U}(4j+2) & \supset & (\text{U}(2j+1) & \supset & \text{Sp}(2j+1) & \supset & \cdots & \supset & \text{SO}(3)) & \otimes & \text{SU}_T(2) \\ \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow \\ [1^n] & & [h] & & [\sigma] & & & & J & & T \end{array}, \quad (16)$$

where $[h]$ and $[\sigma]$ are Young tableaux associated with $\text{U}(2j+1)$ and $\text{Sp}(2j+1)$. In general, $2j+1$ labels are needed to characterize an IR of $\text{U}(2j+1)$, $[h] = [h_1, h_2, \dots, h_{2j+1}]$, and $j+1/2$ labels are needed for an IR of $\text{Sp}(2j+1)$, $[\sigma] = [\sigma_1, \sigma_2, \dots, \sigma_{j+1/2}]$. To ensure overall antisymmetry under $\text{U}(4j+2)$, the Young tableaux of $\text{U}(2j+1)$ and $\text{U}_T(2)$ must be conjugate, that is, one is obtained from the other by interchanging rows and columns. Since the Young tableau associated with $\text{U}_T(2)$ is determined by the nucleon number n and the total isospin T as $[n/2+T, n/2-T]$, the Young tableau of $\text{U}(2j+1)$ must therefore be

$$[h] = \overbrace{[2, 2, \dots, 2]}^{n/2-T}, \overbrace{[1, 1, \dots, 1]}^{2T}. \quad (17)$$

Since an IR of $U(2j + 1)$ has at most $2j + 1$ labels, it follows that $n/2 + T \leq 2j + 1$. Furthermore, all non-zero labels in $[\sigma]$ must be either 2 or 1 and the Young tableau of $Sp(2j + 1)$ must therefore be of the form

$$[\sigma] = \overbrace{[2, 2, \dots, 2]}^{v/2-t} \overbrace{[1, 1, \dots, 1]}^{2t}. \quad (18)$$

The IR of $Sp(2j + 1)$ is thus characterized by *two* labels: the seniority v and the ‘reduced isospin’ t . The former has the same interpretation as in the like-nucleon case while the latter corresponds to the isospin of the nucleons which are not in pairs coupled to $J = 0$.

The group-theoretical analysis is considerably more complex here than in the case of identical nucleons and, in addition, for each value of j one is faced with a different reduction problem associated with $U(2j + 1) \supset Sp(2j + 1) \supset SO(3)$. It is therefore advantageous to go over to a quasi-spin formulation of the problem and, as was shown by Helmers [6], this is possible for whatever value of the intrinsic quantum number of the particles (which is $t = 1/2$ for nucleons). If the pairing interaction is assumed to be isospin invariant, it is the same in the three $T = 1$ channels, neutron–neutron, neutron–proton and proton–proton, and Eq. (5) can be generalized to

$$\hat{V}' = -g \sum_{\mu} \hat{S}_{+,\mu} \hat{S}_{-,\mu} = -g \hat{S}_+ \cdot \hat{S}_-, \quad (19)$$

where the dot indicates a scalar product in isospin. In terms of the nucleon creation operators $a_{jm_jtm_t}^\dagger$, which now carry also isospin indices (with $t = 1/2$), the pair operators are

$$\hat{S}_{+,\mu} = \frac{1}{2} \sum_j \sqrt{2j + 1} (a_{jt}^\dagger \times a_{jt}^\dagger)_{0\mu}^{(01)}, \quad \hat{S}_{-,\mu} = (\hat{S}_{+,\mu})^\dagger, \quad (20)$$

where the coupling refers to angular momentum and to isospin. The index μ (isospin projection) distinguishes neutron–neutron ($\mu = +1$), neutron–proton ($\mu = 0$) and proton–proton ($\mu = -1$) pairs. There are thus three different pairs with $J = 0$ and $T = 1$ and they are related through the action of the isospin raising and lowering operators \hat{T}_\pm . By considering the commutation relations between the different operators, a closed algebraic structure is obtained, generated by the pair operators $\hat{S}_{\pm,\mu}$, the number operator \hat{n} and the isospin operators \hat{T}_\pm and \hat{T}_z . The quasi-spin algebra of neutrons and protons in degenerate j shells turns out to be $SO(5)$, by virtue of which the hamiltonian (19) is analytically solvable [19, 20].

A further generalization is possible in LS coupling. For a neutron and a proton there exists a different paired state with *parallel* spins. The most general pairing interaction for a

system of neutrons and protons is therefore of the form

$$\hat{V}'' = -g\hat{S}_+ \cdot \hat{S}_- - g'\hat{P}_+ \cdot \hat{P}_-, \quad (21)$$

where the pair operators are defined as

$$\begin{aligned} \hat{S}_{+,\mu} &= \sqrt{\frac{1}{2}} \sum_{\ell} \sqrt{2\ell+1} (a_{\ell st}^{\dagger} \times a_{\ell st}^{\dagger})_{00\mu}^{(001)}, & \hat{S}_{-,\mu} &= (\hat{S}_{+,\mu})^{\dagger}, \\ \hat{P}_{+,\mu} &= \sqrt{\frac{1}{2}} \sum_{\ell} \sqrt{2\ell+1} (a_{\ell st}^{\dagger} \times a_{\ell st}^{\dagger})_{0\mu 0}^{(010)}, & \hat{P}_{-,\mu} &= (\hat{P}_{+,\mu})^{\dagger}, \end{aligned} \quad (22)$$

where $a_{\ell m_{\ell} s m_s t m_t}^{\dagger}$ creates a nucleon in the shell ℓ with projection m_{ℓ} , spin projection m_s and isospin projection m_t . The hamiltonian (21) contains two parameters g and g' , the strengths of the isovector and isoscalar components of the pairing interaction. While in the previous case the single strength parameter g just defines an overall scale, this is no longer true for a generalized pairing interaction and different solutions are obtained for different ratios g/g' .

In general, the eigenproblem associated with the interaction (21) can only be solved numerically; for specific choices of g and g' the solution of \hat{V}'' can be obtained analytically [21, 22]. A closed algebraic structure is obtained, formed by the pair operators (22), their commutators, the commutators of these among themselves, and so on until closure is attained. The quasi-spin algebra in this case turns out to be $\text{SO}(8)$, with 28 generators, consisting of the pair operators $\hat{S}_{\pm,\mu}$ and $\hat{P}_{\pm,\mu}$, the number operator \hat{n} , the spin and isospin operators \hat{S}_{μ} and \hat{T}_{μ} , and the Gamow–Teller-like operator $\hat{Y}_{\mu\nu}$, which is a vector in spin and isospin. The symmetry character of the hamiltonian (21) is obtained by studying the subalgebras of $\text{SO}(8)$. Of relevance are the subalgebras $\text{SO}_T(5) \equiv \{\hat{S}_{\pm,\mu}, \hat{n}, \hat{T}_{\mu}\}$, $\text{SO}_T(3) \equiv \{\hat{T}_{\mu}\}$, $\text{SO}_S(5) \equiv \{\hat{P}_{\pm,\mu}, \hat{n}, \hat{S}_{\mu}\}$, $\text{SO}_S(3) \equiv \{\hat{S}_{\mu}\}$ and $\text{SO}(6) \equiv \{\hat{S}_{\mu}, \hat{T}_{\mu}, \hat{Y}_{\mu\nu}\}$, which can be placed in the following lattice of algebras:

$$\text{SO}(8) \supset \left\{ \begin{array}{c} \text{SO}_S(5) \otimes \text{SO}_T(3) \\ \text{SO}(6) \\ \text{SO}_T(5) \otimes \text{SO}_S(3) \end{array} \right\} \supset \text{SO}_S(3) \otimes \text{SO}_T(3). \quad (23)$$

By use of the explicit form of the generators of $\text{SO}(8)$ and its subalgebras, and their commutation relations [22], the following relations can be shown to hold:

$$\hat{S}_+ \cdot \hat{S}_- = \frac{1}{2} \hat{C}_2[\text{SO}_T(5)] - \frac{1}{2} \hat{C}_2[\text{SO}_T(3)] - \frac{1}{8} (2\Omega - \hat{n})(2\Omega - \hat{n} + 6),$$

$$\begin{aligned}\hat{S}_+ \cdot \hat{S}_- + \hat{P}_+ \cdot \hat{P}_- &= \frac{1}{2}\hat{C}_2[\text{SO}(8)] - \frac{1}{2}\hat{C}_2[\text{SO}(6)] - \frac{1}{8}(2\Omega - \hat{n})(2\Omega - \hat{n} + 12), \\ \hat{P}_+ \cdot \hat{P}_- &= \frac{1}{2}\hat{C}_2[\text{SO}_S(5)] - \frac{1}{2}\hat{C}_2[\text{SO}_S(3)] - \frac{1}{8}(2\Omega - \hat{n})(2\Omega - \hat{n} + 6),\end{aligned}\quad (24)$$

with $\Omega = \sum_\ell (2\ell + 1)$ and where $\hat{C}_n[G]$ is the n^{th} -order Casimir operator of the algebra G . This shows that the interaction (21) in the three cases (i) $g = 0$, (ii) $g' = 0$ and (iii) $g = g'$, can be written as a combination of Casimir operators of algebras belonging to a chain of *nested* algebras of the lattice (23). They are thus the dynamical symmetries of the SO(8) model.

The nature of ‘SO(8) superfluidity’ can be illustrated in the specific example of the ground state of even–even $N = Z$ nuclei. In the SO(6) limit of the SO(8) model the exact ground-state solution can be written as [23]

$$\left(\hat{S}_+ \cdot \hat{S}_+ - \hat{P}_+ \cdot \hat{P}_+\right)^{n/4} |0\rangle. \quad (25)$$

This shows that the superfluid solution acquires a *quartet* structure in the sense that it reduces to a condensate of bosons each of which corresponds to four nucleons. Since the boson in (25) is a scalar in spin and isospin, it can be thought of as an α particle; its orbital character, however, might be different from that of an actual α particle. A quartet structure is also present in the two SO(5) limits of the SO(8) model, which yields a ground-state wave function of the type (25) with either the first or the second term suppressed. A reasonable *ansatz* for the $N = Z$ ground-state wave function of the SO(8) pairing interaction (21) with arbitrary strengths g and g' is therefore

$$\left(\cos\theta \hat{S}_+ \cdot \hat{S}_+ - \sin\theta \hat{P}_+ \cdot \hat{P}_+\right)^{n/4} |0\rangle, \quad (26)$$

where θ is a parameter that depends on the ratio g/g' . The condensate (26) of α -like particles provides an excellent approximation to the $N = Z$ ground state of the pairing hamiltonian (21) for any combination of g and g' [23]. It should nevertheless be stressed that, in the presence of both neutrons and protons in the valence shell, the pairing hamiltonian (21) is *not* a good approximation to a realistic shell-model hamiltonian which contains an important quadrupole component.

These results can be generalized to the case of several non-degenerate shells. In fact, the Richardson equations (14) are valid for the quasi-spin symmetry SU(2) but they are known for any Lie algebra [24]. Closed solutions have been obtained for a system of neutron and

protons with a pairing interaction of pure isovector character and of equal isovector and isoscalar strength, based on the SO(5) and the SO(6) quasi-spin algebras, respectively [25, 26].

V. BOSE–EINSTEIN CONDENSATES OF ATOMS WITH SPIN

In this section the concept of seniority is illustrated with an application to the physics of cold atoms. If atoms in a Bose–Einstein condensate (BEC) are trapped by optical means [27], their hyperfine spins (or spins) are not frozen in one particular direction but are essentially free but for their mutual interactions. As a result, the atoms do not behave as scalar particles but each of the components of the spin is involved in the formation of the BEC. This raises interesting questions concerning the structure of the condensate and how it depends on the spin-exchange interactions between the atoms.

Such questions were addressed in a series of theoretical papers by Ho and co-workers [28] who obtained solutions based on a generating function method. In the case of spin-1 atoms the problem of quantum spin mixing was analyzed by Law *et al.* [29] who proposed an elegant solution based on algebraic methods. It is shown here that an exact solution is also available for the spin value $f = 2$ (for any number of atoms n) which allows the analytic determination of the structure of the ground state of the condensate. This was simultaneously and independently pointed out in Refs. [30, 31].

We consider a one-component dilute gas of trapped bosonic atoms with arbitrary (integer) hyperfine spin f . In second quantization the hamiltonian of this system has a one-body and a two-body piece that can be written as (in the notation of Ref. [29])

$$\mathcal{H} = \sum_m \int \hat{\Psi}_m^\dagger \left(-\frac{\nabla^2}{2M_a} + V_{\text{trap}} \right) \hat{\Psi}_m d^3x + \sum_{m_i} \Omega_{m_1 m_2 m_3 m_4} \int \hat{\Psi}_{m_1}^\dagger \hat{\Psi}_{m_2}^\dagger \hat{\Psi}_{m_3} \hat{\Psi}_{m_4} d^3x, \quad (27)$$

where $\hbar = 1$, M_a is the mass of the atom, and $\hat{\Psi}_m$ and $\hat{\Psi}_m^\dagger$ are the atomic field annihilation and creation operators associated with atoms in the hyperfine state $|fm\rangle$ with $m = -f, \dots, +f$, the possible values of all summation indices in (27). The trapping potential V_{trap} is assumed to be the same for all $2f + 1$ components. According to the assumptions outlined in Ref. [29], the atomic field creation and annihilation operators at zero temperature can be approximated by $\hat{\Psi}_m^\dagger \approx b_m^\dagger \phi(\vec{x})$, $\hat{\Psi}_m \approx b_m \phi(\vec{x})$, $m = -f, \dots, +f$, where $\phi(\vec{x})$ is a single wave function (independent of m) and b_m and b_m^\dagger are annihilation and creation

operators, satisfying the usual boson commutation rules. In this approximation the entire hamiltonian (27) can be rewritten as

$$\mathcal{H} \approx \hat{H} \equiv \epsilon b^\dagger \cdot \tilde{b} + \frac{1}{2} \sum_F \nu_F (b^\dagger \times b^\dagger)^{(F)} \cdot (\tilde{b} \times \tilde{b})^{(F)}, \quad (28)$$

where the coefficients ϵ and ν_F are related to those in the original hamiltonian (27) and with $\tilde{b}_m \equiv (-)^{f-m} b_{-m}$.

Exactly solvable hamiltonians with rotational or SO(3) invariance are now found by the determination of all Lie algebras G satisfying $U(2f+1) \supset G \supset SO(3)$. The canonical reduction of $U(2f+1)$ is of the form as encountered by Racah (see Sect. I),

$$U(2f+1) \supset SO(2f+1) \supset SO(3), \quad (29)$$

defining a class of solvable hamiltonians of the type

$$\hat{H}' = a_1 \hat{C}_1[U(2f+1)] + a_2 \hat{C}_2[U(2f+1)] + b \hat{C}_2[SO(2f+1)] + c \hat{C}_2[SO(3)], \quad (30)$$

where a_1 , a_2 , b , and c are numerical coefficients. The solvability properties of the original hamiltonian (28) now follow from a simple counting argument. For atoms with spin $f = 1$ the solvable hamiltonian (30) has three coefficients a_1 , a_2 , and c [since $SO(2f+1) = SO(3)$] while the general hamiltonian (28) also contains three coefficients ϵ , ν_0 , and ν_2 . (Note that the coupling of two spins to odd F is not allowed in the approximation of a common spatial wave function, so no ν_1 term occurs.) They can be put into one-to-one correspondence. For atoms with spin $f = 2$ both the solvable and the general hamiltonian contain four coefficients (a_1 , a_2 , b , and c versus ϵ , ν_0 , ν_2 , and ν_4) which also can be put into one-to-one correspondence. Hence the general hamiltonian (28) is solvable for $f \leq 2$. The same counting argument shows that it is no longer solvable for $f > 2$.

The case of interacting $f = 1$ atoms was discussed by Law *et al.* [29] who identified the existence of two possible condensate ground states: one with all atoms aligned to maximum spin $F = n$ and a second with pairs of atoms coupled to $F = 0$. Whether the condensate is aligned or paired depends on a single interaction parameter. With the technique explained above, the phase diagram for atoms with spin $f = 2$ can also be derived. The results are exact and valid for arbitrary n . The entire spectrum is determined by the eigenvalue expression together with the necessary branching rules. In particular, the allowed values of total spin F for a given seniority v are derived from the $SO(5) \supset SO(3)$ branching rule given by $F = 2\tau, 2\tau - 2, 2\tau - 3, \dots, \tau + 1, \tau$ with $\tau = v, v - 3, v - 6, \dots$ and $\tau \geq 0$.

It is now possible to determine all possible ground-state configurations of the condensate and their quantum numbers ν_0 and F_0 [30]. The character of the ground state does not depend on the coefficients a_i since the first two terms in the expression (30) give a constant contribution to the energy of all states. Although this contribution is dominant, the spectrum-generating perturbation of the hamiltonian is confined to the last two terms and depends solely on the coefficients b and c which are related to the original interactions ν_F according to $b = (-7\nu_0 + 10\nu_2 - 3\nu_4)/70$ and $c = (-\nu_2 + \nu_4)/14$. The phase diagram displays a richer structure than in the $f = 1$ case. There is an aligned phase where the seniority is maximal, $\nu_0 = n$, and all spins are aligned, $F_0 = 2n$. Secondly, there is a low-seniority (paired) and consequently low-spin phase. For even n , this corresponds to $(\nu_0, F_0) = (0, 0)$. The aligned and paired phases are also encountered for interacting $f = 1$ atoms. For $f = 2$ a third phase occurs characterized by high seniority (*i.e.*, unpaired) and low total spin, $(\nu_0, F_0) = (n, 2\delta)$ with $\delta = 0$ or 1.

Since the hamiltonian (30) is solvable for $f = 2$, all eigenstates, and in particular the three different ground states, can be determined analytically. The general expressions given by Chacón *et al.* [32] reduce to

$$\begin{aligned} |v = n, F = M_F = 2n\rangle &\propto (d_{+2}^\dagger)^n |0\rangle, \\ |v = 0, F = M_F = 0\rangle &\propto (d^\dagger \cdot d^\dagger)^{n/2} |0\rangle, \\ |v = n, F = M_F = 0\rangle &\propto ((a^\dagger \times a^\dagger)^{(2)} \cdot a^\dagger)^{n/3} |0\rangle, \end{aligned} \quad (31)$$

where the $f = 2$ atoms are denoted as d bosons. In the second of these expressions it is assumed that n is even and in the third that $n = 3k$; other cases are obtained by adding a single boson d^\dagger or a $d^\dagger \cdot d^\dagger$ pair. The a^\dagger are the so-called traceless boson operators [32] which are defined as $a_m^\dagger = d_m^\dagger - d^\dagger \cdot d^\dagger (2\hat{n} + 5)^{-1} \tilde{d}_m$ (see also Chapt. 8 of Ref. [33]). The wave functions (31) are the *exact* finite- n expressions for the eigenstates of the hamiltonian (30). Since in the large- n limit the traceless boson operators a_m^\dagger become identical to d_m^\dagger , one arrives at a simple interpretation of the three types of configurations: (i) spin-aligned, (ii) condensed into *pairs of atoms* coupled to $F = 0$, and (iii) condensed into *triplets of atoms* coupled to $F = 0$.

In conclusion, the consideration of seniority is crucial in obtaining results concerning Bose–Einstein condensates consisting of atoms with spin. Since all eigenstates of interacting atoms with spin $f \leq 2$ are known analytically, this opens up the possibility to study the

relaxation properties of such condensates using their exact, macroscopic wave functions. In addition, preliminary studies indicate that seniority can be exploited even when $f > 2$. These problems are currently under investigation [34].

This paper is dedicated to the memory of Marcos Moshinsky. The two years I have spent in Mexico as a visitor and the many hours with Marcos as a teacher, were crucial to my formation as a physicist. Without him I never could have written this paper.

-
- [1] G. Racah, *Phys. Rev.* **63**, 367 (1943).
 - [2] This is a rather loose statement in words. The precise definition of seniority involves a property of vanishing n -to- $(n - 2)$ coefficients of fractional parentage.
 - [3] G. Racah, *Phys. Rev.* **76**, 1352 (1949).
 - [4] B.G. Wybourne, *Symmetry Principles and Atomic Spectroscopy*, Wiley-Interscience, New York, 1970.
 - [5] A.K. Kerman, *Ann. Phys. (NY)* **12**, 300 (1961).
 - [6] K. Helmers, *Nucl. Phys.* **23**, 594 (1961).
 - [7] I. Talmi, *Simple Models of Complex Nuclei*, Harwood, Chur, 1993.
 - [8] D.J. Rowe and G. Rosensteel, *Phys. Rev. Lett.* **87**, 172501 (2001).
 - [9] G. Rosensteel and D.J. Rowe, *Phys. Rev. C* **67**, 014303 (2003).
 - [10] J.N. Ginocchio and W.C. Haxton, in *Symmetries in Science VI*, edited by B. Gruber and M. Ramek, Plenum, New York, 1993.
 - [11] I. Talmi, *Nucl. Phys. A* **172**, 1 (1971).
 - [12] R.W. Richardson, *Phys. Lett.* **3**, 277 (1963)
 - [13] M. Gaudin, *La Fonction d'Onde de Bethe* Masson, Paris, 1983.
 - [14] J. Dukelsky, S. Pittel, and G.Sierra, *Rev. Mod. Phys.* **76**, 1 (2004).
 - [15] F. Pan, J.P. Draayer, and W.E. Ormand, *Phys. Lett. B* **422**, 1 (1998).
 - [16] A.B. Balantekin and Y. Pehlivan, *Phys. Rev. C* **76**, 051001(R) (2007).
 - [17] G. Racah, *L. Farkas Memorial Volume*, Research council of Israel, Jerusalem, 1952, p. 294.
 - [18] B.H. Flowers, *Proc. Roy. Soc. (London) A* **212**, 248 (1952).
 - [19] K.T. Hecht, *Phys. Rev.* **139**, B794 (1965); *Nucl. Phys. A* **102**, 11 (1967); *Nucl. Phys. A* **493**, 29 (1989).

- [20] J.N. Ginocchio, *Nucl. Phys.* **74**, 321 (1965).
- [21] B.H. Flowers and S. Szpikowski, *Proc. Phys. Soc.* **84**, 673 (1964).
- [22] S.C. Pang, *Nucl. Phys. A* **128**, 497 (1969).
- [23] J. Dobeš and S. Pittel, *Phys. Rev. C* **57**, 688 (1998).
- [24] M. Asorey, F. Falceto, and G. Sierra, *Nucl. Phys. B* **622**, 593 (2002).
- [25] J. Dukelsky, V.G. Gueorguiev, P. Van Isacker, S. Dimitrova, B. Errea, and S. Lerma H., *Phys. Rev. Lett.* **96**, 072503 (2006).
- [26] S. Lerma H., B. Errea, J. Dukelsky, and W. Satuła, *Phys. Rev. Lett.* **99**, 032501 (2007).
- [27] D.M. Stamper-Kurn, M.R. Andrews, A.P. Chikkatur, S. Inouye, H.-J. Miesner, J. Stenger, and W. Ketterle, *Phys. Rev. Lett.* **80**, 2027 (1998).
- [28] T.-L. Ho, *Phys. Rev. Lett.* **81**, 742 (1998); T.-L. Ho and S. Yip, *Phys. Rev. Lett.* **82**, 247 (1999); T.-L. Ho and L. Yin, *Phys. Rev. Lett.* **84**, 2302 (2000).
- [29] C.K. Law, H. Pu, and N.P. Bigelow *Phys. Rev. Lett.* **81**, 5257 (1998).
- [30] P. Van Isacker and S. Heinze, *J. Phys. A* **40**, 14811 (2007).
- [31] S. Uchino, T. Otsuka, and M. Ueda, *Phys. Rev. A* **78**, 023609 (2008).
- [32] E. Chacón, M. Moshinsky, and R.T. Sharp, *J. Math. Phys.* **17**, 668 (1976).
- [33] A. Frank and P. Van Isacker, *Algebraic Methods in Molecular and Nuclear Structure Physics* Wiley, New York, 1994.
- [34] P. Van Isacker and S. Heinze, to be published.