

Detection and Classification of Local Ca²⁺ Release Events in Cardiomyocytes Using 3D-UNet Neural Network

Prisca Dotti^{1,2}, Pablo Márquez Neila¹, Miguel Fernandez-Tenorio², Radoslav Janicek², Marcel Wullschlegler², Till Meyer zu Westram¹, Raphael Sznitman¹, Marcel Egger²

Affiliations:

¹AIMI, ARTORG Center, University of Bern, Bern, Switzerland

²Department of Physiology, University of Bern, Bern, Switzerland

Correspondence to prisca.dotti@unibe.ch

u^b

UNIVERSITÄT
BERN

Introduction

Global Ca²⁺ increase in the cytosol of cardiomyocytes is crucial for the contraction of the heart. Malfunctioning of proteins involved in this process can trigger local events (e.g., sparks and puffs) and global events (e.g., waves). These are thought to be involved in the development of arrhythmia. Therefore, it is important to detect and classify local Ca²⁺ release events. We present a novel approach, based on a 3D U-Net architecture, to perform these tasks in a fully automated fashion. We employed data obtained with fast xyt confocal imaging of cardiomyocytes where such subcellular Ca²⁺ events are manually annotated and trained the neural network to infer comparable segmentation as output. Despite the relatively small amount of available data and the challenges that it exhibits, we obtained qualitatively promising results.

Challenging Ca²⁺ imaging data

The main task is the detection of three types of SR-Ca²⁺ release events in atrial cardiomyocytes:

- **Ca²⁺ sparks:** coordinated opening of cluster of 6 to 20 ryanodine receptors (RyRs) triggered by calcium ions [2].




Figure 1: Signal over time of a sample Ca²⁺ spark (left). Annotation mask used for training, plotted Ca²⁺ spark is in the red box (top right). Slice of sample movie, the yellow circle denotes the region over which temporal data is averaged (bottom right).

- **Ca²⁺ puffs:** opening of 20 to 35 coordinated inositol trisphosphate receptors (InsP3Rs) [3].




Figure 2: Signal over time of a sample Ca²⁺ puff (left). Annotation mask used for training, plotted Ca²⁺ puff is in the red box (top right). Slice of sample movie, the yellow circle denotes the region over which temporal data is averaged (bottom right).

- **Ca²⁺ waves:** propagation of Ca²⁺ sparks along the length of the cell [2].




Figure 3: Signal over time of a sample Ca²⁺ wave (left). Propagation of Ca²⁺ wave in time, different colors indicate time from the start of the event. Dark purple represents the origin of the event and yellow its end (top right). Slice of sample movie, the yellow circle denotes region over which temporal data is averaged (bottom right).

- Simple thresholds on measures such as amplitude, rise and decay times, etc., are not sufficient to separate the three classes of events.
- **Crosstalks are possible between Ca²⁺ sparks and Ca²⁺ puffs** [1], making the classification task even more challenging.

The dataset consists of 43 xyt image series. Each sample contains from 500 to 1900 frames (512x64 pixels), recorded with a frequency of images between 137.5 Hz and 169.5 Hz. Pharmacology has been utilized to stimulate InsP3Rs activity in a subset of the samples, for this reason, data present different types of noise.

Method

Data segmentation and labelling




Figure 4: Data were manually annotated, with the help of a semi-automatic preprocessing of the samples.

3D U-Net input and architecture




Figure 5: 3D U-Net architecture: the input is an extract of the input image series, and the output consists of 4 image series, representing the probability distribution for each class (events and background).

We split each sample into overlapping chunks of 256 frames, with a step of 32 frames. Data augmentation is done by flipping the samples vertically and/or horizontally. The neural network architecture is based on the standard U-Net architecture which has been adapted to handle 3-dimensional data (fig. 5).

Experiments and results

We trained the neural network for 100K epochs on the Lovász-Softmax loss [4], using the Adam optimizer.




Figure 6a: average IoU (Jaccard Score) and Matthews correlation coefficient for Ca²⁺ puff and Ca²⁺ wave classes, with respect to detection threshold.




Figure 6b: average metrics for Ca²⁺ spark class, with respect to detection threshold.




Figure 7: Example of the model's raw predictions. Left column: sample frame (top) with corresponding annotated Ca²⁺ wave, in green (center) and output of the network's softmax layer, for the Ca²⁺ waves class (bottom). Right column: sample frame (top) with corresponding annotated Ca²⁺ sparks, in orange, and Ca²⁺ puff, in yellow (center) and output of the network's softmax layer, for the Ca²⁺ sparks class (bottom).

Conclusion

- We present a novel approach to the detection and classification of local Ca²⁺ release events in cardiomyocytes.
- The trained network architecture provides a tool to solve this problem in a fully automated way, without manually processing xyt image series.
- Machine learning can help the detection and the categorization of local Ca²⁺ release.
- Further steps include the identification of the events in the prediction, and training with different types of neural network models.

References

- [1] M. Wullschlegler, J. Blanch, and M. Egger, "Functional local crosstalk of inositol 1,4,5-trisphosphate receptor- and ryanodine receptor-dependent Ca²⁺ release in atrial cardiomyocytes," *Cardiovascular Research*, vol. 113, pp. 542–552, 02 2017.
- [2] H. Cheng, W. Lederer, and M. Cannell, "Calcium sparks: elementary events underlying excitation-contraction coupling in heart muscle," *Science*, vol. 262, no. 5134, pp. 740–744, 1993.
- [3] I. Parker and Y. Yao, "Regenerative Release of Calcium from Functionally Discrete Subcellular Stores by Inositol Trisphosphate," *Proceedings of the Royal Society of London Series B*, vol. 246, pp. 269–274, Dec. 1991.
- [4] M. Berman and M. B. Blaschko, "Optimization of the Jaccard index for image segmentation with the Lovász hinge," *CoRR*, 2017.