

REVIEW PAPER

Nanofoms of essential metals: from hormetic phytoeffects to agricultural potential

Zsuzsanna Kolbert^{1,*}, Réka Szöllösi¹, Andrea Rónavári², and Árpád Molnár¹

¹ Department of Plant Biology University of Szeged, Közép fasor 52, Szeged H6726, Hungary

² Department of Applied and Environmental Chemistry, University of Szeged, Rerrich Béla tér 1, Szeged H6720, Hungary

* Correspondence: ordogne.kolbert.zsuzsanna@szte.hu

Received 13 August 2021; Editorial decision 8 December 2021; Accepted 26 January 2022

Editor: Ann Cuypers, Hasselt University, Belgium

Abstract

Vital plant functions require at least six metals (copper, iron, molybdenum, manganese, zinc, and nickel), which function as enzyme cofactors or inducers. In recent decades, rapidly evolving nanotechnology has created nanofoms of essential metals and their compounds (e.g. nZnO, nFe₂O₃) with a number of favourable properties over the bulk materials. The effects of nanometals on plants are concentration-dependent (hormesis) but also depend on the properties of the nanometals, the plant species, and the treatment conditions. Here, we review studies examining plant responses to essential nanometal treatments using a (multi)omics approach and emphasize the importance of gaining a holistic view of the diverse effects. Furthermore, we discuss the beneficial effects of essential nanometals on plants, which provide the basis for their application in crop production as, for example, nanopriming or nanostimulator agents, or nanofertilizers. As lower environmental impact and increased yield can be achieved by the application of essential nanometals, they support sustainable agriculture. Recent studies have actively examined the utilization of green-synthesized metal nanoparticles, which perfectly fit into the environmentally friendly trend of future agriculture. Further knowledge is required before essential nanometals can be safely applied in agriculture, but it is a promising direction that is timely to investigate.

Keywords: Hormesis, nanofertilization, nanometals, nanopriming, nutrient deficiency, omics.

Introduction

Essential elements are indispensable for the vegetative growth and reproduction of plants and are not replaceable by another element. These elements are directly involved in the metabolism forming structural or functional components of plant cells. Moreover, inadequate availability of essential elements causes damage to the plant. These criteria are met by certain metals, such as copper (Cu), iron (Fe), molybdenum (Mo), manganese (Mn), zinc (Zn), and nickel

(Ni), which are essential trace components of plant structures (Barker, 2015).

Land plants have specific transport systems to absorb essential metal ions effectively from the environment, appropriately distribute them in plant organs and tissues, and sequester them within cells (Andresen *et al.*, 2018). The transport of trace metal ions across membranes requires energy derived directly from ATP hydrolysis (primary active transport) or provided by the

metal transporter due to the symport or antiport of other ions (e.g. H^+ ; secondary active transport). Primary active transport of essential metal ions is mediated by P2A and P1B-type (also called CPx-type) ATPases (Williams and Mills, 2005), while glutathione- or phytochelatin-bound metal ion ligands are transported by ABC proteins (Dahuja *et al.*, 2021). Most ABC transporters are not specifically involved in metal translocation but are involved in other transport routes (e.g. malate transport, indole-acetic acid transport; Dahuja *et al.*, 2021). Secondary active transport systems associated with metal uptake into the cytoplasm and intracellular translocation are natural resistance-associated macrophage proteins (NRAMPs), ZRT/IRT-like proteins (ZIPs), and cation diffusion facilitators (CDFs) (Hall and Williams, 2003; Lira-Morales *et al.*, 2019). The latter group of transporters are mainly embedded in the vacuolar or Golgi membranes and function as metal-proton antiporters. Additional types of transporters involved in metal translocation are cation/proton exchangers (CAXs) found in the tonoplast (Pittman and Hirschi, 2016). Additionally, endocytosis and exocytosis may be involved in metal uptake and export from the cytoplasm (Andresen *et al.*, 2018).

In plant cells, essential metals can be present in diverse biologically active forms such as hydrated ions, cofactors in proteins, nucleic acid-bound forms, and forms associated with amino acids, glutathione, nicotinamine, citrate, and malate (Juárez-Maldonado *et al.*, 2018). Among their chemical properties, redox activity has significant relevance in influencing the specific biological function of the essential metals. Due to their different oxidation states, Cu and Fe catalyse oxido-reduction reactions in active centres of enzymes, and thus they are involved in electron transport and cellular redox control.

As the cofactor of plastocyanin, Cu has a unique role in photosynthetic electron transport (Droppa and Horváth, 1990). Cu is involved also in the mitochondrial electron transport chain as a cofactor of cytochrome oxidase, which contains also Fe (in the form of haem) as a supporter of its function. Cu also has a role in reactive oxygen species (ROS) detoxification, since it activates Cu/Zn superoxide dismutase (SOD) isoenzymes and ascorbate oxidase (Alscher *et al.*, 2002; De Tullio *et al.*, 2013). There are several other Cu-dependent oxidases, such as laccase, polyphenol oxidase, and amine oxidase, all of which are involved in wound healing and thus in plants' defence against pathogens (Li *et al.*, 2019). Moreover, Cu is found in the active site of ethylene receptors and thus has a relevant role in perception of the hormone ethylene in plant cells (Li *et al.*, 2019; Schott-Verdugo *et al.*, 2019).

The antioxidant enzyme SOD has an isoform that functions in the presence of Fe. The biological roles of Cu/ZnSODs and FeSODs are different, since FeSODs control the level of ROS as signal molecules (Alscher *et al.*, 2002). Alternative oxidase, involved in respiration, is also a protein that binds haem-Fe. Furthermore, there are multi-haem proteins belonging to the P450 superfamily. These enzymes control redox reactions and convert various types of secondary metabolites, such as amines,

fatty acids, and steroids. In addition, multiple steps of lignin biosynthesis involve the action of P450 superfamily enzymes. Fe is involved directly in the light reactions of photosynthesis, since it is a substituent of cytochrome b_{559} in the PSII reaction centre and it forms Fe_4S_4 clusters in PSI reaction centres. Additionally, the cytochrome b_6f complex contains both haem and Fe-S clusters. In the respiratory electron transport chain, Fe-S clusters can be found in complex I and II, and in the cytochrome bc_1 complex. Nitrate and sulfate metabolisms also require available Fe, since the enzymes nitrate reductase (NR), nitrite reductase (NiR), and sulfite oxidase (SO) contain haem-Fe (Vigani and Murgia, 2018).

NR and SO enzymes contain also Mo in the form of an organic cofactor, molybdopterin, which is involved in substrate binding and reduction. Aldehyde oxidases (acting in abscisic acid and indoleacetic acid biosynthesis) and xanthine oxidase are additional enzymes that function in the presence of Mo, and these enzymes contain a Fe_2S_2 cluster as well (Mendel and Hänsch, 2002).

The major role of Mn in plants is related to the fact that it is a central substituent of the water-splitting complex in the PSII reaction centre. Furthermore, ROS-detoxifying MnSODs are present in mitochondria and in peroxisomes. Mn is also bounded by germin and germin-like proteins, which are located in the cell walls and catalyse the production of hydrogen peroxide (H_2O_2) from oxalate. H_2O_2 contributes to pathogen defence as a signal, as an antimicrobial agent, and as an inducer of lignification. There are several enzymes (e.g. decarboxylases and dehydrogenases in the tricarboxylic acid cycle; phenylalanine ammonia lyase in the shikimic acid pathway of secondary metabolite synthesis) that do not bind Mn but are activated in the presence of Mn (Schmidt and Husted, 2019).

The biological functions and roles of Zn differ from those of Cu and Fe, since Zn is not a redox-active metal. Zn does not catalyse redox reactions but it acts as a Lewis base at the active sites of enzymes or is a structural component of proteins. As a cofactor, Zn is present in all six enzyme classes. Carbonic anhydrase is a very important Zn-dependent plant enzyme, which catalyses the interconversion of carbon dioxide and bicarbonate. Besides Cu, Fe, and Mn, Zn is also involved in ROS detoxification due to its binding to the active site of Cu/ZnSOD (Castillo-Gonzales *et al.*, 2018). There are multiple phosphatase enzymes that contain Zn. The active sites of alkaline phosphatases contain one magnesium (Mg) ion (Mg^{2+}) and two Zn^{2+} , while purple acid phosphatases contain an active site with one Fe^{3+} and one Zn^{2+} bound. Due to its regulatory role in phosphatases, Zn is involved in phosphate nutrition (Olczak *et al.*, 2003). Another interesting enzyme is S-nitrosogluthathione reductase, which regulates nitric oxide metabolism and signalling, and which contains catalytic and structural Zn ions (Lindermayr, 2018). Zinc-finger transcription factors bind Zn and undergo conformational changes leading to altered DNA-binding activity. This means that Zn

controls gene expression via the regulation of non-protein transcription factors (Takatsuji, 1998).

Ni is a very special metal, since the only known plant enzyme that requires Ni for its function is urease. Urea is the end-product of protein and ureide degradation, and urease detoxifies urea, catalysing the recycling of nitrogen (N) (Fabiano *et al.*, 2015).

Collectively, plant cells contain at least six types of metals that are essential for basic life functions. Essential metals are part of the metallome, which can be defined as the sum of the organic and inorganic metal forms in the cell, changes in which affect the physiological processes of plants (Singh and Verma, 2018).

Size matters: beneficial properties of nanoforms

Nanotechnology has become a dynamically emerging industry, with recent manufacturing developments in science and technology leading to the generation of materials of different shapes and sizes, with nanosized materials being among them. These advancements provide the basis for further design to create unique properties targeted toward specific applications (Khot *et al.*, 2012).

These nanosized materials, the so-called nanomaterials and nanoparticles (NPs), which have sizes typically below 100 nm, refer to a colloidal particle system that has received much attention in relation to chemical and biological applications because of their unique structure and tunable physico-chemical and biological properties, such as electrical and thermal conductivity, light absorbance, and catalytic activity, as well as antimicrobial activity, resulting in greater performance compared with their bulk counterparts (Fig. 1). These versatile properties can change as a function of the size and shape of the material. The dissolution, surface reactivity, and aggregation states of nanomaterials play key roles in their fate, lifetime, behaviour, and interactions in environments, often with global complex effects.

Metals and metal oxides in nanosized form can exist naturally in the environment; they are generated in nature either by biological species or through anthropogenic activities. They can also be manufactured (the so-called engineered nanomaterials) by several routes including chemical, physical, and biological methods (e.g. chemical reduction, photochemical methods, microwave processing, laser ablation, and grinding). The synthesis of NPs by chemical and physical means is more frequent, and the use of toxic and expensive chemicals in these processes limits their applications in daily life, human health, and agriculture-related applications. Therefore, the biogenic approaches, which utilize microbial or plant sources, have emerged recently as novel and potential methods for producing NPs. The rapid development of the synthesis procedures brought a significant increase in the number of nanomaterials

with desired properties being manufactured (Gebre and Sendeku, 2019).

Nanometals have a metal core composed of inorganic metal or metal oxide that is usually covered with a thin oxide shell. They can be produced with different morphologies, such as NPs, nanofibers, nanowires, nanotubes, or nanosheets, possessing different kinds of promising features, such as size-dependent qualities, high surface-to-volume ratio, specific surface plasma resonance, increased optical and magnetic properties, and enhanced antimicrobial activity (Jagdish *et al.*, 2018). Due to their reduced molecular size and increased surface area, and also because of changed interactions between molecules, nanomaterials have attributes that may be novel and often differ significantly from those of the bulk material, and which are not characteristics on the micro- or macroscopic scale, or in ionic form.

It is well known that the physico-chemical features of NPs are determined by a number of factors, such as NP size and morphology, or by the nature of the capping materials covering the particle surface (Rónavári *et al.*, 2017). Therefore, proper selection of synthesis methods including adequate reducing and stabilizing agents is crucial to achieve the desired particle properties. As metal NPs are highly reactive, and rapidly react with oxygen or biomolecules in living systems, surface modification is frequently applied to prevent aggregation. These surface adjustments can be achieved by the use of stabilizing agents, coating, or functionalizing (e.g. with oleic acid, sodium citrate, polymers, or biomolecules), promoting the dispersion of NPs and uniform particle size distribution. Various biomolecules (e.g. proteins, polysaccharides, and peptides) have been also used in the biosynthesis of NPs. These molecules interact with the NPs and are able to adsorb on to NP surfaces, establishing the so-called biomolecular corona (Bélteky *et al.*, 2021). This corona serves as a kind of organic complex on the surface of the NP that can improve the NP stability; moreover, it significantly determines and could improve the properties, behaviour, and evoked responses of NPs in the environment and living systems (e.g. enhanced particle release, less toxicity). Thus, the characteristics and fate of metal NPs can be designed and controlled on demand by using different biomolecules during their synthesis, as a function of the specific purpose.

Behaviour of essential nanometals on root/leaf surfaces and in plant cells

The uptake and transport of nanometals is a complex and in some cases contradictory process. To enter plant tissues, NPs need to breach multiple barriers that protect cells from hazards. These protective layers vary across species, tissues, and environmental conditions (Schwab *et al.*, 2016). In case of the root system, the first line of defence is a layer of rhizosphere bacteria and fungi. Symbiotic microenvironments influence the mobility of nanometals in soil, affecting their




Fig. 1. Unique properties of nanometals compared with the bulk form, and the phytoeffects and application of nanometals in plant cultivation. The nanoforms of essential metals possess greater surface area and hardness, specific optical, magnetic, and electrical properties, and show antimicrobial effects. Essential nanometals promote growth/yield and induce the synthesis of phytochemicals and protection against abiotic and biotic stressors. The positive effects of essential nanometals on the physiology of plants can be utilized during agricultural approaches such as nanoprimering, nanostimulation, nanofertilization, and nanopesticide application. For further details refer to the text.

uptake. For example, in ZnO and CuO NP-treated mung bean, *Pseudomonas chlororaphis* O6 affected the uptake of essential metals (Dimkpa et al., 2015). It is known that mycorrhizae can improve heavy metal tolerance in plants and increase the soil porosity in response to FeO NPs (Feng et al., 2013). The second physiological barrier near plant roots consists of mucilage and exudates (M/E) excreted into the soil. These biomolecules have a role in sensing and protecting the root system on the outer cell surface (McNear, 2013). M/E can reversibly absorb various NPs near the root cap and root hairs (Aubert

et al., 2012; Zhu et al., 2012; Lee et al., 2013). M/E has a role in acidifying the root microenvironment, which could intensify the dissolution of nanometals (Lv et al., 2015). It is often reported that nanometals in the roots are smaller in diameter than those in the treatment solution, and in some cases they are completely dissolved (Lv et al., 2015). The effectiveness of M/E in alleviating NP stress is variable; it is more pronounced in the case of small and positively charged nanomaterials (Zhu et al., 2012). The aerial plant parts are protected by a lipophilic cuticle barrier that consists of epicuticular wax crystals, which

create a robust and almost impenetrable layer (Albersheim, 2011). It is highly unlikely that hydrophilic NPs can cross this barrier; however, thin permeable regions with uptake or secretory functions are suggested to function as uptake sites. Beyond the root system, nanomaterials can enter the plant body via the opened stomata, where the water solubility of ZnO and MgO resulted in better translocation to the roots (Wang *et al.*, 2013). Plant cell walls are protective layers of fibres and other composites with relatively continuous porosity (Schwab *et al.*, 2016). The pore size of the cell walls (defined as the space between the cell wall components within the wall matrix) is mostly between 0.6 nm and 4.8 nm (Eichert and Goldbach, 2008; Kurczyńska *et al.*, 2021), which is favourable for extremely small neutrally charged NPs. There is experimental evidence that nanometals such as Fe₃O₄-NP (18 nm) or ZnO NP (8 nm) enter the cytoplasm (Molnár *et al.*, 2020; Iannone *et al.*, 2021), which can be made possible by the NP-induced modification of the chemical (macromolecule composition) and physical (pore size) properties of the cell wall, as suggested by Kurczyńska *et al.* (2021). The Casparian strip is the major barrier to apoplastic transport, and it has been suggested that NPs can enter the pericycle via the disruption of this barrier in early lateral root development (Lv *et al.*, 2015). Metallic NPs are usually charged, making them unable to cross the plasma membrane by diffusion. However, it is documented clearly that metal NPs enter the cells (reviewed by Bhirde *et al.*, 2011; Lv *et al.*, 2019), suggesting possible uptake mechanisms. It has been documented that NP endocytosis can occur via multiple pathways (Onelli *et al.*, 2008; Palocci *et al.*, 2017); however, results with essential nanometals are scarce. Moreover, NP binding to carrier proteins or ion channels has also been proposed as an NP uptake mechanism (Brandenberger *et al.*, 2010; Shukla *et al.*, 2016). Symplastic transport between cells via plasmodesmata, which are 20–50 nm in diameter, has also been proposed. Small particles of ~3 nm can be transported via plasmodesmata between cells (Dietz and Herth, 2011; recently reviewed by Kurczyńska *et al.*, 2021). Furthermore, NPs are assumed to influence the size exclusion limit of plasmodesmata, allowing bigger NPs to travel through them (Kurczyńska *et al.*, 2021). Interorganellar translocation of NPs has been documented in various plant species. For instance, FeO NPs were taken up by pumpkin roots and transferred to the leaves via the xylem (Zhu *et al.*, 2008; reviewed by Peng *et al.*, 2020).

In the past, the detection of nanomaterials in plant tissues provided contradictory results, partly due to the different detection techniques, nanomaterial transformation processes, nanomaterial sizes and properties, and experimental designs used in studies (Montes *et al.*, 2017; Wojcieszek *et al.*, 2020). Nowadays, numerous methods are being used in nanomaterials research, which will be summarized briefly below. Synchrotron-based X-ray fluorescence mapping and X-ray absorption spectroscopy is one of the newest methods in this field. It can provide details about the elemental composition, location, and chemical properties of nanomaterials. Due to the nature of

this method, it can detect biotransformation of nanomaterials, making it an excellent tool to visualize the behaviour of NPs in live tissues (Castillo-Michel *et al.*, 2017). Inductively coupled plasma-mass spectrometry (ICP-MS) is able to detect metals in tissues in the extremely low ng l⁻¹ range. The technique is limited due to the sample preparation method, since it can measure metal content only from acid-decomposed samples. A new technique, single-particle ICP-MS, has been developed, which is able to measure dissolved particles, and can measure each NP individually (Wojcieszek *et al.*, 2020). It is able to detect NP alterations, which is crucial information in samples. Imaging methods used in visualizing the uptake and localization of nanomaterials in tissues include transmission and scanning electron microscopy (TEM and SEM), matrix-assisted laser desorption/ionization (MALDI), and laser ablation (LA) ICP-MS (Montes *et al.*, 2017; Wojcieszek *et al.*, 2020). The most prevalent technique is TEM, which can provide information about the localization, size, and aggregation of nanomaterials in cells. It has a high spatial resolution (<1 nm); however, the interpretation of the data requires multiple control samples. To evaluate the behaviour of nanomaterials in tissues, a combination of various methods is recommended to obtain detailed results.

It has been well documented that the synthesized engineered form of NPs changes when they are released into the environment. Nanomaterials are transformed by natural organic matter in ways including dissolution, chelation, precipitation, and reactions with various functional groups (Maurer-Jones *et al.*, 2013). Additionally, plant biochemical pathways have a tendency to modify internalized nanomaterials. In aqueous solutions, the stability of nanomaterials is influenced heavily by dissolution and aggregation, which reduce surface size. Ion release from metal NPs is considered to be a quick process in the case of Cu (Conway *et al.*, 2015) and Zn (David *et al.*, 2012) NPs. Dissolution of Zn ions from ZnO NPs is hypothesized to be one of the main pathways of NP toxicity (Kouhi *et al.*, 2014). In realistic conditions, natural organic matter can limit NP aggregation rates, thus stabilizing the solution (Liu *et al.*, 2012). Different soil microenvironments also transform nanomaterials. In clay soils, reducing conditions are present, whereas in sandy soils oxidative transformations are favoured (Zhang *et al.*, 2020). On leaf surfaces in the presence of oxygen, redox-active metals such as Cu NPs can receive an oxide coating (Tani *et al.*, 2018). Additionally, ZnO NPs have been reported to be transformed into other inorganic (Zn-nitrate, Zn-phosphate) and organic (Zn-citrate) forms by desert plants (De la Rosa *et al.*, 2011). Biomolecular coatings on NPs called coronas (as described above) have a significant effect on NP properties. The chemical composition of the corona can influence the uptake (Francia *et al.*, 2019), dissolution (Collin *et al.*, 2016), and spatial distribution of NPs. In the case of pumpkin xylem fluid, the corona mainly consists of selectively bound proteins on the surface of CuO NPs (Borgatta *et al.*, 2021). It is hypothesized that the

positively charged NP surface interacts with partially negatively charged functional groups in proteins. Interestingly, it has been reported that metal ions can transform into NPs that are naturally in certain plant tissues (Kuppusamy *et al.*, 2016).

Taken together, the uptake, translocation, and transformation of (essential) nanometals in biological systems is still not understood fully. Analysing the transformed NPs in plant tissues has several methodological difficulties; however, detailed knowledge of these processes is of paramount importance.

The hormetic effects of essential nanometals in plants: an omics approach

The effect of essential nanometals on plants depends on various factors, including the characteristics of the nanometal and also of the interacting plant partner (Fig. 2). The physico-chemical properties of nanometals, such as their hydrodynamic size, morphology, crystal structure, agglomeration, surface charge, and modification, influence the degree of their effects on plants. From the plant side, the species dependence of the effects of nanometals has been described, and the metal accumulation capacity and tolerance of the plant species also determine the effects of the nanometal. The evoked response to the presence of nanometals may also depend on the developmental stage

of the plant. From the methodological point of view, the duration of exposure to nanometals and the way in which they are administered also influence their effects on plants (Fig. 2).

Literature data agree that plants respond to the presence of essential nanometals in a concentration-dependent way, which—using a toxicological term—is called hormesis (Agathokleous *et al.* 2019b). The response of plants to metallic micronutrients and nanometals depends on the concentration in such a way that stimulation occurs in a lower concentration range, whereas higher doses exert adverse effects on plants (Fig. 3).

In their work evaluating the relevant literature, Agathokleous *et al.* (2019a) revealed that the extent of the hormetic plant responses to nanomaterials (not only to nanometals) depends on the size and type of the nanomaterial, since different nanomaterials caused different hormetic effects when applied at the same concentration under the same experimental conditions. The amplitude of the hormetic response proved to depend also on whether the nanomaterial contains essential or non-essential nutrients. Based on the analysis, element release from the NPs does not seem to be the major mechanism responsible for the stimulatory effect of low NP concentrations (Agathokleous *et al.* 2019a).

According to the holistic view, plant responses to essential nanometals are the sum of changes at the genome, transcriptome, proteome, metabolome, ionome, and phenome




Fig. 2. Factors determining plant responses to nanometals and the different levels at which plant responses can be examined. Among the characteristics of nanometals, their type, dosage, size, agglomeration, crystal structure, and surface charge are the most important determining factors. The response to essential nanometals also depends on the species, metal tolerance capacity, and developmental stage of the plant. Additionally, the method of application (e.g. foliar spray, irrigation, hydroponics) and the period of exposure to the nanometal also influence its effect. Plants show multi-level responses to nanometal exposure. Genomics, transcriptomics, proteomics, metabolomics, and phenomics studies provide a holistic view about the complex effects of essential nanometals on plants.




Fig. 3. Hormetic effect of increasing doses of essential nanometals on plants. At lower concentrations, nanometals promote plant biomass production and physiological processes, whereas higher nanometal concentrations exert inhibitory effects.

levels (Fig. 2). Few studies have evaluated transcriptome-level changes in plants treated with essential nanometals (Table 1). For example, Landa *et al.* (2017) observed that exposure to CuO NPs (10 mg l^{-1}) altered the expression of 922 genes (mainly involved in stress responses and Cu binding/transport), whereas Cu ion treatment resulted in the up-regulation of 482 genes in *Arabidopsis thaliana* L. roots. Moreover, the similar transcription profiles of NP- and ion-exposed plants indicates that the release of Cu^{2+} ions from CuO NPs can be considered as the main contributor to phytotoxicity. More recently, 2270 and 4264 genes were found to be differentially expressed in response to exposure to 100 mg l^{-1} and 1000 mg l^{-1} CuO-NPs, respectively, in lettuce. The affected genes belonged to the ABC transporter family and heavy metal-associated isoprenylated plant proteins (HIPPs), which are involved in endocytosis, metal ion binding, and transport (Xiong *et al.*, 2021). Moreover, genes associated with cell wall biogenesis, antioxidant processes, and photosynthesis were also affected by CuO NP treatment of lettuce (Xiong *et al.*, 2021). Using metabolomics platforms, 65 metabolites, including jasmonates, scopoletin, and glucosinolate derivatives, were detected in leaves of nano CuO-treated *Arabidopsis* (Chavez Soria *et al.*, 2019). Targeted analysis of the jasmonic acid (JA) pathway revealed that CuO NP up-regulates JA precursors and down-regulates JA levels, indicating the involvement of JA-associated defence signalling in plant responses to CuO NPs. As for proteome-level changes induced by CuO NPs, the abundance of proteins involved in glycolysis and the tricarboxylic acid cycle increased in stress-tolerant wheat (*Triticum aestivum* L.) varieties, while photosynthesis- and tetrapyrrole synthesis-related proteins were down-regulated upon nano CuO exposure (Yasmeen *et al.*, 2018). Using a

gel-free proteomic approach in grains of CuO NP-exposed wheat, elevated levels of proteins associated with starch degradation and glycolysis have been detected (Yasmeen *et al.*, 2017). Multiple studies revealed that nano ZnO extensively modifies the gene expression profiles of *Arabidopsis* and tomato (*Solanum lycopersicum* L.), mainly up-regulating biotic- and abiotic stress-related genes (Landa *et al.*, 2012; Wan *et al.*, 2019; Sun *et al.*, 2020). Furthermore, ZnO NPs induced changes in the secondary metabolism and increased the amino acid and sugar contents in the leaves and roots of tomato (Sun *et al.*, 2020). In response to nano ZnO exposure of soybean (*Glycine max* L.), 298 root proteins showed specific modifications, whereas in the shoot the expression of only 36 proteins was altered exclusively by nano ZnO (Hossain *et al.*, 2016). Using gel-free proteomics, Yasmeen *et al.* (2016) found that Fe NP treatment affected proteins mainly associated with photosynthetic processes in wheat. Moreover, an enhanced abundance of light-reaction-related proteins was associated with salt tolerance. In Fe NP-exposed wheat seeds, Yasmeen *et al.* (2017) described the accumulation of proteins involved in starch degradation, glycolysis, and the tricarboxylic acid cycle. In a recent metabolomics study, 21 and 53 dysregulated metabolites were detected in molybdenum trioxide (MoO_3) NP-treated corn (*Zea mays* L.) and wheat leaves, respectively. Notably, more metabolic pathways were affected by the NPs in wheat than in corn. The untargeted analyses were completed by the analysis of individual groups of metabolites and, for example, asparagine, fructose, reduced glutathione, and mannose were found to be modulated by MoO_3 NPs in both corn and wheat (Huang *et al.*, 2021).

Promoting effects of essential nanometals on non-metal-accumulator plants and their practical application in plant cultivation

Hormetic effects of essential nanometals on germination, vegetative growth, and yield

Since seed germination is an early and sensitive phase of plant ontogenesis, it is important to determine the influencing effects of different nanomaterials (including essential nanometals) on it, especially in crops. Thus, the concentration-dependent hormetic effect of essential nanometals on germination and seedling development has been revealed by several authors in the past years, and has been recently reviewed by Szöllősi *et al.* (2020). More recent literature shows a trend for authors to prefer biological, plant-based NP production over chemical synthesis. For example, ultra-small ($\sim 6.6 \text{ nm}$) CuO NPs derived from plant-based synthesis (green tea extract) at a lower concentration positively affect the germination and seedling growth of lettuce, whereas elevated doses of nano CuO (nCuO; hereafter, nanoforms of essential metals and their compounds are indicated with the prefix n) have an adverse effect. The toxicity of CuO NPs was shown to be associated with inhibited antioxidant

Table 1. Transcriptomic, proteomic, and metabolomic studies in plants treated with essential nanometals

Essential nanometal	Plant species	Nanometal treatment	Approach/technology	Most relevant effects	Reference
Transcriptomics					
nCuO	<i>Arabidopsis thaliana</i> (Col-0), root	10 mg l ⁻¹ nCuO (<50 nm) in hydroponics	Microarray/qRT-PCR	922 regulated genes, e.g. superoxide dismutase (<i>CSD1</i>), copper chaperone (<i>CCS1</i>), copper transporter (<i>COPT2</i>), laccase (<i>LAC3</i>)	Landa <i>et al.</i> , (2017)
nCuO	<i>Lactuca sativa</i>	100 or 1000 mg l ⁻¹ CuO NP (~40-200 nm) via foliage	RNA sequencing/qRT-PCR	2270 (100 mg l ⁻¹ CuO NP) and 4264 (1000 mg l ⁻¹ CuO NP) modified genes, e.g. cellulose synthase A catalytic subunit 4 (<i>CESA4</i>), photosystem I chlorophyll <i>a/b</i> -binding protein 3 (<i>LHCA3</i>), chlorophyll <i>a/b</i> -binding protein 3C (<i>CAB3C</i>), photosystem I reaction centre subunit (<i>psaK</i> , <i>psaD</i> , <i>psaL</i>), and photosystem II reaction centre protein (<i>psbB</i> and <i>psbZ</i>) up-regulated, photosystem I chlorophyll <i>a/b</i> -binding protein 3 (<i>LHCA3</i>), chlorophyll <i>a/b</i> -binding protein 3C (<i>CAB3C</i>), photosystem I reaction centre subunit [<i>psaK</i> , <i>psaD</i> (ncbi_111898767), <i>psaL</i>], and photosystem II reaction centre protein (<i>psbB</i> and <i>psbZ</i>) down-regulated.	Xiong <i>et al.</i> , (2021)
nZnO	<i>Arabidopsis thaliana</i> Col-0, root	100 ppm ZnO NP (<100 nm) (or TiO NP (<150 nm) or fullerene soot (>7% fullerene) in hydroponics	Microarray/qRT-PCR	660 up-regulated genes (involved in abiotic and biotic stress responses), 826 down-regulated genes (involved in cell organization and biogenesis, including translation, nucleosome assembly, and microtubule-based process) by ZnO NPs.	Landa <i>et al.</i> , (2012)
nZnO	<i>Arabidopsis thaliana</i> Col-0, whole seedlings	100 or 200 mg l ⁻¹ ZnO NP (20-45 nm) in agar-solidified medium	RNA sequencing/qRT-PCR	A total of 1024 genes were up-regulated and 447 genes were down-regulated in nZnO-treated seedlings. After 3 d of recovery, 71% of DEGs had returned to normal levels in nZnO-treated seedlings.	Wan <i>et al.</i> , (2019)
nZnO	<i>Solanum lycopersicum</i> , leaf, root	Foliar spraying with ZnO NPs (20 and 100 mg l ⁻¹)	RNA sequencing/qRT-PCR	808 up-regulated and 103 down-regulated genes in ZnO NP-treated tomato leaves (e.g. nutrient, amino acid and sugar transporters, genes involved in sugar metabolism), and 395 up-regulated and 1127 down-regulated genes in tomato roots (e.g. element transporters, cutin, wax, suberin synthesis, flavonoid biosynthesis, sugar metabolism).	Sun <i>et al.</i> , (2020)
Proteomics					
nFe	<i>Triticum aestivum</i> seedlings (Pakistan-13, NARC-11 varieties), shoot	5 ppm Fe NP (20 nm) via root system	Gel-free/label-free proteomics	The abundance of proteins related to photosynthesis (e.g. ribulose biphosphate carboxylase small chain clone 512, ribulose biphosphate carboxylase/oxygenase activase A, phosphoglycerate kinase) and proteins involved in amino acid metabolism (e.g. ketol acid reductoisomerase, probable LL diaminopimelate aminotransferase) was decreased in the drought-tolerant variety. The abundance of proteins involved in photosynthesis (e.g. ribulose biphosphate carboxylase small chain clone 512, phosphoglycerate kinase) was increased by Fe NP, while amino acid metabolism-related proteins (e.g. glutamate glyoxylate aminotransferase 1, ketol acid reductoisomerase) decreased in the salt-tolerant variety. Of photosynthesis-related proteins, light reaction was enhanced in the salt-tolerant variety compared with the drought-tolerant variety on exposure to Fe NPs.	Yasmeen <i>et al.</i> , (2016)
nCuO or nFeO	<i>Triticum aestivum</i> (galaxy-13, NARC-11, Pakistan-13 varieties), grain	20, 25, 35, or 40 ppm CuO NP (15–30 nm) or FeO NP (20–30 nm) in soil	Proteomics/gel-free	In 25 ppm CuO-treated galaxy-13, Pakistan-13, and NARC-11 wheat varieties, 58, 121, and 25 proteins, respectively, were changed in abundance. Glycolysis and protein degradation-related proteins were increased by both nanometals.	Yasmeen <i>et al.</i> , (2017)

Table 1. Continued

Essential nanometal	Plant species	Nanometal treatment	Approach/technology	Most relevant effects	Reference
nCuO	<i>Triticum aestivum</i> seedlings (Pakistan-13, NARC-11 varieties), shoot	1, 5, 10, or 50 ppm CuO NP (<50 nm) in solution	Proteomics/gel-free	Abundance of proteins related to glycolysis and the tricarboxylic acid cycle was increased by CuO NPs. Proteins related to photosynthesis and tetrapyrrole synthesis were decreased by CuO NPs in both varieties.	Yasmeen <i>et al.</i> , (2018)
Metabolomics nZnO	<i>Glycine max.</i> , leaf, root		Metabolomics	In nZnO-exposed root, 104 changed proteins associated with secondary metabolism, cell organization, and hormone metabolism were detected. In the leaf, 16 common proteins were significantly changed in NP-exposed soybean, predominantly associated with photosystem and protein degradation.	Hossain <i>et al.</i> , (2016)
nCuO	<i>Arabidopsis thaliana</i> (Col-0), root, leaf, flowering shoot	10 mg l ⁻¹ CuO NP (490 ± 70 nm) in hydroponics	LC-QToF-MS and LC-Orbitrap-MS untargeted metabolomics platforms	65 metabolites altered by CuO NPs (e.g. jasmonates, scopoletin and glucosinolate derivatives). Targeted analysis: up-regulation of JA precursors (12-OPDA and dinor-12-OPDA), and down-regulation of the final product (JA).	Soria <i>et al.</i> , (2019)
nMoO ₃	<i>Triticum aestivum</i> , <i>Zea mays</i> , leaf	100 or 500 mg l ⁻¹ MoO ₃ NPs (~375–399 nm) in vermiculite-grown plants	UHPLC, LC-MS/MS metabolomics platform	21 dysregulated metabolites in corn leaves and 53 in wheat leaves. Five more metabolomic pathways were perturbed in wheat leaves compared with corn leaves. Targeted analysis: the amounts of asparagine, fructose, reduced glutathione, and mannose were re-programmed in both corn and wheat root and leaf.	Huang <i>et al.</i> , (2021)

defence and intensified S-nitrosothiol signalling (Pelegriño *et al.*, 2020). Similarly, plant-extract-derived, phytochemical-capped FeO NPs were used to improve the germination capacity of rice (*Oryza sativa* L.), whereas the same concentrations of the ionic form (FeSO₄) had an inhibitory effect (Afzal *et al.*, 2021). The germination-promoting effect of nFeO was due to the enhancement of water uptake, alpha-amylase activity, soluble sugar content, and antioxidant enzyme activities. The observed nFeO-induced ROS production may be involved in cell wall loosening and endosperm weakening. Moreover, the contents of macro- and micronutrients in the endosperm of the nFeO-treated rice seeds increased, supporting that FeO NPs can be used effectively as a 'nano-nutrient' for efficient germination and growth (Afzal *et al.*, 2021). In another recent study, MnO NPs were synthesized using onion (*Allium cepa* L.) bulb extract and were applied for seed priming of watermelon (*Citrullus lanatus* L.) genotypes (Kasote *et al.*, 2021). In comparison with the bulk forms (KMnO₄ and Mn₂O₃), the seed treatment with nMnO proved to be less toxic. Changes in the leaf metabolome of seedlings showed a clear MnO NP effect only in case of the triploid line. Furthermore, the levels of phytohormones such as abscisic acid, gibberellic acid, salicylic acid, and JA changed in a genotype-dependent manner in MnO NP-treated watermelon, which emphasizes the genotype dependence of plant responses to nanometals (Kasote *et al.*, 2021). Similarly, the effect of nFe and nCu on seedling development varied depending on the genotype of Scots pine

(*Pinus sylvestris* L.) (Kalyakina *et al.*, 2021). A hormetic effect of Mo NPs on seedling development has been described, without a significant effect on germination of rice (Adhikari *et al.*, 2013). In a recent study, rice seedlings were treated with high doses of chemically synthesized Mo NPs (α-MoO₃ and MoS₂) and decreases in pigment content and antioxidant activities were described, indicating Mo NP-induced oxidative stress in rice seedlings. Furthermore, the authors concluded that, based on the poor translocation and low accumulation without any notable effect on seedling growth, a concentration of MoS₂ NPs of 100 ppm appears to be a promising treatment for future application during rice cultivation (Sharma *et al.*, 2021).

Far fewer studies have dealt with the later developmental stages of plants, for example, flowering, and yield in connection to essential nanometals. A study found that treatment of soybean seeds with Fe, Cu, or Co NPs (zerovalent) slightly induced germination and, when used in a field experiment, increased chlorophyll content and other parameters such as number of nodules per root, number of pods per plant, pod weight, and 1000-grain weight; nCo exerted the most efficient promoting effect, followed by Cu and Fe NPs (Ngo *et al.*, 2014). In carrot, the highest rate of root and shoot growth induction and yield was achieved by a combined treatment with ZnO and FeO NPs (Elizabeth *et al.*, 2017). In a recent field study, the combined foliar application of ZnSO₄ and nZnO produced the highest yield and improved nutrient content of rice compared with ZnSO₄-fertilized plants, indicating that the application of

ZnO NPs as additives in fertilizers is beneficial in agricultural practice (Elshayb *et al.*, 2021). Similarly, the application of nFe on greenhouse-grown tomato significantly enhanced growth and yield, and the promoting effect of nFe proved to be more intense than that of conventional Fe fertilizers (El-Desouky *et al.*, 2021).

Most of the studies on the effects of essential nanometals on plant yield are descriptive, without revealing the biochemical and molecular mechanisms explaining the observed effects. Future studies should include an omics approach in order to understand the responses of developing plants to essential nanometals at different levels.

Seed priming by essential nanometals for inducing germination capacity and vegetative and reproductive growth seems to be a promising way to increase the effectiveness (in terms of cost, time, quality, and quantity) of crop cultivation. The effects, however, depend on several factors, making the development of general treatment protocols difficult. In the case of seed nanopriming, the use of plant-synthesized NPs has become more and more desirable; thus, 'green nanopriming' will surely be a focus of future research (Song and He, 2021).

Essential nanometals induce phytochemical synthesis in plants

Beyond affecting primary metabolism and consequently promoting growth and development, nanometal treatments elicit the production of secondary metabolites in diverse plant species. More than 5000 phytochemicals have been identified in fruits, vegetables, and grains. Among the most valuable phytochemicals are vitamin C, folate, provitamin A, potassium (K), calcium (Ca), Mg, flavonoids, phenolic acids, alkaloids, carotenoids and fibres, which have a wide range of applications in cosmetics, the food industry, pharmacology, and medicine due to their beneficial effects. According to the very recent review of Rivero-Montejo *et al.* (2021), among the essential nanometals, mostly nCu and nZn as biostimulants have been examined so far. Nanostimulation by the application of Cu or Zn NPs increases the total phenol, flavonoid, rebaudioside A, stevioside, thymol, and carvacrol contents of diverse plant species (Juarez-Maldonado *et al.*, 2018, Rivero-Montejo *et al.*, 2021). More recently, some studies have applied essential nanometals as biostimulants in plants exposed to abiotic stress. For example, foliar application of ZnO NP on gold-of-pleasure plant (*Camelina sativa L.*) exposed to salt stress increased the total phenol content and the levels of anthocyanins, carotenoids, Ca, Zn, and phosphorus (P) compared with plants exposed to salt stress alone (Hezaveh *et al.*, 2020). In another study, bell pepper (*Capsicum annuum L.*) exposed to salt stress was treated with selenium (Se), silicon (Si), or Cu NPs, and in each case it was found that the NPs stimulated ascorbate peroxidase, glutathione peroxidase, catalase, and phenylalanine ammonia lyase activities and increased the content of phenols, flavonoids, glutathione, β -carotene,

and yellow carotenoids in the fruits, suggesting that the application of Se, Si, or Cu NPs to bell pepper plants during conditions of high salinity is an efficient way to increase the content of bioactive compounds in fruits (González-García *et al.*, 2021).

For this subfield, in addition to observing changes in compound composition, it is also necessary to understand the molecular mechanisms behind the effects of essential nanometals. Nevertheless, nanobiostimulation is a promising approach because low doses of NPs can trigger the accumulation of beneficial plant compounds, thus making cultivation cost-effective. As during nanostimulation nanometals may be introduced directly into products for human consumption, further efforts should be made to assess the toxicological risks.

Essential nanometals intensify biotic and abiotic stress tolerance in plants: focusing on nutrient deficiency

In association with global climate change and with the increasing consumption of plant-based food, it is becoming increasingly important to intensify the resilience of (crop) plants to environmental stresses (Kilian *et al.*, 2021). The living environment (biotic factors) exerts notable impacts on the physiological processes of plants. Viruses, viroids, phytoplasma, bacteria, fungi, oomycetes, nematodes, insects, and weeds cause major problems in crop production. According to certain estimates, yield losses of major crops due to pests range from 20% to 40% or even more than 50% in certain cases (Pestovsky and Martinez-Antonio, 2017), causing considerable financial damage for farmers (Hajong *et al.*, 2019). Therefore, the introduction of new approaches such as the utilization of nanomaterials, including nanometals, in agricultural practices is urgently needed. Crop yield loss can be reduced by detecting infections as soon as possible; thus, an important area is the development and application of NPs as sensors in crop production (Li *et al.*, 2020). Among the nanometals, mostly nCuO, nZnO, nFe₂O₃, nFe₃O₄, and nNiO have potential as nanopesticides in disease management practices against viruses, bacteria, fungi, and oomycetes. The current interest and significance of this topic is indicated by the large number of detailed literature reviews that have been published recently (e.g. Djiwanti and Kausik, 2019; Hajong *et al.*, 2019, de Oliveira, 2021; Farooq *et al.*, 2021).

Besides biotic factors, non-living environmental conditions (abiotic stress factors), such as water or nutrient limitations, heat, cold, salinity, and UV radiation, reduce the average yield of most major crop plants by more than 50% (Hirayama and Shinozaki, 2010). Based on the experimental data accumulated in the past years, it can be stated that nanomaterials, including nanometals, provide a promising strategy to address these environmental challenges that plants face. This topic is a focus of several recent reviews (Kahn and Upadhyaya, 2019; Zhao *et al.*, 2020; Singh *et al.*, 2021; Thounaojam *et al.*, 2021).

One of the significant abiotic stressors influencing crop quantity and quality is the lack or inadequate accessibility of essential nutrient elements in soils. Plants require optimal availability of at least 14 macro- (N, P, K, S, Ca, Mg) and micronutrients (Cu, Zn, Mn, B, Mo, Si, Ni, and Cl) (Kirkby, 2012). Particularly in developing countries, low soil fertility, reduced availability of mineral nutrients in soils, improper nutrient management strategies, and the lack of plant genotypes possessing tolerance to nutrient limitations collectively lead to food insecurity, malnutrition, and degradation of the ecosystem. About 50% of the world's population is endangered by an inadequate micronutrient supply, highlighting the necessity of implementing new strategies (Kumari *et al.*, 2021).

Plant breeding is a good option to develop plant genotypes with improved potential to acclimatize to nutrient-deficient conditions (Kumari *et al.*, 2021). Another option is to increase the available element content of the soil by fertilization; however, the application of conventional fertilizers has multiple drawbacks. Most importantly, the nutrient use efficiencies of conventional fertilizers are only around 30–35% for N, 18–20% for P, and 35–40% for K, meaning that the applied nutrients are utilized by plants with low efficiency. Moreover, the large-scale application of conventional fertilizers causes irreversible damage to the soil structure, mineral cycles, and soil microbiome, as well as to plants (Mohammadi and Sohrabi, 2012; Solanki *et al.*, 2015; Riaz *et al.*, 2020). Nanofertilizers release nutrients slowly and steadily, providing balanced nutrition for crops without causing damage to soils, microbes, or plants (Elemike *et al.*, 2019; Mittal *et al.*, 2020). Using nanoforms of fertilizers reduces nutrient loss from the soils; thus, this is an 'eco-friendly' way of land use that supports sustainable agriculture (Subramanian *et al.*, 2015). However, despite the significant practical potential of nanofertilizers, the amount of laboratory data about the effects of NPs and nanometals on element-deficient plants is surprisingly low.

Inadequate Zn availability for crops is a relatively common problem worldwide. Bala *et al.* (2019) applied ZnO NP treatments (at 0.5, 1, or 5 g l⁻¹) to the leaves of rice grown in soil with adequate or low Zn supply. The ZnO NPs could be detected on the leaf surface near to the stomata by SEM, indicating that one putative route for internalization of NPs is via stomata. The foliar application of ZnO (5 g l⁻¹) has positive effect on the soil microbiome, possibly due to the increased Zn ion concentration of the roots. For the Zn-deficient plants, ZnO NP treatment improved vegetative growth and yield parameters (e.g. tiller number, 1000-grain weight), enhanced Zn content, and altered levels of nutrients in the root, shoot, and grain. These results indicate that foliar nZnO fertilization successfully reverts the symptoms of Zn deficiency in rice (Bala *et al.*, 2019).

The recent work of Sun *et al.* (2020) revealed that the application of nZnO to tomato plants through the foliage increases Fe accumulation and efficiently improves tolerance of Fe deficiency due to decreasing Fe-deficiency-induced oxidative damage and enhancing the contents of essential metal

elements. Additionally, transcriptomic and metabolomic analyses indicated that the expression of genes involved in the antioxidant system, transport, carbon and N metabolism, and secondary metabolism was increased by ZnO NP spraying, possibly contributing to the Fe-deficiency tolerance of tomato (Sun *et al.*, 2020).

The efficiency of nCuO as a fertilizer has also been studied; however, those experiments did not involve element-deficient plants. Soil-grown Chinese scallions (*Allium fistulosum* L.) were treated with nCuO, bulk CuO, and Cu in ionic form (CuSO₄). The nanoform of CuO proved to be the most efficient in increasing the contents of Ca, Fe, or Mg, as well as alliin (Wang *et al.*, 2020). Similar results were obtained by Kohatsu *et al.* (2021) in a study in which the foliar application of biogenic CuO NPs triggered the accumulation of K, Na, S, Ag, Cd, Cr, Cu, and Zn in lettuce leaves without causing necrosis, while CuSO₄ caused the most intense changes in the mineral element content of lettuce and caused necrotic damage. These findings indicate that nCuO is a more effective fertilizer than the other forms of Cu for lettuce cultivation (Kohatsu *et al.*, 2021).

Peanut (*Arachis hypogaea* L.) is sensitive to limited Fe supply, and therefore nFe fertilizers have great potential for this crop. In the study of Rui *et al.* (2016), Fe₂O₃ NPs increased growth and biomass production, and the SPAD values (a measure of chlorophyll content) of peanut plants. Moreover, nFe₂O₃-induced changes in the levels of certain phytohormones [abscisic acid, gibberellic acid (GA₄₊₇, GA₃), zeatin riboside, dihydrozeatin riboside] and antioxidant activities (peroxidase, SOD, catalase) were associated with the growth induction. Additionally, nFe treatment efficiently increased the Fe content of peanut plants, indicating that nFe₂O₃ is a promising nanonutrient for peanut plants, although the putative amelioration of damage in Fe-deficient plants would have provided additional relevant data (Rui *et al.*, 2016).

Not all types of nFe can be effective as a source of Fe for plants. A recent study indicated that despite the fact that supermagnetic iron oxide NPs (SPIONs, ~12.5 nm) internalize in the roots, this treatment alone is not effective in supplying Fe to summer squash (*Cucurbita pepo* L.); however, the authors suggest that applying SPIONs together with Fe(III-EDTA) can be beneficial in the fertilization of *Cucurbita* spp. (Tombuloglu *et al.*, 2019).

The phyto-effect of nFe also depends on the plant species and their Fe-acquisition strategies. Both cucumber and maize used Fe and P derived from FePO₄ NPs more efficiently than when supplied as bulk material, but cucumber mainly used FePO₄ NPs as a source of P, while maize preferred FePO₄ NPs as a source of Fe. Interestingly, cucumber roots were not able to take up the nFe, possibly due to agglomeration. These comparative results clearly indicate that the nutrient utilization of roots exposed to FeSO₄ NPs is affected by species-specific metabolic responses (Sega *et al.*, 2020).

It is also true for NPs studied as nanofertilizers that the effects on the plant may differ depending on the method of

application. One of the few studies examining nMn found that foliar exposure resulted in higher shoot and grain Mn contents, lower soil nitrate-N, and higher soil and shoot P compared with soil application of nMn in wheat plants. These results indicate that applying nMn via the foliage can exert more intense effects on nutrient composition of wheat. Additionally, exposure to the nanoform of Mn via the soil could mildly affect plants, differing from exposure to bulk or ionic Mn (Dimpka *et al.*, 2018).

The above examples show that, with few exceptions, essential nanometals are better utilized by plants and have a more positive effect on plant biomass production, as well as more efficiently alleviating nutrient deficiency, than bulk materials. One of the reasons behind this could be that due to their nanometric size, a greater amount of NPs may reach the root surface compared with the bulk forms. Moreover, NPs may be more rapidly and more efficiently dissolved due to their large surface-to-volume ratio. Additionally, the literature clearly points out that the beneficial effects of essential nanometals depend on factors such as the form, concentration, and route of administration. Furthermore, the plant species and its element acquisition strategy may influence the observed effects of nanofertilizers (Kalwani *et al.*, 2022).

Conclusions and future perspectives

Plants use Cu, Fe, Mo, Mn, Zn, and Ni as essential metals for operating vital physiological processes, such as photosynthesis, cellular respiration, and antioxidant protection. Nanoforms of these metals possess a number of favourable properties over bulk materials, such as a large surface area and superior electrical, optical, magnetic, and biological properties.

Essential nanometals come into contact with plant leaves and roots and enter the tissues through diverse pathways. In the case of roots, soil microbes and root exudates may help or limit NP uptake, while in the case of leaves opened stomata or thin permeable regions of the epidermis can serve as entry points for NPs. At the cellular level, the movement of NPs is limited by the cell wall pore size and the Casparian strip, and the uptake of NPs into the cytoplasm may occur by endocytosis or interaction with carriers and channels. Between the cytoplasm of adjacent cells, NPs may travel through the plasmodesmata.

The detection of nanometals in plant tissues faces technical difficulties, so it is recommended to use several complementary methods (e.g. TEM, ICP-MS). Biotransformation of nanometals may take place in the plant tissues, with the ionic form of the metal being released from the nanometal taken up by the plant. Our knowledge about the internalization, behaviour, and interactions of essential nanometals in plant cells is incomplete, so more studies are needed to examine the transport mechanisms and biomolecular interactions from a physico-chemical point of view.

A more actively studied topic is the effect of essential nanometals on plants, which is concentration dependent

and can therefore be considered as hormesis. Metal ions also exert hormetic effects on plants, but in the case of nanometals there are some special influencing factors, for example, the NP size. Furthermore, the physico-chemical properties of the nanometals, the properties of the plant species, and the conditions of the plant treatments affect the hormetic effect. All of these factors indicate that the effects of essential nanometals on plants can be highly variable among experimental systems, which makes it difficult to draw general conclusions and points out the necessity of further tests in different experimental setups. To get a holistic view, plant responses need to be examined with (multi) omics approaches. From the data available so far for nZn, nFe, nCu, and nMo, a picture is beginning to emerge that processes mainly involved in stress responses are activated at the genome, proteome, and metabolome levels, with more significant changes in the root than in the shoot. These data should be completed by the examination of nMn and nNi using (multi) omics approaches.

The beneficial effects of essential nanometals include promoting germination and vegetative and reproductive development, promoting the synthesis of phytochemicals (e.g. secondary metabolites), and enhancing abiotic and biotic stress tolerance. In most cases, laboratory or field studies do not cover the molecular details, and thus plant physiological responses need to be examined in more depth in the future.

The multifaceted effects of essential nanometals provide the basis for their application in plant cultivation as priming agents, nanofertilizers, nanopesticides, or nanoherbicides. This review has focused on the effects of nanometals in ameliorating nutrient deficiency in plants. In many cases, the effects of nanometals as nanofertilizers are tested on healthy plants, with far fewer studies dealing with amelioration of metal deficiency, which would, however, be required. The use of essential nanometals in sustainable agriculture is a desirable strategy, as efficient cultivation can be achieved with less environmental impact. Also fitting into the environmentally friendly view, the biogenic approaches utilizing microbe or plant sources for producing NPs have become increasingly popular in recent studies. At the same time, we should consider the risks (both environment- and health-related) during the application of nanomaterials. Although we are still in the early stages of understanding the molecular-level responses of plants to essential nanometals, we can state that the effects exerted by essential nanometals are more beneficial than those of the bulk forms, indicating the potential of their application in future agricultural practices.

Author contributions

ZSK: conceptualization; ZSK: visualization; ZSK, RSZ, AR, and AM: writing—original draft, writing—review & editing.

Funding

The work in the Kolbert lab is financed by the National Research, Development and Innovation Office (grant no. 135303).

References

- Adhikari T, Kundu S, Subba Rao A.** 2013. Impact of SiO₂ and Mo nano particles on seed germination of rice (*Oryza sativa* L.) International Journal of Agricultural Food Science and Technology **4**, 809–816.
- Afzal S, Sharma D, Singh NK.** 2021. Eco-friendly synthesis of phytochemical-capped iron oxide nanoparticles as nano-priming agent for boosting seed germination in rice (*Oryza sativa* L.). Environmental Science and Pollution Research **28**, 40275–40287.
- Agathokleous E, Feng ZZ, Iavicoli I, Calabrese EJ.** 2019a. The two faces of nanomaterials: A quantification of hormesis in algae and plants. Environmental International **131**, 105044.
- Agathokleous E, Kitao M, Calabrese EJ.** 2019b. Hormesis: a compelling platform for sophisticated plant science. Trends in Plant Science **4**, 318–327.
- Albersheim P.** 2011. Plant cell walls: from chemistry to biology. New York: Garland Science.
- Alscher RG, Erturk N, Heath LS.** 2002. Role of superoxide dismutases (SODs) in controlling oxidative stress in plants. Journal of Experimental Botany **53**, 1331–1341.
- Andresen E, Peiter E, Küpper H.** 2018. Trace metal metabolism in plants. Journal of Experimental Botany **69**, 909–954.
- Aubert T, Burel A, Esnault MA, Cordier S, Grasset F, Cabello-Hurtado F.** 2012. Root uptake and phytotoxicity of nanosized molybdenum octahedral clusters. Journal of Hazardous Materials **219**, 111–118.
- Bala R, Kalia A, Dhaliwal SS.** 2019. Evaluation of efficacy of ZnO nanoparticles as remedial zinc nanofertilizer for rice. Journal of Soil Science and Plant Nutrition **19**, 379–389.
- Barker AV, Pilbeam DJ.** 2015. Handbook of plant nutrition. Boca Raton: CRC Press.
- Béteky P, Rónavári A, Zakupszky D, Boka E, Igaz N, Szerencsés B, Pfeiffer I, Vágvolgyi C, Kiricsi M, Kónya Z.** 2021. Are smaller nanoparticles always better? Understanding the biological effect of size-dependent silver nanoparticle aggregation under biorelevant conditions. International Journal of Nanomedicine **16**, 3021–3040.
- Bhirde A, Xie J, Swierczewska M, Chen X.** 2011. Nanoparticles for cell labeling. Nanoscale **3**, 142–153.
- Borgatta JR, Lochbaum CA, Elmer WH, White JC, Pedersen JA, Hamers RJ.** 2021. Biomolecular corona formation on CuO nanoparticles in plant xylem fluid. Environmental Science **8**, 1067–1080.
- Brandenberger C, Cliff MJD, Gehr P, Muhlfeld C, Rothen-Rutishauser B, Stone V.** 2010. Intracellular imaging of nanoparticles: Is it an elemental mistake to believe what you see? Part Fibre Technology **7**, 15.
- Castillo-González J, Ojeda-Barrios D, Hernández-Rodríguez A, González-Franco AC, Robles-Hernández L, López-Ochoa GR.** 2018. Zinc metalloenzymes in plants. Interciencia **43**, 242–248.
- Castillo-Michel HA, Larue C, Pradas del Real AE, Cotte M, Sarret G.** 2017. Practical review on the use of synchrotron based micro- and nano-X-ray fluorescence mapping and X-ray absorption spectroscopy to investigate the interactions between plants and engineered nanomaterials. Plant Physiology and Biochemistry **110**, 13–32.
- Chavez Soria NG, Bisson MA, Atila-Gokcumen GE, Aga DS.** 2019. High-resolution mass spectrometry-based metabolomics reveal the disruption of jasmonic pathway in *Arabidopsis thaliana* upon copper oxide nanoparticle exposure. Science of the Total Environment **693**, 133443.
- Collin B, Tsyusko OV, Starnes DL, Unrine JM.** 2016. Effect of natural organic matter on dissolution and toxicity of sulfidized silver nanoparticles to *Caenorhabditis elegans*. Environmental Science **3**, 728–736.
- Conway JR, Adeleye AS, Gardea-Torresdey J, Keller AA.** 2015. Aggregation, dissolution, and transformation of copper nanoparticles in natural waters. Environmental Science and Technology **49**, 2749–2756.
- Dahuja A, Kumar RR, Sakhare A, Watts A, Singh B, Goswami S, Sachdev A, Praveen S.** 2021. Role of ATP-binding cassette transporters in maintaining plant homeostasis under abiotic and biotic stresses. Physiologia Plantarum **171**, 785–801.
- David CA, Galceran J, Rey-Castro C, Puy J, Companys E, Salvador J, Monné J, Wallace R, Vakourov A.** 2012. Dissolution kinetics and solubility of ZnO nanoparticles. Journal of Physical Chemistry C **116**, 11758–11767.
- De la Rosa G, Lopez-Moreno ML, Hernandez-Viescaz J, Montes MO, Peralta-Videa JR, Gardea-Torresdey JL.** 2011. Toxicity and bio-transformation of ZnO nanoparticles in the desert plants *Prosopis juliflora-velutina*, *Salsola tragus* and *Parkinsonia florida*. International Journal of Nanotechnology **8**, 492–506.
- de Oliveira JL.** 2021. Nano-biopesticides: present concepts and future perspectives in integrated pest management. In Jogaiah S, Singh HB, Fernandes Fraceto L, de Lima R, eds. Advances in nano-fertilizers and nano-pesticides in agriculture. Duxford: Woodhead Publishing, 1–27.
- De Tullio M, Guether M, Balestrini R.** 2013. Ascorbate oxidase is the potential conductor of a symphony of signaling pathways, Plant Signaling & Behavior **8**, 3, e23213.
- Dietz KJ, Herth S.** 2011. Plant nanotoxicology. Trends in Plant Science **16**, 582–589.
- Dimkpa CO, McLean JE, Britt DW, Anderson AJ.** 2015. Nano-CuO and interaction with nano-ZnO or soil bacterium provide evidence for the interference of nanoparticles in metal nutrition of plants. Ecotoxicology **24**, 119–129.
- Dimkpa CO, Singh U, Adisa IO, Bindraban PS, Elmer WH, Gardea-Torresdey JL, White JC.** 2018. Effects of manganese nanoparticle exposure on nutrient acquisition in wheat (*Triticum aestivum* L.). Agronomy **8**, 158.
- Djiwanti SR, Kaushik S.** 2019. Nanopesticide: future application of nanomaterials in plant protection. In: Prasad R. ed. Plant nanobionics. Vol. 2, Approaches in nanoparticles, biosynthesis, and toxicity. Cham: Springer, 255–298.
- Droppa M, Horváth G.** 1990. The role of copper in photosynthesis. Critical Review in Plant Sciences **9**, 111–123.
- Eichert T, Goldbach HE.** 2008. Equivalent pore radii of hydrophilic foliar uptake routes in stomatal and astomatous leaf surfaces – further evidence for a stomatal pathway. Physiologia Plantarum **132**, 491–502.
- El-Desouky HS, Islam KR, Bergefurd B, Gao G, Harker T, Abd-El-Dayem H, Ismail F, Mady M, Zewail RMY.** 2021. Nano iron fertilization significantly increases tomato yield by increasing plants' vegetable growth and photosynthetic efficiency. Journal of Plant Nutrition **44**, 1649–1663.
- Elemike EE, Uzoh IM, Onwujiwe DC, Babalola OO.** 2019. The role of nanotechnology in the fortification of plant nutrients and improvement of crop production. Applied Sciences **9**, 499.
- Elizabeth A, Bahadur V, Misra P, Prasad VM, Thomas T.** 2017. Effect of different concentrations of iron oxide and zinc oxide nanoparticles on growth and yield of carrot (*Daucus carota* L.). Journal of Pharmacognosy and Phytochemistry **6**, 1266–1269.
- Elsabay OM, Farroh KY, Amin HE, Atta AM.** 2021. Green synthesis of zinc oxide nanoparticles: fortification for rice grain yield and nutrients uptake enhancement. Molecules **26**, 584.
- Fabiano CC, Tezotto T, Favarin JL, Polacco JC, Mazzafera P.** 2015. Essentiality of nickel in plants: a role in plant stresses. Frontiers in Plant Science **6**, 754.
- Farooq T, Adeel M, He Z, Umar M, Shakoor N, da Silva W, Elmer W, White JC, Rui Y.** 2021. Nanotechnology and plant viruses: an emerging disease management approach for resistant pathogens. ACS Nano **15**, 6030–6037.
- Feng YZ, Cui XC, He SY, Dong G, Chen M, Wang JH, Lin X.** 2013. The role of metal nanoparticles in influencing arbuscular mycorrhizal fungi effects on plant growth. Environmental Science and Technology **47**, 9496–9504.

- Francia V, Yang K, Deville S, Reker-Smit C, Nelissen I, Salvati A.** 2019. Corona composition can affect the mechanisms cells use to internalize nanoparticles, *ACS Nano* **13**, 11107–11121.
- Gebre SH, Sendeku MG.** 2019. New frontiers in the biosynthesis of metal oxide nanoparticles and their environmental applications: an overview. *SN Applied Sciences* **1**, 928.
- González-García Y, Cárdenas-Álvarez C, Cadenas-Pliego G, Benavides-Mendoza A, Cabrera-de-la-Fuente M, Sandoval-Rangel A, Valdés-Reyna J, Juárez-Maldonado A.** 2021. Effect of three nanoparticles (Se, Si and Cu) on the bioactive compounds of bell pepper fruits under saline stress. *Plants* **10**, 217.
- Hajong M, Devi NO, Debbarma M, Majumder D.** 2019. Nanotechnology: an emerging tool for management of biotic stresses in plants. In: Prasad R, ed. *Plant nanobionics*. Vol. 2, Approaches in nanoparticles, biosynthesis, and toxicity. Cham: Springer, 299–335.
- Hall JL, Williams LE.** 2003. Transition metal transporters in plants. *Journal of Experimental Botany* **54**, 2601–2613.
- Hezaveh TA, Rahmani F, Alipour H, Pourakbar L.** 2020. Effects of foliar application of ZnO nanoparticles on secondary metabolite and microelements of *Camelina (Camelina sativa L.)* under salinity stress. *Journal of Stress Physiology & Biochemistry* **16**, 54–69.
- Hirayama T, Shinozaki K.** 2010. Research on plant abiotic stress responses in the post-genome era: past, present and future. *The Plant Journal* **61**, 1041–1052.
- Hossain Z, Mustafa G, Sakata K, Komatsu S.** 2016. Insights into the proteomic response of soybean towards Al₂O₃, ZnO, and Ag nanoparticles stress. *Journal of Hazardous Materials* **304**, 291–305.
- Huang X, Cervantes-Avilés P, Li W, Keller AA.** 2021. Drilling into the metabolomics to enhance insight on corn and wheat responses to molybdenum trioxide nanoparticles. *Environmental Science and Technology* **55**, 13452–13464.
- Iannone MF, Groppa MD, Zawoznik MS, Coral DF, van Raap MBF, Benavides MP.** 2021. Magnetite nanoparticles coated with citric acid are not phytotoxic and stimulate soybean and alfalfa growth. *Ecotoxicology and Environmental Safety* **211**, 111942.
- Jagadish K, Shiralgi Y, Chandrashekar BN, Dhananjaya BL, Srikantswamy S.** 2018. Ecofriendly synthesis of metal/metal oxide nanoparticles and their application in food packaging and food preservation. In: Grumezescu AM, Holban AM, eds. *Impact of nanoscience in the food industry*. London: Academic Press, 197–216.
- Juárez-Maldonado A, González-Morales S, Cabrera-De la F, Medrano-Macías J, Benavides-Mendoza A.** 2018. Nanometals as promoters of nutraceutical quality in crop plants. In: Grumezescu AM, Holban AM, eds. *Impact of nanoscience in the food industry*. London: Academic Press, 277–310.
- Kalwani M, Chakdar H, Srivastava A, Pabbi S, Shukla P.** 2022. Effects of nanofertilizers on soil and plant-associated microbial communities: emerging trends and perspectives. *Chemosphere* **287**, 132107.
- Kalyakina RG, Ryabinina ZN, Rjabuchina MV.** 2021. Morphobiological response of different genotypes of *Pinus sylvestris* to the influence of iron and copper nanoparticles. *Nanotechnology in Russia* **16**, 110–114.
- Kasote DM, Lee JHJ, Jayaprakasha GK, Patil BS.** 2021. Manganese oxide nanoparticles as safer seed priming agent to improve chlorophyll and antioxidant profiles in watermelon seedlings. *Nanomaterials* **11**, 1016.
- Khan Z, Upadhyaya H.** 2019. Impact of nanoparticles on abiotic stress responses in plants: an overview. In: Tripathi DK, Ahmad P, Sharma S, Chauhan DK, Dubey NK, eds. *Nanomaterials in plants, algae and microorganisms*. London: Academic Press, 305–322.
- Khot LR, Sankaran S, Maja JM, Ehsani R, Schuster EW.** 2012. Applications of nanomaterials in agricultural production and crop protection: a review. *Crop Protection* **35**, 64–70.
- Kilian B, Dempewolf H, Guarino L, Werner P, Coyne C, Warburton ML.** 2021. *Crop Science* special issue: Adapting agriculture to climate change: a walk on the wild side. *Crop Science* **61**, 32–36.
- Kirkby E.** 2012. Introduction, definition and classification of nutrients. In: Marschner S, ed. *Marschner's mineral nutrition of higher plants*, 3rd edn. London: Elsevier, 3–5.
- Kohatsu MY, Pelegrino MT, Monteiro LR, et al.** 2021. Comparison of foliar spray and soil irrigation of biogenic CuO nanoparticles (NPs) on elemental uptake and accumulation in lettuce. *Environmental Science and Pollution Research* **28**, 16350–16367.
- Kouhi SM, Lahouti M, Ganjeali A, Entezari MH.** 2014. Comparative phytotoxicity of ZnO nanoparticles, ZnO microparticles, and Zn²⁺ on rapeseed (*Brassica napus L.*): Investigating a wide range of concentrations. *Toxicology and Environmental Chemistry* **96**, 861–868.
- Kumari A, Sharma B, Singh BN, Hidangmayum A, Jatav HS, Chandra K, Singhal RK, Sathyanarayana E, Patra A, Mohapatra KK.** 2021. Physiological mechanisms and adaptation strategies of plants under nutrient deficiency and toxicity conditions. In: Aftab T, Roychoudhury A, ed. *Plant perspectives to global climate changes*. London: Elsevier, 173–194.
- Kuppusamy P, Yusoff MM, Govindan N.** 2016. Biosynthesis of metallic nanoparticles using plant derivatives and their new avenues in pharmacological applications—an updated report. *Saudi Pharmaceutical Journal* **24**, 473–484.
- Kurczyńska E, Godel-Jędrychowska K, Sala K, Milewska-Hendel A.** 2021. Nanoparticles—plant interaction: what we know, where we are? *Applied Sciences* **11**, 5473.
- Landa P, Dytrych P, Prerostova S, Petrova S, Vankova R, Vanek T.** 2017. Transcriptomic response of *Arabidopsis thaliana* exposed to CuO nanoparticles, bulk material, and ionic copper. *Environmental Science and Technology* **51**, 10814–10824.
- Landa P, Vankova R, Andrlouva J, Hodek J, Marsik P, Storchova H, White JC, Vanek T.** 2012. Nanoparticle-specific changes in *Arabidopsis thaliana* gene expression after exposure to ZnO, TiO₂, and fullerene soot. *Journal of Hazardous Materials* **241–242**, 55–62.
- Lee S, Kim S, Kim S, Lee I.** 2013. Assessment of phytotoxicity of ZnO NPs on a medicinal plant, *Fagopyrum esculentum*. *Environmental Science and Pollution Research* **20**, 848–854.
- Li C, Li Y, Ding C.** 2019. The role of copper homeostasis at the host-pathogen axis: from bacteria to fungi. *International Journal of Molecular Sciences* **20**, 175.
- Li Z, Yu T, Paul R, Fan J, Yang Y, Wei Q.** 2020. Agricultural nanodiagnosics for plant diseases: recent advances and challenges. *Nanoscale Advances* **2**, 3083–3094.
- Lindermayr C.** 2018. Crosstalk between reactive oxygen species and nitric oxide in plants: key role of S-nitrosoglutathione reductase. *Free Radical Biology and Medicine* **122**, 110–115.
- Lira-Morales JD, Varela-Bojórquez N, Montoya-Rojo MB, Sañudo-Barajas JA.** 2019. The role of ZIP proteins in zinc assimilation and distribution in plants: current challenges. *Czech Journal of Genetics and Plant Breeding* **55**, 45–54.
- Liu J, Legros S, Ma G, Veinot JGC, von der Kammer F, Hofmann T.** 2012. Influence of surface functionalization and particle size on the aggregation kinetics of engineered nanoparticles. *Chemosphere* **87**, 918–924.
- Lv J, Christie P, Zhang S.** 2019. Uptake, translocation, and transformation of metal-based nanoparticles in plants: recent advances and methodological challenges. *Environmental Science: Nano* **6**, 41–59.
- Lv JT, Zhang SZ, Luo L, Zhang J, Yang K, Christie P.** 2015. Accumulation, speciation and uptake pathway of ZnO nanoparticles in maize. *Environmental Science* **2**, 68–77.
- Maurer-Jones M, Gunsolus IL, Murphy CJ, Haynes, CL.** 2013. Toxicity of engineered nanoparticles in the environment. *Analytical Chemistry* **85**, 3036–3049.
- McNear DH.** 2013. The rhizosphere – roots, soil and everything in between. *Nature Education Knowledge* **4**, 1. <http://www.nature.com/scitable/knowledge/library/the-rhizosphere-roots-soil-and-67500617>
- Mendel RR, Hänsch R.** 2002. Molybdoenzymes and molybdenum cofactor in plants. *Journal of Experimental Botany* **53**, 1689–1698.

- Mittal D, Kaur G, Singh P, Yadav K and Ali SA.** 2020. Nanoparticle-based sustainable agriculture and food science: recent advances and future outlook. *Frontiers in Nanotechnology* **2**, 579954.
- Mohammadi K, Sohrabi Y.** 2012. Bacterial biofertilizers for sustainable crop production: a review. *APRN Journal of Agricultural and Biological Science* **7**, 237–308.
- Molnár Á, Papp M, Kovács DZ, et al.** 2020. Nitro-oxidative signaling induced by chemically synthesized zinc oxide nanoparticles (ZnO NPs) in *Brassica* species. *Chemosphere* **251**, 126419.
- Montes A, Bisson MA, Gardella JA, Aga DS.** 2017. Uptake and transformations of engineered nanomaterials: critical responses observed in terrestrial plants and the model plant *Arabidopsis thaliana*. *Science of the Total Environment* **607–608**, 1497–1516.
- Ngo QB, Dao TH, Nguyen HC, Tran XT, Nguyen TV, Khuu TD, Huynh TH.** 2014. Effects of nanocrystalline powders (Fe, Co and Cu) on the germination, growth, crop yield and product quality of soybean (Vietnamese species DT-51). *Advances in Natural Sciences: Nanoscience and Nanotechnology* **5**, 015016.
- Olczak M, Morawiecka B, Watorek W.** 2003. Plant purple acid phosphatases – genes, structures and biological function. *Acta Biochimica Polonica* **50**, 1245–1256.
- Onelli E, Prescianotto-Baschong C, Caccianiga M, Moscatelli A.** 2008. Clathrin-dependent and independent endocytic pathways in tobacco protoplasts revealed by labelling with charged nanogold. *Journal of Experimental Botany* **59**, 3051–3068.
- Palocci C, Valletta A, Chronopoulou L, Donati L, Bramosanti M, Brasili E, Baldan B, Pasqua G.** 2017. Endocytic pathways involved in PLGA nanoparticle uptake by grapevine cells and role of cell wall and membrane in size selection. *Plant Cell Reports* **36**, 1917–1928.
- Pelegrino MT, Kohatsu MY, Seabra AB, et al.** 2020. Effects of copper oxide nanoparticles on growth of lettuce (*Lactuca sativa* L.) seedlings and possible implications of nitric oxide in their antioxidative defense. *Environmental Monitoring and Assessment* **192**, 232.
- Peng C, Tong H, Shen C, Sun L, Yuan P, He M, Shi J.** 2020. Bioavailability and translocation of metal oxide nanoparticles in the soil-rice plant system. *Science of the Total Environment* **713**, 136662.
- Pestovsky YS, Martinez-Antonio A.** 2017. The use of nanoparticles and nanoformulations in agriculture. *Journal of Nanoscience and Nanotechnology* **17**, 8699–8730.
- Pittman JK, Hirschi KD.** 2016. CAX-ing a wide net: cation/H⁺ transporters in metal remediation and abiotic stress signalling. *Plant Biology* **18**, 741–749.
- Riaz U, Mehdi SM, Iqbal S, Khalid HI, Qadir AA, Anum W, Ahmad M, Murtaza G.** 2020. Bio-fertilizers: eco-friendly approach for plant and soil environment. In: Hakeem K., Bhat R., Qadri H. eds. *Bioremediation and biotechnology*. Cham: Springer, 189–213.
- Rivero-Montejo SDJ, Vargas-Hernandez M, Torres-Pacheco I.** 2021. Nanoparticles as novel elicitors to improve bioactive compounds in plants. *Agriculture* **11**, 134.
- Rónavári A, Kovács D, Igaz N, Vágvölgyi C, Boros IM, Kónya Z, Pfeiffer I, Kiricsi M.** 2017. Biological activity of green-synthesized silver nanoparticles depends on the applied natural extracts: a comprehensive study. *International Journal of Nanomedicine* **12**, 871–883.
- Rui M, Ma C, Hao Y, et al.** 2016. Iron oxide nanoparticles as a potential iron fertilizer for peanut (*Arachis hypogaea*). *Frontiers in Plant Science* **7**, 815.
- Schmidt SB, Husted S.** 2019. The biochemical properties of manganese in plants. *Plants* **8**, 381.
- Schott-Verdugo S, Müller L, Classen E, Gohlke H, Groth G.** 2019. Structural model of the ETR1 ethylene receptor transmembrane sensor domain. *Scientific Reports* **9**, 8869.
- Schwab F, Zhai G, Kern M, Turner A, Schnoor JL, Wiesner MR.** 2016. Barriers, pathways and processes for uptake, translocation and accumulation of nanomaterials in plants – critical review. *Nanotoxicology* **10**, 257–278.
- Sega D, Baldan B, Zamboni A, Varanini Z.** 2020. FePO₄ NPs are an efficient nutritional source for plants: combination of nano-material properties and metabolic responses to nutritional deficiencies. *Frontiers in Plant Science* **11**, 586470.
- Sharma PK, Raghubanshi AS, Shah K.** 2021. Examining the uptake and bioaccumulation of molybdenum nanoparticles and their effect on antioxidant activities in growing rice seedlings. *Environmental Science and Pollution Research* **28**, 13439–13453.
- Shukla PK, Misra P, Kole C.** 2016. Uptake, translocation, accumulation, transformation and generational transmission of nanoparticles in plants. In: Kole C, Kumar DS, Khodakovskaya MV, eds. *Plant nanotechnology*. Cham: Springer International Publishing, 183–218.
- Singh A, Tiwari S, Pandey J, Lata C, Singh IK.** 2021. Role of nanoparticles in crop improvement and abiotic stress management. *Journal of Biotechnology* **337**, 57–70.
- Singh V, Verma K.** 2018. Metals from cell to environment: connecting metallomics with other omics. *Open Journal of Plant Science* **3**, 001–014.
- Solanki P, Bhargava A, Chhipa H, Jain N, Panwar J.** 2015. Nano-fertilizers and their smart delivery system. In: Rai M., Ribeiro C., Mattoso L., Duran N, eds. *Nanotechnologies in food and agriculture*. Cham: Springer, 81–101.
- Song K, He X.** 2021. How to improve seed germination with green nanoprimer. *Seed Science and Technology* **49**, 81–92.
- Subramanian KS, Manikandan A, Thirunavukkarasu M, Rahale CS.** 2015. Nano-fertilizers for balanced crop nutrition. In: Rai M., Ribeiro C., Mattoso L., Duran N, eds. *Nanotechnologies in food and agriculture*. Cham: Springer, 69–80.
- Sun L, Wang Y, Wang R, Wang R, Zhang P, Ju Q, Xu J.** 2020. Physiological, transcriptomic, and metabolomics analyses reveal zinc oxide nanoparticles modulate plant growth in tomato. *Environmental Science: Nano* **7**, 3587.
- Szöllösi R, Molnár Á, Kondak S, Kolbert Z.** 2020. Dual effect of nanomaterials on germination and seedling growth: stimulation vs. phytotoxicity. *Plants* **9**, 1745.
- Takatsujii H.** 1998. Zinc-finger transcription factors in plants. *Cellular and Molecule Life Science* **54**, 582–596.
- Tani T, Kan R, Yamano Y, Uchida T.** 2018. Stabilization of Ag nanostructures by tuning their Fermi levels. *Japanese Journal of Applied Physics* **57**, 055001.
- Thounaojam TC, Thounaojam TM, Upadhyaya H.** 2021. Role of zinc oxide nanoparticles in mediating abiotic stress responses in plant. In: Abd-Elisalam KA, ed. *Zinc-based nanostructures for environmental and agricultural applications*. Amsterdam: Elsevier, 323–337.
- Tombuloglu H, Slimani Y, Tombuloglu G, Demir Korkmaz A, Baykal A, Almeseiere M, Ercan I.** 2019. Impact of superparamagnetic iron oxide nanoparticles (SPIONs) and ionic iron on physiology of summer squash (*Cucurbita pepo*): a comparative study. *Plant Physiology and Biochemistry* **139**, 56–65.
- Vigani G, Murgia I.** 2018. Iron-requiring enzymes in the spotlight of oxygen. *Trends in Plant Science* **23**, 874–882.
- Wan J, Wang R, Wang R, Ju Q, Wang Y, Xu J.** 2019. Comparative physiological and transcriptomic analyses reveal the toxic effects of ZnO nanoparticles on plant growth. *Environmental Science and Technology* **53**, 4235–4244.
- Wang Y, Deng C, Cota-Ruiz K, et al.** 2020. Improvement of nutrient elements and allicin content in green onion (*Allium fistulosum*) plants exposed to CuO nanoparticles. *Science of the Total Environment* **725**, 138387.
- Wang WN, Tarafdar JC, Biswas P.** 2013. Nanoparticle synthesis and delivery by an aerosol route for watermelon plant foliar uptake. *Journal of Nanoparticle Research* **15**, 1417.
- Williams LE, Mills RF.** 2005. P_{1B}-ATPases – an ancient family of transition metal pumps with diverse functions in plants. *Trends in Plant Science* **10**, 491–502.

- Wojcieszek J, Jiménez-Lamana J, Ruzik L, Szpunar J, Jarosz M.** 2020. To-do and not-to-do in model studies of the uptake, fate and metabolism of metal-containing nanoparticles in plants. *Nanomaterials* **10**, 1480.
- Xiong T, Zhang S, Kang Z, Zhang T, Li S.** 2021. Dose-dependent physiological and transcriptomic responses of lettuce (*Lactuca sativa* L.) to copper oxide nanoparticles—insights into the phytotoxicity mechanisms. *International Journal of Molecular Sciences* **22**, 3688.
- Yasmeen F, Iqbal Raja N, Abdul Razzaq A, Komatsu S.** 2017. Proteomic and physiological analyses of wheat seeds exposed to copper and iron nanoparticles. *Biochimica et Biophysica Acta – Proteins and Proteomics* **1865**, 28–42.
- Yasmeen F, Raja NI, Ilyas N, Komatsu S.** 2018. Quantitative proteomic analysis of shoot in stress tolerant wheat varieties on copper nanoparticle exposure. *Plant Molecular Biology Reporter* **36**, 326–340.
- Yasmeen F, Raja NI, Razzaq A, Komatsu S.** 2016. Gel-free/label-free proteomic analysis of wheat shoot in stress tolerant varieties under iron nanoparticles exposure. *Biochimica et Biophysica Acta – Proteins and Proteomics* **1864**, 1586–1598.
- Zhang P, Guo Z, Zhang Z, Fu H, White JC, Lynch I.** 2020. Nanomaterial transformation in the soil–plant system: implications for food safety and application in agriculture. *Small* **16**, 2000705.
- Zhao L, Lu L, Wang A, Zhang H, Huang M, Wu H, Xing B, Wang Z, Ji R.** 2020. Nano-biotechnology in agriculture: use of nanomaterials to promote plant growth and stress tolerance. *Journal of Agricultural and Food Chemistry* **68**, 1935–1947.
- Zhu H, Han J, Xiao JQ, Jin Y.** 2008. Uptake, translocation, and accumulation of manufactured iron oxide nanoparticles by pumpkin plants. *Journal of Environmental Monitoring* **10**, 713–717.
- Zhu ZJ, Wang HH, Yan B, Zheng H, Jiang Y, Miranda OR, Rotello VM, Xing B, Vachet RW.** 2012. Effect of surface charge on the uptake and distribution of gold nanoparticles in four plant species. *Environmental Science and Technology* **46**, 12391–12398.