

Health of African Refugee Children Outside Africa: A Scoping Review

**Bukola Salami*, Shirley Mogale, Folakemi Ojo, Michael Kariwo, Jill Thompson,
Philomina Okeke-Ihejirika, Sophie Yohani**

***Corresponding Author: Bukola Salami, Bukola.salami@ualberta.ca**

Bukola Salami, RN, BScN, MN, PhD

Associate Professor

Faculty of Nursing

University of Alberta

11405 87 Avenue

Edmonton, Alberta, T6G 1C9

Phone: 7802481801

Fax: (780) 492-2551

Email: Bukola.salami@ualberta.ca

Dr. Ramadimetja Shirley Mogale, RN, PhD

Professor and Head of Department Nursing Science

University of Pretoria

8-38 HW Snyman Building North

Phone: +27123563179

Fax +27123561366

Email: shirley.mogale@up.ac.za

Folakemi Ojo, BSc, MPH

Research Assistant

Faculty of Nursing

University of Alberta

11405 87 Avenue

Edmonton, Alberta. T6G 1C9

Email: foluwakuyide@yahoo.com

Phone: 780 248 1801

Fax: 780 492 2551

Michael Kariwo, PhD

Research Coordinator

Faculty of Nursing

University of Alberta

11405 87 Avenue

Edmonton, Alberta

T6G 1C9

Email: mkariwo@ualberta.ca

Phone: 780 248 1801

Fax: 780 492 2551

Jill Thompson, PhD

Senior Lecturer

Health Sciences School

The University of Sheffield

Barber House Annexe

3a Clarkehouse Road

Sheffield

S10 2LA

Jill.thompson@sheffield.ac.uk

Philomina Okeke-Ihejirika, PhD

Full Professor

Faculty of Arts

University of Alberta

1-13 Assiniboia Hall

Edmonton, Alberta

Phone: 780318 4785

Email: pokeke@ualberta.ca

Sophie Yohani, PhD., R Psych.

Associate Professor

Department of Educational Psychology

University of Alberta

Edmonton, Alberta

Canada, T6G 2G5

Tel: (780) 492-1164

Fax: (780) 492-1318

Email: sophie.yohani@ualberta.ca

Declarations of Conflict of Interest

None

Funding Acknowledgement

This work was supported by the Worldwide Universities Network Research Development Fund.

The funder has no role in the conduct of the review and/or preparation of the article.

Author Contributions

Bukola Salami: Conceptualization, Methodology, Validation, Funding Acquisition, Writing Original Draft, Review and Editing, Supervision, Project Administration

Shirley Mogale: Writing Original Draft, Review and Editing

Folakemi Ojo: Formal Analysis, Writing Original Draft

Michael Kariwo: Review and Editing, Supervision, Project Administration

Jill Thompson: Funding Acquisition, Review and Editing, Conceptualization

Philomina Okeke-Ihejirika: Funding Acquisition, Review and Editing, Conceptualization

Sophie Yohani: Funding Acquisition, Review and Editing, Conceptualization

Highlights

- Areas of published research include infectious disease, mental health and nutrition
- African child refugees living outside Africa experience poor mental and physical health
- Experiences during the pre-migration and postmigration stage influence their health
- Income and gender influence the mental health of African refugee children living outside Africa

Abstract

Problem: Africa is the largest source continent of refugee children. However, we found no published synthesis of the literature on the health of African refugee children outside Africa. Conducting a review of the literature on this particular population will help illuminate the particular contextual health issues faced by African child refugees who live outside Africa. The

purpose of this review is to synthesize what is known from the existing literature regarding the health of sub-Saharan African refugee children who live outside Africa.

Methods: We completed a scoping review of the published literature. We included articles published in English with a focus on the health of sub-Saharan African refugee children living outside Africa. We excluded studies of refugees in Africa as the living conditions of these refugees, most of whom reside in camps, are very different from those outside Africa. Using relevant keywords, we searched 10 databases to identify and screen 6,602 articles after duplicates were eliminated.

Sample: A total of 20 studies were included in this review.

Findings: Published research articles on sub-Saharan African child refugees living outside Africa focus on infectious diseases, mental health and neurodevelopmental disorders, food insecurity and psychosocial adjustment, physical health (including obesity), and health promotion strategies. This population is characterized by a high rate of infectious diseases (e.g., malaria), obesity, and mental health problems, especially post-traumatic stress disorder (PTSD).

Conclusions and Implications: To attend to the health needs of sub-Saharan African refugee children who live outside Africa, interventions should address pre-migration factors as well as post-migration factors (including income and community belonging) while employing a strengths-based perspective.

Keywords: Africa; African; Child; Migrant; Refugee

Introduction

Africa is a leading source continent for child refugees (International Organization for Migration, 2020). According to the UNHCR (2020), “refugees are people who have fled war, violence, conflict or persecution and have crossed an international border to find safety in another country” (Paragraph 1). African child refugees are a growing population in destination countries. Between 2006 and 2016, the number of African refugee children more than doubled (UNICEF, 2016). Of the ~26 million refugees in 2016, 52% were under the age of 18 years (World Migration Report, 2020). While the majority of child migrants, including refugees, remain in Africa (30%), some migrate outside of Africa. Addressing the health of African child refugees will contribute to meeting global health targets, including the United Nations’ Sustainable Development Goals (United Nations Development Program, 2020).

Migrant and refugee well-being is envisioned in several of the Sustainable Development Goals, including Target 10.2: “By 2030, empower and promote the social, economic and political inclusion of all, irrespective of age, sex, disability, race, ethnicity, origin, religion or economic or other status” (United Nations Development Program, 2020). However, migrants and refugees continue to experience several social and health challenges in destination countries, including exploitation, abuse, trafficking, and poor socio-economic integration (International Organization for Migration, 2020). Refugee children face physical and mental health challenges, rendering them vulnerable to developmental challenges that further impact their future health outcomes (Borrego et al., 2019). Despite the existing, albeit sparse, evidence about the large number of African child refugees (International Organization for Migration, 2020) and the general challenges they face, we did not find any synthesis of evidence on this topic, especially for those who live outside Africa. Conducting a review of the literature on this unique population will help illuminate the particular contextual health issues faced by African child refugees who

live outside Africa. We specifically focus on sub-Saharan African refugee children living outside Africa, as the experience of these migrants and historical context is different from those from northern parts of Africa. We define sub-Saharan Africa as countries that are predominantly inhabited by Black populations in African and largely does not include countries in North Africa. Our definition includes 49 of African's 54 countries excluding Algeria, Egypt, Libya, Morocco, and Tunisia. Our definition of sub-Saharan Africa includes Somalia, Sudan and Djibouti which some authors have excluded from such definitions in the past.

Methodology

We used a scoping review methodology to map or summarize the range of evidence in the field to convey breadth and depth. Specifically, we utilized Arksey and O'Malley's (2005) approach, which follows five key stages: 1) identify the research question; 2) identify relevant studies; 3) study selection; 4) charting the data; and 5) collating, summarizing, and reporting the results. Scoping reviews differ from systematic reviews in that the goal is not to assess the quality of studies, although this may be done by some reviewers. Rather, scoping reviews "aim to map rapidly the key concepts underpinning a research area and the main sources and types of evidence available" (Arksey & O'Malley, 2005, p. 194), which is the approach taken here. To map out evidence in the field, scoping reviews deal with a greater range of study designs and methodologies.

Identify Research Question

Our research question was: What is known from the existing literature on the health of sub-Saharan African refugee children who live outside Africa?

Identify Relevant Studies

In line with Arksey and O'Malley (2005), we completed a search of multiple databases. We included articles published in English with a focus on sub-Saharan African first-generation refugee children and health. By sub-Saharan African refugee children, we mean children who were born in sub-Saharan Africa and later migrated to other countries outside Africa (a definition is provided in the introduction). Thus, we excluded children who were not born in Africa but were children of African migrants. We also excluded children from north African countries such as Libya, Egypt, Morocco, Tunisia, and Algeria as the culture and histories of these countries are more similar to those of the Middle East than countries south of the Sahara. We excluded studies of refugees in Africa as the living conditions of these refugees, most of whom reside in camps, are very different from those who live outside Africa. We also excluded literature reviews, systematic reviews, epidemiological studies (where sub-Saharan African refugees were only a variable and not the sole focus), and studies in which results from African refugee children were mixed with other children. While we excluded reviews from our inclusion, we searched their reference lists and included articles cited therein in our work. Epidemiological studies that provided robust information were included; however, epidemiological studies where sub-Saharan African refugees outside Africa were only a variable and where no full results or discussion were provided were excluded. We limited studies to those published from 2000 to 2019.

Based on the inclusion and exclusion criteria, a librarian assisted with developing a search of 10 electronic databases: CINAHL, MEDLINE, Embase, Global Health, Psych Info, Cochrane Library, SocIndex, Sociological Abstract, ProQuest Dissertations & Theses Global, and Child Development and Adolescent Studies. Search terms included a combination of words related to Africa (including names of sub-Saharan African countries), child, and migrant or refugee. The search resulted in total of 12,720 articles, with 6,602 remaining after duplicates

were excluded. Articles retrieved were exported to www.covidence.org for further selection by research assistants. We also reviewed reference lists of included articles. (Please see PRISMA diagram for number of articles at each stage of the selection process)

Study Selection

Six graduate students supported the selection of articles due to the large number returned from the initial search. The abstract and title of each article were reviewed by two of the graduate students, all of whom received training prior to the search and met intermittently with the lead author throughout selection to ensure consistency in the application of inclusion criteria. Two research assistants then read the full articles to determine relevance to inclusion criteria. Figure 1 is a Prisma flow diagram for the scoping review process.

Charting Data

A research assistant supported the extraction of data (see Appendix Table 1 for some of the information extracted from the articles).

Data Summary and Analysis

We completed a numerical analysis of articles that met our inclusion criteria. We identified the country of origin, research methods, sample size, and health focus. We also organized our results thematically according to the health condition of interest. We also considered consistency and contradictions across studies as well as research gaps. This scoping review did not include a quality appraisal of the studies.



PRISMA Flow Diagram – Health of African Refugee Children Outside Africa: A Scoping Review




Fig. 1. PRISMA flow diagram – health of African refugee children outside Africa: a scoping review.

Table 1 Studies on African Refugees Outside Africa Included in the Scoping Review

Author(s)	Research Question/ Purpose	Methodology and Method	Summary of Findings
1. Alasagheirin, M. H., & Clark, M. K. (2018)	To study the physical growth, bone growth, body composition, metabolic risks, physical activity, and food security of Sudanese children living in the United States.	Quantitative, Cross-sectional study.	About 1/3 of children had low bone mineral (BMC and BMD). Higher prevalence rate of obesity (32%). There was an association between gender and increase in body mass index (BMI) among African refugee children studied.
2. Caron, R. M., Tshabangu-Soko, T., & Finefrock, K. J. (2013).	To describe the agency' role in the settlement process; to examine communication methods used in public health system; and to develop effective health communication strategies that assist key refugee resettlement agencies, which comprise part of the public health system, in building the community's capacity to manage childhood lead poisoning.	Qualitative: structured interviews	Childhood lead poisoning was associated with a lack of a government regulations and education of African refugee populations; ineffective communication methods by settlement agencies.
3. Ellis, B. H., MacDonald, H. Z., Lincoln, H. Z., & Cabral, H. J. (2008)	To examine relations between trauma exposure, post-resettlement stressors, perceived discrimination, and mental health symptoms in Somali adolescent refugees resettled in the U.S.	Community-based participatory with quantitative design	PTSD symptoms showed positive correlations with depressive symptoms, trauma exposure, post-war hardships, acculturative hassles, and perceived discrimination.
4. Ellis, H. B., Lincoln, A. K., Charney, M. E., Ford-Paz, R., Benson, M., & Strunin, L. (2010).	To examine the utility of the Gateway Provider Model in understanding service utilization and pathways to help for Somali refugee adolescents resettled in the Northeastern United States.	Community-based participatory with quantitative design	Only 10% of parents report seeking help for mental health issues; 2% seek help through prayer. However, 3% screened clinically positive for depression while 20% screened positive for PTSD.
5. Geltman, P. L., Grant-Knight, W., Mehta, S. D., Lloyd-Travaglini, C., Lustig, S., Landgraf, J. M., Wise, P. H. (2005).	To assess the functional and behavioral health of unaccompanied Sudanese refugee minors approximately 1 year after resettlement in the United States.	Quantitative	20% of the minors had a diagnosis of posttraumatic stress disorder and were more likely to have worse scores on all the Child Health Questionnaire subscales. Social isolation and history of personal injury were associated with PTSD.

6. Geltman, P. L., Grant-Knight, W., Ellis, H., & Landgraf, J. M. (2007).	To assess whether mental health counseling and other health services were associated with functional health outcomes of unaccompanied Sudanese refugee minors in the U.S.	Descriptive Survey in a quantitative study.	20% of the sample surveyed was diagnosed for PTSD. The use of a variety of health services was high. Ninety-six percent reported having a routine source of care. The Child Health Questionnaire (CHQ) revealed a high prevalence of seeking medical care for somatic complaints and health problems often associated with behavioral or emotional disorders.
7. Griffith, M., Mellor, D., Green, J., & Renzaho, A. M. (2014)	To examine migration- and socioeconomic-related influences on obesity among African migrant adolescents in Melbourne, Australia.	Quantitative: used standardized measures including the acculturation scale	Gender and parental BMI were linked with increase in adolescent BMI. Boys had lower BMI than girls.
8. Hadley, C., Zodhiates, A., & Sellen, D. W. (2006).	To assess the occurrence and severity of food insecurity, and examine associations between food insecurity and measures of socio-economic status and indicators of acculturation.	Ethnography study in a Qualitative research.	Food insecurity was indicated in approximately half of households (53%). Food insecurity was associated with measures of socio-economic status such as income, employment status and participation in the food stamp program. Measures of acculturation such as difficulty in the shopping environment and language difficulty were also associated with the occurrence and severity of food insecurity, as well as time lived in the USA.
9. Herve, K., Vargas, D., Klesges, L., Fischer, P. R., Trippel, S., & Juhn, Y. J. (2009).	To determine the role of age at the time of arrival in the subsequent risk of becoming overweight; and to assess the impact of age and weight at arrival on the rate of subsequent weight changes.	Quantitative: a retrospective cohort study	Children who were overweight at arrival in the U.S. remained at an increased risk of overweight. Older children showed a tendency to gain weight more rapidly and to be at risk of overweight, in comparison with the younger children.
10. Huemer, J., Karnik, N., Voelkl-Kernstock, S., Granditsch, E., Plattner, B., Friedrich, M., & Steiner, H. (2011).	Hypotheses a) the prevalence of PTSD and related internalizing psychopathology would be higher when compared to age appropriate norms and results among accompanied refugee minors; b) positive quality of life in the assessed youths would be lower than in age matched norms, given their dire psychosocial living circumstances; and c) war exposure would predict higher rates of PTSD symptoms after controlling for age and gender.	Quantitative used 5 standardized measures: Youth self report; UCLA PTSD; Mini -International Neuropsychiatric Interview; Facts about You; and Scales for Children Afflicted by war	Girls are more susceptible to PTSD than boys when controlling for age. There was an association between the incidence of PTSD and exposure to war or war-related event incidence rate of posttraumatic stress disorders among African refugees.
11. Huemer, J., Volkl-Kernstock, S., Karnik,	To explore whether high levels of protective intra-individual processes	Quantitative: theoretical model of resilience	Compared to the unaccompanied African refugees, study shows high level of repressive defensiveness.

N., Denny, K. G., Granditsch, E., Mitterer, M., Humphreys, K., Plattner, B., Friedrich, M., Shaw, R. J., & Steiner, H. (2013)	such as repression are present in unaccompanied refugee minors (URMs) when compared to a sample of age-matched norms; and to explore the role of other personality factors.		High levels of resilience. Reduction in self - reported satisfaction with life.
12. Lincoln, A. K., Lazarevic, V., White, M. T., & Ellis, B. H. (2016)	1. Are acculturative hassles associated with worse PTSD symptoms and depression symptoms? Is there a moderating effect for gender on this relationship? 2. Does the degree to which Somali refugee adolescents experience acculturative hassles differ by acculturation? 3. Does the association between acculturative hassles and mental health (PTSD and depression symptoms) differ by acculturation style?	Community-based participatory research in mixed method study.	The severity of acculturative hassles was associated with higher levels of both PTSD symptoms and depression symptoms. There was no moderating effect of gender in the data obtained. The severity of acculturative hassles varies with acculturative style. The effect of hassles is more detrimental to marginalized participants than other acculturative groups.
13. Maroushek, S. R., Aguilar, E. F., Stauffer, W., & Abd-Alla, M. D. (2005).	To determine the prevalence of malaria among asymptomatic children from West Africa.	Retrospective Chart review, Quantitative study	The prevalence rate of malaria was 60% among those who were asymptomatic.
14. Mellor, D., Renzaho, A., Swinburn, B., Green, J., & Richardson, B. (2012).	To investigate the relationship between parenting style and family functioning, and BMI among adolescent migrants and refugees from African countries.	Quantitative	There was an association between gender and increase in BMI among African refugee children studied. Girls' BMI was on average slightly higher than that of boys. Lack of parenting supervision accounts for significant changes in BMI.
15. Miller-Gairy, S., & Mofya, S. (2015).	To examine the elements of culture and tradition that affect perceptions and expectations of Somali refugee mothers regarding autism spectrum disorder (ASD).	Case study using ethnography in qualitative study design	Etiology and pathology of ASD was not well understood because of cultural and traditional beliefs. Communication also affected Somali refugee mother's perception of ASD. Lack of access to healthcare resource by children because of misunderstanding between health care providers and mothers.
16. Ndao, M., Bandyayera, E., Kokoskin, E., Diemert, D., Gyorkos, T. W., MacLean, J. D., St. John, R., & Ward. B. J. (2005).	To investigate the performance of the health care system in the diagnosis and management of imported malaria.	Quantitative and clinical	Small outbreaks of malaria were missed regularly. Therefore, there is a need for optimal malaria diagnosis and treatment for refugee children from the malaria-endemic region.

17. Völkl-Kernstock, S., Karnik, N., Mitterer-Asadi, M., Granditsch, E., Steiner, N., Friedrich, M. H., & Huemer, J. (2014).	To investigate African unaccompanied refugee minors living in Austria for PTSD prevalence and related symptoms, comorbidity, demographics, and coping strategies.	Quantitative: used standardized measures	Girls were more likely to develop PTSD, which was significantly correlated with single war-related traumatic events.
18. Weine, S. M., Ware, N., Tugenberg, T., Hakizimana, L., Dahnweih, G., Currie, M., Wagner, M., & Levin, E. (2013).	To characterize the patterns of psychosocial adjustment among adolescent African refugees in U.S. resettlement.	Longitudinal study employing ethnography in a mixed methods design	Identified means of promoting psychosocial adjustment as finances, social support network, family engagement, adherence to cultural values, literacy, and participation in religious activities.
19. Weine, S. M., Ware, N., Hakizimana, L., Tugenberg, T., Currie, M., Dahnweih, G., Wagner, M., & Wulu, J. (2014).	1. What protective agents and protective resources appear to contribute to psychosocial well-being in adolescent refugees? 2. What are the protective mechanisms through which these protective resources appear to contribute to psychosocial well-being in adolescent refugees?	Longitudinal study employing ethnography in a qualitative methods design	Eight family and community capacities were identified that appeared to promote psychosocial well-being in the adolescent refugees. These included: 1) finances for necessities; 2) English proficiency; 3) social support networks; 4) engaged parenting; 5) family cohesion; 6) cultural adherence and guidance; 7) educational support; and 8) faith and religious involvement. Protective mechanisms were 1) relational (supporting, connecting, belonging); 2) informational (informing, preparing); and 3) developmental (defending, promoting, adapting).
20. Wilson, A., & Renzaho, A. (2013).	To investigate the differences in acculturation experiences between parent and adolescent refugees from the Horn of Africa in Melbourne, Australia; and to explore food beliefs and perceived health risks from an intergenerational perspective.	Exploratory design in a Qualitative study	There were differences between parents and adolescents in relation to lifestyle, diet and physical activity, and views on health consequences of their changed diets. Three distinct themes emerged: (i) acculturation; (ii) food beliefs and perceived health consequences; and (iii) parental feeding practices.

Results

After screening, 20 articles were ultimately included in the review, of which 13 used quantitative methods, five used qualitative methods, and two used mixed methods. The authors of these articles predominantly obtained data from children, followed by their parents. The studies included were conducted in the United States (n=13), Australia (n=3), Austria (n=3), and Canada (n=1). Participant countries of origin included Nigeria, Gambia, Ghana, Niger, Guinea-Bissau, Mali Eritrea, Kenya, Ethiopia, Burundi, Democratic Republic of Congo, Rwanda, Tanzania, Sudan, Somalia, Liberia, and Mali. Sample size ranged from 19 to 101 in the qualitative studies, from 41 to 304 in the quantitative studies, and from 73 to 114 in the mixed-method studies. Sampling methods included convenience sampling (n=4), purposeful sampling (n=4), and snowball sampling (n=1); eleven articles did not specify. The age of participants ranged from 6 months to 18 years. Data sources included children (n=15), parents, caretakers, and adolescents (n=5), and parents only (1). The studies reported on five broad themes: infectious diseases (n=2), mental health and neurodevelopmental disorders (n=9), food insecurity and psychosocial adjustment (n=3), physical health (n=4), and health promotion strategies (n=2).

Infectious Diseases

Two studies (both quantitative) investigated infectious diseases (specifically malaria) among sub-Saharan African refugee children outside Africa. Ndao et al. (2005) in a quantitative study in Canada found 47% of African refugee children with malaria were asymptomatic. In the other quantitative study conducted among African refugee children in the United States, Maroushek et al. (2005) found 29% of children with malaria were asymptomatic. The findings from these two studies suggest malaria is regularly missed and therefore indicate the need for

optimal malaria screening, diagnosis, and treatment for refugee children from malaria-endemic regions.

Mental Health and Neurodevelopmental Disorders

Nine studies focused on mental health disorders among sub-Saharan African refugee children outside Africa, of which seven were quantitative (Ellis et al., 2008, 2010; Geltman et al., 2005, 2008; Huemer et al., 2011, 2013; Volkl-Kernstock et al., 2014), one was a mixed study (Lincoln et al., 2016), and one was qualitative (Miller-Gairy & Mofya, 2015). The main focus of studies was post-traumatic stress disorder (PTSD), depression, and autism spectrum disorder (ASD). Huemer et al. (2011) in a study on psychopathology in African unaccompanied refugee minors in Austria found girls to be more susceptible to PTSD than boys after controlling for age. Similarly, Volkl-Kernstock et al. (2014) in a quantitative study among unaccompanied African refugee minors in Austria reported that girls were more likely to develop PTSD than boys.

Several studies found the incidence rate of PTSD among African refugee children outside Africa to be between 17 and 20% (Ellis et al., 2010; Geltman et al., 2008; Huemer et al., 2011; Lincoln et al., 2016). Ellis et al. (2008) conducted a quantitative study on the mental health of Somali adolescent refugees in the United States and reported the severity of acculturative stress is associated with higher levels of PTSD symptoms. This result corroborates findings from a study by Geltman et al. (2005) of unaccompanied Sudanese refugee minors in United States that found social isolation and a history of personal injury were associated with PTSD. Four studies found an association between the incidence of PTSD and exposure to war or war-related events (Ellis et al., 2008; Geltman et al., 2005; Huemer et al., 2011; Volkl-Kernstock et al., 2014). In these studies, resettlement stress and acculturation were associated with depression (Lincoln et al., 2016). Huemer and colleagues' (2013) study on unaccompanied minors in Australia used a

theoretical model of resilience and had similar findings: unaccompanied minors have increased distress and depression. Finally, an ethnographic study on the perceptions and expectations of ASD among Somali refugee mothers in Georgia in the United States (Miller-Gairy & Mofya, 2015) found communication was the main barrier faced by children with ASD with respect to access to health care services.

Food insecurity and psychosocial adjustment

One mixed-method and two qualitative studies investigated food insecurity and psychosocial adjustment among sub-Saharan African refugee children outside Africa. The qualitative study by Hadley et al. (2007) used an ethnographic approach to study food insecurity among West African refugees resettled in the United States. Their results indicated measures of socioeconomic status are associated with food security and psychosocial adjustment. Wilson & Renzaho's (2015) qualitative study identified intergenerational difference in acculturation experience and food beliefs among East African refugees in Australia. The study by Weine et al. (2013) conducted among Liberian and Burundi refugee children using family eco-developmental theory found that finances, social support network, family engagement, adherence to cultural values, literacy, and participation in religious activities are means of promoting psychosocial adjustment.

Physical health

Four studies (all quantitative) focused on the physical health of sub-Saharan African refugee children living outside Africa, including obesity, skeletal growth, body composition, and metabolic risk. Hervey et al. (2009) in their study of Somali refugee children found 2% of the sample of 69 were overweight. Mellor et al. (2012) found the rate of overweight was 19.0% and

rate of obesity was 1% amongst Australian sub-Saharan African refugee children. However, Alasagheirin & Clark's (2018) study of 64 children found a higher prevalence rate of obesity (32%). Two studies reported an association between sex and increase in body mass index (BMI) among sub-Saharan African refugee children living outside Africa. Mellor et al. (2012) found the BMI for girls was on average slightly higher than for boys. Griffith et al. (2014) in a study on obesity among sub-Saharan African migrant adolescents in Australia reported sex and parental BMI were linked with an increase in adolescent BMI, with boys having a lower BMI than girls. However, Hervey et al.'s (2009) study found female Somali refugee children after arrival in the United States without a comorbid condition and those who never attended school were less likely than their counterparts to become overweight.

Health promotion strategies

Two qualitative studies investigated health promotion strategies among sub-Saharan African refugee children living outside Africa. In their study on adolescent refugees in the United States, Weine et al. (2014) used family eco-developmental theory to identify eight themes of family and community capacities as a means of promoting psychosocial well-being in adolescent refugees. This study also identified nine protective mechanisms as the processes for developing adolescent refugees' competencies and behaviors. Caron et al. (2013) conducted a qualitative study on lead poisoning among African refugee children in a Somali refugee resettlement community in the United States. Childhood lead poisoning was associated with government policies on African refugee education, ineffective communication methods, and lack of adequate education about this public health problem.

Discussion

We sought to synthesize the literature on the health of sub-Saharan African child refugees living outside the continent. Major areas of published research identified in our review on the health of sub-Saharan African refugee children living outside Africa include infectious disease, mental health, food insecurity, physical health (including obesity), and health promotion strategies. By and large, the literature in this field shows poor health status among sub-Saharan African child refugees living outside Africa. Our review is particularly important given that Africa contributes 25% of child refugees and migrants under the age of 18 years (International Organization for Migration, 2020). Given the lack of previous reviews on African refugee children living outside Africa, below we situate our findings within the broader literature on refugees, including non-African refugees and African adult refugees.

Our review found a high rate of reported post-traumatic stress among refugee children, consistent with other reviews of the literature that, for example, show refugees who have been affected by war and violence are vulnerable to poor mental health outcomes including post-traumatic stress (Colucci et al., 2014; Guruge & Butt, 2015). A systematic review of Karen refugees that included 18 studies shows they experience a high rate of anxiety, depression, and PTSD (Hoffman & Robertson, 2016). Similarly, a review on Sudanese refugees, mostly adults, found high rates of post-traumatic stress and indicated post-migration stressors, such as family stressors, have more impact on mental health than past experience of trauma. While we found limited data on the resilience displayed by sub-Saharan African refugee children living outside Africa, researchers report that refugee children generally have good resilience and function well even in the face of substantial adversities (Hodes & Vostanis, 2019). Given our current review did not identify studies that focus on the resilience of sub-Saharan African refugee children living outside Africa, this remains a gap that needs to be examined. Supporting refugees'

resilience and using a strengths-based approach can help with their healing path (Marshall et al., 2016); this is therefore an important consideration for sub-Saharan African refugee children living outside Africa.

Better understanding of the health experience of refugees requires consideration of the trajectory of the phases refugees experience, including the pre-migration or pre-flight phase, migration or flight phase, and post-migration or resettlement phase (Colucci et al., 2014; Guruge & Butt, 2015; Lustig et al., 2004; Marshall et al., 2016). Our review found that experiences during all stages of the migration process, including pre-migration trauma, migration as an unaccompanied minor, and socioeconomic context, may influence the mental and physical health of sub-Saharan African refugee children living outside Africa. During the pre-flight stage, refugees may have experienced or witnessed several atrocities including rape and murder of loved ones. In addition, children might have been witness to front-line combat, which can expose them to trauma and physical injuries (Lustig et al., 2004). During the flight stage, refugee children may experience separation from parents and become unaccompanied minors. They may also feel depersonalized in refugee camps and lack basic resources that are essential to mental and physical health. Lack of basic resources, such as water, and improper sanitation can increase their risk of contracting infectious diseases. Challenges with access to food can contribute to malnutrition. Some children are also detained or face a lengthy asylum process. Upon arrival in destination countries, these children may face challenges with settlement including access to health services, discrimination, poverty, and lack of social support (Guruge & Butt, 2015). Refugees often face many hurdles related to accessing healthcare in destination countries, including poor economics and unemployment, cultural differences, language barriers, and legal barriers. To attend to the mental health needs of refugee children, health professionals, including nurses, should pay attention to pre-migration factors as well as post-migration factors (including

income) from a strengths-based perspective (Hodes & Vostanis, 2019). Group interventions may also be effective for refugee children but, unfortunately, our research found limited intervention studies on sub-Saharan African refugee children living outside Africa. This indicates the need for more research on culturally appropriate interventions to improve the health of sub-Saharan African refugee children living outside the continent.

Based on our review, several social factors including income and sex influence the mental health of sub-Saharan African refugee children living outside Africa. Previous research on the mental health of immigrant and refugee children found female youths experience more mental health problems than male youths (Guruge & Butt, 2015), which is consistent with our findings. Also, the importance of community belonging and having a strong social support network has been established as a contributor to immigrant mental health in Canada (Salami et al., 2017). Research on Karen refugees shows that reformation of community and building a social network contribute to better mental health (Hoffman and Robertson, 2016). Creating natural environments (e.g., during delivery of parenting programs) that foster community belonging and improve social support networks can assist in improving the mental health of sub-Saharan African refugee children in Western host societies (Salami et al., 2019). Future research should shed light on the influence of the social determinants of mental health for sub-Saharan African refugee children in Western host societies.

Our review found several physical health problems among sub-Saharan African refugee children outside the continent, including high rates of infectious disease (especially malaria) and obesity. Similarly, Hoffman and Robertson (2016) note that refugee adults experience acute health conditions including infectious diseases (including tuberculosis and hepatitis) and chronic health conditions (such as hypertension) (Hoffman & Robertson, 2016). In two chart audits at a refugee clinic in Australia, researchers found the refugees were at high risk of being diagnosed

with vitamin D deficiency, tuberculosis infection, *Helicobacter pylori* infection, schistosomiasis, hepatitis B carrier status, dental disease, and anaemia (Johnston et al., 2012; Mutch et al., 2012). Two-thirds of children had at least one pathological condition that was not reported in their medical history (Mutch et al., 2012). Similar findings have been reported in studies in Canada (Salehi et al., 2015). A major contributing factor to health problems is pre- and post-migration socioeconomic conditions. Our review identified that refugee children in low-income households experience food insecurity, which may influence their nutritional status. Health practitioners should ensure a complete physical and mental assessment of refugees to address and detect these priority health issues. Addressing the social and economic conditions including access to food that may contribute to the health of refugees is also important. In addition, our review indicates sex differences in the rate of overweight among sub-Saharan African refugees living outside Africa, but the evidence in this regard is mixed. Thus, more research is needed in this field.

Our numerical summary further identifies gaps in the literature. Thirteen of the 20 studies reviewed were quantitative, which indicates the need for more qualitative work in this area. Regional disparities in the research are also evident. The majority (n=13) of the studies were done in the United States with others from Canada (n=1), Austria (n=3), and Australia (n=3). Gaps in interventions on key health issues under the thematic areas observed are also apparent. In particular, health interventions on infectious diseases, nutrition, and mental health issues are needed along with follow-up studies to investigate the association between the exposure of children to the risk factors observed and the development of adverse health outcomes.

References

- Alasagheirin, M. H., & Clark, M. K. (2018). Skeletal growth, body composition, and metabolic risk among Sudanese immigrant children. *Public Health Nursing, 35*(2), 91-99.
<http://dx.doi.org/10.1111/phn.12386>
- Arksey, H., & O'Malley, L. (2005). Scoping studies: towards a methodological framework. *International Journal of Social Research Methodology, 8*(1), 19-32.
- Borrego Jr., J., Ortiz-González, E., & Gissandaner, T. D. (2019). Ethnic and cultural considerations. In S. N. Compton, M. A. Villabø, & H. Kristensen (Eds.), *Pediatric Anxiety Disorders* (pp. 461-497). Academic Press.
- Caron, R. M., Tshabangu-Soko, T., & Finefrock, K. J. (2013). Childhood lead poisoning in a Somali refugee resettlement community in New Hampshire. *Journal of Community Health, 38*(4), 660-669. <http://dx.doi.org/10.1007/s10900-013-9661-5>
- Colucci, E., Szwarc, J., Minas, H., Paxton, G., & Guerra, C. (2014). The utilisation of mental health services by children and young people from a refugee background: a systematic literature review. *International Journal of Culture and Mental Health, 7*(1), 86-108.
- Ellis, B. H., MacDonald, H. Z., Lincoln, H. Z., & Cabral, H. J. (2008) Mental health of Somali adolescent refugees: the role of trauma, stress, and perceived discrimination. *Journal of Consulting and Clinical Psychology, 76*(2), 184-193. <http://dx.doi.org/10.1037/0022-006X.76.2.184>
- Ellis, H. B., Lincoln, A. K., Charney, M. E., Ford-Paz, R., Benson, M., & Strunin, L. (2010). Mental health service utilization of Somali adolescents: religion, community, and school as gateways to healing. *Transcultural Psychiatry, 47*(5), 789-811.
<http://dx.doi.org/10.1177/1363461510379933>

- Geltman, P. L., Grant-Knight, W., Mehta, S. D., Lloyd-Travaglini, C., Lustig, S., Landgraf, J. M., & Wise, P. H. (2005). The lost boys of Sudan: functional and behavioral health of unaccompanied refugee minors resettled in the United States. *Archives of Pediatrics and Adolescent Medicine*, *159*(6), 585-591.
- Geltman, P. L., Grant-Knight, W., Ellis, H., & Landgraf, J. M. (2008). The lost boys of Sudan: use of health services and functional health outcomes of unaccompanied refugee minors resettled in the U.S. *Journal of Immigrant and Minority Health*, *10*(5), 389-396.
<http://dx.doi.org/10.1007/s10903-007-9110-6>
- Griffith, M., Mellor, D., Green, J., & Renzaho, A. M. (2014). Migration-related influences on obesity among sub-Saharan African migrant adolescents in Melbourne, Australia. *Nutrition and Dietetics*, *71*(4), 252-257. <http://dx.doi.org/10.1111/1747-0080.12135>
- Guruge, S., & Butt, H. (2015). A scoping review of mental health issues and concerns among immigrant and refugee youth in Canada: looking back, moving forward. *Canadian Journal of Public Health*, *106*(2), e72-e78.
- Hadley, C., Zodhiates, A., & Sellen, D. W. (2007). Acculturation, economics and food insecurity among refugees resettled in the USA: a case study of West African refugees. *Public Health Nutrition*, *10*(4), 405-412. <http://dx.doi.org/10.1017/S1368980007222943>
- Hervey, K., Vargas, D., Klesges, L., Fischer, P. R., Trippel, S., & Juhn, Y. J. (2009). Overweight among refugee children after arrival in the United States. *Journal of Health Care for the Poor and Underserved*, *20*(1), 246-256. <http://dx.doi.org/10.1353/hpu.0.0118>
- Hodes, M., & Vostanis, P. (2019). Practitioner review: mental health problems of refugee children and adolescents and their management. *Journal of Child Psychology and Psychiatry*, *60*(7), 716-731.

- Hoffman, S. J., & Robertson, C. L. (2016). A systematic review of Karen refugee health. *International Journal of Migration, Health and Social Care*, 12(1), 1-15.
- Huemer, J., Karnik, N., Voelkl-Kernstock, S., Granditsch, E., Plattner, B., Friedrich, M., & Steiner, H. (2011). Psychopathology in African unaccompanied refugee minors in Austria. *Child Psychiatry and Human Development*, 42(3), 307-319.
<http://dx.doi.org/10.1007/s10578-011-0219-4>
- Huemer, J., Volkl-Kernstock, S., Karnik, N., Denny, K. G., Granditsch, E., Mitterer, M., Humphreys, K., Plattner, B., Friedrich, M., Shaw, R. J., & Steiner, H. (2013) Personality and psychopathology in African unaccompanied refugee minors: repression, resilience and vulnerability. *Child Psychiatry Human Development*, 44(1), 39-50.
<http://dx.doi.org/10.1007/s10578-012-0308-z>
- International Organization for Migration. (2020). Migration data portal. Retrieved from <https://migrationdataportal.org/themes/child-and-young-migrants>
- International Organization for Migration. (2020). World Migration Report. Geneva, Switzerland: International Organization for Migration.
- Johnston, V., Smith, L., & Roydhouse, H. (2012). The health of newly arrived refugees to the Top End of Australia: results of a clinical audit at the Darwin Refugee Health Service. *Australian Journal of Primary Health*, 18(3), 242-247.
- Lincoln, A. K., Lazarevic, V., White, M. T., & Ellis, B. H. (2016). The impact of acculturation style and acculturative hassles on the mental health of Somali adolescent refugees. *Journal of Immigrant and Minority Health*, 18(4), 771-778.
<http://dx.doi.org/10.1007/s10903-015-0232-y>

- Lustig, S. L., Kia-Keating, M., Knight, W. G., Geltman, P., Ellis, H., Kinzie, J. D., Keane, T., & Saxe, G. N. (2004). Review of child and adolescent refugee mental health. *Journal of the American Academy of Child & Adolescent Psychiatry*, 43(1), 24-36.
- Maroushek, S. R., Aguilar, E. F., Stauffer, W., & Abd-Alla, M. D. (2005). Malaria among refugee children at arrival in the United States. *The Pediatric Infectious Disease Journal*, 24(5), 450-452. <http://dx.doi.org/10.1097/01.inf.0000160948.22407.0d>
- Marshall, E. A., Butler, K., Roche, T., Cumming, J., & Taknint, J. T. (2016). Refugee youth: a review of mental health counselling issues and practices. *Canadian Psychology / Psychologie canadienne*, 57(4), 308-319.
- Mellor, D., Renzaho, A., Swinburn, B., Green, J., & Richardson, B. (2012). Aspects of parenting and family functioning associated with obesity in adolescent refugees and migrants from African backgrounds living in Australia. *Australian and New Zealand Journal of Public Health*, 36(4), 317-324. <http://dx.doi.org/10.1111/j.1753-6405.2012.00894.x>
- Miller-Gairy, S., & Mofya, S. (2015). Elements of culture and tradition that shape the perceptions and expectations of Somali refugee mothers about autism spectrum disorder. *International Journal of Child and Adolescent Health*, 8(4), 425-438.
- Mutch, R. C., Cherian, S., Nemba, K., Geddes, J. S., Rutherford, D. M., Chaney, G. M., & Burgner, D. P. (2012). Tertiary paediatric refugee health clinic in Western Australia: analysis of the first 1026 children. *Journal of Paediatrics and Child Health*, 48(7), 582-587.
- Ndao, M., Bandyayera, E., Kokoskin, E., Diemert, D., Gyorkos, T. W., MacLean, J. D., St. John, R., & Ward, B. J. (2005). Malaria “epidemic” in Quebec: diagnosis and response to imported malaria. *Canadian Medical Association Journal*, 172(1), 46-50. <http://dx.doi.org/10.1503/cmaj.1031862>

- Salami, B., Yaskina, M., Hegadoren, K., Diaz, E., Meherali, S., Rammohan, A., & Ben-Shlomo, Y. (2017). Migration and social determinants of mental health: results from the Canadian Health Measures Survey. *Canadian Journal of Public Health, 108*(4), e362-367.
- Salami, B., Salma, J., Hegadoren, K., Meherali, S., Kolawole, T., & Diaz, E. (2019). Sense of community belonging among immigrants: perspective of immigrant service providers. *Public Health, 167*, 28-33.
- Salehi, L., Lofters, A. K., Hoffmann, S. M., Polsky, J. Y., & Rouleau, K. D. (2015). Health and growth status of immigrant and refugee children in Toronto, Ontario: a retrospective chart review. *Paediatrics & Child Health, 20*(8), e38-42.
- UNICEF (2016). Uprooted. The Growing Crisis for Refugee and Migrant Children.
https://www.unicef.org/publications/files/Uprooted_growing_crisis_for_refugee_and_migrant_children.pdf
- United Nations Development Program (UNDP). (2020). Sustainable Development Goals.
Retrieved from <https://www.undp.org/content/undp/en/home/sustainable-development-goals/goal-10-reduced-inequalities/targets.html>
- United Nations High Commissioner for Refugees (UNHCR). (2020). What is a refugee.
Retrieved from <https://www.unhcr.org/what-is-a-refugee.html>
- Völkl-Kernstock, S., Karnik, N., Mitterer-Asadi, M., Granditsch, E., Steiner, N., Friedrich, M. H., & Huemer, J. (2014). Responses to conflict, family loss and flight: posttraumatic stress disorder among unaccompanied refugee minors from Africa. *Neuropsychiatry, 28*, 6-11. <http://dx.doi.org/10.1007/s40211-013-0094-2>
- Weine, S. M., Ware, N., Tugenberg, T., Hakizimana, L., Dahnweih, G., Currie, M., Wagner, M., & Levin, E. (2013). Thriving, managing, and struggling: a mixed methods study of

- adolescent African refugees' psychosocial adjustment. *Adolescent Psychiatry (Hilversum)*, 3(1), 72-81.
- Weine, S. M., Ware, N., Hakizimana, L., Tugenberg, T., Currie, M., Dahnweih, G., Wagner, M., Polutnik, C., & Wulu, J. (2014). Fostering resilience: Protective agents, resources, and mechanisms for adolescent refugees' psychosocial well-being. *Adolescent Psychiatry (Hilversum)*, 4(4), 164–176. <http://dx.doi.org/10.2174/221067660403140912162410>.
- Wilson, A., & Renzaho, A. (2015). Intergenerational differences in acculturation experiences, food beliefs and perceived health risks among refugees from the Horn of Africa in Melbourne, Australia. *Public Health Nutrition*, 18(1), 176–188. <http://dx.doi.org/10.1017/S1368980013003467>