



Calhoun: The NPS Institutional Archive
DSpace Repository

Faculty and Researchers

Faculty and Researchers' Publications

2002-09

Near Shore Wave Processes

Thornton, Edward B.; Stanton, Timothy P.

<http://hdl.handle.net/10945/69023>

This publication is a work of the U.S. Government as defined in Title 17, United States Code, Section 101. Copyright protection is not available for this work in the United States.

Downloaded from NPS Archive: Calhoun



Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.

Dudley Knox Library / Naval Postgraduate School
411 Dyer Road / 1 University Circle
Monterey, California USA 93943

<http://www.nps.edu/library>

Near Shore Wave Processes

Edward B. Thornton
Oceanography Department
Naval Postgraduate School
Monterey, CA 93943-5000
phone: (831) 656-2847 fax: (831) 656-2712 email: thornton@nps.navy.mil

Timothy P. Stanton
phone: (831) 656-3144 email: stanton@nps.navy.mil

Award #'s: N0001402WR20188, N0001402WR20376, N0001402WR20153
<http://www.oc.nps.navy.mil/~thornton/>
<http://www.oc.nps.navy.mil/~stanton/>
<http://www.frf.usace.army.mil/SandyDuck/SandyDuck.stm>

LONG-TERM GOALS

Long-term goals are to predict the wave-induced three-dimensional velocity field and induced sediment transport over arbitrary bathymetry in the near shore given the offshore wave conditions.

OBJECTIVES

The interrelationship of wave-induced hydrodynamic and sediment processes over the vertical and morphologic processes at the bed are measured and modeled. The primary mechanism for changes in momentum flux that drive near shore hydrodynamics is due to the dissipation by breaking waves, the processes of which are poorly understood. To improve our understanding of breaking waves, the dissipation associated with bubble injection is measured along with the velocity fields over the vertical. Bottom boundary layer measurements are obtained to determine bottom stress and dissipation. Sediment transport is measured in response to the measured mean longshore and cross-shore currents, wave velocities and induced stresses. The small-scale morphology, which acts as hydraulic roughness for the mean flows and perturbs the velocity-sediment fields, is measured as a function of time and over large areas to examine cross-shore and alongshore variation.

APPROACH

A combined experiment (SteepBeach/RIPEX) investigating wave transformation over a steep beach (high Iribarren number) and rip currents was conducted April/May 2001 in Southern Monterey Bay as a pilot experiment to test instrumentation for NCEX. The large-scale morphology is a barred shoreline incised by deep rip channels spaced 100-200 m apart. Offshore of the bar the slope is 1:20, and the beach slope is steep at 1:5. An unexpectedly large variation in incident waves occurred during the experiment owing to a series of storms, resulting in waves up to 4 m at breaking. Comprehensive wave and current data were obtained for the first time on a steep beach and for rip currents, and these data greatly expand the range of measured beach processes. The experiment was a collaborative effort lead by Thornton and Stanton of the Naval Postgraduate School, PhD student MacMahan (University of Florida), and Lippmann (Ohio State University).

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 30 SEP 2002		2. REPORT TYPE		3. DATES COVERED 00-00-2002 to 00-00-2002	
4. TITLE AND SUBTITLE Near Shore Wave Processes				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Oceanography Department,,Naval Postgraduate School,,Monterey,,CA, 93943				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Long-term goals are to predict the wave-induced three-dimensional velocity field and induced sediment transport over arbitrary bathymetry in the near shore given the offshore wave conditions.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

The comprehensive Delft3D morphodynamic model developed by Delft Hydraulics is being assessed by comparing with field data. This work is lead by Reniers (National Research Council post-doc) with participation of NPS graduate students. In addition, process models for breaking waves, momentum mixing due to the interaction of longshore and cross-shore vertical mean profiles, and bottom shear stress enhanced by the form drag of bedforms and by turbulence of wave breaking are compared with observations, which are the focus of Stanton, Thornton and their students. Both linear and nonlinear (Boussinesq) wave models (collaborative with Karakiewicz and PhD student Morichon, U. of Quebec) are considered.

WORK COMPLETED

Work completed includes and analysis of the SteepBeach/RIPEX data, improved techniques for determining reflection, and modeling the hydrodynamics and large-scale morphodynamics in the nearshore with an advanced Delft3D model.

- 1) The generation of infragravity waves by directionally spread short waves incident on an alongshore uniform beach was investigated using an advanced version of Delft3D. The infragravity wave response is examined using linear shallow water equations, taking into account the presence of bottom friction, set-up, rollers and the longshore current. The infragravity energy density spectrum and surface elevation at the infragravity band time scale is generated by summing all difference interaction frequency pairs of the input short-wave directional spectra obtained. Model results are compared with the data from the Delilah experiment (Reniers et.al., 2002a), and the results of the SteepBeach/RIPEX experiment (Reniers et.al., 2002b)
- 2) The breaking wave parameterization used in the Battjes and Janssen (1978) surf zone wave transformation model incorporated in Delft3D calibration was extended to include a larger range of field data including longer period swell (Morris et.al., 2002).
- 3) A new technique for determining wave reflection in the nearshore using PUV measurements as an extension of the work by Dixon et.al. (1995) was developed (Thornton et.al., 2002)
- 4) Data obtained during the SteepBeach/RIPEX was analyzed. Pulsations of the rip currents at infragravity frequencies were found to be the result of wave group forcing (MacMahan et.al., 2002).
- 5) Significant very low frequency energy (VLF's) with periods greater than 5 minutes outside the region of zero-mode edge waves was found and shown to be attributed to directionally broad wave group forcing and/or instabilities of the rip current itself, resulting in large-scale vorticies (MacMahan et. al., 2002).
- 6) The morphodynamics of rip current systems on an initially alongshore uniform barred beach with directionally broad spectral wave forcing with a mean angle of normal incidence is initially perturbed by VLF's and/or self organized motions that determine the alongshore length scale of the rip channels The rip channels are enhanced by a positive feedback of the wave group forcing and rip currents themselves. The alongshore scale of the rip channel spacing is shown to be a function of the directional bandwidth. (Reniers et.al., 2002c).

7) The Delft3D morphodynamic model was assessed by comparing model output with data from Duck94 experiment.

RESULTS

The morphodynamic response of an embayed beach induced by wave groups generated by a directionally broad wave spectrum with mean angle normally incident is examined with a numerical model (Reniers, et.al. 2002). The model utilizes the nonlinear shallow water equations to phase resolve the mean and infragravity motion in combination with an advection-diffusion equation for sediment transport. Starting with an initially alongshore uniform barred beach, the bathymetry evolves to the shoals cut by quasi-periodic rip channels. Without directional spreading, the smallest alongshore separation is obtained and the beach response is self-organizing in nature. Introducing directional spreading results in a limited range of preferred spacing between rip channels, qualitatively similar to observations (Figure 1). The hypothesized correlation between the observed rip spacing and wave group forced edge waves over the initially alongshore uniform bathymetry is not found. However, there is a correlation between the alongshore lengths of the wave-group induced quasi-steady flow circulations (very-low frequency oscillations) and the rip current spacing. This suggests that the scouring associated with the flow circulations of the initial wave groups triggers the development of rip channels via a positive feedback mechanism in which the small scour holes start attracting more and more discharge.

A linear version of the above model was used to generate infragravity energy density spectra and compared with data from the Delilah experiment. Two mechanisms responsible for the generation of infragravity waves are considered: the release of the bound infragravity waves associated with changes in the spatial variation of the incident shore wave energy, and the forcing of trapped waves by obliquely incident directionally spread short waves. Typically, 80% of the infragravity wave height variability is explained by the model, of which 30% or less is due to bound infragravity waves (Reniers, et.al., 2002b).

Numerical model computations of infragravity motions are compared with measurements obtained during the RIP-current EXperiment (RIPEX) in concert with the Steep Beach Experiment (SBE). The experiments were performed at Sand City, Monterey Bay, CA, during the spring of 2001. The nearshore bathymetry was made up of shore-connected shoals incised by relatively narrow rip-channels spaced approximately 125 m apart. The comparison considers a 24 day period during which significant changes in both the offshore wave climate and nearshore bathymetry occurred. Analysis from the alongshore array (MacMahan et.al., 2002) indicates that there is significant energy in the cross-shore infragravity velocities, and that there is little alongshore spatial variation, even in the presence of rip channels. Rip current pulsations at the infragravity band frequencies were found linked to the infragravity motions of the bound and free long waves, as opposed to forcing by dynamic hydraulic head (wave set-up) associated with incoming short-wave groups with preferred drainage through the rip channels. Reniers et.al., (2002c) utilized a non-linear shallow water wave model operating on the time-scale of wave groups, to compare with measurements to examine the effects of the complex bathymetry on the infragravity motions (Figure 2). The temporal variation in infragravity conditions during the experiment is strong, with computational results typically explaining 75 % of the observed infragravity motions within the nearshore. In contrast to the temporal variation, the alongshore spatial variation in infragravity intensity during the experiment is generally small, even though the underlying bathymetry shows strong depth variations.

IMPACT/APPLICATIONS

On the basis of Delft3D hydrodynamic model comparisons with comprehensive nearshore field data acquired over two decades funded in all or part by ONR, it is recommended the U.S. Navy adopt Delft3D as an operational surf model.

TRANSITIONS

It is recommended on the basis of comparisons with comprehensive nearshore field data that the Delft3D hydrodynamics model be modified to include roller dynamics and then be adopted by the U.S. Navy as an operational surf model.

RELATED PROJECTS

1. Results of process modeling obtained on this project are being applied to nearshore modeling efforts under the following programs: Modeling Wave Dissipation within the Wave Boundary Layer (ONR), and Development and Verification of a Comprehensive Community Model for Physical Processes in the Nearshore (NOPP).
2. Collaborative modeling and data comparisons of breaking waves using Boussinesq equations is being performed by PhD students at the U of Quebec under co-direction with Barbara Boczar-Karakiewicz.

REFERENCES

Battjes, J. A. and J.P.F.M. Janssen, 1978, Energy loss and set-up due to breaking of random waves, Proc. 16th Int. Conf. on Coastal Eng., ASCE, 569-587.

Dickson, W.S., T.H.C. Herbers and E.B. Thornton, 1994, "Wave reflection from a breakwater", J. Waterway, Port, Coastal and Ocean Engineering, 121 (5), 262-268.

MacMahan, J, A.J.H.M. Reniers, E.B. Thornton, and T.P. Stanton, 2002, Infragravity Motions on a Complex Beach, Part 1: Observations, submitted to J. Geophysical Research.

Morris, B., E.B. Thornton, and A. Reniers, 2002, Calibration of a random wave model extended to long period swell, accepted J. Coastal Engineering.

Reniers, A.J.H.M., J.A. Roelvink, and E.B. Thornton, 2002a, Morphodynamic modeling of an embayed beach under wave group forcing, submitted to J. Geophysical Research.

Reniers, A.J.H.M., A. van Dongeren, J. Battjes, and E.B. Thornton, 2002b, Linear modelling of infragravity waves during Delilah, in press, J. Geophysical Research.

Reniers, A.J.H.M, J. MacMahan, J, E.B. Thornton, and T.P. Stanton, 2002c, Infragravity Motions on a Complex Beach, Part 2: Modeling, submitted to J. Geophysical Research.




Figure 1. Model predicted average spacing between rip channels, L_r , for one-meter wave height with mean wave direction normal to shore varies between 150 m for unidirectional waves (zero directional spreading factor, DSPR) to 350 m for DSPR of 2° and decreases thereafter for increasing DSPR, showing that the rip current spacing is bounded and in qualitative agreement with observations. The bars indicate the standard deviation of an ensemble of 8 simulations.




Figure 2. Predicted wave heights (left panel), radiation stress and pressure gradient forcing (middle panel) and currents (right panel) overlying bathymetry and denoting location of sensors during SteepBeach/Ripex experiment. A rip current at 120m alongshore and very low frequency oscillations at 20 m alongshore are evident. Good comparisons with data are obtained.

PUBLICATIONS

Reniers, A.J.H.M., A. van Dongeren, J. Battjes, and E.B. Thornton, 2002b, Linear modelling of infragravity waves during Delilah, in press, J. Geophysical Research.

Gallagher, E.L., and E.B. Thornton, 2002, Megaripples in the surf zone and their relation to steady flow dunes, accepted in Marine Geology.

Gallagher, E.L., E.B. Thornton and T.P. Stanton, 2002, Sand bed roughness in the nearshore, accepted J. Geophys. Res.

Morris, B., E.B. Thornton, and A. Reniers, 2002, Calibration of a random wave model extended to long period swell, accepted J. Coastal Engineering.

Saulter, A.N., P.E. Russell, J.R. Miles, and E.L. Gallagher, 2001, Observations of Bed Level Change in a Satruated Surf Zone, submitted to J. Geophysical Research.

Feddersen, F. E.L. Gallagher, R.T.Guza, and S. Elgar, 2001, The Drag Coefficient in the Nearshore, submitted to J. Geophysical Research.

Chen, Q., J.T. Kirby, R.A. Dalrymple, F. Shi, and E.B. Thornton, 2001, Boussinesq Modeling of Longshore Currents, submitted to J. Geophysical Research.

Reniers, A.J.H.M., E.B. Thornton and T.C. Lippmann, 2001, Effects of Alongshore Non-uniformities on longshore currents measured in the field, submitted to the J. Geophysical Research.

Reniers, A.J.H.M., J.A. Roelvink, and E.B. Thornton, 2002, Morphodynamic modeling of an embayed beach under wave group forcing, submitted to J. Geophysical Research.

MacMahan, J, A.J.H.M. Reniers, E.B. Thornton, and T.P. Stanton, 2002, Infragravity Motions on a Complex Beach, Part 1: Observations, submitted to J. Geophysical Research.

Reniers, A.J.H.M, J. MacMahan, J, E.B. Thornton, and T.P. Stanton, 2002, Infragravity Motions on a Complex Beach, Part 2: Modeling, submitted to J. Geophysical Research.

NON-REFEREED PUBLICATIONS

Conference Proceedings

Van Dongeren, A., A. Reniers, 2001, Nonlinear modeling of infragravity wave response under field conditions, Proc. Coastal Dynamics '01, ASCE, 283-292.

Reniers, A., G. Symonds and E. Thornton, 2001, Modelling of rip currents during RDEX, Proc. Coastal Dynamics '01, ASCE, 493-499.

Lippmann, T.C., T.H.C. Herber, and E.B. Thornton, 2001, Observations of infragravity waves in the nearshore, Proc. Coastal Dynamics '01, ASCE, 55-61.

Morichon, D., B. Boczar-Karakiewicz, and E.B. Thornton, 2001, Boussinesq wave model compared with field data, 2001, Proc. Coastal Dynamics '01, ASCE, 365-372.

Morris, B., E.B. Thornton and A. Reniers, 2001, Nearshore wave and current predictions compared with field observations, Proc. Coastal Dynamics '01, ASCE, 788-797.

Foster, D.L., T. Stanton, K. Andersen, J. Fredsoe, and E. Thornton, 2001, Model-data comparisons of velocity and suspended sediment in a wave dominated environment, Proc. Coastal Dynamics '01, ASCE, 751-758.

Conference Presentations:

Reniers, A., and E.B. Thornton, Directional properties of infragravity waves in the surfzone, Waves '01, San Francisco, July 2001.

Morichon, D., E.B. Thornton, and B. Boczar-Krakiewicz, Boussinesq model for breaking wave height distributions, mean currents and morphology compared with field data, Waves '01, San Francisco, July 2001.

Thornton, E. B., My Historical Perspective on the ONR Coastal Science Program and the Kinder Years, American Geophysical Union Fall Mtg., San Francisco, Dec. 2001.

Reniers, A., E.B. Thornton, T. Stanton, and J. MacMahan, Modeling of Rip-Currents during RIPEX, American Geophysical Union Fall Mtg., San Francisco, Dec. 2001.

Stanton, T. P., Turbulent Stresses and Shear Production in Bottom Boundary Layers in the Surf Zone and Inner Shelf, American Geophysical Union Fall Mtg., San Francisco, Dec. 2001.

Foster, D.L., T.P. Stanton, and J. Fredsoe, Model-Data Comparisons of Velocity and Suspended Sediment, American Geophysical Union Fall Mtg., San Francisco, Dec. 2001.

Natoo, P, D.L. Foster, and T.P. Stanton, Evaluation of Three Bed Load Transport Models, American Geophysical Union Fall Mtg., San Francisco, Dec. 2001.

Reniers, A., E.B. Thornton, T.P. Stanton, and J. MacMahan, RIPEX: Rip-current pulsation modeling, 28th International Conf. Coastal Engineering , Cardiff, Wales, UK, July 2002.

MacMahan, J., E.B. Thornton, T.P. Stanton, A. Reniers and R. Dean, RIPEX: Rip-current pulsation measurements, 28th International Conf. Coastal Engineering , Cardiff, Wales, UK, July 2002.

Foster, D., P. Natoo, T. Stanton, J. Fredsoe, and E. Thornton, Evaluation of three bed load transport models, 28th International Conf. Coastal Engineering , Cardiff, Wales, UK, July 2002.

Thornton, E., T. Stanton, and A. Reniers, Wave reflection on a steep beach, 28th International Conf. Coastal Engineering , Cardiff, Wales, UK, July 2002.

Thesis Directed:

Martin, Steve, Vortex Ripple Morphology Using Dune2D Model, M.S. Thesis, Naval Postgraduate School, March 2001.

Morris, Bruce, Nearshore Wave and Current Dynamics, Ph.D. Thesis, Naval Postgraduate School, September 2001, 89 pp.

Cutshaw, Charles, Verification of a one-dimensional surf prediction model for steep beach conditions, M.S. Thesis, Naval Postgraduate School, June 2002, 25 pp.

Egley, Lora, An application of LIDAR to examine erosion in Southern Monterey Bay during the 1997-98 El Nino, M.S. Thesis, Naval Postgraduate School, March 2002, 55 pp.

Welsch, Charlotte A., Assessment of Delft3D morphologic model during Duck94, M.S. Thesis, Naval Postgraduate School, September 2002.