



Calhoun: The NPS Institutional Archive
DSpace Repository

Faculty and Researchers

Faculty and Researchers' Publications

1999-09

Collaborative Research Projects in Support of FNMOOC Operational Mission

Elsberry, Russell L.

<http://hdl.handle.net/10945/69525>

This publication is a work of the U.S. Government as defined in Title 17, United States Code, Section 101. Copyright protection is not available for this work in the United States.

Downloaded from NPS Archive: Calhoun



Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.

Dudley Knox Library / Naval Postgraduate School
411 Dyer Road / 1 University Circle
Monterey, California USA 93943

<http://www.nps.edu/library>

Collaborative Research Projects in Support of FNMOC Operational Mission

Russell L. Elsberry
Department of Meteorology
Naval Postgraduate School
589 Dyer Rd., Room 254
Monterey, CA 93943-5114

Phone: (831) 656-2373 Fax: (831) 656-3061 e-mail: elsberry@met.nps.navy.mil
Award #: N0001499WR30003

LONG-TERM GOAL

The long-term goal is to improve the prediction of tropical cyclone track and structure so that warnings to the Fleet units afloat and ashore are optimized.

OBJECTIVES

(1) Improve tropical cyclone track prediction by demonstrating that tailored objective aids within the framework of the Systematic Approach to Tropical Cyclone Track Forecasting will provide better guidance than existing objective aids. (2) Improve understanding of tropical cyclone structure including rapid intensification and extratropical transition.

APPROACH

(1) Utilize the Systematic Approach framework to demonstrate the feasibility of optimizing track forecast guidance. (2) Use the ONR-sponsored Tropical Cyclone Motion (TCM-90) field experiment data set to analyze the environmental factors leading to intensity and structure changes in Supertyphoon Flo. (3) Utilize a new data base of tropical cyclones undergoing extratropical transition in the western North Pacific to document the characteristics and the physical processes involved.

WORK COMPLETED

(1) A simple statistical-synoptic technique for tropical cyclone track forecasting to 72 h in the western North Pacific has been derived and demonstrated to have considerable skill. (2) An exhaustive study of the TCM-90 data set related to the structure changes of Supertyphoon Flo has been completed with the publication of the Ph.D. dissertation of David Titley (CDR, USN).

RESULTS

(1) A journal article by Chen et al. (1999) documents the simple statistical-synoptic technique. For a statistical-synoptic equation set derived and tested with tropical cyclones in the Standard/Dominant Ridge pattern/region for the entire 72-h forecast interval, the improvement relative to the operational CLIPER is 13% after only 12 h and increases to 24% at 72 h. For the Poleward/Poleward-oriented pattern/region, the improvement relative to the operational CLIPER is about 14% after 12 h, and is about 11% at 72 h. Given a perfect knowledge of the type and timing of these two transitions, the appending of a composited post-transition track can lead to about 50% reduction relative to the 72-h

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 30 SEP 1999	2. REPORT TYPE	3. DATES COVERED 00-00-1999 to 00-00-1999			
4. TITLE AND SUBTITLE Collaborative Research Projects in Support of FNMOC Operational Mission		5a. CONTRACT NUMBER			
		5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)		5d. PROJECT NUMBER			
		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Postgraduate School, Department of Meteorology, 589 Dyer Rd., Room 254, Monterey, CA, 93943		8. PERFORMING ORGANIZATION REPORT NUMBER			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	4	

operational CLIPER errors if the transition occurs early in the forecast interval. This result is a validation of the environmental structure change focus of the Systematic Approach.

Elsberry et al. (1999a) and Boothe et al. (1999) describe a statistical post-processing of the Naval Operational Global Atmospheric Prediction System (NOGAPS) tropical cyclone track forecasts. Zonal and meridional adjustments to the NOGAPS-predicted track positions are derived from a 1992-96 sample of 152 tropical cyclones in the western North Pacific. The primary objective is to improve the early track errors that arise from an offset in the initial position in the NOGAPS model relative to an updated position that will be known 6 h after the 00 UTC and 12 UTC synoptic times, when the forecasters first make use of the NOGAPS predictions. Early track discrepancies relative to a persistence of past motion orientation that may arise from an incorrect analysis or a model systematic bias are also corrected by the post-processing technique. Validations with three independent data sets illustrate the likely usefulness of the post-processing technique throughout the Northern Hemisphere, even though the development sample contained only western North Pacific cyclones during 1992-96. Application to the 1997 western North Pacific tropical cyclones results in improvements relative to the unadjusted NOGAPS forecast errors of 50% at 12 h and 12% at 36 h. Corresponding percentage improvements for a 1995-97 sample of NOGAPS forecasts of Atlantic tropical cyclones are 38% and 14%. In the eastern/central North Pacific, the corresponding percentage improvements in the mean adjusted NOGAPS track error are 50% and 17%.

An overview of the Systematic Approach and its likely global applicability has been prepared for the H. Riehl memorial volume (Elsberry et al. 1999b). This overview brings together the original concepts developed for western North Pacific tropical cyclones with the experiences gained in applications in the eastern/central North Pacific and in the Southern Hemisphere. Three of the four original synoptic patterns apply in all these regions. However, each basin has a special synoptic pattern: monsoon Gyre (G) in the western North Pacific, upper-tropospheric Low (L) in the eastern/central North Pacific, and High-amplitude (H) in the Southern Hemisphere. It is important that all tropical cyclone tracks may be classified into just four synoptic patterns in each basin, so that a consistent "storyline" can be used to describe each track. Extension to the Atlantic hurricanes is in progress, and cyclones in the North Indian Ocean will be classified in the future.

A broader perspective on progress and perspectives for providing early (and more accurate) warnings of tropical cyclones was prepared (Elsberry 1999) for an international conference in Potsdam, Germany during September 1998. This conference was both a retrospective on progress during the International Decade for Natural Disaster Reduction and a planning for future activities. In addition to providing the keynote address on tropical cyclones, a session was organized that brought together forecasters and emergency managers from around the globe. Four of these presentations were edited for inclusion in the proceedings volume.

As co-chair of the U. S. Weather Research Program Hurricane Landfall Workshop in Miami, Florida, during November 1997, a Workshop report (Elsberry and Marks 1999) was prepared. This report summarizes and ranks opportunities in research and observations in five topics related to hurricane landfall on the U. S. coastline from Texas to Maine. These topics are track prediction, outer and inner (including intensity) structure changes, modifications of circulation as the hurricane crosses coastline, quantitative precipitation estimation and prediction, and social-economic impact assessments. This workshop report will serve as the basis for developing a five-year research plan beginning in October 1999.

(2) Titley (1998) developed a Multi-Quadric (MQ) analysis and compared it with the TCM-90 four-dimensional data assimilation (4DDA) analyses of the evolution of Typhoons Flo and Ed. The MQ analysis is shown to provide an alternate, plausible depiction of the tropical atmosphere. These analyses are then used to analyze the physical processes leading to rapid intensification of Flo, but not of Ed. Significant results (Titley and Elsberry 1999) include: (i) existence of a “cyclonic wind burst” extending beyond 1000 km radius at 200 mb during a pre-conditioning period prior to the rapid intensification of Flo; (ii) Flo had developed a warmer core near the tropopause, and was less stable than Ed in the lower- and mid- troposphere prior to the beginning of rapid intensification; (iii) a strong sustained eddy flux convergence (EFC) in the upper troposphere during the pre-conditioning period had the role of a catalyst, since the EFC forcing was significantly reduced during the rapid intensification; (iv) vertical wind shear was not an inhibiting factor in the intensification, rather it appeared to decrease as a result of the rapid intensification; (v) Supertyphoon Flo rapidly decreased in intensity while over warm water and while large EFC forcing aloft was still present, presumably because the interaction with the midlatitude circulation also produced a mid-tropospheric EFC that led to an outward secondary circulation that tends to spin down the mid-tropospheric vortex.

(3) The so-called extratropical transition of a tropical cyclone to an extratropical cyclone is not well understood because it involves complex interactions on various horizontal and vertical scales and over a range of time scales. Our recent progress and journal submissions on extratropical transition are described in the reports of P. A. Harr and E. A. Ritchie.

TRANSITIONS

The feasibility of the statistical-synoptic track prediction technique will be tested as part of a 6.4 project funded by SPAWAR and offered to JTWC for operational testing. The post-processing technique for NOGAPS will be re-derived for the combined data sets throughout the Northern Hemisphere (and separately for the Southern Hemisphere). A version for the Advanced Tropical Cyclone Forecast (ATCF) workstation will be tested as part of the SPAWAR project and provided to JTWC, Pearl Harbor, and Norfolk centers.

REFERENCES

- Boothe, M. A., R. L. Elsberry, and G. A. Ulses, 1999: Statistical post-processing of NOGAPS track forecasts. *Preprints, 23rd Conf. Hurr. Trop. Meteor.*, Amer. Meteor. Soc., Boston, MA 02108, 1061-1064.
- Chen, J.-M., R. L. Elsberry, M. A. Boothe, and L. E. Carr, III, 1998: A simple statistical-synoptic track prediction technique for western North Pacific tropical cyclones. *Mon. Wea. Rev.*, **127**, 89-102.
- Elsberry, R. L., 1999: Track forecast guidance improvements for early warnings of tropical cyclones. *Proceedings, Early Warning Conference-98* (in press).
- Elsberry, R. L., and L. E. Carr, III, 1999: Consensus of dynamical tropical cyclone track forecasts. *Mon. Wea. Rev.*, (submitted).
- Elsberry, R. L., and F. D. Marks, Jr., 1999: The Hurricane Landfall Workshop Report. *Bulletin American Meteorological Society*, **80**, 683-685.

- Elsberry, R. L., M. A. Boothe, G. A. Ulses, and P. A. Harr, 1999: Statistical post-processing of NOGAPS tropical cyclone track forecasts. *Mon. Wea. Rev.*, **127**, 1912-1919.
- Elsberry, R. L., L. E. Carr, III, and M. A. Boothe, 1999b: Progress toward a generalized description of the environment structure contribution to tropical cyclone track types. *Meteor. Atmos. Phys.*, 93-116.
- Titley, D. W., 1998: Intensification and structure change of Supertyphoon Flo as related to the large-scale environment. Ph.D. Dissertation, Naval Postgraduate School, Monterey, CA 93943-5114, 389 pp.
- Titley, D. W., and R. L. Elsberry, 1999: On rapid intensity changes in tropical cyclones: A case study of Supertyphoon Flo during TCM-90. *Mon. Wea. Rev.*, accepted.
- Titley, D. W., and R. L. Elsberry, 1999: Rapid intensification of tropical cyclones: A case study of Supertyphoon Flo during TCM-90. *Preprints 23rd Conf. Hurr. Trop. Meteor.*, Amer. Meteor. Soc., Boston, MA 02108, 588-591.

PUBLICATIONS

- Boothe, M. A., R. L. Elsberry, and G. A. Ulses, 1999: Statistical post-processing of NOGAPS track forecasts. *Preprints, 23rd Conf. Hurr. Trop. Meteor.*, Amer. Meteor. Soc., Boston, MA 02108, 1061-1064.
- Chen, J.-M., R. L. Elsberry, M. A. Boothe, and L. E. Carr, III, 1998: A simple statistical-synoptic track prediction technique for western North Pacific tropical cyclones. *Mon. Wea. Rev.*, **127**, 89-102.
- Elsberry, R. L., 1999: Track forecast guidance improvements for early warnings of tropical cyclones. *Proceedings, Early Warning Conference-98* (in press).
- Elsberry, R. L., and F. D. Marks, Jr., 1999: The Hurricane Landfall Workshop Report. *Bulletin American Meteorological Society*, **80**, 683-685.
- Elsberry, R. L., M. A. Boothe, G. A. Ulses, and P. A. Harr, 1999a: Statistical post-processing of NOGAPS tropical cyclone track forecasts. *Mon. Wea. Rev.*, **127**, 1912-1919.
- Elsberry, R. L., L. E. Carr, III, and M. A. Boothe, 1999b: Progress toward a generalized description of the environment structure contribution to tropical cyclone track types. *Meteor. Atmos. Phys.*, 93-116.
- Titley, D. W., and R. L. Elsberry, 1999: On rapid intensity changes in tropical cyclones: A case study of Supertyphoon Flo during TCM-90. *Mon. Wea. Rev.*, (accepted).
- Titley, D. W., and R. L. Elsberry, 1999: Rapid intensification of tropical cyclones: A case study of Supertyphoon Flo during TCM-90. *Preprints 23rd Conf. Hurr. Trop. Meteor.*, Amer. Meteor. Soc., Boston, MA 02108, 588-591.