

# Identification of QTLs Controlling Resistance to Anthracnose Disease in Water Yam (*Dioscorea alata*)

Paterne Angelot Agre <sup>1</sup>, Kwabena Darkwa <sup>2</sup>, Bunmi Olasanmi <sup>3</sup>, Olufisayo Kolade <sup>1</sup>, Pierre Mournet <sup>4,5</sup>, Ranjana Bhattacharjee <sup>1</sup>, Antonio Lopez-Montes <sup>6</sup>, David De Koeber <sup>7</sup>, Patrick Adebola <sup>1</sup>, Lava Kumar <sup>1</sup>, Robert Asiedu <sup>1</sup> and Asrat Asfaw <sup>1,\*</sup>

<sup>1</sup> International Institute of Tropical Agriculture (IITA), Ibadan 5320, Nigeria; P.Agre@cgiar.org (P.A.A.); O.Kolade@cgiar.org (O.K.); R.Bhattacharjee@cgiar.org (R.B.); p.adebola@cgiar.org (P.A.); L.Kumar@cgiar.org (L.K.); R.Asiedu@cgiar.org (R.A.)

<sup>2</sup> Savanna Agricultural Research Institute, Tamale NT0000 - NT2701, Ghana; kwabenadarkwa@gmail.com

<sup>3</sup> Department of Agronomy, University of Ibadan, Ibadan 200284, Nigeria; bunminadeco@yahoo.com

<sup>4</sup> Centre de Coopération Internationale en Recherche Agronomique pour le Développement Montpellier, 34398 Montpellier, France; pierre.mournet@cirad.fr

<sup>5</sup> Amélioration Génétique et Adoption des Plants Méditerranéennes et Tropical AGAP, Université de Montpellier, 34398 Montpellier, France

<sup>6</sup> International Trade Centre (ITC), Addison House International Trade Fair Center, FAGE, Accra GA145, Ghana; mijuamarel@gmail.com

<sup>7</sup> Agriculture and Agri-Food Canada, Fredericton, NB 20280, Canada; david.dekoeber@aaagr.gc.ca (D.D.K.)

\* Correspondence: A.Amele@cgiar.org; Tel.: +234-8039-784-287

**Citation:** Agre, P.A.; Darkwa, K.; Olasanmi, B.; Kolade, O.; Mournet, P.; Bhattacharjee, R.; Lopez-Montes, A.; De Koeber, D.; Adebola, P.; Kumar, L.; Asiedu, R.; Asfaw, A. Identification of QTLs Controlling Resistance to Anthracnose Disease in Water Yam (*Dioscorea alata*). *Genes* **2022**, *13*, 347. <https://doi.org/10.3390/genes13020347>

Academic Editor: Anna M. Mastrangelo

Received: 17 January 2022

Accepted: 11 February 2022

Published: 14 February 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** Anthracnose disease caused by a fungus *Colletotrichum gloeosporioides* is the primary cause of yield loss in water yam (*Dioscorea alata*), the widely cultivated species of yam. Resistance to yam anthracnose disease (YAD) is a prime target in breeding initiatives to develop durable-resistant cultivars for sustainable management of the disease in water yam cultivation. This study aimed at tagging quantitative trait loci (QTL) for anthracnose disease resistance in a bi-parental mapping population of *D. alata*. Parent genotypes and their recombinant progenies were genotyped using the Genotyping by Sequencing (GBS) platform and phenotyped in two crop cycles for two years. A high-density genetic linkage map was built with 3184 polymorphic Single Nucleotide Polymorphism (NSP) markers well distributed across the genome, covering 1460.94 cM total length. On average, 163 SNP markers were mapped per chromosome with 0.58 genetic distances between SNPs. Four QTL regions related to yam anthracnose disease resistance were identified on three chromosomes. The proportion of phenotypic variance explained by these QTLs ranged from 29.54 to 39.40%. The QTL regions identified showed genes that code for known plant defense responses such as GDSL-like Lipase/Acylhydrolase, Protein kinase domain, and F-box protein. The results from the present study provide valuable insight into the genetic architecture of anthracnose resistance in water yam. The candidate markers identified herewith form a relevant resource to apply marker-assisted selection as an alternative to a conventional labor-intensive screening for anthracnose resistance in water yam.

**Keywords:** *Dioscorea* spp.; greater yam; genetic map; marker–trait association; linkage analysis

## 1. Introduction

Yam (*Dioscorea* spp.) is a multi-species monocotyledonous crop widely grown in the tropics and subtropics [1]. It is the most valuable crop in West Africa, where its cultivation began 11,000 years ago [2]. Of the over 600 yam species, water yam (*D. alata*) is an extensively cultivated species worldwide [3]. In Africa, white Guinea yam (*D. rotundata*) is the most cultivated yam species, followed by water yam [3]. Yam production in West Africa is mainly by smallholder farmers, making it a significant source of farm employment and

income for this group. In addition, yam plays a vital role in traditional medicine and the socio-cultural life of the people as it is involved in many key life ceremonies [4].

Water yam possesses several valuable attributes for cultivation and consumption. These include high multiplication ratio, early vigor for weed smothering, the higher genetic potential for yield (especially under low to average soil fertility), low post-harvest losses, good processing quality, and high nutritional value, including its possession of low glycemic index [5,6]. However, anthracnose disease caused by the *Colletotrichum gloeosporioides* (Penz) is the most limiting factor affecting the productivity of water yam by devastating all parts of the yam plant at every developmental stage, including leaves, stems, tubers, and seeds in many regions of the world [7,8]. Anthracnose causes mild to acute leaf necrosis, premature leaf abscission, and shoot die-back [9]. Severe infections result in defoliation, leaving naked, black, and drying vines [10]. Yield losses from the disease of up to 90% have been reported under severe conditions on different cultivars of water yam in Africa [10–12]. High genetic and pathogenic variances have been reported among isolates of *C. gloeosporioides* from different geographical locations [7,13,14], suggesting a high probability of the geographic variation in strains, some of which could be overcome existing resistance [15].

Cultural control approaches such as the use of disease-free planting materials, adjustment of plant spacing and planting dates, burying infected plant residues in the soil immediately after harvesting, intercropping, crop rotation with non-host crops, and fallowing have been used in other plant pathosystems to reduce pathogen inoculum in the field, delay disease onset, or slow disease progress [16,17]. Nonetheless, these disease management practices have not been effective for controlling anthracnose disease in water yam or result in a substantial increase in tuber yield [8], especially in disease-endemic areas. Additionally, biological control to impede or out-compete the multiplication and spread of virulent *C. gloeosporioides* strain in yam fields has been limited [18]. Chemical control can be an effective disease management approach. Still, most yam producers are smallholder growers and may not have the prerequisite technical support and finance to afford the use of fungicides [19].

Furthermore, inappropriate use of fungicides could potentially result in the development of resistant *C. gloeosporioides* strains to systemic fungicides [20] as well as detrimental environmental effects. Therefore, the best control option is developing and deploying cultivars with durable resistance to anthracnose. Substantial progress has been made to develop anthracnose resistant water yam varieties at the International Institute of Tropical Agriculture (IITA), Nigeria, and national agricultural research systems in West Africa and elsewhere through conventional breeding using phenotypic observations [3,21]. Anthracnose-resistant cultivars of yam such as TDa1425 and TDr2040 were identified at IITA [22]. In India, laboratory and field investigations also found highly resistant *D. alata* lines [23]. However, this effort is arduous and considerably slow due to the crop's heterozygous and vegetatively propagated nature [24]. Genomics-informed breeding techniques such as molecular marker-assisted breeding and genomic selection would accelerate efforts in introgressing anthracnose resistance into preferred genetic backgrounds [3].

Earlier investigations on anthracnose disease in water yam showed that resistance is likely dominant and quantitatively inherited [25]. Efforts have also been made to identify QTL controlling yam anthracnose disease (YAD) using low-throughput molecular markers and less dense or unsaturated genetic maps such as Amplified fragment length polymorphism (AFLP) markers [5,26] and Expressed Sequence Tag – Simple Sequence Repeats (EST-SSRs) [27–29]. Prospects for locating additional QTLs and applying molecular breeding methods in water yam improvement programs are up-and-coming, mainly due to advances in next-generation sequencing and the recent development of the reference genome sequence of *D. rotundata* and *D. alata*. Therefore, it is imperative to apply new molecular tools to develop additional genomic resources from different genetic backgrounds to facilitate the breeding for anthracnose resistance in water yam. The objective of this study was to develop a SNP-based genetic linkage map and identify QTL for anthracnose

disease resistance in a diploid bi-parental mapping population of *D. alata*. It assessed the QTL presence, positions, the effects of QTL alleles, and the underlying putative genes in the QTL vicinities responsible for anthracnose resistance in water yam.

## 2. Materials and Methods

### 2.1. Plant Materials

An F<sub>1</sub> recombinant clonal population of 204 individuals derived from a single cross of TDa0500015 × TDa9900048 was used for this study. TDa0500015 (diploid) is a female breeding line showing a tolerant reaction to yam anthracnose disease, while TDa9900048 (diploid) is a male breeding line expressing a susceptible response. The two parents and their F<sub>1</sub>-derived recombinant clonal progenies, along with a highly susceptible cultivar (TDa92-2), were field-phenotyped in two cropping cycles for two seasons (2017 and 2018) at IITA, Ibadan research farm in Nigeria. The field experiment was carried out using a partial replicated design of three plants per genotype in 1 × 1 m planting spacing during the main rainy seasons when anthracnose incidence and severity are high. Genotypes with high plant numbers were planted in 2 replications, and susceptible reference cultivar TDa92-2 was planted as a spreader row between blocks and around the field.

### 2.2. Phenotyping

Anthracnose disease severity was scored at two months after planting and after that, fortnightly till six months. Severity was scored by visual assessment of the relative area of plant tissue affected by anthracnose using a 1–5 severity rating scale. Where, 1 = No visible symptoms of anthracnose disease or infection spot on the leaf surface; 2 = Few anthracnose spots or symptoms on 1–25% of the plant (i.e., one or two spots of less than 1 cm diameter width, and dry tissue on the leaf surface); 3 = Anthracnose symptoms covering 26–50% of the plant (i.e., one or two spots of more than 1 cm diameter width, and dry tissue on the leaf surface, small dark and no dried spots with more than 1 cm width are present); 4 = Symptoms on >50% of the plant (i.e., coalesced spots with dry tissue and covering a significant proportion of the leaf surface, areas with less than 1 cm width coalesce to more prominent spots and yellowing of green tissue is intense around the spots areas); and 5 = Severe necrosis and death of the plant (i.e., coalesced spots with dry tissue more than 1.5 cm in diameter and covering a significant proportion of the leaf surface and yellowing of the green tissue is generalized in the leaf blade) [30]. The time series severity score was recorded on individual plants in a plot. The mean anthracnose severity for a genotype in a plot was estimated by summing severity scores >1 in a plot divided by the total number of symptomatic plants.

The area under the disease progression curve (AUDPC) was estimated from the mean disease severity scores of a genotype per plot using the trapezoidal method [31]. This method discretizes the time variable and calculates the average disease intensity between each pair of adjacent time points.

$$AUDPC = \sum_{i=1}^n \left( \frac{y_i + y_{i+1}}{2} \right) (t_{i+1} - t_i) \quad (1)$$

where n = total number of observations, y<sub>i</sub> = disease severity at the t<sub>i</sub> observation, and t = time at the t<sub>i</sub> observation.

### 2.3. Genotyping

Young fresh leaf samples were collected from the 207 genotypes (204 recombinant progenies, the two parents and a check variety) and immediately dipped in dry ice. The leaves were stored at −80 °C before lyophilization. Lyophilized leaf samples were sent to CIRAD-France for DNA extraction, library construction, and Genotyping by Sequencing

(GBS). DNA extraction and Genotyping by Sequencing (GBS) were performed as described in Cormier et al. [32]. GBS libraries were constructed as described by Elshire et al. [33] using PstI-MseI restriction enzymes. Sequencing was conducted on an Illumina HiSeq 3000 system Montpellier, France (150 bp, single-end reads) at the GeT-PlaGe platform in Toulouse, France.

#### 2.4. Data Analyses

##### 2.4.1. Phenotype Data

Anthraxnose severity score data collected at different times during the crop's growth period were converted to AUDPC for quantitative comparison over the years. The area under disease progress curve data was subjected to mixed model analysis using lme4 library package implemented in R [34].

$$Y_{ijk} = \mu + \beta_i + R_{ij} + G_k + (\beta_i \times G_k) + e_{ijkm} \quad (2)$$

where  $Y_{ijk}$  = phenotypic value,  $\mu$  = overall phenotypic mean,  $\beta_i$  = effect of year  $i$ ,  $R_{ij}$  = effect of block  $j$  in year  $i$ ,  $G_k$  = effect of genotype  $k$ ,  $(\beta_i \times G_k)$  = effect of interaction between year  $i$  and genotype  $k$ , and  $e_{ijkm}$  = residual. Block within-year effects were added to the model as a random variable to remove the spatial variation within the trial field. Broad sense heritability was estimated from the model to assess the proportion of phenotypic variation in the data set due to genetic effects. Phenotypic BLUE (Best Linear Unbiased Estimator) values of un-shrunken means for QTL analysis were extracted for the years and over the years.

##### 2.4.2. SNP Calling and Quality Assessment

Raw data were first filtered using a pipeline described in Scarcelli et al. [35]. Demuladapt (<https://github.com/Maillol/demuladapt> accessed on 12 March 2020) was used for demultiplexing. Cutadapt 1.2.1 [36] was used to remove the adaptors and low-quality bases read with a mean quality score <30 using a free perl script [https://github.com/SouthGreenPlatform/arcad-hts/blob/master/scripts/arcad\\_hts\\_2\\_Filter\\_Fastq\\_On\\_Mean\\_Quality.pl](https://github.com/SouthGreenPlatform/arcad-hts/blob/master/scripts/arcad_hts_2_Filter_Fastq_On_Mean_Quality.pl) accessed on 11 March 2020. For the final SNP calling, GATK was used while mapping was performed using default options of Burrows-Wheeler Aligner (BWA) [37] using the *D. alata* reference genome. The SNP quality assessment was performed using vcftools [38] and plink [39]. SNPs with low MAF <0.05 and low depth sequencing <5 were removed. This retained 7583 SNPs out of the raw 15,936 SNPs called. For the missing point, SNP markers and genotypes with high missing information >20% were removed as well.

##### 2.4.3. Genetic Map Construction

Linkage analysis was conducted using MAPpoly package [40] in the R environment [34]. A series of filtering steps were applied using all segregating markers polymorphic in at least one of the parents to construct an integrated genetic map. Chi-square ( $\chi^2$ ) test was conducted to calculate the marker segregation ratio and exclude markers showing significant segregation distortion from map construction. The  $p$ -value threshold to assess the significant marker segregation distortion was set using the filter\_segregation function as implemented in MAPpoly package. To construct linkage groups, the pairwise recombination fraction and LOD matrices between markers retained after the segregation test were calculated using the function est\_pairwise\_rf in MAPpoly package. Linkage grouping was then performed using an initial LOD value of >6 obtained from  $\alpha$ thres function in MAPpoly. The LOD value of 6.0 that established known linkage groups was then chosen as the significance criterion for multipoint linkage testing. First, for the genetic map construction, marker loci were partitioned primarily into linkage groups (LGs). Secondly, the modified logarithm of odds (MLOD) scores between markers were calculated to further confirm the robustness of markers for each LGs. Markers with MLOD scores <6 were filtered

out prior to ordering. Thirdly, the highMap strategy described by Liu et al. [41] was utilized to order the Bin markers and correct genotyping errors within and between LGs. Genetic recombination fraction (RF) was estimated for the retained SNP markers to confirm the non-switch of alleles from one LG to another using “est.rf” function implemented in R/QTL [42]. To confirm the well ordering SNP markers across LG, the recombination fraction against the LOD score was then plotted and the graph was viewed using ggplot2 R package. The final GM was then constructed using R/QTL2 [42] and viewed in Linkage-MapView.

#### 2.4.4. QTL Analysis

The QTL analysis was performed with mean trait value over years and linkage map constructed from the 159 recombinant clones using the Composite Interval Mapping (CIM) method in R/QTL2 package [42]. A forward and backward simple stepwise regression was run to select background markers with a significant level of  $P < 0.05$ . The threshold levels to declare significant QTLs were empirically determined through 1000 permutations of the data, which maintained a chromosome-wise Type I error rate of 0.05 [43] with a fixed LOD of 4 as a minimum threshold of declaring a SNP marker linked with the YAD.

The location of a QTL was described according to its LOD peak location. The proportion of phenotypic variance accounted for by each detected QTL was estimated by a single-factor analysis of variance using the General Linear Model. The additive (Add) and dominance (Dom) effects and the proportion of phenotypic variation explained (PVE%) by each QTL were estimated using the “fitqtl” function in R/QTL. The sign of the additive effect of each QTL was used to identify the origin of the favorable alleles. A simple mixed model was implemented in lmer4 package to estimate QTL interaction/environment using the identified QTL by considering the year and the SNP marker as fixed effect, while the genotypes were considered as a random effect. Markers linked with the yam anthracnose disease were then placed in the respective chromosome, and their position was viewed using Qtl/jittermap. For the gene mining, the related putative genes associated with SNP markers were searched within the upstream and downstream locations of the QTL generic feature format (GFF3) of the reference genome of *D. alata*. Functions of the different genes associated with the identified QTL were determined using the public database Interpro, European Molecular Biology Laboratory-European Bioinformatics Institute (EMBL-EBI).

### 3. Results

#### 3.1. Phenotypic Variability

Significant differences ( $p < 0.05$ ) were observed for the reaction of the progenies to YAD in both years (Table 1). The mean squares for the year and genotype-by-year interaction effects were highly significant ( $p < 0.01$ ). The disease pressure was higher in 2018 compared to 2017. The area under disease progression curve (AUDPC) estimates ranged from 210.0 to 397.5 with an average of 245.5 in 2017, while the range was from 233.4 to 482.1 with an average of 299.8 in 2018. None of the recombinant clones demonstrated immune or highly resistant (mean severity score of 1, equivalent to AUDPC value <105) or highly susceptible (mean severity score of 5, the equivalent of AUDPC > 525) reaction to natural field infestation by anthracnose disease. However, most of the genotypes (67–92%) expressed moderate resistance to anthracnose. Broad sense heritability was high (70.64%).

**Table 1.** Mean squares and heritability estimate for yam anthracnose disease severity in the mapping population.

| Trait | Mean Squares |            |                 | CV (%) | Broad Sense Heritability (%) |
|-------|--------------|------------|-----------------|--------|------------------------------|
|       | Genotype     | Year       | Genotype × Year |        |                              |
| AUDPC | 163.01 *     | 2190.01 ** | 3371.8 ***      | 17.6   | 70.64                        |


AUDPC: area under disease progression curve; \*, \*\*, \*\*\* significance at 0.05, 0.01, and 0.001 *p*-values, respectively; CV: coefficient of variation.

### 3.2. SNP Filtering

Total of 15,936 SNP markers were identified in the parental individuals and mapping population. Filtering for minor allele frequency (MAF < 0.05), low depth sequencing (< 5), and polymorphism between the parents TDa0500015 and TDa9900048 reduced the number of SNPs to 7,583 SNPs (47.6% of the raw SNPs identified). Further filtering for 20% missing data (both for the SNP and genotype) (Figure S1) and segregation distortion with chi-square test (Figure S2) identified 3257 informative markers and 179 individuals out of 204 progenies with good coverage for linkage map construction. Pairwise recombination fractions calculated for all informative markers showed high SNP markers ordering across the different linkage groups (Figure S3).

### 3.3. Linkage Mapping

A genetic map was constructed that covered all 20 linkage groups of the water yam genome (Figure 1) with a total genetic distance of 1460.94 cM. The marker order on the linkage map showed perfect genetic recombination (Figure S3) as the recombination fraction of the mapped SNP markers on linkage groups displayed a perfect alignment with no half circles between the recombination fraction and the LOD score (Figure S4). The linkage map had an average of 163 markers per linkage group or chromosome, with the highest SNP markers mapped on linkage 5. Linkage groups 6, 5, and 2 were the longest with 109.52, 109.19, and 109.17 cM, respectively, while linkage group 19 was the shortest with 33.08 cM (Table 2). The genetic map intervals ranged from 2.03 and 9.07 cM on chromosomes 19 and 16, respectively (Table 2).



**Figure 1.** Genetic linkage map showing the SNP distribution across the 20 yam chromosomes. Each vertical line represents a yam chromosome with genetic distance in Kosambi units (cM).

**Table 2.** Distribution of SNP markers and linkage group length (cM) in each chromosome.

| Chromosomes | Number of SNPs | Chromosome Length (cM) | Average SNP Distance | Maximum Gap |
|-------------|----------------|------------------------|----------------------|-------------|
| Chr1        | 80             | 80.93                  | 1.95                 | 5.00        |
| Chr2        | 84             | 109.17                 | 1.28                 | 6.01        |
| Chr3        | 115            | 64.63                  | 0.56                 | 4.48        |
| Chr4        | 520            | 92.32                  | 0.16                 | 8.26        |
| Chr5        | 199            | 109.19                 | 0.50                 | 7.71        |
| Chr6        | 191            | 109.52                 | 0.57                 | 5.54        |
| Chr7        | 127            | 59.84                  | 0.48                 | 5.26        |
| Chr8        | 200            | 91.39                  | 0.48                 | 4.44        |
| Chr9        | 124            | 77.12                  | 0.62                 | 5.80        |
| Chr10       | 104            | 57.13                  | 0.55                 | 5.52        |
| Chr11       | 85             | 55.95                  | 0.66                 | 7.94        |
| Chr12       | 125            | 83.22                  | 0.65                 | 4.30        |
| Chr13       | 116            | 95.55                  | 0.78                 | 5.47        |

|              |             |                 |      |      |
|--------------|-------------|-----------------|------|------|
| Chr14        | 208         | 51.447          | 0.25 | 4.40 |
| Chr15        | 180         | 34.70           | 0.18 | 2.39 |
| Chr16        | 139         | 77.16           | 0.61 | 9.07 |
| Chr17        | 208         | 67.44           | 0.31 | 3.95 |
| Chr18        | 129         | 75.71           | 0.54 | 7.23 |
| Chr19        | 111         | 33.09           | 0.28 | 2.03 |
| Chr20        | 139         | 35.48           | 0.23 | 3.41 |
| <b>Total</b> | <b>3184</b> | <b>1,460.98</b> |      |      |

### 3.4. QTL Identification


The QTLs detected on three chromosomes out of the 20 are presented in Table 3 and Figure 2. Two significant QTLs were detected on chromosome 7 at position 10.60 cM (between 10.596 and 19.217 cM). This QTL (Qyad-7-1) had a LOD score of 4.51 and accounted for 33.7% of the total phenotypic variation in anthracnose severity score (Table 3, Figure 2). The second QTL located on chromosome 7 (Qyad-7-2) was at position 19.21 cM (between 10.596 and 19.218 cM) at LOD score of 5.28 and accounted for 29.54% of the total phenotypic variation in anthracnose severity score. The 3rd significant QTL, Qyad-15, which explained 30.90% of the total phenotypic variance with a LOD score of 4.43 was detected at 28.80 cM on chromosome 15. The QTL on chromosome 18 (Qyad-18) was at position 61.4 cM (between 61.345 and 61.432 cM) at LOD score of 4.65 and contributed 39.40% of the total phenotypic variance. For the four markers associated with the YAD, the favorable alleles were contributed by TDa0500015 tolerant to the YAD.

**Table 3.** Summary of significant QTLs detected for yam anthracnose disease resistance in water yam.

| Markers  | Chr | Pos (cM) | LOD  | Add/Dom | CI. Low | CI. High | R <sup>2</sup> (%) | Putative Genes                  |
|----------|-----|----------|------|---------|---------|----------|--------------------|---------------------------------|
| Qyad-7-1 | 7   | 10.60    | 4.51 | -2.56   | 10.596  | 19.217   | 33.7               | DRNTG_08663.1                   |
| QTL-7-2  | 7   | 19.21    | 5.28 | -5.98   | 10.596  | 19.218   | 29.54              | DRNTG_08664.1,<br>DRNTG_23336.1 |
| Qyad-15  | 15  | 28.80    | 4.43 | -10.12  | 10.171  | 28.817   | 30.90              | DRNTG_14305.1                   |
| Qyad-18  | 18  | 61.4     | 4.65 | -3.48   | 61.345  | 61.432   | 39.40              | DRNTG_18245.1,<br>DRNTG_29617.1 |

Chr: chromosome; pos: position; LOD: logarithm of odds score; CI: confidence interval; R<sup>2</sup>: % variation explained; Add: additive; Dom: dominance.





**Figure 2.** Genetic map of water yam showing significant QTLs associated with yam anthracnose disease resistance. Only those chromosomes where significant QTL is located are displayed. The identified QTLs are highlighted in blue on each chromosome.

The QTL region linked to YAD resistance on chromosome 7 has known genes in plant biotic stress such as DRNTG\_08663.1 (GDSL-like Lipase/Acylhydrolase), DRNTG\_08664.1 (Protein kinase domain), and DRNTG\_23336.1 (Table S1). Additionally, the regions within the Qyad-15 locus were related to the N-terminal  $\alpha/\beta$  domain gene DRNTG\_14305.1. The vicinity of Qyad-18 showed genes that code for ANTH domain Putative clathrin assembly protein (DRNTG\_18245.1) and WD domain-WD40 repeat-containing protein (DRNTG\_29617.1) (Table S1).

Interaction among the four QTLs related to YAD resistance revealed significant ( $p < 0.05$ ) QTL by QTL interaction for Qyad-7-1 and Qyad-15, Qyad7-2 and Qyad-18. In contrast, no significant variation was observed among all other QTL combinations (Table 4).

Of the four QTLs related to YAD resistance, three were stable over the years and showed non-significant QTL by environment interaction (Table 5).

**Table 4.** Interactions among the detected QTL.

| Marker Interactions                 | df | MS     | p-Value  | Adjusted Squared | R- |
|-------------------------------------|----|--------|----------|------------------|----|
| Qyad-7-1: QTL-7-2                   | 1  | 55.9   | 0.835    | 0.04147          |    |
| Qyad-7-1: Qyad-15                   | 1  | 5303.5 | 0.0456 * | 0.02544          |    |
| Qyad7-1: Qyad-18                    | 1  | 155.2  | 0.734    | -0.0002131       |    |
| QTL-7-2: Qyad-15                    | 1  | 2580.7 | 0.158    | 0.04395          |    |
| Qyad-7-2: Qyad-18                   | 1  | 6341.0 | 0.026 *  | 0.06074          |    |
| Qyad-15: Qyad-18                    | 1  | 1408.4 | 0.309    | -0.01079         |    |
| Qyad-7-1: QTL-7-2: Qyad-15: Qyad-18 | 3  | 1247.7 | 0.068    | 0.04413          |    |

df: degree of freedom; MS: mean square; \* statistical significance at p-value 0.05.


**Table 5.** QTL by environment analysis considering the major QTL.

|                 | Sum Sq | Mean Sq | F Value    |
|-----------------|--------|---------|------------|
| Year            | 174125 | 87062   | 370.9525   |
| Qyad-7-1        | 12.25  | 12.25   | 0.0002 *** |
| QTL-7-2         | 101    | 101     | 0.0003 *** |
| Qyad-15         | 129    | 129     | 0.0001 *** |
| Qyad-18         | 2.6    | 2.6     | 0.0002 *** |
| Year × Qyad-7-2 | 275    | 112     | 0.789ns    |
| Year × QTL-7-2  | 342    | 78      | 0.02 *     |
| Year × Qyad-15  | 278    | 110     | 0.226ns    |
| Year × Qyad-18  | 178    | 89      | 0.567ns    |


MS: mean square; TPVE: total phenotypic variation estimation; ns: non-significant; \*, \*\*\* statistical significance at p values 0.05 and 0.001, respectively.

### 3.5. Marker Effect


The presence of allele T for loci Qyad-7-1 and Qyad-7-2 appeared to lower the AUDPC score in the evaluated population, while the presence of the alleles C tended to increase the disease score, especially with Qyad-7-2 with  $p$ -value = 0.03 (Figures 3 and 4). For QTLs detected on chromosomes 15 and 18, allele A of the variant AG/GA was associated with a higher AUDPC value while allele G linked with the lower AUDPC value in the population (Figures 5 and 6).




**Figure 3.** The boxplot showing the effect of the different alleles (variants) of Qyad-7-1 on the AUDPC values. The letters on the X-axis represent alleles (CC, CT, and TT).



**Figure 4.** The boxplot displaying the effect of the different alleles (variants) of Qyad-7-2 on the AUDPC estimates. The letters on the X-axis represent alleles (CC, CT, and TT).



**Figure 5.** Comparison of the effects of the different alleles (variants) of Qyad-15 on the AUDPC estimates in the study population. The letters on the X-axis represent alleles (AA, AG, and GG), \*\*\* statistical significance at p values 0.0001 while ns is non-significant



**Figure 6.** Comparisons of allelic effects of the QTL Qyad-18 on AUDPC estimates in the study population. The letters on the X-axis represent alleles (AG and GG), \*\* and \*\*\* are statistical significance at p values 0.05 and 0.001

#### 4. Discussion

This study selected two parents based on their responses to yam anthracnose disease and created their F<sub>1</sub>-derived recombinant clonal population to assess the functional association of anthracnose resistance and genetic markers using the QTL mapping approach. The recombinant clonal population showed a differential response to the disease-causing organism over the two-year evaluation period. The recombinant population showed quantitative tolerance with a continuous distribution from resistance to the susceptible range with substantial skewness towards resistance. However, no immune or highly resistant clones were identified. Instead, a large number of the clones expressed tolerance reaction to YAD field infestation. The heritability estimate in the present study was high, indicating the proportion of phenotypic variance that is genetic. Similarly, Petro et al. [26] and Bhattacharjee et al. [29] reported high heritability estimates for YAD in water yam.

In an earlier effort, Cormier et al. [32] constructed a high-density genetic map of *D. alata* using 1,579 polymorphic SNP markers with a consensus map length of 2613.5 cM. However, our genetic linkage map was built using 3184 SNPs from the GBS platform that spanned a total length of 1460.94 cM representing the most saturated and accurate genetic map for *D. alata* to date. Genetic linkage maps of water yam were also developed using EST-SSRs [29] and AFLPs [5,26]. The genetic linkage map presented in this report will offer a unique opportunity for qualitative and quantitative trait analysis in water yam.

Three studies have been conducted to map QTLs controlling resistance to anthracnose in water yam [5,26,29]. The study by Mignouna et al. [5] and Petro et al. [26] utilized AFLP maps and identified one and nine QTLs, respectively, for anthracnose resistance, explaining 10% and 26–74% of the total phenotypic variation. Bhattacharjee et al. [29] utilized an EST-SSR genetic map for their study and identified a major QTL on linkage group 14, explaining 69% of the total phenotypic variance. Even though the previous studies ordered markers on 20 linkage groups, the absence of a standard genetic map and the different marker systems makes it difficult to compare the location of the detected QTLs in these studies. In the present study, four QTLs located on three chromosomes, accounting for 29.54–39.4% of the total phenotypic variation in the trait, were identified. QTL interaction over the years revealed the stability of three QTL and indicated their potential to predict the specific effect of variation in strain or intensity of strains of *C. gloeosporioides* infestation over the years during the field experimentation.

Furthermore, gene annotation in the QTL regions related to YAD resistance showed known genes that code for plant defense mechanisms. Notably, the region composed of Qyad-7-1 shows the GDSL-like Lipase/Acylhydrolase gene that is reported to regulate systemic resistance to *Alternaria brassicicola* in Arabidopsis [44,45]. Hong et al. [46] also found this gene involved in the defense against drought and *Xanthomonas campestris* pv. Vesicatoria in pepper. Additionally, the protein kinase domain is involved in regulating the resistance against bacterial blight (*Xanthomonas oryzae*) in rice [47] and resistance to the necrotrophic fungal pathogen *Plectosphaerella cucumerina* in Arabidopsis [48] is also present within the QTL region related to YAD in our study. Moreover, the QTL loci on chromosomes 15 and 18 showed the ANTH domain associated with defense against *Pseudomonas syringae* in *Nicotiana benthamiana* and Arabidopsis [49], and the WD domain enhanced the resistance to anthracnose leaf blights in maize caused by *Colletotrichum sublineolum* [50,51]. The F-box protein found within the QTL region of chromosome 15 was reported to be involved in cell death and defense response during the pathogen recognition of *Pseudomonas syringae* and Tobacco mosaic virus in tomato and tobacco [52]. The N-terminal domain within the flanking sequence of the QTL region was involved in the resistance to the downy mildew pathogen *Hyaloperonospora arabidopsidis* in Arabidopsis [53]. Therefore, enough evidence exists to confirm that the genes within the flanks of the significant QTLs for anthracnose disease resistance discovered in this study code for response to plant biotic stress.

## 5. Conclusions

This study developed a highly saturated and accurate genetic linkage map for water yam. The linkage mapping approach used in this study identified and mapped QTLs linked to yam anthracnose disease. The QTL regions identified in this study showed six already known genes involved in plant defense. Our results are valuable tools for developing water yam cultivars with quantitative resistance to anthracnose disease. However, these QTLs need to be validated in different genetic backgrounds and environments to be routinely applied in marker-assisted selection in water yam breeding programs in Africa.

**Supplementary Materials:** The following supporting information can be downloaded at: [www.mdpi.com/article/10.3390/genes13020347/s1](http://www.mdpi.com/article/10.3390/genes13020347/s1), Table S1: List of the putative genes identified in the vicinity of the QTL associated with yam anthracnose disease; Figure S1: SNP quality assessment for missing markers and genotypes (missing threshold = 20%) showing informative markers (the left figure) and genotypes with good coverage (right figure) for linkage map construction. The blue horizontal dotted lines represent the threshold; Figure S2: Evaluation of polymorphic SNP markers in the biparental population for segregation distortion using chi-square test; Figure S3: Recombination fraction of genotyped GBS markers. A pairwise comparison of recombination fractions after SNP markers were ordered across the 20 chromosomes. Recombination fractions are in the upper left triangle, the LOD scores are in the lower right triangle, and the red corresponds to the large LOD or a small recombination fraction; Figure S4: Plot of recombination fraction against the LOD score showing perfect alignment as no half circles were observed.

**Author Contributions:** A.A., A.L.-M., R.A., B.O., and D.D.K. designed the project; A.A. and A.L.-M. developed the mapping population; K.D., P.A.A. and O.K. executed the field phenotyping with contribution from A.A.; P.A.A. and A.A. performed phenotypic analysis; P.M. managed the genotyping for the sequencing; P.A.A. managed the genotypic data and performed the analysis; P.A.A. and K.D. wrote the first draft with contributions from A.A. and B.O., R.A.; L.K., R.A. and P.A. reviewed and edited the manuscript; A.A.; R.B. supervised the study. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the Bill and Melinda Gates Foundation (BMGF) through the AfricaYam project (OPP1052998). The African Union Commission funded Kwabena Darkwa at the Pan African University, Institute of Life and Earth Sciences (PAULESI), University of Ibadan.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Data available at <https://yambase.org/breeders/trial/1047> and <https://figshare.com/account/home>.

**Acknowledgments:** Special thanks to the yam breeding team at IITA for their contribution to the mapping population development and field phenotyping. The yam research at IITA is supported by the Bill and Melinda Gates Foundation (BMGF) and CGIAR Research Program on Roots, Tubers, and Bananas (RTB).

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Asiedu, R.; Sartie, A. Crops that feed the world 1. Yams. *Food Secur.* **2010**, *2*, 305–315.
2. Scarcelli, N.; Cubry, P.; Akakpo, R.; Thuillet, A.C.; Obidiegwu, J.; Baco, M.N.; Otoo, E.; Sonké, B.; Dansi, A.; Djedatin, G.; et al. Yam genomics supports West Africa as a major cradle of crop domestication. *Sci. Adv.* **2019**, *5*, 1–8.
3. Darkwa, K.; Olasanmi, D.; Asiedu, R.; Asfaw, A. Review of empirical and emerging breeding methods and tools for yam (*Dioscorea* spp.) improvement: Status and prospects. *Plant Breed.* **2020**, *139*, 474–497.
4. Obidiegwu, J.E.; Akpabio, E.M. The geography of yam cultivation in southern Nigeria: Exploring its social meanings and cultural functions. *J. Ethn. Foods* **2017**, *4*, 28–35.
5. Mignouna, H.; Mank, R.; Ellis, T.; Van den Bosch, N.; Asiedu, R.; Abang, M. and Peleman, J. A genetic linkage map of water yam (*Dioscorea alata* L.) based on AFLP markers and QTL analysis for anthracnose resistance. *Theor. Appl. Genet.* **2002**, *105*, 726–735.
6. Baah, F.D.; Maziya-Dixon, B.; Asiedu, R.; Oduro, I.; Ellis, W.O. Nutritional and biochemical composition of *D. alata* (*Dioscorea* spp.) tubers. *J. Food Agric. Environ.* **2009**, *7*, 373–378.

7. Abang, M.M.; Fagbola, O.; Smalla, K.; Winter, S. Two genetically distinct populations of *Colletotrichum gloeosporioides* Penz. causing anthracnose disease of yam (*Dioscorea* spp.). *J. Phytopathol.* **2005**, *153*, 137–142.
8. Ntui, V.O.; Uyoh, E.A.; Ita, E.E.; Markson, A.-A. A.; Tripathi, J.N.; Okon, N.I.; Akpan, M.O.; Phillip, J.O.; Brisibe, E.A.; Ene-Obong, E.; et al. Strategies to combat the problem of yam anthracnose disease: Status and prospects. *Mol. Plant Pathol.* **2021**, *22*, 1302–1314.
9. Abang, M.M.; Green, K.R.; Wanyera, N.W.; Iloba, C. Characterization of *Colletotrichum gloeosporioides* Penz. from yam (*Dioscorea* spp.) in Nigeria. In *Root Crops in the 21st Century, Proceedings of the 7th Triennial Symposium of the International Society for Tropical Root Crops—Africa Branch, Cotonou, Bénin, 11–17 October 1998*; ISTRC, Acra, pp. 613–615.
10. Akem, C.N. Yam die-back and its principal cause in the yam belt of Nigeria. *Pak. J. Biol. Sci.* **1999**, *2*, 1106–1109.
11. Appiah-Kubi, Z.; Apetorgbor, A.K.; Moses, E.; Appiah-Kubi, D. Farmers Knowledge of Anthracnose Disease of Cassava and Yam in Four Ecological Zones in Ghana. *Greener J. Agric. Sci.* **2015**, *5*, 204–209.
12. Nwadi, C.O.; Augusto, J.; Bhattacharjee, R.; Atehnkeng, J.; Lopez-Montes, A.; Onyeka, T.J.; Kumar, P.L.; Asiedu, R.; Bandyopadhyay, R. Comparative reliability of screening parameters for anthracnose resistance in water yam (*Dioscorea alata*). *Plant Dis.* **2017**, *101*, 209–216.
13. Frezal, L.; Jacqua, G.; Neema, C. Study of the genetic diversity of *Colletotrichum gloeosporioides*, causative agent of anthracnose of white yam in Guadeloupe: In Proceedings of the 7th International Conference on Plant Diseases, Tours, France, 3–5 December 2003.
14. Abang, M.M.; Hoffmann, P.; Winter, S.; Green, K.R.; Wolf, G.A. Vegetative compatibility among isolates of *Colletotrichum gloeosporioides* from yam (*Dioscorea* spp.) in Nigeria. *J. Phytopathol.* **2004**, *152*, 21–27.
15. McDonald, B.A.; Linde, C. The population genetics of plant pathogens and breeding strategies for durable resistance. *Euphytica* **2002**, *124*, 163–180.
16. Simons, S.A.; Green, K.R. Epidemiology of yam anthracnose: Sources of inoculum. In Proceedings of 4th International Conference of Plant Protection, Kuala Lumpur, Malaysia, 28–31 March 1994; pp. 67–69.
17. Reis, E.M.; Casa, R.T.; Blum, M.M.; dos Santos, H.P.; Medeiros, C.A. Effects of cultural practices on the severity of leaf blotches of wheat and their relationship to the incidence of pathogenic fungi in the harvested seed. *Fitopatol. Bras.* **1997**, *22*, 407–412.
18. Palaniyandi, S.A.; Yang, S.H. Cheng, J.H.; Meng, L.; Suh, J.W. Biological control of anthracnose (*Colletotrichum gloeosporioides*) in yam by *Streptomyces* sp. MJM5763. *J. Appl. Microbiol.* **2001**, *111*, 443–455.
19. Hepperly, P.; Vazquez, F. Resistance and scouting in the control of yam anthracnose of the winged yam (*Dioscorea alata*), In Proceedings of 25th Annual Meeting of the Caribbean Food Crops Society Gosier, Guadeloupe, France, 1–6 July 1989; pp. 587–596.
20. Bayart, J.D.; Pallas, B. Tolerance of yam anthracnose of a benzimidazole. Results of the first study conducted in Guadeloupe. *Phytoma* **1994**, *461*, 37–40.
21. Lebot, V.; Abraham, K.; Kaoh, J.; Rogers, C.; Molisalé, T. Development of anthracnose resistant hybrids of the greater yam (*Dioscorea alata* L.) and interspecific hybrids with *D. nummularia* Lam. *Genet. Resour. Crop. Evol.* **2019**, *66*, 871–883.
22. Popoola, A.R.; Adedibu, B.O.; Ganiyu, S.A. Rapid assessment of resistance to tissue-cultured water yam (*Dioscorea alata*) and white yam (*Dioscorea rotundata*) to anthracnose (*Colletotrichum gloeosporioides* Penz.). *Arch. Phytopathol. Plant Prot.* **2013**, *46*, 663–669.
23. Arya, R.S.; Sheela, M.N.; Jeeva, M.L.; Abhilash, P.V. Identification of host plant resistance to anthracnose in greater yam (*Dioscorea alata* L.). *Int. J. Curr. Microbiol. Appl. Sci.* **2019**, *8*, 1690–1696.
24. Asiedu, R.; Ng, N.G.; Bai, K.V.; Ekanayake, I.J.; Wanyera, N.M. Genetic improvement. In *Food Yams: Advances in Research*; IITA and NCRI: Ibadan, Nigeria, 1998; pp. 63–104.
25. Mignouna, H.D.; Abang, M.M.; Green, K.R.; Asiedu, R. Inheritance of resistance in water yam (*Dioscorea alata*) to anthracnose (*Colletotrichum gloeosporioides*). *Theor. Appl. Genet.* **2001**, *103*, 52–55.
26. Petro, D.; Onyeka, T.J.; Etienne, S.; Rubens, S. An intraspecific genetic map of water yam (*Dioscorea alata* L.) based on AFLP markers and QTL analysis for anthracnose resistance. *Euphytica* **2001**, *179*, 405–416.
27. Narina, S.S.; Buyyarapu, R.; Kottapalli, K.R.; Sartie, A.M.; Ali, M.I.; Robert, A.; Hodeba, M.J.D.; Sayre, B.; Scheffler, B.E. Generation and analysis of expressed sequence tags (ESTs) for marker development in yam (*Dioscorea alata* L.). *BMC Genom.* **2011**, *12*, 100.
28. Sasaki, C.A.; Bhattacharjee, R.; Scheffler, B.E.; Asiedu, R. Genomic resources for water yam (*Dioscorea alata* L.): Analyses of EST-sequences, de novo sequencing and GBS libraries. *PLoS ONE* **2015**, *10*, e0134031.
29. Bhattacharjee, R.; Nwadi, C.O.; Sasaki, C.A.; Paterna, A.; Scheffler, B.E.; Augusto, J.; Lopez-Montes, A.; Onyeka, J.D.; Kumar, P.L.; Bandyopadhyay, R. An EST-SSR based genetic linkage map and identification of QTLs for anthracnose disease resistance in water yam (*Dioscorea alata* L.). *PLoS ONE* **2018**, *13*, e0197717.
30. Asfaw, A. Standard Operating Protocol for Yam Variety Performance Evaluation Trial; IITA: Ibadan, Nigeria, 2016; pp. 1–27.
31. Campbell, C.L.; Madden, L.V. Introduction to plant disease epidemiology, John Wiley and Sons: New York, NY, USA, 1990.
32. Cormier, F.; Lawac, F.; Maledon, E.; Gravillon, M.C.; Nudol, E.; Mournet, P.; Vignes, H.; Arnau, G. A reference high-density genetic map of greater yam (*Dioscorea alata* L.). *Theor. Appl. Genet.* **2019**, *132*, 1733–1744.

33. Elshire, R.J.; Glaubitz, J.C.; Sun, Q.; Poland, J.A.; Kawamoto, K.; Buckler, E.S.; Mitchell, S.E. A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PLoS ONE* **2011**, *6*, e19379.
34. R Core Team. *R: A Language and Environment for Statistical Computing*; R Foundation for Statistical Computing: Vienna, Austria, 2017. Available online: <https://www.R-project.org/> (accessed 22 June 2021).
35. Scarcelli, N.; Mariac, C.; Couvreur, T.L.; Faye, A.; Richard, D.; Sabot, F.; Berthouly-Salazar, C.; Vigouroux, Y. Intra-individual polymorphism in chloroplasts from NGS data: Where does it come from and how to handle it? *Mol. Ecol. Resour.* **2016**, *16*, 434–445.
36. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J.* **2011**, *17*, 10–12.
37. Sarah, G.; Homa, F.; Pointet, S.; Contreras, S.; Sabot, F.; Nabholz, B.; Santoni, S.; Sauné, L.; Ardisson, M.; Chantret, N.; et al. A large set of 26 new reference transcriptomes dedicated to comparative population genomics in crops and wild relatives. *Mol. Ecol. Resour.* **2017**, *17*, 565–580.
38. Danecek, P.; Auton, A.; Abecasis, G.; Albers, C.A.; Banks, E.; DePristo, M.A.; Handsaker, R.E.; Lunter, G.; Marth, G.T.; Sherry, S.T.; et al. The variant call format and VCFtools. *Bioinformatics* **2011**, *27*, 2156–2158.
39. Purcell, S.; Neale, B.; Todd-Brown, K.; Thomas, L.; Ferreira, M.A.; Bender, D.; Maller, J.; Sklar, P.; De Bakker, P.I.; Daly, M.J.; et al. PLINK: A tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* **2007**, *81*, 559–575.
40. Mollinari, M.; Olukolu, B.A.; Pereira, G.D.; Khan, A.; Gemenet, D.; Yencho, G.C.; Zen, Z.B. Unraveling the hexaploid sweet-potato inheritance using ultra-dense multilocus mapping. *G3-Genes Genom. Genet.* **2020**, *10*, 281–292.
41. Liu, D.; Ma, C.; Hong, W.; Huang, L.; Liu, M.; Liu, H.; Zheng, H. Construction and analysis of high-density linkage map using high-throughput sequencing data. *PLoS ONE*, **2014**, *9*, e98855.
42. Broman, K.W.; Gatti, D.M.; Simecek, P.; Furlotte, N.A.; Prins, P.; Sen, S.; Yandell, B.S.; Churchill, G.A. R/qtl2: Software for mapping quantitative trait loci with high-dimensional data and multiparent populations. *Genetics* **2019**, *211*, 495–502.
43. Churchill, G.A.; Doerge, R.W.; Empirical threshold values for quantitative trait mapping. *Genetics* **1994**, *138*, 963–971.
44. Ling, H.; Zhao, J.; Zuo, K.; Qiu, C.; Yao, H.; Qin, J.; Sun, X.; Tang, K. Isolation and expression analysis of a GDSL-like lipase gene from Brassica napus L. *J. Biochem. Mol. Biol.* **2006**, *39*, 297.
45. Kwon, S.J.; Jin, H.C.; Lee, S.; Nam, M.H.; Chung, J.H.; Kwon, S.S.I.; Ryu, C.M.; Park, O.K. GDSL lipase-like 1 regulates systemic resistance associated with ethylene signaling in Arabidopsis. *Plant J.* **2009**, *58*, 235–245.
46. Hong, J.K.; Choi, H.W.; Hwang, I.S.; Kim, D.S.; Kim, H.K.; Choi, D.S.; Kim, Y.J.; Hwang, B.K. Function of a novel GDSL-type pepper lipase gene, CaGLIP1, in disease susceptibility and abiotic stress tolerance. *Planta* **2008**, *227*, 539–558.
47. Sun, X.; Cao, Y.; Yang, Z.; Xu, C.; Li, X.; Wang, S.; Zhang, Q. Xa26, a gene conferring resistance to Xanthomonas oryzae pv. oryzae in rice, encodes an LRR receptor kinase-like protein. *Plant J.* **2004**, *37*, 517–527.
48. Llorente, F.; Alonso-Blanco, C.; Sánchez-Rodríguez, C.; Jorda, L.; Molina, A. ERECTA receptor-like kinase and heterotrimeric G protein from Arabidopsis are required for resistance to the necrotrophic fungus Plectosphaerella cucumerina. *Plant J.* **2005**, *43*, 165–180.
49. Mbengue, M.; Bourdais, G.; Gervasi, F.; Beck, M.; Zhou, J.; Spallek, T.; Bartels, S.; Boller, T.; Ueda, T.; Kuhn, H.; et al. Clathrin-dependent endocytosis is required for immunity mediated by pattern recognition receptor kinases. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, 11034–11039.
50. Ibraheem, F.; Gaffoor, I.; Chopra, S. Flavonoid phytoalexin-dependent resistance to anthracnose leaf blight requires a functional yellow seed1 in Sorghum bicolor. *Genetics* **2010**, *184*, 915–926.
51. Ibraheem, F.; Gaffoor, I.; Tan, Q.; Shyu, C.R.; Chopra, S. A sorghum MYB transcription factor induces 3-deoxyanthocyanidins and enhances resistance against leaf blights in maize. *Molecules* **2015**, *20*, 2388–2404.
52. van den Burg, H.A.; Tsitsigiannis, D.I.; Rowland, O.; Lo, J.; Rallapalli, G.; MacLean, D.; Takken, F.L.; Jones, J.D. The F-box protein ACRE189/ACIF1 regulates cell death and defense responses activated during pathogen recognition in tobacco and tomato. *Plant Cell* **2008**, *20*, 697–719.
53. Takemoto, D.; Rafiqi, M.; Hurley, U.; Lawrence, G.I.; Bernoux, M.; Hardham, A.R.; Ellis, J.G.; Dodds, P.N.; Jones, D.A. N-terminal motifs in some plant disease resistance proteins function in membrane attachment and contribute to disease resistance. *Mol. Plant Microbe Interact.* **2012**, *25*, 379–392.