



Title	Body Weight Loss After Orthognathic Surgery: Comparison Between Postoperative Intermaxillary Fixation with Metal Wire and Elastic Traction, Factors Related to Body Weight Loss
Author(s)	Ooi, K.; Inoue, N.; Matsushita, K.; Yamaguchi, H.; Mikoya, T.; Kawashiri, S.; Tei, K.
Citation	Journal of maxillofacial and oral surgery, 20(1), 95-99 https://doi.org/10.1007/s12663-019-01318-6
Issue Date	2021-03
Doc URL	http://hdl.handle.net/2115/84252
Rights	This is a post-peer-review, pre-copyedit version of an article published in Journal of maxillofacial and oral surgery. The final authenticated version is available online at: https://doi.org/10.1007/s12663-019-01318-6
Type	article (author version)
File Information	body weight loss (JMFOS) 2019.pdf



[Instructions for use](#)

Body weight loss after orthognathic surgery: comparison between postoperative intermaxillary fixation with metal wire and elastic traction, factors related to body weight loss

K Ooi^{1,2}, N Inoue³, K Matsushita², H Yamaguchi², T Mikoya², S Kawashiri¹, K Tei²

¹ Department of Oral and Maxillofacial Surgery, Graduate School of Medical Science, Kanazawa University, 13-1 Takaramachi, Kanazawa, Ishikawa, 920-8641, Japan

² Oral and Maxillofacial Surgery, Department of Oral Patho-biological Science, Graduate School of Dental Medicine, Hokkaido University, Kita 13 Nishi 7 Kita-ku, Sapporo, Hokkaido 060-8586, Japan

³ Gerodontology, Department of Oral Health Science, Graduate School of Dental Medicine, Hokkaido University, Kita 13 Nishi 7 Kita-ku, Sapporo, Hokkaido 060-8586, Japan

Corresponding author at:dental medicine,

Kazuhiro Ooi, D.D.S.,Ph.D.

Department of Oral and Maxillofacial Surgery, Kanazawa University

Graduate School of Medical Science

13-1 Takaramachi, Kanazawa, Ishikawa 920-8641, Japan

E-mail address : k-ooi@staff.kanazawa-u.ac.jp

Conflict of interest

The authors declare that they have no conflicts of interest.

Role of the funding source

There was no source of funding for this research.

Key words: body weight loss; orthognathic surgery; intermaxillary fixation with metal wire; elastic traction

Short title: Body weight loss after orthognathic surgery

Abstract

The aim of this study was to compare body weight loss between postoperative intermaxillary fixation with metal wire and elastic traction, and to investigate factors related to body weight loss after orthognathic surgery. Subjects were 59 patients with dentofacial deformity, comprising 31 patients treated with intermaxillary fixation (IMF) and 28 patients treated with elastic traction without intermaxillary fixation (ELT) just after surgery. Body weight loss was measured at 1 week (T1) and 2 weeks (T2) after surgery. Body weight loss was compared between IMF and ELT, and factors related to body weight loss were statistical analyzed. Body weight loss ratio was significantly increased in IMF (2.6%) rather than in ELT (1.4%) at T1, but only tend to be increased in both groups at T2, showing no statistical difference. Body weight loss ratio was significantly increased at T2 compared to T1 in both groups. Body weight loss was significantly greater at T2 than at T1. Both IMF and ELT cause body weight loss after orthognathic surgery, but IMF causes body weight loss earlier than ELT and increased early body weight loss increases continuous body weight loss after orthognathic surgery.

Introduction

Nutritional management is one of the basic cornerstones to the medical treatment of all diseases. Nutrition can affect postoperative morbidity in surgical patients^{1, 2}, and insufficient food intake can decrease therapeutic effects. In the perioperative period, poor nutrition is thought to adversely affect wound healing^{3, 4} and postoperative infections⁵. Preventing deterioration of nutritional status in the perioperative period is desirable. In particular, patients following orthognathic surgery are unable to eat a normal diet for 6-8 weeks due to trismus and postoperative occlusal management. If postoperative intermaxillary fixation with metal wire (IMF) is performed just after surgery, patients can only take food orally by liquid diet or by nasogastric tube feeding until the IMF is removed. IMF is one of the modalities used before the utilization of miniplates to stabilize and promote the healing of fractured facial bones in cases of maxilla-mandibular trauma or orthognathic surgery. Recently, elastic traction (ELT) has been mainly used for occlusal management without IMF according to development of the miniplates osteosynthesis method. No reports appear to have compared body weight loss between postoperative occlusal management with IMF or ELT, although body weight loss has been noted as one of the major side effects of IMF in patients with

mandibular fractures⁶ or orthognathic surgery⁷. This may eventually lead to impaired bone and wound healing and thus to a deterioration in overall functional recovery⁸. The literature lacks systematic documentation of the weight loss experienced by patients during the first few weeks or months or of the time required for postoperative recovery following orthognathic surgery. The perioperative period of orthognathic surgery is thus suboptimal for nutrition, but the details remain unclear. In addition, few reports have investigated factors related to nutritional status after orthognathic surgery. Knowledge about nutritional status and related factors in the early postoperative period following orthognathic surgery is important for perioperative management of orthognathic surgery. The aim of this study was therefore to compare body weight loss between IMF and ELT and to investigate factors related to body weight loss after orthognathic surgery.

Materials and methods

Subjects were 59 patients with dentofacial deformity who underwent bilateral sagittal splitting ramus osteotomy (BSSRO). Of these, 31 patients were treated with intermaxillary fixation using metal wire (IMF), and 28 patients were treated with elastic

traction without IMF (ELT) after surgery (Fig. 1). IMF was done for 2 weeks only in IMF group. Patients were able to have food without elastics during only meal time in ELT group. This study performed retrospectively. ELT group was subsequently treated after IMF group was treated. No significant differences in clinical status was seen between the two groups (Table 1). No cases showed postoperative nausea or vomiting. All patients consumed food at nearly 35 kcal/kg body weight from 1 day after surgery. A liquid diet was ingested orally or through a nasogastric tube in IMF during 2 weeks only. In ELT, the liquid diet was ingested orally until 5 days after surgery, then paste or a soft diet was started depending on the state of elastics removal for each meal. Age and body mass index (kg/m^2) were assessed as physical factors, while operation time, blood loss, and amount of mandibular movement were assessed as operative stress factors, and body weight loss, total protein, serum albumin were assessed as nutritional state factors at 1 week postoperatively (T1) and 2 weeks postoperatively (T2).

Body weight loss ratio ($(\text{preoperative body weight} - \text{postoperative body weight} / \text{preoperative body weight} \times 100 (\%))$) was compared between IMF and ELT, and factors related to body weight loss were statistically analyzed (Prism 7 GraphPad software, San

Diego, CA). This study was approved by the Hokkaido University Research Ethics Committee (Ref. No. 010-0285).

Results

Body weight loss ratio was significantly higher in IMF (2.6%) than in ELT (1.4%) at T1, but was increased with no significant difference between groups at T2. A body weight loss ratio $\geq 5\%$ was observed in 5 patients in both groups at T2, but in only 1 patient in IMF at T1 (Fig. 2). Body weight loss ratio was significantly increased at T2 compared to T1 in both groups (Table 2). Body weight loss ratio at T2 compared to that at T1 (Fig. 3).

Discussion

The evidence for nutritional support during the perioperative period has been reviewed, and recommendations have been made regarding when nutritional support is most useful and when it may be counterproductive⁷. Past studies have focused on the relationship between the preoperative nutritional state and postoperative complications.

The importance of postoperative nutritional changes has rarely been discussed, particularly in oral and maxillofacial surgery. Nutrition affects the defense systems of the body in many ways. More than 60 years ago, Studley⁹ reported that postoperative infection-related morbidity and mortality increased when patients lost $\geq 20\%$ of their preoperative body weight. When 35–40% of body weight is lost, death is generally inevitable¹⁰. Recent studies on nutrition and wound healing have focused on the relationship between specific nutrients and wound healing¹¹. Nutrition directly influences the wound-healing process, since malnutrition decreases anabolism and depresses immunity¹². Animal studies have shown that 6 weeks of nutritional deficiency reduces wound strength and depresses both cellular and humoral immunity^{13,14}. Clinically, these deficits manifest as delayed healing, wound infection, and prolonged rehabilitation¹⁵. These indicate that nutrition is a key factor in the development of postoperative complications. Tadano¹⁶ reported that administration of non-nutrient solutions via gastrostomy under TPN accelerated jejunal anastomotic healing in comparison with TPN alone with a fasting period, and provided comparable effects to early enteral feeding. The mechanical loading under such stretch conditions contributed

to the upregulation of de novo type I collagen and type III collagen synthesis by rat fibroblasts derived from the gastrointestinal tract. These results indicate that nutrition is important for immunity and wound healing in animal models.

Body weight offers a simple, easily measured, and important index of nutrition.

Nutrition affects the defense systems of the body in many ways. Body weight was reduced by a mean of 3.5 kg in patients with complications and 2.1 kg in patients with no complications following oral and maxillofacial malignancy¹⁷. This indicates that nutrition is a key factor in the development of postoperative complications.

Intermaxillary fixation with metal wire is used for reliable occlusal management just after orthognathic surgery or maxilla-mandibular fracture reduction. Long-term trismus due to this method often causes body weight loss. Worrall¹⁸ reported no significant difference in weight loss or body composition changes between plated and non-plated groups at 1 week postoperatively, but the IMF group had lost significantly more weight (mean, 4.5 kg) than the plated group (mean, 1.1 kg) by 6 weeks postoperatively.

Cawood¹⁹ reported that patients whose mandibular fractures were treated by internal fixation without IMF lost less weight and regained their preoperative weight faster than

those with fractures treated by IMF alone. In a series of 100 cases, the plated group lost less weight (mean, 3 kg) in the first postoperative week than the IMF group (mean, 5 kg) and had regained their preoperative weight within 4 weeks. In these studies, intermaxillary fixation was shown to cause an undernourished state. However, no mention was made of the quantity of nutrient intake in the intermaxillary fixation and non-intermaxillary fixation groups in either study. Generally, fewer calories are present in fluid food than in solid food, because fluid food is included more water. Fluid food with a higher caloric content is needed to provide the same content as solid food. For the purpose of preventing marked differences in nutritional intake, 35 kcal/kg was provided to both groups in our study. No significant difference in body weight loss ratio was evident between groups at 2 weeks after surgery, suggesting the effectiveness of this approach. The stress of IMF in the initial stage up to 1 week may have been greater than that of ELT, because body weight loss at 1 week was significantly large in IMF than in ELS. In addition, body weight loss ratio $\geq 5\%$ was observed in 5 patients in both groups at 2 weeks after surgery, but only 1 patient with IMF at 1 week after surgery.

This indicates that elastic traction carries similar levels of nutritional stress to IMF by 2

weeks, whereas ELS offers advantages over IMF by 1 week after surgery. Both diet and conversation are difficult due to trismus for a certain period of time after orthognathic surgery, and this is thought to be a strong stressor. These factors should therefore be taken into account when determining the nutrient dose. In recent years, the need for intermaxillary fixation has decreased with miniplate fixation, but undernutrition appears avoidable by providing sufficient nutrients when intermaxillary fixation is needed for abnormal fractures. Many hospitals maintain a close relationship with dietetics departments, whose staff provide invaluable support and advice both before and after surgery. However, despite dietary advice and support, many patients continue to lose weight during the 6- to 8-week period with jaws immobilized by intermaxillary fixation or while eating a soft diet to protect the internal titanium mini plates.

Patients who undergo orthognathic surgery should be counseled on the importance of maintaining an adequate diet throughout the healing period, regardless of intermaxillary fixation.

References

1. Guo CB, Ma DQ, Zhang KH (1994) Applicability of general nutritional status score to patients with oral and maxillofacial malignancies. *Int J Oral Maxillofac Surg* 23:167–9.
2. Howard L, Ashley C (2003) Nutrition in the perioperative patient. *Annu Rev Nutr* 23:263–82.
3. Albina JE (1994) Nutrition and wound healing. *J Parenter Enteral Nutr* 18:367–76.
4. Williams JZ, Barbul A (2003) Nutrition and wound healing. *Surg Clin North Am* 83:571–96.
5. Felblinger DM (2003) Malnutrition, infection, and sepsis in acute and chronic illness. *Crit Care Nurs Clin North Am* 15:71–8.
6. Hayter JP, Cawood JI (1993) The functional case for miniplates in maxillofacial surgery. *Int J Oral Maxillofac Surg* 22 : 91-6.
7. Behbehani F, Al-Aryan H, Al-Attar A, Al-Hamad N (2006) Perceived effectiveness and side effects of intermaxillary fixation for diet control. *Int J Oral Maxillofac Surg* 35 : 618-23.
8. Kuvat SV, Güven E, Hoccoqlu E, Basaran K, Marsan G, Cura N, Emekli U (2010) Body fat composition and weight changes after double-jaw osteotomy. *J Craniofac Surg* 21:1516-8.
9. Studley HD (1936) Percentage of weight loss: basic indicator of surgical risk in patients with chronic peptic ulcer. *JAMA*;106:458–60.
10. World Health Organization. Obesity (1998) Preventing and managing the global epidemic. Report of a WHO consultation on obesity. Geneva, Switzerland : World Health Organization.
11. Williams JZ, Barbul A (2003) Nutrition and wound healing. *Surg Clin North Am* 83:571–96.
12. Chandra RK (1995) Nutrition and immunity in elderly in the elderly: clinical significance. *Nutrition Review*; 53:S80 –S3.
13. Ruberg I (1984) Role of nutrition in wound healing. *Surg Clin North Am* 64: 721.

14. Law DK, Dudrick SJ, Abdou XI (1974) The effects of dietary protein depletion on immunocompetence. *Ann Surg* 179: 168.
15. Leigh J (1986) Postoperative care. III: Rowe KL, Li Williams J, eds. *Maxillofacial Injuries*. Edinburgh: Churchill Livingstone 701.
16. Tadano S, Terashima H, Fukuzawa J, Mastuo R, Ikeda O, Ohkohchi N (2011) Early Postoperative Oral Intake Accelerates Upper Gastrointestinal Anastomotic Healing in the Rat Model. *Journal of Surgical Research* 169: 202–208.
17. Bozzetti F (1989) Effects of artificial nutrition on the nutritional status of cancer patients. *J Parenter Enteral Nutr*;13:406–20.
18. S. F. Worrall (1994) Changes in weight and body composition after orthognathic surgery and jaw fractures: a comparison of miniplates and intermaxillary fixation. *Br J Oral Maxillofac Surg* 32: 289-292.
19. Cawood JJ (1985) Small plate osteosynthesis of mandibular fractures. *Br J Oral Maxillofac Surg* 23: 77.

Table1 : Clinical status of the patients with dentofacial deformity treated by bilateral sagittal splitting ramus osteotomy

	IMF	ELT	P value
Case (number)	31	28	
Age	23 (17~33)	24 (17~40)	0.2947
BMI	21.0 (15.1~26.6)	21.0 (17.7~28.0)	0.9624
Body weight (kg)	53.0 (33~73)	53.8 (43.5~71.5)	0.6655
Laboratory data			
Total protein (g/dl)	7.4 (6.5~8.3)	7.2 (5.9~7.9)	0.1691
Total albumin (mm)	4.6 (4.1~5.2)	4.5 (3.9~4.9)	0.0899

IMF : titanium mini plates with 2 weeks intermaxillary fixation using metal wire

ELT: titanium mini plates fixation with elastic traction

BMI: body mass index


mean (range)

* p<0.05 t test


Table 2. Factors related body weight loss at 2 week after bilateral sagittal splitting ramus osteotomy (n=59)

	R square	P value
Physical factors		
Age	0.0348	0.1570
Body mass index (BMI)	0.0001	0.9836
Operative stress factors		
Operation time (min)	0.0003	0.8983
Blood loss (g)	0.0376	0.1408
Mandibular movement (mm)	0.0028	0.6924
Nutritional state factors		
Body weight loss (1 week after surgery)	0.3697	0.0001 *
Laboratory data (change from pre-operation)		
total protein	0.0940	0.0203
serum albumin	0.0450	0.1273

* p<0.05 Spearman`s test




IMF



ELT

IMF : titanium mini plates fixation with intermaxillary fixation using metal wire
ELT: titanium mini plates fixation with elastic traction

Fig.1 Occlusal management after orthognathic surgery



IMF : titanium mini plates fixation with intermaxillary fixation using metal wire
 ELT: titanium mini plates fixation with elastic traction
 T1: 1 week after surgery
 T2 : 2 weeks after surgery

Fig.2 Body weight loss ratio at 1 and 2 weeks after bilateral sagittal splitting ramus osteotomy in IMF and ELT




Fig.3 Relationship between body weight loss ratio at 2 weeks after surgery and 1 week after surgery