



Title	Brown bear digging for cicada nymphs : a novel interaction in a forest ecosystem
Author(s)	Tomita, Kanji; Hiura, Tsutom
Citation	Ecology, 101(3), e02899 https://doi.org/10.1002/ecy.2899
Issue Date	2020-03
Doc URL	http://hdl.handle.net/2115/86161
Rights	This is the peer reviewed version of the following article: Tomita, K., and T. Hiura. 2020. Brown bear digging for cicada nymphs : a novel interaction in a forest ecosystem. Ecology, 101(3): e02899, which has been published in final form at https://doi.org/10.1002/ecy.2899 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley 's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited.
Type	article (author version)
File Information	TH -2019-Ecology.pdf



[Instructions for use](#)

1 Article type : The Scientific Naturalist

2 Running Head: The Scientific Naturalist

3 **Brown Bear Digging for Cicada Nymphs: a novel interaction in a forest ecosystem**

4

5 KANJI TOMITA^{1,3*} and TSUTOM HIURA²

6

7 ¹Graduate School of Environmental Science, Hokkaido University, N10 W5, Kita-ku,

8 Sapporo, Hokkaido 060 0810, Japan

9 ²Field Science Center for Northern Biosphere, Hokkaido University, N9 W9, Kita-ku,

10 Sapporo, Hokkaido 060 0809, Japan

11

12 *** Corresponding author:**

13 KANJI TOMITA

14 ³E-mail: ktomita38@gmail.com

15

16 **Key words:** *Ursus arctos*; feeding habit; dietary change; bioturbation; social learning;

17 omnivory; novel species interaction; ecosystem engineer

18

19 The brown bear (*Ursus arctos*) is one of the most widespread large carnivores in
20 the northern hemisphere. Their diets are highly diverse and depend on resource availabilities
21 across regions and seasons (Bojarska and Selva 2012). Brown bears often consume starchy
22 plant roots and insects by digging into the soil (Mattson et al. 1991; Tardiff and Stanford
23 1998). In the Shiretoko World Heritage, Hokkaido, northern Japan (44° N, 145° E), where the
24 density of brown bears is high (Kohira et al. 2009), these animals have recently been
25 observed digging for cicada nymphs (*Lyristes bihamatus*) during summer (Fig. 1a, b). This
26 behavior has not been previously reported, even though some other mammalian species
27 forage for cicada nymphs (Hahus and Smith 1990). In this article, we show some preliminary
28 results of our field survey on this novel interaction between brown bear and cicada. We then
29 discuss the reasons why brown bears forage on cicada nymphs and the possible ecological
30 consequences of their digging in the forest ecosystem.

31 Wildlife managers first observed brown bears digging for cicada nymphs in 2000 in
32 the study area. In fact, cicada nymphs have not been reported in scats of brown bears
33 collected in the summer of mid-1980s in the study area (Fig.2; Yamanaka and Aoi 1988).
34 This earlier study along with ours allows us to determine whether the proportion of cicadas
35 and other foods in the diet has changed during the past 30 years. In 2018, we evaluated the
36 composition of brown bear scats collected in the same season and region as the earlier study.

37 By comparing scat compositions between the mid-1980s and the 2018, we determined the
38 changes in brown bears diets in the study area during the past 30 years. The proportion of
39 cicada nymphs was estimated 14.3%, suggesting that brown bears consume cicada nymphs at
40 a certain rate in summer. Final instar nymphs are highly nutritious and finish their
41 development until emergence in shallow soil during summer, and so bears can easily dig for
42 them (Hayashi and Saisho 2011). Although herbaceous plants comprised the highest
43 proportion in scats in both periods, we found that the proportion of herbaceous plants in scats
44 in 2018 is half that recorded in mid-1980s (Fig.2).

45 Evidence of brown bear digging was frequently observed in the larch (*Larix*
46 *kaempferi*) plantations, whereas we could not find evidence of digging in the Sakhalin spruce
47 (*Picea glehnii*) plantations and the natural mixed forests. In August 2017, when final instar
48 cicada nymphs completed emergence, we created 10 survey plots of 100 m² area in each
49 forest type and counted the number of exuviae of *L. bihamatus* nymphs as a proxy of their
50 abundance in each plot. The largest number of the *L. bihamatus* exuviae occurred in the larch
51 plantations (mean \pm SE / 100m² = 153.9 \pm 15.6). The number of *L. bihamatus* exuviae in the
52 Sakhalin spruce plantations and the natural mixed forests was 8.7 \pm 3.0 and 4.5 \pm 2.3,
53 respectively. These data indicate that brown bears can most efficiently forage on cicada
54 nymphs by searching intensively in the larch plantation where the nymphs aggregate at high

55 density.

56 We set up 8 camera traps in the larch plantations where brown bears dug the
57 previous year to evaluate how many brown bears dug for cicada nymphs, by counting the
58 minimum number of brown bears based on useful features for individual identification (e.g.,
59 color, body size, and family structure). As a result, we captured 112 videos wherein bears
60 were recorded to be digging and detected a minimum number of 11 individuals; 3 adult bears
61 were females with cub(s), 2 were solitary adult female bears, and 2 were subadult bears. ([See](#)
62 [Videos S1 and S2 for two examples.](#))

63 Why have brown bears begun foraging on cicada nymphs since 2000? One possible
64 reason may be that overgrazing by sika deer (*Cervus nippon yesoensis*) has altered the diet of
65 brown bears by reducing the available herbaceous plants for bears. An index of population
66 density of sika deer by spotlight surveys has revealed a remarkable increase from 1 deer/km
67 in the late 1980s to 20 deer/km in the early 2000s in the study area (Kaji et al. 2006). Because
68 herbaceous plant species preferred by sika deer were partially in common with the species
69 upon which brown bears foraged in summer during the 1980s (Kaji 1988), the availability of
70 herbaceous plants for bears during summer may have decreased in the study area. This is also
71 supported by the results of our scat analysis (Fig. 2). Therefore, brown bears may have
72 foraged on cicada nymphs as an alternative food resource to herbaceous plants.

73 Our camera-traps revealed that 3 female brown bears with cub(s) dug for cicada
74 nymphs in the study area. Possibly, the digging behavior of brown bears will propagate
75 through the bear population via social learning from mother bear to cub, because bears likely
76 acquire foraging behavior through learning from their mothers during at least the first year of
77 their life (Gilbert 1999; Hopkins III 2013). Moreover, because home ranges of female brown
78 bears tend to be fixed in proximity to that of their mother's (Zedrossor et al. 2007), female
79 bears acquiring the digging behavior will remain in the same population. Thus, in the future,
80 the number of brown bears that dig for cicada nymphs will probably increase via social
81 learning in the study area.

82 Bioturbation by digging mammals is a well-studied example of ecosystem
83 engineering that alters habitat structure for other species and soil nutrient dynamics (Tardiff
84 and Stanford 1998; Meysman et al. 2006; Mallen-Cooper et al. 2019). We observed the area
85 of a dug patch was often over 100 m², and there were at least several dozens of the patches.
86 Furthermore, individual bear apparently dug up a large amount of soil according to
87 observation by our camera traps. Thus, bioturbation by brown bear digging may have
88 significant engineering effects in the forest ecosystem. Additionally, in the study area brown
89 bears may have exerted these effects since 2000 when they started digging for cicada
90 nymphs. In other words, the novel interaction between native species may have generated

91 novel bioturbation in the forest ecosystem.

92 As the number of brown bears that dig for cicada nymphs increases, the ecological
93 importance of the brown bear as an ecosystem engineer will increase in the forest ecosystem.

94 Finally, we propose a hypothesis that propagation of the digging behavior via social learning
95 might strengthen their ecological effects of bioturbation. This hypothesis may shed light on
96 linkage between social learning and ecosystem engineering (i.e. bioturbation). Testing this
97 hypothesis requires data on (1) population trends of digging bears; (2) their kinship based on
98 fecal DNA which can evaluate the relationship between kinship and diet; (3) temporal
99 dynamics in spatial patterns of brown bear digging; and (4) engineering effects of the digging
100 on other organisms and soil nutrient dynamics.

101 **Acknowledgments**

102 We thank members of Shiretoko Nature Foundation for providing valuable information
103 about brown bear digging for cicada nymphs.

104

105 **Literature cited**

- 106 Bojarska, K., & Selva, N. 2012. Spatial patterns in brown bear *Ursus arctos* diet: the role of
107 geographical and environmental factors. *Mammal Review* 42: 120-143.
- 108 Gilbert, B. K. 1999. Opportunities for social learning in bears. Pages 225-235 in H. O. Box
109 and K. R. Gibson editors *Mammalian social learning: comparative and ecological*
110 *perspectives*. Cambridge University Press, Cambridge, United Kingdom.
- 111 Hahus, S. C., & Smith, K. G. 1990. Food habits of *Blarina*, *Peromyscus*, and *Microtus* in
112 relation to an emergence of periodical cicadas (*Magicicada*). *Journal of Mammalogy*
113 71: 249-252.
- 114 Hayashi, M., & Saisho, Y. 2011. The Cicadidae of Japan. Seibundo-shinkosha, Tokyo, 1-224.
115 [in Japanese]
- 116 Hopkins III, J. B. 2013. Use of genetics to investigate socially learned foraging behavior in
117 free-ranging black bears. *Journal of Mammalogy*, 94: 1214-1222.
- 118 Kaji, K., 1988. Sika deer. Pages 155-180 in Ohtaishi N and Nakagawa H, editors. *Animals of*
119 *Shiretoko*. Hokkaido University Press, Sapporo, Japan Pages 155–180 [in Japanese
120 with an English abstract]
- 121 Kaji, K., Okada, H., Kohira, M., M. Yamanaka. 2006. The Shiretoko sika deer herd. Pages
122 221-224. in D. R. McCullough, K. Kaji and M. Yamanaka, editors. *Wildlife in*
123 *Shiretoko and Yellowstone National Parks. Lessons in Wildlife Conservation from Two*
124 *World Heritage Sites*. Shiretoko Nature Foundation, Syari, Japan.
- 125 Kohira, M., Okada, H., Nakanishi, M., & Yamanaka, M. 2009. Modeling the effects of
126 human-caused mortality on the brown bear population on the Shiretoko Peninsula,
127 Hokkaido, Japan. *Ursus* 20: 12-22.
- 128 Mallen-Cooper, M., Nakagawa, S., & Eldridge, D. J. 2019. Global meta-analysis of soil-
129 disturbing vertebrates reveals strong effects on ecosystem patterns and processes.
130 *Global Ecology and Biogeography* 28: 661-679.

- 131 Mattson, D. J., Gillin, C. M., Benson, S. A., & Knight, R. R. 1991. Bear feeding activity at
132 alpine insect aggregation sites in the Yellowstone ecosystem. *Canadian Journal of*
133 *Zoology* 69: 2430-2435.
- 134 Meysman, F. J., Middelburg, J. J., & Heip, C. H. 2006. Bioturbation: a fresh look at Darwin's
135 last idea. *Trends in Ecology & Evolution* 21: 688-695.
- 136 Tardiff, S. E., & Stanford, J. A. 1998. Grizzly bear digging: effects on subalpine meadow
137 plants in relation to mineral nitrogen availability. *Ecology* 79: 2219-2228.
- 138 Yamanaka, M., & Aoi, T. 1988. Brown bear. Pages 181-223 in Ohtaishi N. and Nakagawa H.
139 editors. *Animals of Shiretoko*. Hokkaido University Press, Sapporo, Japan. [in Japanese
140 with an English abstract]
- 141 Zedrosser, A., Støen, O. G., Sæbø, S., & Swenson, J. E. 2007. Should I stay or should I go?
142 Natal dispersal in the brown bear. *Animal Behaviour* 74: 369-376.
- 143

144 **Figure legends**

145 Figure 1: (a) Female brown bear with two cubs dig for cicada nymphs in a larch plantation.

146 (b) Brown bear scat containing final instar *Lyristes bihamatus* nymphs. Scale bar: 50 mm

147 (photo credit: Shiretoko Nature Foundation)

148 Figure 2: Scat composition of brown bears in visually estimated percent volumes for major

149 categories, in 1985-1986 (left) and 2018 (right). Data of the scat composition in 1985-1986

150 from Yamanaka and Aoi (1988).

151

152



