


Cite this article as: Rubino AS, De Santo LS, Pisano A, Mauro M, Benussi S, Borghetti V *et al.* Cardiac surgery practice during the COVID-19 outbreak: a multicentre national survey. *Eur J Cardiothorac Surg* 2021;59:901–7.

## Cardiac surgery practice during the COVID-19 outbreak: a multicentre national survey

Antonino Salvatore Rubino <sup>a†</sup>, Luca Salvatore De Santo<sup>a†</sup>, Antonio Pisano<sup>b</sup>, Michele di Mauro<sup>c</sup>,  
Stefano Benussi<sup>d</sup>, Valentino Borghetti<sup>e</sup>, Alessandro Castiglioni<sup>f</sup>, Luigi Chiariello<sup>g</sup>, Andrea Colli<sup>h</sup>,  
Antonio De Bellis<sup>i</sup>, Carlo Maria De Filippo<sup>j</sup>, Ruggero De Paulis<sup>k</sup>, Giuseppe Di Benedetto<sup>l</sup>, Marco Di Eusanio<sup>m</sup>,  
Giuseppe Faggian<sup>n</sup>, Brenno Fiorani<sup>o</sup>, Pasquale Antonio Fratto<sup>p</sup>, Angelo Giuseppe Giuffrida<sup>q</sup>,  
Mattia Glauber<sup>r</sup>, Gabriele Iannelli<sup>s</sup>, Severino Iesu<sup>t</sup>, Ugolino Livi<sup>u</sup>, Gianluca Martinelli<sup>v</sup>, Massimo Massetti<sup>w</sup>,  
Pasquale Mastroberoberto<sup>x</sup>, Lorenzo Menicanti<sup>y</sup>, Giuseppe Minniti<sup>z</sup>, Fabio Miraldi<sup>aa</sup>, Gianfranco Montesi<sup>ab</sup>,  
Francesco Musumeci<sup>ac</sup>, Francesco Nicolini<sup>ad</sup>, Carlo Pace Napoleone<sup>ae</sup>, Paolo Panisi<sup>af</sup>, Aniello Pappalardo<sup>ag</sup>,  
Francesco Patanè<sup>ah</sup>, Temistocle Ragni<sup>ai</sup>, Mauro Rinaldi<sup>aj</sup>, Salvatore Tribastone<sup>ak</sup>, Michele Triggiani<sup>al</sup>,  
Francesco Paolo Tritto<sup>am</sup>, Carlo Zebele<sup>an</sup>, Alessandro Parolari<sup>ao</sup>, Gino Gerosa<sup>ap†</sup> and Marisa De Feo<sup>af</sup>,  
for the Italian Society for Cardiac Surgery Task Force on COVID-19 Pandemic<sup>††</sup>

- <sup>a</sup> Department of Translational Medicine, Cardiac Surgery Unit, University of Campania, Luigi Vanvitelli, AORN dei Colli, Cardiac Surgery, Vincenzo Monaldi Hospital, Naples, Italy  
<sup>b</sup> Cardiac Anaesthesia and Intensive Care Unit, AORN Dei Colli, Monaldi Hospital, Naples, Italy  
<sup>c</sup> Cardiology and Cardiac Surgery, API “Madonna del Ponte”, Lanciano, Italy  
<sup>d</sup> Cardiac Surgery Unit, Spedali Civili, Brescia, Italy  
<sup>e</sup> Cardiac Surgery Unit, Azienda Ospedaliera Santa Maria, Terni, Italy  
<sup>f</sup> Cardiac Surgery Unit, San Raffaele Hospital, Milan, Italy  
<sup>g</sup> Cardiac Surgery Unit, Clinica Mediterranea, Naples, Italy  
<sup>h</sup> Cardiac Surgery Unit, Azienda Ospedaliero-Universitaria Pisana, Pisa, Italy  
<sup>i</sup> Cardiac Surgery Unit, Casa di Cura San Michele, Maddaloni, Caserta, Italy  
<sup>j</sup> Cardiac Surgery Unit, Gemelli Molise, Fondazione Giovanni Paolo II, Campobasso, Italy  
<sup>k</sup> Cardiac Surgery Unit, European Hospital, Rome, Italy  
<sup>l</sup> Cardiac Surgery Unit, Clinica Pineta Grande, Castel Volturno, Caserta, Italy  
<sup>m</sup> Cardiac Surgery Department, Lancisi Cardiovascular Center—OORR—Polytechnic University of Marche—School of Medicine, Ancona, Italy  
<sup>n</sup> Cardiac Surgery Unit, Azienda Ospedaliera Universitaria Integrata, University of Verona, Verona, Italy  
<sup>o</sup> Cardiac Surgery Unit, AORNAS “San Giuseppe Moscati”, Avellino, Italy  
<sup>p</sup> Cardiac Surgery Unit, Grande Ospedale Metropolitano “Bianchi-Melacrino-Morelli”, Reggio Calabria, Italy  
<sup>q</sup> Cardiac Surgery Unit, AOU Policlinico-Vittorio Emanuele, Catania, Italy  
<sup>r</sup> Cardiac Surgery Unit, Istituto Clinico Sant’Ambrogio, Milan, Italy  
<sup>s</sup> Cardiac Surgery Unit, University of Naples Federico II, Naples, Italy  
<sup>t</sup> Cardiac Surgery Unit, A.O. San Giovanni di Dio e Ruggi d’Aragona, Salerno, Italy  
<sup>u</sup> Cardiac Surgery Unit, Azienda Ospedaliero-Universitaria S. Maria della Misericordia, Udine, Italy  
<sup>v</sup> Cardiac Surgery Unit, Clinica San Gaudenzio, Gruppo Policlinico di Monza, Novara, Italy  
<sup>w</sup> Cardiac Surgery Unit, Policlinico “A. Gemelli”, Università Cattolica del Sacro Cuore, Rome, Italy  
<sup>x</sup> Cardiac Surgery Unit, Università degli Studi Magna Graecia, Catanzaro, Italy  
<sup>y</sup> Cardiac Surgery Unit, Centro “E. Malan”, Milan, Italy  
<sup>z</sup> Cardiac Surgery Unit, Ospedale di Treviso, Treviso, Italy  
<sup>aa</sup> Cardiac Surgery Unit, Università La Sapienza, Rome, Italy  
<sup>ab</sup> Cardiac Surgery Unit, Università Policlinico “Le Scotte”, Siena, Italy  
<sup>ac</sup> Cardiac Surgery Unit, Azienda Ospedaliera San Camillo-Forlanini, Rome, Italy  
<sup>ad</sup> Cardiac Surgery Unit, University of Parma, Parma, Italy  
<sup>ae</sup> Azienda Ospedaliera Città della Salute e della Scienza di Torino, Presidio Ospedale Infantile Regina Margherita, Torino, Italy  
<sup>af</sup> Cardiac Surgery Unit, Humanitas Gavazzeni, Bergamo, Italy  
<sup>ag</sup> Cardiac Surgery Unit, Ospedali Riuniti di Trieste, Italy

<sup>†</sup> The first two authors contributed equally to the study.

<sup>††</sup> The last two authors supervised equally to this paper as co-senior authors.


<sup>†††</sup> Members of the Italian Society for Cardiac Surgery Task Force on COVID-19 Pandemic are listed in the Supplementary Material.

Presented at the 34th Annual Meeting of the European Association for Cardio-Thoracic Surgery, Barcelona, Spain, 8–10 October 2020.

- <sup>ah</sup> Cardiac Surgery Unit, Azienda Ospedaliera Papardo, Messina, Italy  
<sup>ai</sup> Cardiac Surgery Unit, Ospedale R. Silvestrini, Perugia, Italy  
<sup>aj</sup> Cardiac Surgery Unit, Azienda Ospedaliera Città della Salute e della Scienza di Torino, Presidio Molinette, Università di Torino, Torino, Italy  
<sup>ak</sup> Cardiac Surgery Unit, Centro Cuore Morgagni, Pedara, Catania, Italy  
<sup>al</sup> Cardiac Surgery Unit, Ospedale Giuseppe Mazzini, Teramo, Italy  
<sup>am</sup> Cardiac Surgery Unit, AORNAS SS. Anna e Sebastiano, Caserta, Italy  
<sup>an</sup> Cardiac Surgery Unit, Casa di Cura Montevergine, Mercogliano (AV), Italy  
<sup>ao</sup> Department of University Cardiac Surgery and Translational Research, IRCCS Policlinico S. Donato, University of Milan, Milan, Italy  
<sup>ap</sup> Department of Cardiac Surgery, University of Padua, Padua, Italy

\* Corresponding author. Department of Translational Medical Sciences, University of Campania Luigi Vanvitelli, Cardiac Surgery, Vincenzo Monaldi Hospital, AORN dei Colli, Via Leonardo Bianchi 5, 80131 Naples, Italy. Tel/fax: +39-0815464594; e-mail: antoninosalvatore.rubino@unicampania.it (A.S. Rubino).

Received 24 June 2020; received in revised form 7 October 2020; accepted 18 October 2020



## Abstract

**OBJECTIVES:** Healthcare systems worldwide have been overburdened by the coronavirus disease 2019 (COVID-19) outbreak. Accordingly, hospitals had to implement strategies to profoundly reshape both non-COVID-19 medical care and surgical activities. Knowledge about the impact of the COVID-19 pandemic on cardiac surgery practice is pivotal. The goal of the present study was to describe the changes in cardiac surgery practices during the health emergency at the national level.

**METHODS:** A 26-question web-enabled survey including all adult cardiac surgery units in Italy was conducted to assess how their clinical practice changed during the national lockdown. Data were compared to data from the corresponding period in 2019.

**RESULTS:** All but 2 centres (94.9%) adopted specific protocols to screen patients and personnel. A significant reduction in the number of dedicated cardiac intensive care unit beds (-35.4%) and operating rooms (-29.2%), along with healthcare personnel reallocation to COVID departments (nurses -15.4%, anaesthesiologists -7.7%), was noted. Overall adult cardiac surgery volumes were dramatically reduced (1734 procedures vs 3447;  $P < 0.001$ ), with a significant drop in elective procedures [580 (33.4%) vs 2420 (70.2%)].

**CONCLUSIONS:** This national survey found major changes in cardiac surgery practice as a response to the COVID-19 pandemic. This experience should lead to the development of permanent systems-based plans to face possible future pandemics. These data may effectively help policy decision-making in prioritizing healthcare resource reallocation during the ongoing pandemic and once the healthcare emergency is over.

**Keywords:** Cardiac surgery • COVID-19 pandemic • Waiting list • Healthcare resources • Prioritization •

**ABBREVIATIONS**

COVID-19	Coronavirus disease 2019
ICU	Intensive care unit
IQR	Interquartile range
LOS	Length of stay
PPE	Personal protective equipment
SARS-CoV-2	Severe acute respiratory syndrome coronavirus-2

**INTRODUCTION**


The coronavirus disease 2019 (COVID-19) pandemic is the world's largest infectious disease crisis in the last 100 years. Italy was the first European country affected and has the third highest number of casualties after the UK and Spain. Despite several attempts to control the spread of infection, the prevalence rose significantly and led to a nationwide lockdown on 9 March 2020 that ended only on 4 May 2020 [1]. The availability of inpatient beds in Italian hospitals is chronically lower than the European average (262.5 per 100 000 inhabitants vs 372.2 per 100 000, respectively) [2]. As the cases of acute respiratory failure requiring intensive care unit (ICU) admission surged, a dramatic imbalance between the healthcare needs of the population and the actual availability of acute/critical care resources became evident. In this 'disaster medicine' scenario, the principle of proportionality of care has driven resource allocation, leaving many patients, especially those with other diseases, without adequate care [3], and leading to a possible increase in non-COVID-19-related deaths as a further dramatic consequence of the pandemic [4, 5]. Remodelling of therapeutic algorithms for non-COVID-19 medical and surgical care has therefore become mandatory. Knowledge about the impact of the COVID-19 pandemic on cardiac surgery practice is pivotal to define potential improvement strategies in case of similar scenarios in the future.

In fact, the risk of overlooking severely ill patients and postponing life-saving treatments was high during the central phase of the epidemic. Moreover, quantifying the backlog of deferred cases might help estimate the need for increasing cardiac surgery volumes during the COVID-19 recovery period (phase 2 and over) [6]. The goal of the present study was to describe the changes in cardiac surgery practices during the health emergency at the national level.

**MATERIALS AND METHODS**

On 24 April 2020, a 26-question survey (Supplementary Material, File S1) was sent by e-mail to a total of 99 Italian adult cardiac surgery centres identified from the Italian Society for Cardiac Surgery mailing list. In view of the rapidly evolving situation and given the need for timely presentation of results, we predetermined to close the survey on 20 May 2020.

The topics investigated by the questionnaire included the redistribution of dedicated healthcare resources; the modalities for screening surgical candidates and for active surveillance of healthcare workers; the availability of adequate personal protective equipment (PPE); the number, urgency status, type of surgical

**Hospital Activity During Lockdown  
by resource allocation**

**Figure 1:** Proportion of centres stratified by resource reallocation to treat COVID-19.

procedure, patterns of referral and discharge and the length of hospital stay of all consecutive patients referred for surgery during the lockdown period. These data were compared with those from the equivalent period in 2019.

**Data collection**

Each centre retrieved data from their internal records or from the Department of Management. The completed questionnaires were submitted to the coordinating unit at the Department of Translational Medical Sciences, University of Campania, Naples, which was in charge of data collection and analysis.

**Statistical analyses**

Categorical variables are presented as numbers and percentages, continuous data as median, interquartile range (IQR) and maximum value. Data were compared using the  $\chi^2$  test, the paired-sample *t*-test and the Kruskal-Wallis test. Statistical significance was set at an alpha level of 0.05. All statistical analyses were performed using SAS version 9.4 (SAS Institute, Cary, NC, USA).

**RESULTS**

Thirty-nine national cardiac surgery centres out of 99 (39.4%) agreed to participate in the survey and returned the completed questionnaire. Of these, 15 were academic hospitals (38.5%), 15 were public hospitals (38.5%) and 9 were private clinics (23.1%). Fourteen centres (35.9%) were in northern Italy, 10 (25.6%) in

**Table 1:** Reallocation of healthcare resources in the 3 macro areas

Beds	Northern Italy (n = 12) <sup>a</sup>	Central Italy (n = 8) <sup>a</sup>	Southern Italy (n = 10) <sup>a</sup>	P-value
ICU	-45.1% (20.0–81.0%; max 100%)	-30.0% (0–71.3%; max 100%)	-21.2% (0–56.9%; max 100%)	0.50
Operating room	-41.7% (0–74.3%; max 100%)	-12.5% (0–50.0%; max 100%)	-16.7% (0–50.0%; max 50.0%)	0.60
Ward	-35.4% (2.3–84.9%; max 100%)	-41.7% (6.3–55.4%; max 66.7%)	-8.3% (0–35.8%; max 50.0%)	0.18
Healthcare professionals	Northern Italy (n = 10) <sup>a</sup>	Central Italy (n = 5) <sup>a</sup>	Southern Italy (n = 6) <sup>a</sup>	P-value
Nurses	-25.7% (4.6–67.9%; max 100%)	-28.6% (7.7–40.8%; max 41.7%)	-1.8% (0–15.0%; max 40.0%)	0.18
Anaesthesiologists	-23.8% (0–63.8%; max 100%)	0% (0–36.9%; max 37.5%)	-3.8% (0–16.7%; max 33.3%)	0.43
Cardiac surgeons	0% (0–37.5%; max 77.8%)	0% (0–3.8%; max 7.7%)	0% (0–5.0%; max 20.0%)	0.39
Cardiologists	0% (0–51.6%; max 87.5%)	0	0% (0–7.1%; max 28.6%)	0.20
Perfusionists	0% (0–25.0%; max 50.0%)	0	0% (0–12.5%; max 50.0%)	0.42

Results are expressed as median (interquartile range; maximum) percentage reduction, comparing availability of resources during lockdown to the corresponding period in 2019.

<sup>a</sup>The number of centres experiencing resource reallocation.

ICU: intensive care unit.

central Italy and 15 (38.6%) in southern Italy. Thirty out of 39 (76.9%) were hub hospitals for COVID-19 patients.

### Screening protocols and availability of personal protective equipment

All but 2 centres (94.9%) adopted specific protocols for screening patients admitted to the hospital. Naso- and oropharyngeal swabs for the detection of severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) infection were the most frequently performed preoperative screening tests for surgical candidates: 34 out of 39 centres (87.2%) performed swabs alone (30 centres) or in combination with serum tests (4 centres). The rapid test was the preferred screening test in 2 centres (5.1%), whereas titration of serum antibodies against SARS-CoV-2 was used in 1 centre (2.6%).

Surveillance protocols for healthcare professionals, including self-assessment, recording body temperature, rapid tests, swabs, antibody titre or various combinations of these, were implemented in all but 2 centres.

Use of PPE was mandatory in all centres and was perceived as sufficiently provided in 27 centres (69.2%), temporarily inadequate in 10 (25.6%) and chronically inadequate in 2 (5.1%).

### Resource allocation

Hospital resources were rationed to cope with the national outbreak and to provide adequate regional support. Figure 1 shows the proportion of centres stratified based on whether their hospital resources were reallocated during the lockdown.

The ICUs normally admitting patients after cardiac surgical procedures were the facilities mostly involved, with a median bed reduction of -35.4% compared to 2019 standards (IQR 0–68.8%, maximum 100%, 106 unit reduction out of 189), followed by operating rooms (-29.2%, IQR 0–50%, maximum 100%, 35-unit reduction out of 64) and wards (-28.1%, IQR 0–50%, maximum 100%, 250 unit reduction out of 524). Twenty-one centres (53.8%) experienced a reduction in personnel. In particular, the largest reallocation involved nurses (-15.4%, IQR 0–40.0%, maximum 100%, 149 unit reduction out of 543), followed by anaesthesiologists (-7.7%, IQR 0–36.9%, maximum 100%, 39 unit

reduction out of 101), cardiac surgeons (0%, IQR 0–13.8%, maximum 77.8%, 17 unit reduction out of 50), cardiologists (0%, IQR 0–6.7%, maximum 87.5%, 71 unit reduction out of 145) and perfusionists (0%, IQR 0–0%, maximum 50%, 6 unit reduction out of 23), which were reallocated to the emergency departments dealing with COVID-19 patients. Such redistribution, although not statistically significant, was most evident in northern Italy (Table 1).

Hospital pathways were differentiated to account for suspected or confirmed SARS-CoV-2-positive patients. Thirty-seven centres (94.9%) had a dedicated area for triage (22 outside the ward and 15 inside). Twenty-three centres (59.0%) also had an operating room dedicated to suspected/confirmed COVID-19 patients (15 inside the operating block, 8 outside). Thirty-four centres (87.2%) had dedicated ICU beds available for patients with positive or ongoing screening tests (18 centres in the general ICU, 16 centres inside the cardiac surgery ICU).

The admittance of relatives was forbidden in 27 centres (69.2%) and limited in 11 (28.2%). When relatives were not allowed to visit patients, updates were provided by telephone in 21 cases, whereas the remaining centres allowed a consultation in dedicated spaces. When admittance was limited, consultations with the referring physician were available in dedicated spaces preoperatively, at the end of the operation or during the entire hospital stay until discharge.

### Referral, operative volumes and surgical procedures

Patterns of referral during the pandemic significantly changed compared to those during the equivalent period in 2019 (Table 2). In particular, we observed a significant reduction in patients' admittance from the waiting lists (22.5% during lockdown vs 43.3% during the same time period in 2019) in favour of increased referral by the emergency contact (12.0% vs 9.1%), in-hospital consultations (32.7% vs 25.4%) or from peripheral hospitals (20.5% vs 16.5%).

All centres experienced a significant drop in operative procedures during the lockdown compared to 2019 (1734 procedures vs 3447;  $P < 0.001$ ), with a median reduction in cardiac surgery case volume of 53.5% (IQR 23.7–81.4%). Notably, the percentage



**Table 2:** Details of patterns of referrals and surgical volumes recorded during the lockdown compared to the same period in 2019 (absolute numbers and percentage)

	Lockdown 2020	2019	P-value
Referral, n (%)			<0.001
Emergency contact	161 (12.0)	221 (9.1)	
Internal to the hospital	438 (32.7)	619 (25.4)	
Peripheral hospitals	275 (20.5)	402 (16.5)	
Waiting list	301 (22.5)	1057 (43.3)	
Other	63 (4.7)	39 (1.6)	
Unknown	102 (7.6)	102 (4.2)	
Procedural volume, n (%)	1734	3447	<0.001
Elective	580 (33.4)	2420 (70.2)	
Urgent	955 (55.1)	832 (24.1)	
Emergency	166 (9.6)	155 (4.5)	
Salvage	33 (1.9)	40 (1.2)	

of elective surgery cases fell significantly in favour of urgent and emergency interventions, with 22 centres (56.4%) having stopped elective scheduling.

The distribution of aetiologies (Table 3) and the types of surgical procedures (Table 4) were different between the 2 study periods. During lockdown, we observed a percentage increase in admissions for acute coronary and aortic syndromes and for heart failure, with a consequent relative increase in coronary artery bypass graft surgery and either open or endovascular aortic procedures, whereas the percentage of valve procedures decreased significantly. Conversely, the occurrences of endocarditis or of complications of acute myocardial infarction and the number of transcatheter aortic valve implant procedures were similar.

### Length of hospital stay and patterns of discharge

Twelve centres (30.8%) reported an increase in the average post-operative length of stay (LOS) compared to the same period in 2019 ( $13.6 \pm 3.8$  vs  $8.8 \pm 2.9$  days;  $P < 0.001$ ). During the lockdown, 19 centres (48.7%) did not change their pattern of discharge, maintaining the usual standard of care. On the other hand, 16 centres (41.0%) preferably discharged home and 3 centres (7.7%) continued to send patients directly to a rehabilitation facility.

## DISCUSSION

This multicentre national survey clearly shows the unprecedented impact on cardiac surgery practice of the current COVID-19 pandemic. Excluding those few countries (such as China, Vietnam, Singapore and Canada), which faced the 2003 severe acute respiratory syndrome outbreak, most Western countries were found woefully unprepared both culturally and in terms of facilities and equipment [7, 8]. As mentioned previously, the number of hospital beds in Italy is chronically lower than the European average and has been further reduced by healthcare cost reduction policies since 2012 [2].

### Resource allocation

Usually, cardiac surgery is the single largest user of ICU beds. Contingent policy responses to the COVID-19 outbreak had to be organized along 3 key priorities: staff, supplies and facilities.

**Table 3:** Details of aetiologies at hospital admittance during lockdown compared to the same period of 2019 (absolute numbers and percentage)

Aetiologies	Lockdown 2020, n (%)	2019, n (%)
Valvulopathies	391 (22.2)	1094 (30.9)
Acute coronary syndromes	348 (19.8)	602 (17)
Chronic coronary artery disease	360 (20.4)	802 (22.7)
Myocardial infarction complications	13 (0.7)	19 (0.5)
Endocarditis	76 (4.3)	158 (4.5)
Acute aortic syndromes	91 (5.2)	100 (2.8)
Prosthetic failure	30 (1.7)	40 (1.1)
Heart failure	84 (4.8)	110 (3.1)
Rescue	45 (2.6)	46 (1.3)
Other	324 (18.4)	567 (16.0)

**Table 4:** Details of surgical procedures performed during lockdown compared to the same period in 2019 (absolute numbers and percentages)

Surgical procedures	Lockdown 2020, n (%)	2019, n (%)
CABG	598 (35.4)	1178 (32.8)
Valve replacement	289 (17.1)	649 (18.1)
Valve repair	126 (7.5)	399 (11.1)
Aortic/arch	149 (8.8)	212 (5.9)
Combined	207 (12.2)	522 (14.5)
GUCH	2 (0.1)	15 (0.4)
Heart transplant	19 (1.1)	16 (0.4)
VAD	14 (0.8)	66 (1.8)
ECMO-post-cardiotomy	10 (0.6)	7 (0.2)
ECMO-HF	14 (0.8)	13 (0.4)
ECMO-respiratory	32 (1.9)	6 (0.2)
TAVI	128 (7.6)	271 (7.5)
Endovascular	19 (1.1)	20 (0.6)
Infected device	10 (0.6)	38 (1.1)
Tumours	13 (0.8)	25 (0.7)
Tamponade	40 (2.0)	81 (2.3)
Miscellaneous	20 (1.2)	74 (2.1)

CABG: coronary artery bypass graft; ECMO: extracorporeal membrane oxygenation; GUCH: grown-up congenital heart disease; HF: heart failure; TAVI: transcatheter aortic valve implantation; VAD: ventricular assist device.

Accordingly, this survey showed a significant reduction in the number of dedicated cardiac operating rooms and ICU beds along with personnel reallocation to other departments, with peaks mostly confined to northern Italy, whose regions were the first to cope with the COVID-19 outbreak. The observed reallocation is consistent with that described by a recent UK nationwide survey [9]. The effects of these rationalization strategies and the response of the cardiac surgery system are authoritatively described in a recent paper by Belluschi *et al.* [10] about the Lombardy reorganization model. As these systems-based changes were evolving, guidelines with adequate specificity to address the complexity of decision-making for safely and effectively performing and/or deferring cardiac surgery were unavailable or, at best, under development [9, 11–13]. Nevertheless, the response of the Italian cardiac surgery community was prompt, and all centres developed programme-specific policies [14].

## Screening protocols

As shown in this survey, preoperative screening for SARS-CoV-2 infection and algorithms for active surveillance of healthcare providers significantly differed from one institution to another. PPE has been an important and emotive subject during the current pandemic: Appropriate use in some clinical settings is mandatory to reduce the risk of viral transmission, but a critical shortage has been reported worldwide [15]. Limited availability of PPE has been experienced in variable degrees in Italian surgical centres, with up to 30% experiencing a critical scarcity at some time during the outbreak. A similar situation has been reported in other countries [9, 16, 17]. Modalities for access of patients' relatives to surgical wards and triage algorithms for suspected COVID-19 cardiac surgical candidates were modelled on local resource availability and varied widely. Notably, hospital LOS and the patterns of discharge were also affected by the COVID-19 pandemic. The competition for diagnostics, the scarcity of blood products, the need to maximize patient health status before discharge in times of limited access to cardiac surgery outpatient clinics, along with the scarcity of rehabilitation facilities, are among the factors that synergistically contributed to these practice changes. Usually, resource availability is not factored in the decision-making process and the choices of an individual clinical or surgical case [18].

## Surgical activity

As elegantly shown in the few available surveys, indications for cardiac surgical procedures varied widely from one country to another and even within the same nation according to local resource availability, severity of viral outbreak and local expertise [9, 16, 17]. Although no official consensus statements were published by either academic or professional cardiac surgical societies, our survey reveals an almost uniform approach to indications for surgical procedures in Italy. It largely complies with that forwarded by the reorganization statement published by the Lombardy region [10]. When looking at the nationwide case mix during the lockdown, 4 points are worthy of a comment. First, although coronary artery bypass graft surgery for acute coronary syndrome accounted for more than 30% of all performed procedures, its absolute number significantly and worryingly dropped (by nearly 50%). This unexpected finding mirrors that reported by an Italian nationwide cardiologists' survey showing a dramatic reduction in hospital admissions for acute myocardial infarction during the lockdown and an increase in inherent mortality and complication rates [19]. Second, the number of transcatheter aortic valve implant procedures appeared to be less affected by the COVID-19 outbreak. Again, this pattern is consistent with that reported by recently published surveys and probably reflects a change in the treatment algorithm for severe aortic stenosis in times of limited healthcare resources [4, 9, 16, 17]. Third, surprisingly, the number of heart transplants appeared to be substantially unaffected by the pandemic. Recent data from the Italian Ministry of Health show that, by May 2020, despite a 4.5% decrease in available donors due to the reduced number of ICU beds, the number of oppositions to organ donations during the COVID-19 outbreak dropped from 33.1% to 25.3%, and the total number of solid organ transplants is currently 1.45% lower than that observed in 2019 [20]. Notably, as reported by the Italian National Organ Transplant Centre, the

number of organ transplants in Italy was significantly higher than those in the USA, Spain and France during the most critical phase of the surge [20]. Fourth, as expected, the number of extracorporeal membrane oxygenation systems implanted for respiratory failure significantly increased 10-fold (in percentages of procedures performed), which corroborates the commitment of the cardiac surgical programmes to supporting the care of critically ill COVID-19 patients [21].

The dramatic reduction in the total adult cardiac surgery case load also deserves a comment. This finding is consistent with the findings of recently published international surveys [16, 17]. Most centres restricted cardiac surgery activity to urgent/emergency cases, and 10.2% cancelled all cases including emergencies. Such a decrease has led to obvious positive effects. Indeed, it contributed to spare the limited hospital resources, prevented the in-hospital spread of SARS-CoV-2 infection and limited the risk of performing surgery on patients with asymptomatic infection during the incubation period. The drawbacks of such practice patterns are less clear now, but the outlook is not optimistic. In fact, the current mortality rate while waiting for elective cardiac surgical procedures is still relevant [22, 23]. Moreover, an increase in the number of deaths has previously been reported by healthcare systems in the aftermath of natural disasters [24, 25]. In fact, the clearance of the backlog of cardiac surgical cases is complicated by the exhaustion of supplies and resources from the pandemic and the competition with similar needs by other medical and surgical subspecialties [6]. In this respect, our findings may effectively help policy decision-making in prioritizing healthcare resource reallocation once the healthcare emergency is over [26].

## Limitations

This study has several limitations. First, only 39.4% out of 99 centres participated in the survey, which might imply an issue of representativeness. Nevertheless, participating units were evenly distributed over the nation to account for possible differences pertaining to the spread of infection and regional availability of healthcare resources. Besides, the response rate is comparable to those reported in contemporary cardiological and surgical surveys on the practice pattern during the pandemic [9, 16, 17, 19]. Second, this survey is a snapshot of a rapidly evolving situation, focused on a restricted time span, within a context of limited resources and unsettled practice guidelines during an unprecedented pandemic. The descriptive nature of the study prevents us from making any inferences from the observed results. Third, the survey design implies an inherent potential for subjectivity. Fourth, we have assumed that historical rates of surgical procedures might provide a valuable benchmark to quantify the surgical backlog and thus intensify the need for a post-pandemic increase in the volume of cardiac surgical procedures.

## CONCLUSIONS

This nationwide survey describes the major changes in adult cardiac surgery practice imposed by the COVID-19 pandemic. The cardiac surgical network responded promptly and effectively despite the severe shortage of healthcare resources in terms of facilities, staff and PPE and despite the absence of specific guidelines. In this respect, this experience should lead to the development of

permanent systems-based plans to face possible future pandemics and may effectively help policy decision-making in prioritizing healthcare resource reallocation once this healthcare emergency is over.

## SUPPLEMENTARY MATERIAL

Supplementary material is available at *EJCTS* online.

**Conflict of interest:** none declared.

## Author contributions

**Antonino Salvatore Rubino:** Data curation; Formal analysis; Methodology; Validation; Writing—original draft. **Luca Salvatore De Santo:** Conceptualization; Investigation; Methodology; Validation; Writing—original draft. **Antonio Pisano:** Writing—review & editing. **Michele di Mauro:** Project administration; Validation. **Stefano Benussi:** Data curation; Validation. **Valentino Borghetti:** Data curation; Validation. **Alessandro Castiglioni:** Data curation; Validation. **Luigi Chiariello:** Data curation; Validation. **Andrea Colli:** Data curation; Validation. **Antonio De Bellis:** Data curation; Validation. **Carlo Maria De Filippo:** Data curation; Validation. **Ruggero De Paulis:** Data curation; Validation. **Giuseppe Di Benedetto:** Data curation; Validation. **Marco Di Eusanio:** Data curation; Validation. **Giuseppe Faggian:** Data curation; Validation. **Brenno Fiorani:** Data curation; Validation. **Pasquale Antonio Fratto:** Data curation; Validation. **Angelo Giuseppe Giuffrida:** Data curation; Validation. **Mattia Glauber:** Data curation; Validation. **Gabriele Iannelli:** Data curation; Validation. **Severino Iesu:** Data curation; Validation. **Ugolino Livi:** Data curation; Validation. **Gianluca Martinelli:** Data curation; Validation. **Massimo Massetti:** Data curation; Validation. **Pasquale Mastrobert:** Data curation; Validation. **Lorenzo Menicanti:** Data curation; Validation. **Giuseppe Minniti:** Data curation; Validation. **Fabio Miraldi:** Data curation; Validation. **Gianfranco Montesi:** Data curation; Validation. **Francesco Musumeci:** Data curation; Validation. **Francesco Nicolini:** Data curation; Validation. **Carlo Pace Napoleone:** Data curation; Validation. **Paolo Panisi:** Data curation; Validation. **Aniello Pappalardo:** Data curation; Validation. **Francesco Patané:** Data curation; Validation. **Temistocle Ragni:** Data curation; Validation. **Mauro Rinaldi:** Data curation; Validation. **Salvatore Tribastone:** Data curation; Validation. **Michele Triggiani:** Data curation; Validation. **Francesco Paolo Tritto:** Data curation; Validation. **Carlo Zebele:** Data curation; Validation. **Alessandro Parolari:** Project administration; Validation. **Gino Gerosa:** Supervision. **Marisa De Feo:** Supervision. **Italian Society for Cardiac Surgery Task Force on COVID-19 Pandemic:** Project administration.

## Reviewer information

European Journal of Cardio-Thoracic Surgery thanks Paul Modi, Marko Ivan Turina and the other, anonymous reviewer(s) for their contribution to the peer review process of this article.

## REFERENCES

- Rosenbaum L. Facing Covid-19 in Italy—ethics, logistics, and therapeutics on the epidemic's front line. *N Engl J Med* 2020;382:1873–5.
- Eurostat. Healthcare Resource Statistics—Beds. [https://ec.europa.eu/eurostat/statistics-explained/index.php/Healthcare\\_resource\\_statistics\\_beds](https://ec.europa.eu/eurostat/statistics-explained/index.php/Healthcare_resource_statistics_beds) (22 May 2020, date last accessed).
- Vergano M, Bertolini G, Giannini A, Gristina GR, Livigni S, Mistracchi G *et al.* Clinical ethics recommendations for the allocation of intensive care treatments in exceptional, resource-limited circumstances: the Italian perspective during the COVID-19 epidemic. Version 2. *Crit Care* 2020;24:165.
- Giordano A, Biondi-Zoccai G, Frati G, Bartorelli AL. Management of structural heart disease and acute coronary syndromes in the COVID-19 pandemic. *Curr Atheroscler Rep* 2020;22:29.
- Pisano A, Landoni G, Zangrillo A. Protecting high-risk cardiac patients during the COVID-19 outbreak. *J Cardiothorac Vasc Anesth* 2020;34:1698.
- Salenger R, Etchill EW, Ad N, Matthew T, Alejo D, Whitman G *et al.* The surge after the surge: cardiac surgery post-COVID-19. *Ann Thorac Surg* 2020;110:2020–5.
- Tan Z, Phoon PHY, Zeng LA, Fu J, Lim XT, Tan TE *et al.* Response and operating room preparation for the COVID-19 outbreak: a perspective from the National Heart Centre in Singapore. *J Cardiothorac Vasc Anesth* 2020;34:2331–7.
- Pisano A, Landoni G, Verniero L, Zangrillo A. Cardiothoracic surgery at the time of the coronavirus disease-2019 pandemic: lessons from the East (and From a Previous Epidemic) for Western Battlefields. *J Cardiothorac Vasc Anesth* 2020;34:2338–40.
- Benedetto U, Goodwin A, Kendall S, Uppal R, Akowuah E. A nationwide survey of UK cardiac surgeons' view on clinical decision making during the coronavirus disease 2019 (COVID-19) pandemic. *J Thorac Cardiovasc Surg* 2020;160:968–73.
- Belluschi I, De Bonis M, Alfieri O, Del Forno B, Alamanni F, Polvani G *et al.* First reorganization in Europe of a regional cardiac surgery system to deal with the coronavirus-2019 pandemic. *Eur J Cardiothorac Surg* 2020;21:58:25–29.
- Engelman DT, Lother S, George I, Funk DJ, Ailawadi G, Atluri P *et al.* Adult cardiac surgery and the COVID-19 pandemic: aggressive infection mitigation strategies are necessary in the operating room and surgical recovery. *Ann Thorac Surg* 2020;110:707–11.
- Hassan A, Arora RC, Lother SA, Adams C, Bouchard D, Cook R *et al.* Ramping up the delivery of cardiac surgery during the COVID-19 pandemic: a guidance statement from the Canadian Society of Cardiac Surgeons. *Can J Cardiol* 2020;36:1139–1143.
- Haft JW, Atluri P, Alawadi G, Engelman D, Grant MC, Hassan A *et al.* Adult cardiac surgery during the COVID-19 pandemic: a tiered patient triage guidance statement. *Ann Thorac Surg* 2020;110:697–700.
- Bonalumi G, di Mauro M, Garatti A, Barili F, Gerosa G, Parolari A; for the Italian Society for Cardiac Surgery Task Force on COVID-19 Pandemic. The COVID-19 outbreak and its impact on hospitals in Italy: the model of cardiac surgery. *Eur J Cardiothorac Surg* 2020;57:1025–8.
- Mandrola J. CoVID-19 e dispositivi di protezione individuale: qualcuno di noi morirà per la loro carenza [CoVID-19 and PPE: some of us will die because of the shortage]. *Recenti Prog Med* 2020;111:183.
- Gaudino M, Chikwe J, Hameed I, Robinson NB, Fremes SE, Ruel M. Response of cardiac surgery units to COVID-19: an Internationally-Based Quantitative Survey. *Circulation* 2020;142:300–2.
- Seese L, Aranda-Michel E, Sultan I, Morell VO, Mathier MA, Mulukutla SR *et al.* Programmatic responses to the coronavirus pandemic: a survey of 502 cardiac surgeons. *Ann Thorac Surg* 2020;110:761–3.
- Emanuel EJ, Persad G, Upshur R, Thome B, Parker M, Glickman A *et al.* Fair allocation of scarce medical resources in the time of Covid-19. *N Engl J Med* 2020;382:2049–55.
- De Rosa S, Spaccarotella C, Basso C, Calabrò MP, Curcio A, Filardi PP *et al.* Reduction of hospitalizations for myocardial infarction in Italy in the COVID-19 era. *Eur Heart J* 2020;41:2083–8.
- <http://www.trapianti.salute.gov.it/trapianti/dettaglioComunicatiNotizieCnt.jsp?lingua=italiano&area=cnt&menu=media&sottomenu=news&id=561> (22 May 2020, date last accessed).
- Bartlett RH, Ogino MT, Brodie D, McMullan DM, Lorusso R, MacLaren G *et al.* Initial ELSO Guidance Document: ECMO for COVID-19 patients with severe cardiopulmonary failure. *ASAIO J* 2020;66:472–4.
- Malaisrie SC, McDonald E, Kruse J, Li Z, McGee EC Jr, Abicht TO *et al.* Mortality while waiting for aortic valve replacement. *Ann Thorac Surg* 2014;98:1564–70.
- Seddon ME, French JK, Amos DJ, Ramanathan K, McLaughlin SC, White HD. Waiting times and prioritization for coronary artery bypass surgery in New Zealand. *Heart* 1999;81:586–92.
- Lee EE, Stewart B, Zha YA, Groen TA, Burkle FM Jr, Kushner AL. Surgical care required for populations affected by climate-related natural disasters: a global estimation. *PLoS Curr* 2016;8:ecurrents.dis.e601960a8cd66c3083d160877abfdde4.
- Chan EY, Sondorp E. Medical interventions following natural disasters: missing out on chronic medical needs. *Asia Pac J Public Health* 2007;19:45–51.
- Parolari A, di Mauro M, Bonalumi G, Barili F, Garatti A, Carretta G *et al.* Safety for all: coronavirus disease 2019 pandemic and cardiac surgery: a roadmap to 'phase' 2. *Eur J Cardiothorac Surg* 2020;58:213–16.