



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## Estimating Microbial Growth and Hydrogen Consumption in Hydrogen Storage in Porous Media

**Citation for published version:**

Thaysen, E, McMahon, S, Strobel, GJ, Butler, I, Heinemann, N, Ngwenya, B, Wilkinson, M, Hassanpouryouzband, A, McDermott, C & Edlmann, K 2021, 'Estimating Microbial Growth and Hydrogen Consumption in Hydrogen Storage in Porous Media', *Renewable and Sustainable Energy Reviews*, vol. 151, 111481. <https://doi.org/10.1016/j.rser.2021.111481>

**Digital Object Identifier (DOI):**

[10.1016/j.rser.2021.111481](https://doi.org/10.1016/j.rser.2021.111481)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Renewable and Sustainable Energy Reviews

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# Estimating Microbial Growth and Hydrogen Consumption in Hydrogen Storage in Porous Media

*Eike M. Thaysen<sup>1\*</sup>, Sean McMahon<sup>1,3</sup>, Gion J. Strobel<sup>2</sup>, Ian B. Butler<sup>1</sup>, Bryne Ngwenya<sup>1</sup>, Niklas Heinemann<sup>1</sup>, Mark Wilkinson<sup>1</sup>, Aliakbar Hassanpouryouzband<sup>1</sup>, Chris I. McDermott<sup>1</sup>, Katriona Edlmann<sup>1</sup>*

<sup>1</sup>School of Geoscience, Grant Institute, The King's Buildings, The University of Edinburgh, James Hutton Road, Edinburgh, EH9 3FE, United Kingdom

<sup>2</sup>Department of Petroleum Engineering, Clausthal University of Technology, Germany

<sup>3</sup>School of Physics and Astronomy, James Clerk Maxwell Building, University of Edinburgh, EH9 3FD, United Kingdom

## ABSTRACT

Subsurface storage of hydrogen, e.g. in depleted oil and gas fields (DOGF), is suggested as a means to overcome imbalances between supply and demand in the renewable energy sector. However, hydrogen is an electron donor for subsurface microbial processes, which may have important implications for hydrogen recovery, gas injectivity and corrosion. Here, we review the controls on the three major hydrogen consuming processes in the subsurface, methanogenesis, homoacetogenesis, and sulfate reduction, as a basis to estimate the risk for microbial growth in geological hydrogen storage. Evaluating our data on 42 DOGF showed that five of the fields may be considered sterile with respect to hydrogen-consuming microorganisms due to temperatures >122 °C. Only three DOGF can sustain all of the hydrogen consuming processes, due to either temperature, salinity or pressure constraints in the remaining fields. We calculated a

---

\* = corresponding author email, phone number: [eike.thaysen@ed.ac.uk](mailto:eike.thaysen@ed.ac.uk), +44 7500533239

potential microbial growth in the order of  $1-17 \times 10^7$  cells  $\text{ml}^{-1}$  for DOGF with favorable conditions for microbial growth, reached after 0.1-19 days for growing cells and 0.2-6.6 years for resting cells. The associated hydrogen consumption is negligible to small (<0.01-3.2 % of the stored hydrogen). Our results can help inform decisions about where hydrogen will be stored in the future.

*Keywords:* Hydrogen, underground storage, microbial hydrogen consumption, homoacetogens, methanogens, sulfate reducers

Word count: 9422 words


### **Highlights**

- Review of the most important hydrogen-oxidizing microorganisms in the underground
- Elucidation of the growth criteria for 518 strains of the major hydrogen-oxidizers
- Screening of 42 depleted oil and gas fields (DOGF) for possible microbial growth
- Calculation of the microbial growth and hydrogen consumption in DOGF

### **Abbreviations and units**

SSR	Sulfur species reduction
SSRM	Sulfur species reducing microorganisms
DOGF	Depleted oil and gas fields
EPS	Extracellular polymeric substances
M	Molarity ( $\text{mol L}^{-1}$ )
MPa	Megapascal

# Graphical abstract



## 1. Introduction

Zero-carbon energy generation from renewable sources can help mitigate carbon emissions and abate climate change [1-3]. One of the most significant challenges for renewable energy is the imbalance between supply and demand [3, 4]. The generation of hydrogen (H<sub>2</sub>) via electrolysis of water during periods of renewable energy oversupply and subsequent H<sub>2</sub> storage is one way of overcoming this imbalance, as H<sub>2</sub> can be recovered and used for electricity generation during periods of renewable energy shortage [1, 5]. Subsurface storage of H<sub>2</sub> in salt caverns, depleted gas or oil fields or saline aquifers is being considered as an alternative to expensive purpose-built storage containers [6]. However, the artificial elevation of the H<sub>2</sub> concentration in the subsurface may stimulate the growth of H<sub>2</sub>-oxidizing (hydrogenotrophic) bacteria and archaea, here collectively referred to as microorganisms, with possible adverse implications for gas injectivity and withdrawal via permeability reduction, H<sub>2</sub> volume loss and corrosion of metal infrastructure [4, 7]. Understanding the controls on microbial H<sub>2</sub> metabolism is therefore highly important.

Much of the subsurface is characterized by combinations of elevated temperature [7], high salt concentrations and high pressure [3], reduced void space [8], limited nutrient availability [9] and typically highly reducing conditions [9-11]. The evidence for microbial life at depth is plentiful (e.g. [12-16]). Most microorganisms in nature grow in biofilms attached to surfaces (communities of aggregated microbial cells embedded in a secreted matrix of extracellular polymeric substances (EPS)) [17, 18]. Even small amounts of biofilm can reduce pore throat sizes and increase the flow-path tortuosity, resulting in dramatic decreases in permeability [19].

Hydrogen plays a central role in the energy metabolism of subsurface life [9]. Yet, a quantitative assessment of the consumption of H<sub>2</sub> by deep microbial communities in the context of the global H<sub>2</sub> cycle is lacking [20]. In underground gas storage sites and oil reservoirs the

most abundant H<sub>2</sub>-oxidizers are hydrogenotrophic sulfate reducers, that couple H<sub>2</sub>-oxidation to sulfate reduction to produce hydrogen sulfide (H<sub>2</sub>S); hydrogenotrophic methanogens that reduce carbon dioxide (CO<sub>2</sub>) to methane (CH<sub>4</sub>) by oxidizing H<sub>2</sub>; and homoacetogens that couple H<sub>2</sub> oxidation to carbon dioxide (CO<sub>2</sub>) reduction producing acetate [7, 21-23]. These three groups of microorganisms are, amongst others, implicated in causing subsurface corrosion [7, 22, 24].

Several recent reviews discussed the potentially very important role of microbial activity in geological H<sub>2</sub> storage [6, 7, 23, 25, 26]. Gregory et al. [7] addressed the many possible abiotic and biotic H<sub>2</sub>-producing and H<sub>2</sub>-consuming processes in the subsurface. Dopffel et al. [23] characterized different microbial issues, giving key indicators for the processes, and advised in the monitoring and management of microbial activity in subsurface H<sub>2</sub> storage. All of these efforts lacked a quantitative assessment of the processes of microbial growth and H<sub>2</sub> consumption relevant for H<sub>2</sub> storage. Strobel et al. [26] summarized the concept and potential of underground methanation using experimental data from the Sun Storage project [27]. These authors highlighted controls on the growth of methanogens and changes in gas composition due to methanogenesis, but did not quantify microbial growth. Many studies report changes in gas composition, biofilm growth and clogging near injection wells but hardly any provide quantitative figures on microbial growth or on permeability changes [28].

To date it remains unclear how subsurface microorganisms might react to elevated H<sub>2</sub> concentrations [7] and hence whether microbial growth is a concern for H<sub>2</sub> storage. Even in natural, non-engineered subsurface environments, there is little information on microbial H<sub>2</sub> turnover rates [29] and the behavior and population kinetics of microorganisms are not fully understood [26]. The majority of the available data on microbial H<sub>2</sub> turnover rates come from batch cultures at optimal growth conditions where the kinetics [29], the pace of life [30, 31],

the physiological states and the prominent organisms may differ widely from the subsurface environment [7, 30]. A further complication arises from the fact that many microorganisms in the deep subsurface are not culturable with modern enrichment techniques [12, 32].

In this work, we review the state-of-the-art understanding of the controls of temperature, salinity, pH, pressure and nutrients and water on microbial growth on H<sub>2</sub> in the subsurface, with emphasis on the three major H<sub>2</sub>-consuming processes methanogenesis, sulfate reduction and homoacetogenesis, to determine what reservoir conditions will be unfavorable to microbial activity and as such more suitable sites for long term gas storage operations of 30 years or longer, such as the UK Rough gas storage site.

Physicochemical data from 42 depleted or close to depleted oil and gas fields (DOGF) of the British and Norwegian North Sea and the Irish Sea as well as five H<sub>2</sub> storage test sites provide the base for an evaluation of the number of sites where microbial growth of methanogens, sulfate reducers and homoacetogens can be expected. Using average nutrient contents of the microbial cells and site-specific dissolved ion concentrations, we calculate significant growth and a small H<sub>2</sub> consumption for growth-permitting DOGF.

## **2. State of the art understanding**

### **2.1 Likely microbial hydrogen oxidation in hydrogen storage systems**

Hydrogen oxidizing processes may be ranked according to the magnitude of their H<sub>2</sub> threshold and their standard free energy change ( $\Delta G^{0'}$ ), two useful metrics to compare the likelihood of reactions to take place and the order at which they proceed (Table 1). The H<sub>2</sub> threshold defines the concentration of H<sub>2</sub> below which it is no longer consumed. Given all other factors are at optimum, the microbial population with the lowest H<sub>2</sub> threshold value is expected to be the most successful population in competing for H<sub>2</sub> [33].

**Table 1.** Biotic H<sub>2</sub>-consuming processes ranked according to their free energy yield ( $\Delta G^0$ ) and measured H<sub>2</sub> threshold. Not included are Vanadium, Cobalt, Technetium, Uranium and Selenium reduction, due their limited relevance for H<sub>2</sub> storage. NA= not available.

H <sub>2</sub> - oxidizing process	Reaction (number)	H <sub>2</sub> threshold (nM)	$\Delta G^0$ (KJ mol H <sub>2</sub> <sup>-1</sup> )	Typical ambient [H <sub>2</sub> ] (nmol L <sup>-1</sup> )	Relevance for H <sub>2</sub> storage
<b>Chromate reduction</b>	$\frac{1}{2}H_2 + \frac{1}{3}CrO_4^{2-} + \frac{5}{3}H^+ \rightarrow \frac{1}{3}Cr^{3+} + \frac{4}{3}H_2O$ (1)	<0.1 <sup>[34]</sup>	NA	NA	low
<b>Aerobic hydrogen oxidation (Knallgas)</b>	$H_2 + \frac{1}{2}O_2 \rightarrow H_2O$ (2)	0.051 <sup>[7]</sup>	-237 <sup>[7, 34]</sup>	NA	low
<b>Denitrification</b>	$H_2 + \frac{2}{5}H^+ + \frac{2}{5}NO_3^- \rightarrow \frac{1}{5}N_2 + \frac{6}{5}H_2O$ (3)	<0.05-0.5 <sup>[7]</sup>	-240.1 <sup>[7, 34]</sup> -224 <sup>[4, 35]</sup>	<0.05 <sup>[4, 33, 34]</sup>	low
<b>Halorespiration</b>	$H_2 + \text{halogenated compounds} \rightarrow \text{dehalogenated compounds} + HCl$ (4)	0.05-0.27 <sup>[34]</sup> <0.3 <sup>[36]</sup> 0.27-2 <sup>[7]</sup>	-230 to -187 <sup>[7]</sup>	NA	low
<b>Iron (III) reduction</b>	$H_2 + \text{ferric(oxy)hydroxides} \rightarrow \text{ferrous iron} + H_2O$ (5)	<0.11-0.8 <sup>[34, 36]</sup>	-228.3 <sup>[7, 36]</sup> -182.5 <sup>[34]</sup> -114 <sup>[4]</sup>	0.2 <sup>[4, 33]</sup> 0.2-1 <sup>[34]</sup>	intermediate
<b>Manganese (IV) reduction</b>	$2H_2 + MnO_2 \rightarrow Mn(OH)_2 + 2H_2O$ (6)	<0.05 <sup>[33]</sup>	-163 <sup>[4, 33]</sup>	<0.05 <sup>[4, 33]</sup>	low
<b>Arsenate reduction</b>	$H_2 + HAsO_4^{2-} + 2H^+ \rightarrow H_3AsO_3 + H_2O$ (7)	0.03-0.09 <sup>[34]</sup>	-162.4 <sup>[34]</sup>	0.4-0.7 <sup>[34]</sup>	low
<b>Ammonification</b>	$4H_2 + 2H^+ + NO_3^- \rightarrow NH_4^+ + 3H_2O$ (8)	0.015- 0.025 <sup>[36, 37]</sup>	-150 <sup>[4, 36]</sup>	<0.05 <sup>[4, 33]</sup>	low
<b>Fumarate reduction</b>	$H_2 + \text{fumarate} \rightarrow \text{succinate}$ (9)	0.015 <sup>[36, 37]</sup>	-86.2 <sup>[36]</sup>	NA	low
<b>Hydrogenotrophic sulfate reduction</b>	$4H_2 + SO_4^{2-} + H^+ \rightarrow HS^- + 4H_2O$ (10)	1-15 <sup>[36, 37]</sup>	-38 <sup>[7, 36]</sup> -48 <sup>[34]</sup> -57 <sup>[4]</sup>	1-2 <sup>[4, 33]</sup>	high
<b>Hydrogenotrophic methanogenesis</b>	$4H_2 + CO_2 \rightarrow CH_4 + 2H_2O$ (11)	0.4-95 <sup>[36-38]</sup>	-34 <sup>[4, 36]</sup> -43.9 <sup>[34]</sup>	5-10 <sup>[4, 33]</sup> 7-13 <sup>[34]</sup>	high
<b>Sulfur reduction</b>	$H_2 + S \rightarrow HS^- + H^+$ (12)	2500 <sup>[7]</sup>	-33.1 <sup>[7]</sup>	NA	intermediate
<b>Homoacetogenesis</b>	$4H_2 + 2CO_2 \rightarrow CH_3COOH + 2H_2$ (13)	328-3640 <sup>[36, 37]</sup>	-26 <sup>[4, 36]</sup> -36.1 <sup>[34]</sup>	100- <sup>[4]</sup> , 117-150 <sup>[34]</sup>	high



The  $\Delta G^{0'}$  marks the thermodynamic favorability of a reaction at ambient pressure and temperature, pH 7 and 1 M of all reactants. In oligotrophic (nutrient poor) high pressure and temperature environments, the order of the  $\Delta G^{0'}$  may be used to determine which reaction is more energetically favorable. As can be seen from Table 1, more negative  $\Delta G^{0'}$  values (more available free energy) are generally accompanied by lower  $H_2$  thresholds. Not included in Table 1 are the kinetics which describe the rate of the electron transfer in the redox reaction. Abiotically, most of the  $H_2$ -oxidizing reactions are very slow but mediated by microbial enzymes the processes are catalyzed [35, 39].

The three main microbial processes with implications for  $H_2$  storage, hydrogenotrophic sulfate reduction, hydrogenotrophic methanogenesis (for simplicity from now on just referred to as sulfate reduction, and methanogenesis unless otherwise specified) and homoacetogenesis, require the highest threshold  $[H_2]$  and are among the processes with lowest  $\Delta G^{0'}$  (Table 1). Nevertheless, e.g. sulfate reduction is instantaneous in most geologic settings [40] possibly due to fast kinetics [35] and/or a relatively high availability of sulfate.

Because sulfate reducers may use the same substrates as sulfur reducers (i.e. sulfide and thiosulfate [41, 42]), they are here collectively referred to as sulfur species reducing microorganisms (SSRM) performing sulfur species reduction (SSR). Direct respiration of sulfur is limited by its low solubility ( $1.6 \cdot 10^{-7}$  M) and hence requires cell attachment to the sulfur particle [43]. However, sulfur readily reacts with sulfide formed during the reduction of sulfate to form easily metabolizable polysulphides [43, 44].

Iron (III) reduction relies on the availability of iron oxides and iron-bearing minerals such as smectite and chlorite [45, 46], as well as the availability of organic carbon, since dissimilatory iron reducing bacteria (DIRB) are strict heterotrophs, i.e. synthesize cell carbon from organic compounds [47]. Iron oxides are abundant in many sediments and aquifers [45] but are

typically not available in the carbon-rich oil fields because they have been reduced over millions of years and are not replenished [21]. Meanwhile, bacteria capable of reducing iron are frequently isolated from hydrocarbon-contaminated or oil-associated sites (reviewed in [48]). However, the mere observation of iron reduction by bacteria, which are given a DIRB enrichment medium in the laboratory, does not imply that these bacteria will reduce iron in nature. In addition, cell counts are often low to intermediate (10–100 cells ml<sup>-1</sup>) and may include non-hydrogenotrophs (e.g. [49, 50]). In non-engineered environments rich in Fe oxides and organic carbon, DIRB may have a great advantage over SSRM, methanogens and homoacetogens, due to a very high affinity for H<sub>2</sub> [45]. We evaluate this process as of intermediate relevance for H<sub>2</sub> storage in DOGF.

Many DIRB and a few SSRM can also couple H<sub>2</sub> oxidation to reduction of a variety of other trace metal oxides, e.g. MnO<sub>4</sub><sup>2-</sup>/MnO<sub>2</sub>, CrO<sub>4</sub><sup>2-</sup>, Co, SeO<sub>4</sub><sup>2-</sup>, UO<sub>2</sub><sup>2</sup>, TcO<sub>4</sub><sup>-</sup>, AsO<sub>3</sub><sup>3-</sup>, and VO<sub>4</sub><sup>-</sup> [41, 51, 52]. After Fe, the most abundant metal in sedimentary environments is Mn (~10 % of Fe abundance) [45, 51]. Due to the trace content of these compounds in the environment, their reduction has low relevance for H<sub>2</sub> storage.

Oxygen and nitrate are scarce in the subsurface [11, 21, 53, 54] and aerobic hydrogen oxidation, denitrification and ammonification hence only become significant when contamination of the aquifer occurs, e.g. by drilling fluid [55-57].

Halogenated compounds are common in aquifers, and may arise from contamination or via natural processes in sediment [58, 59]. However, the concentrations of these compounds are extremely low: In aquifers of 170-1000 m depth, chloroflourocarbons reach maximum concentrations of ≤1.1 µg L<sup>-1</sup> [59] and for pristine aquifers 0.003-0.007 µg L<sup>-1</sup> of chlorinated hydrocarbons were measured [58]. We evaluate the relevance of this process to H<sub>2</sub> storage as negligible.

Literature on the importance of anaerobic fumarate respiration using H<sub>2</sub> is scarce. Fumarate may be used as an alternative electron acceptor by SSRM [41, 60] and homoacetogens [61-64]. In the non-engineered subsurface, readily metabolizable organic matter, like fumarate, is rare [65]. Oil fields being rich in organic C compounds may contain more fumarate. Payler et al. [12] confirmed the presence of fumarate reductase, the key enzyme in fumarate reduction, in three out of five metagenomes from subsurface brines within sandstone. However, the metagenomes belonged primarily to non-H<sub>2</sub> utilizing bacteria (*Halorubrum*) and fumarate concentrations were not reported. Acknowledging the lack of data in this field, we evaluate this process as being of low relevance for H<sub>2</sub> storage.

## **2.2 Factors governing microbial growth**

Microbial growth and H<sub>2</sub> consumption rates vary with nutrient availability and environmental variables (e.g. [17, 66].) Each strain is adapted to an optimum set of nutrients and environmental conditions where potentially the greatest growth rates occur. Beyond the optimum conditions, organisms may grow but at reduced rate or they become dormant. In this section, we discuss the requirements for nutrients and water, and the overall impact of temperature, salinity, pH and pressure on the growth of the major microbial H<sub>2</sub>-oxidizers in DOGFs, in the ranges relevant to H<sub>2</sub> storage. The specific activity of microbial strains grown at optimum conditions varies as well (reviewed in [67]) but the elucidation of differences between strains is beyond the scope of this review.

### **2.2.1 Nutrients**

The nutrient requirements of H<sub>2</sub>-oxidizing microorganisms are poorly elucidated. Often, only a limited number of single strains within each diverse metabolic group have been investigated, which are unlikely to be representative of all strains. Below we summarize the few knowns.

Apart from water of sufficient thermodynamic activity (see section 2.2.4), hydrogenotrophs require H<sub>2</sub> as a source of electrons (energy), an electron acceptor and a carbon source for cell division, together with a set of macro and trace elements as well as various organic nutrients [68]. Microorganisms can only access H<sub>2(aq)</sub> and hence the solubility of H<sub>2(g)</sub> is of direct relevance for all H<sub>2</sub>-consuming reactions. Given a gas phase of ~100 % H<sub>2</sub> in an H<sub>2</sub> storage system, the equilibrium solubility of H<sub>2</sub> exceeds the highest threshold value of an H<sub>2</sub>-consuming microorganism of 3.6 μM (Table 1) by ~3 orders of magnitude at ambient pressure and temperature and under static conditions (Fig. A.1a), with further increase at higher pressures (Fig. A.1b and c). While under non-static conditions hydrogenotrophs will consume part of the H<sub>2</sub>, these figures suggest no limitation by the H<sub>2</sub> solubility on microbial growth under H<sub>2</sub> storage conditions.

Elemental requirements include the macro elements C, N, H, P, Ca, Mg, S and Fe (>95 % of the microbial cell dry weight), and the trace elements Co, Mn, Ni, Mo, Cu, Zn, W as well as Se for some metabolic groups [69, 70]. For optimum growth, many microorganisms additionally require different vitamins (e.g. lipoic acid, biotin, riboflavin, folic acid, thiamine, etc.), yeast extract, coenzyme M, aromatic acids and phospholipids or a combination of these (e.g. [8, 63, 71-73]).

Nutrients may be assimilated from the solution or directly from minerals (e.g., [74-77]), the latter being of particular importance in oligotrophic environments [75]. Carbon, sulfur, phosphorous and iron are amongst the key elements released by mineral weathering [75]. The extent to which subsurface microbial communities depend on mineral weathering is unknown [75]. For soils, Huang et al. [78] analyzed that >50 % of the 1100 microbial strains were capable of mineral weathering, as tested by their ability to mineralize biotite.

Microbial cell carbon may be assimilated from CO<sub>2</sub> alone (autotrophy) or from organic carbon compounds (heterotrophy) [79]. Methanogens and homoacetogens can grow autotrophically or heterotrophically, and several can grow mixotrophically (e.g. [64, 80, 81]). SSRM typically grow heterotrophically but some grow autotrophically or mixotrophically [82, 83]. Nitrogen may be assimilated from ammonia and nitrate or by nitrogen-fixation (diazotrophy). Diazotrophy is common amongst SSRM, methanogens and homoacetogens [84-87], though homoacetogens often inhabit ammonia-rich environments [86].

Little is known about the differences in the nutrient requirements on the level of functional groups and the variation in nutrient requirement within a functional group. SSRM have a higher requirement for iron ( $1.8 \times 10^{-6}$  M) than is usually observed for microorganisms [88] while methanogens have a higher requirement for sulfur with optimal levels ranging from 0.03 to 0.79 mM (reviewed in [89]).

Literature on when nutrients become limiting is also scarce. Sulfate reduction may occur down to 5-77  $\mu$ M sulfate [90, 91]. Specific data on the phosphorous requirement of the major H<sub>2</sub>-oxidizing microbial groups are outdated/lacking but research on other extremophiles indicates that phosphorous concentrations as low as 1.7  $\mu$ M may be sufficient for growth [92]. Methanogens of the order *Methanosarcinae* require 29.6 mM Mg for optimum growth and growth ceases at 16.5 mM (reviewed in [89]). When grown under optimum conditions, the growth rate of autotrophs may be limited by the rate of transfer of H<sub>2</sub> and CO<sub>2</sub> from gas to liquid, as was shown for the methanogen *Methanobacterium thermoautotrophicum* [93] and for the sulfate reducers within *Desulfotomaculum sp* [94].

Carbon is unlikely to be limiting in the hydrocarbon-rich DOGF [54, 95, 96] but this is not a given in saline aquifers with no history of oil or gas. Sulfate is present in significant concentrations in most DOGF (Table 2) but H<sub>2</sub> injection can cause sulfate depletion due to

accelerated growth of SSRM [97]. Nitrogen in the form of the preferred N-source, ammonium [98], may be limiting in DOGF [49, 54, 88] but nitrate levels may be elevated [49], often due to contamination by drilling fluid [55-57].


### ***2.2.2 Temperature***

Temperatures of 22.5–80 °C or 20–100 °C have been suggested for H<sub>2</sub> storage based on a recommended depth range of 500- 2000 m for H<sub>2</sub> storage in DOGF and saline aquifers [99-101]. Microorganisms are classified according to their preferred growth temperature: psychrophiles grow optimally below 20 °C, psychotrophs grow optimally at or above 20 °C and may tolerate temperatures below 5 °C, mesophiles grow between 20 and 45 °C, thermophiles grow above 45-50 °C, and hyperthermophiles show optimal growth at temperatures of 80 °C or above [102, 103]. The upper limit for life is 121-122°C [104, 105].

High temperatures alter the energetic properties (e.g., vibrational modes) of biomolecules in their aqueous solvent, change the substrate solubility or viscosity and the ionization of the aqueous medium [106]. Adverse effects of high temperature include DNA denaturing or damage, decreased protein stability, hydrolysis of ATP and ADP, amongst others [104, 106]. The metabolic strategies of thermophiles are highly diverse. For a discussion, the reader is referred to [107].

Thermophiles and hyperthermophiles are challenged by increased reaction rates at elevated temperature which can imply that abiotic reaction rates are so fast that there is no benefit to the microorganism if it catalyzes the reaction [39]. High-temperature-adapted microorganisms are therefore thought to produce enzymes with faster reaction rates [108].

Most cultivated hydrogenotrophic methanogens are mesophiles but known optimal growth temperatures for methanogens range from 15 to 98 °C (Fig. 1a). A considerable number of



**Figure 1.** Distribution of optimum growth temperature, critical growth temperature, optimum pH values and critical salinity for 101-143 methanogens (a-d), 19-88 homoacetogens (e-h) and 165-277 sulfur species reducing microorganisms (SSRM) (i-l). Distributed between the graphs for the different groups of H<sub>2</sub>-oxidizers are the temperatures, pH values and salinities of 42 depleted oil and gas fields (DOGF) and five test sites for H<sub>2</sub> injection. Where ranges of a parameter were given (see Table A.4), the lower end value was plotted.

methanogens favor temperatures above 60 °C (Fig. 1a). The highest temperature that a methanogen was found to grow under is 122 °C (*Methanopyrus kandleri*) (Fig. 1b) [105].

Cultivated SSRM typically have optimum growth temperatures of 20-30 °C or 50-70 °C where sulfur reducing archaea have higher optimum growth temperatures than sulfur and sulfate reducing bacteria. The full range for optimum growth of SSRM spans 10-106 °C (Fig. 1i). The critical temperature for growth of cultivated SSRM is 113 °C (*Pyrolobus fumarii*) [109].

Homoacetogens typically have optimum growth temperatures between 20-30 °C (85 % of the here gathered cultivated strains; Fig. 1e). Thermophilic growth temperatures  $\geq 60$  °C have been reported for eight strains, only (e.g. *Moorella mulderi*, *Thermoanaerobacter kivui*, *Acetogenium kivui*) [110-112]. Corresponding upper limits for growth are 70-72 °C (Fig. 1f) [110-112].

### **2.2.3 Salinity**

The relevant salt concentration range for H<sub>2</sub> storage is 0-5 M NaCl [100], at which highly diverse prokaryote communities can be found [113]. Microorganisms are classified according to their salt tolerance: Non-halophilic microorganisms grow up to 0.2 M NaCl, slight halophile grow at 0.2–0.5 M NaCl, moderate halophile between 0.5–2.5 M NaCl, and extreme halophile that grow best in hypersaline media containing 2.5–5.2 M NaCl [113].

High salt concentrations exert osmotic stress [114], requiring any microorganism living at high salt concentrations to maintain its intracellular environment at least isosmotic with the environment [113]. Two main strategies to achieve osmotic balance exist: the salt-in strategy, and at the exclusion of salt and biosynthesis/accumulation of organic ‘compatible’ solutes [115]. For a discussion of these strategies in relation to different metabolic pathways, the reader is referred to [115]. Commonly, salt tolerance/requirement is enhanced at increased temperatures [113] but there are many examples of mesophilic halophiles.



Most cultivated hydrogenotrophic methanogens favor salt concentrations up to 0.77 M NaCl (the approximate salinity for seawater) but 16 known strains survive under more halophilic conditions. Two extremely halophilic mesophilic hydrogenotrophic methanogens, will tolerate salt concentrations of ~3.3- 3.4 M, *Methanocalculus halotolerans* FRIT [116] and *Methanocalculus natronophilus* [117] (Fig. 1d).

The large majority of cultivated SSRM grow optimally at low salinities between >0-0.4 M. However, fourteen SSRM (all mesophiles) have upper salinity limits for growth of  $\geq 1.7$  M NaCl (Fig. 1h). *Desulfovibrio oxyclinae*, *Desulfohalobium utahense* and *Desulfohalobium retbaense*, have the highest upper salinity limits for growth of 4.0 to 4.2 M NaCl [118-120] (Fig. 1h).

The salt tolerance of homoacetogens is poorly investigated. The majority of cultivated homoacetogens have low optimum salinities of >0-0.4 M NaCl. However, a few strains, i.e. *Natroniella acetigena* and *Acetohalobium arabaticum*, grow optimally around 2.5 M NaCl and will tolerate salinities up to 4.3-4.4 M (Fig. 1h) [121, 122]. The upper growth temperatures for these strains are 42 and 47 °C, respectively [121, 122].

A clear upper salinity limit to microbial activity has not been established [23, 115]. It appears to be the brine composition, rather than the salinity alone, that puts a hard limit on microbial growth [12], see section 2.2.4. Salt tolerances based on activity measurements from natural microbial communities match results from laboratory studies on cultivated microorganisms for most metabolic pathways [115]. For sulphate reduction, however, activity measurements of natural microbial communities (using any available electron donor) indicate an upper salinity limit of 4.7-8.1 M NaCl [123-125].

#### **2.2.4 Brine complexity**

Natural brines contain dissolved ions whose interaction is extremely complex and may cause physicochemical stressors to brine habitability such as low water activity ( $a_w$ ), high ionic strength, chaotropy (ability to disrupt the network of H<sub>2</sub>-bonds between water molecules) or a combination of these [12]. Most bacteria grow well at an  $a_w$  around 0.98 (the approx.  $a_w$  for sea water) but relatively few species can grow at  $a_w$  of 0.96 or lower [126]. Halophilic microorganisms, including halophilic methanogens are one exception; several can grow at  $a_w$  as low as 0.75 [127] in [126]; [128]. Steinle et al. [129] challenged these limits by detecting SSR in a nearly MgCl<sub>2</sub> saturated brine with  $a_w$  of ~0.4.

There are indications of a more important role of chaotropy over  $a_w$  in limiting microbial life [128]. Chaotropic agents include MgCl<sub>2</sub>, CaCl<sub>2</sub>, FeCl<sub>3</sub>, KI, LiBr, LiCl while examples of kosmotropic agents are NaCl, KCl, Na<sub>2</sub>SO<sub>4</sub>, MgSO<sub>4</sub>, K<sub>2</sub>SO<sub>4</sub>, FeSO<sub>4</sub> [130]. As such one may speculate that most subsurface brines due to their dominance of NaCl and richness in sulfate are kosmotropic and albeit also stress-inducing, more permissive of microbial growth [12, 128]. Meanwhile, the interactions between chao- or kosmotropic agents,  $a_w$  and other physicochemical properties of brines may be very complex and are hitherto not understood [12]. The further elucidation of this topic is subject to more research and beyond the scope of this paper.

### **2.2.5 pH**

The brine pH may affect the growth of microorganisms via 1) a direct effect on the growth metabolism, and 2) an effect on the redox reaction. With respect to the former, most methanogens, homoacetogens and SSRM are adapted to a pH of 6.5-7.5 (Fig. 1c, g, k). Most methanogens and SSRM cannot grow outside the pH range 4–9.5 [26, 131, 132] (Fig. 1c, k). Eight known methanogens can endure a critical pH-value of 10 (e.g. the *Methanocalculus natronophilus* and *alkaliphilus* [133]). At the other end of the spectrum, ten known

methanogens can endure acidic conditions of pH 4, e.g. the *Methanoregula boonei*, the *Methanothermococcus okinawensis*, the *Methanosarcina spelaei* and the *Methanocaldococcus bathoardescens* [134-137].

Eighteen known SSRM are adapted to highly alkaline environment >pH 10, e.g. the *Desulfonatronovibrio hydrogenovorans*, the *Desulfurispira natronophila* and the *Desulfovibrio vietnamensis* [138-140]. Thirteen known SSRM grow down to a pH of 4. Nine known SSRM, all of them sulfur reducers, grow down to a pH of 1, e.g. the *Thiobacillus caldus*, the *Sulfolobus acidocaldarius*, the *Acidianus infernus* and *brierleyi*, and the *Stygiolobus azoricus* [141-144].

Six known homoacetogenic strains have high critical pH values of 10.0-10.7, i.e. *Clostridium ultunense*, *Natroniella acetigena*, *Fuchsiella alkaliacetigena* and *ferrireducens*, *Peptostreptococcus productus B-52* and *Moorella sp HUC22-1* [145-149]. The *Clostridium drakai*, *ljungdahlii*, *scatologenes*, *coccoides* and *thermoautotrophicum* are the most acidophilic known strains; they can tolerate pH as low as 3.6-4.5 [150-154].

### **2.2.6 Pressure**

Pressure ranges for H<sub>2</sub> storage of 5-20 MPa [99] or 1-50 MPa [100] have been reported. Life at high pressure requires homeostatic changes [103]. The high pressures encountered in pore spaces in the crust are generally less inhibitory to microbial cellular activity than the high temperatures, partly because of the relatively high osmotic pressure of cytoplasm [102], in particular in thermophiles and hyperthermophiles [39]. Membrane fluidity, and DNA and protein synthesis are among the most pressure-sensitive cellular components and processes [103, 155, 156]. Different adaptive mechanisms and strategies are used by microorganisms to thrive in high-pressure environments, including efficient expression and activity of proteins

used in protein folding complexes (prefoldins), membrane fluidity maintaining, robust biocatalysts [156], and EPS [18] or spore formation [103].

An upper pressure limit to microbial life has not been established [23]. At 30-50 MPa, the growth of various mesophilic, atmospheric-pressure-adapted microorganisms is inhibited [155] whereas pressure effects are generally favorable for the growth of hyperthermophiles; above 100 °C, elevated pressures are required to maintain a liquid environment [104]. Microorganisms that grow optimally at 10 MPa or above are obligate and facultative piezophiles, where the former do not tolerate ambient pressure and the latter do [103]. A recent publication listed all identified piezophiles and grouped them according to their growth temperature optimum [103]. The list of species is rather short (and as we find incomplete despite being published in 2020), possibly due to the fact that, to date, it has not been possible to isolate genes associated with piezophily, so the effects of pressure on any particular organism can only be determined empirically [103, 156]. Empirical efforts however, do not commonly include pressure tolerance in the description of the environmental growth constraints of a microorganism. In addition, most mesophiles and thermophiles from habitats with pressures of <50 MPa will grow in enrichment cultures incubated at atmospheric pressure [32]. The large majority of identified cultivated piezophiles are psychrophiles (27 strains) [103], the relevance of which is low to our study. Only four mesophilic strains were reported, three of them hydrogenotrophic sulfate reducers (the *Desulfovibrio profundus*, *piezophilus*, and *hydrothermalis*), growing optimally at 10-40 MPa [103]. Eight thermophiles were identified, including one hydrogenotrophic methanogen, *Methanococcus thermolithrophicus*, growing optimally at 50 MPa. The hyperthermophilic group hosts the hydrogenotrophic *Methanopyrus kandleri* and *Methanocaldococcus jannaschii* growing optimally at 20 to 75 MPa, respectively. Examples of hydrogenotrophic piezophiles that are not included in [103] are the mesophilic SSRM *Paracoccus pantrotrophus* and *Pseudodesulfovibrio indicus* which growth optimally at

30 and 10 MPa, respectively [157, 158], and the thermophilic SSRM *Piezobacter thermophilus* and *Archaeoglobus fulgidus* TF2 which grow optimally at 30 and 42 MPa, respectively [73, 159].

A temperature dependence of the pressure response was reported for the SSRM *Desulfovibrio indonesiensis* which has similar growth rates at high and ambient pressure 45 °C but reduces its growth rate at 20 °C and 30 MPa relative to at 0.1 MPa [160]. Elevated pressure may increase the maximum growth temperature by 2-12 °C relative to lower pressure (0.1-3 MPa) [104, 105, 161].

### **2.2.7 Inhibitors**

Exposure to hydrogen sulfide, H<sub>2</sub>S, and its bisulfide ion, HS<sup>-</sup>, causes damage to microbial proteins and coenzymes [89, 162]. It remains unclear whether H<sub>2</sub>S or HS<sup>-</sup> is responsible for the toxicity effect but there is general consensus that H<sub>2</sub>S can penetrate the microbial cell membrane more easily than HS<sup>-</sup> [162]. Hydrogen sulfide dissociates with a pK<sub>1</sub> of 6.99 at 10 MPa and 25 °C to form >99 % HS<sup>-</sup> at pH 8.5 [163].

Growth of SSRM and methanogens is adversely affected at concentrations of H<sub>2</sub>S >3.8-4.0 mM [164-166]. At 5.0-6.3 mM H<sub>2</sub>S growth is completely inhibited for SSRM [164, 166], without however stopping all metabolic activity [164]. For methanogens and homoacetogens 3.8-7.5 mM H<sub>2</sub>S and total sulfide concentrations of 3.3 mM, respectively, stop the growth [162, 166]. In systems with circumneutral pH and ferric ion concentrations above 1 mM, the concentrations of H<sub>2</sub>S are predicted to be kept below toxic levels due to its precipitation in makinawite [44].

Carbon dioxide pressure above 1 bar can be toxic for microorganisms as shown for the SSRM *Desulfotomaculum geothermicum* and the methanogen *Methanothermococcus*

*thermolithotrophicus* [167]. For many anaerobes like methanogens and homoacetogens, oxygen is toxic too [62, 102].

Nitrate inhibits homoacetogenesis [168], and ammonium [169] and sulfate inhibit methanogenesis (reviewed in [170]), with minimum inhibitory concentrations varying depending on the environment [166, 169]. For instance, sulfate concentrations as low as  $2 \cdot 10^{-4}$  M were shown to inhibit methanogenesis for 10 hours in lake sediments, possibly by competition with SSRM for available  $H_2$  and C-substrate [165] (see section 2.3.9). Under  $H_2$  storage conditions however, sulfate is likely not to affect methanogenesis, because sulfate inhibition was shown to be reversed by addition of  $H_2$  [165]. For a discussion of an inhibitory effect of  $H_2$ , see section 2.5.

### **2.2.8 Summary of environmental growth constraints**


Acknowledging the lack of data for the pressure sensitivity of many microorganisms [103], and considering a general abundance of nutrients in DOGF (Table 2), we evaluate temperature and salinity as the most crucial environmental factors constraining the growth of homoacetogens, methanogens and SSRM in DOGF. Pressures encountered in the crust are documented to have less effect than temperature on microbial cellular activity, particularly in thermophiles and hyperthermophiles [39, 102]. The pH does not pose a similar constraint to the growth of homoacetogens, methanogens and SSRM because the pH ranges for growth typically span two to three pH units (not shown) and for most species they comprise the typical aquifer pH values of 6-7 [171] (Table A.4). Brine complexity and inhibitors were not included in this analysis due a lack of information on the brine composition of DOGF beyond a limited set of dissolved ions.

Figure 2 shows the critical temperature versus critical salinity for 287 cultivated strains and reveals that salt tolerances up to 1-1.7 M are widely distributed over the entire temperature

**Table 2.** Reservoir conditions for depleted, or soon to be depleted oil and gas fields. Except where otherwise indicated, the data are from [172].

⊗= reference [171]. The salinity was calculated from the chloride concentration and the concentrations of dissolved N<sub>2</sub> was estimated from the mol percentage in the gas phase, neglecting any effect of salinity. NA= not analyzed. See Table A.4 for extended data.

Field name	Area (Km <sup>2</sup> )	P (MPa)	Temp (°C)	Salinity (M)	pH	HCO <sub>3</sub> <sup>-</sup> (mM)	N <sub>2</sub> (mM)	SO <sub>4</sub> <sup>2-</sup> (mM)	K <sup>+</sup> (mM)	Ca <sup>+2</sup> (mM)	Mg <sup>+2</sup> (mM)	P (mM)	Na <sup>+</sup> (mM)	Cl <sup>-</sup> (mM)	Fe <sup>+2</sup> (mM)	Organic acids (mM)
<b>Frigg</b>	100	19.5	61	0.07-0.53	6.5-7.4	16.3	0.4	NA	26.3-31.2	0.4-2.0	1.9-7.1	NA	75.2-534.8	58.7-490.3	0.04-0.27	NA
<b>Hamilton</b>	15	9.6	30	1.59-4.18	5.8	4.8	2.1	0.6-7.4	8.4-29.7	72.8-720.0	19.5-37.6	0.012-0.028	1354.8-2210.9	1453.3-3700.7	4.03	NA
<b>Barque</b>	36	26.0	79	4.83	4.7	0.3	0.8	3.5	42.2	535.0	156.8	NA	2920.4	4405.4	2.15	NA
<b>Hamilton North</b>	8	10.5	30	2.93	7.9	11.0	2.3	23.1	18.8	13.6	13.6	NA	2640.9	2662.9	NA	NA
<b>Miller</b>	45	49.3	121	1.61	7.2	NA	0.6	0.0	41.6	30.0	NA	NA	1358.7	1471.9	0.02	NA
<b>Beryl</b>	49	36.0	101	1.88	6.1	5.6	0.4	0.0	20.8	90.0	NA	NA	1469.6	1717.9	0.05	1.9
<b>Judy</b>	NA	46.9	137	0.14-0.15	6.8	8.4	0.6	6.4	2.9	4.5	NA	0.002	117.4	131.7	0.11	NA
<b>(Andrew 1)</b>																
<b>Amethyst</b>	97	27.9	88	4.45	5.6	1.0	1.6	3.7	33.2	521.5	148.5	0.452	2673.9	4064.6	2.51	NA
<b>Rhyl</b>	NA	14.9	36	5.80	5.5	13.5	2.8	14.0	62.4	147.0	21.2	0.031	4777.0	5297.9	0.81	>1.2
<b>Dalton</b>	NA	28.8	91	0.26	5	0.9	1.0	1.8	15.6	5.5	5.5	NA	189.1	237.0	0.00	NA
<b>Davy</b>	6	28.2	88	3.87	6.8	6.5	NA	7.0	219.2	15.6	10.7	0.155	818.3	1142.7	0.66	NA
<b>Veslefrikk<sup>⊗</sup></b>	NA	29.8-35.0	67-114	0.29-0.72	6.5	8.4-17.2	NA	0.1-0.15	NA	NA	NA	NA	298.0-666.0	281.0-745.0	NA	2.2-8.1
<b>Average</b>						7.9	1.1	5.2	44.7	166.8	42.2	0.113	1473.7	1857.4	0.97	3.3



**Figure 2.** Critical temperature (without salinity stress) versus critical salinity (without temperature stress) for methanogens, homoacetogens and SSRM.

range while salt tolerances  $>1.7$  M are mainly found at a critical temperature tolerances of 40-50 °C. Hence, from the point of view of minimizing microbial impacts on  $H_2$  storage, sites with temperatures  $>55$  °C and salinities  $>1.7$  M are preferred.

Growth of all cultivated strains in the investigated microbial groups occurs up to 72 °C (Fig. 1). Above 72 °C, known homoacetogens will not grow, and at 80-94 °C sulfate reducers cease to grow. Thirty-six cultivated SSRM and eleven methanogens have optimum growth temperature of  $\geq 80$  °C (Fig.1a and g) and will still grow, albeit at reduced rate, beyond their optimum temperatures. The maximum growth temperature for known methanogens and sulfur reducers is 122 °C and 113 °C, respectively. The upper limits for salinity and pH that allow growth of all cultivated strains of the major groups of  $H_2$ -oxidizing microorganisms are 3 M NaCl and pH 10.2, respectively. The upper pressure limit for most mesophiles is 30-50 MPa.

### 2.3 Growth regulation by competition and syntrophy



Homoacetogenic bacteria are ubiquitous in anaerobic sediments [63, 173] and often co-exist with SSRM and methanogens [15, 174], as revealed by a combination of molecular (16S RNA gene sequences) and culturing (e.g. metabolites, radiotracer) techniques. Few habitats have been identified in which homoacetogens compete with other H<sub>2</sub>-consumers (culturing studies) [173, 175]. Exceptions include a low-temperature and low-salinity petroleum reservoir where homoacetogens dominated over methanogens and SSRM (molecular study) [60], a granite groundwater at 400 m depth where cell numbers of methanogens and homoacetogens were balanced (molecular study) [81], and subsurface marine sediments where mixotrophic homoacetogenesis outperformed methanogenesis (culturing study) [80].

Kinetic advantages of SSRM and methanogens (i.e. a higher affinity for H<sub>2</sub>, expressed as a low Michaelis-Menten constant,  $K_M$ , or Monod half saturation constant,  $K_S$  (H<sub>2</sub> concentration at which growth rate reaches half maximum growth rate), and a higher maximum growth/reaction rate,  $V_{max}$  or  $\mu_{max}$  for Michaelis-Menten kinetics and Monod kinetics, respectively) were proposed as the underlying cause for the few examples of the poor competitiveness of homoacetogens [176]. Very limited information on the H<sub>2</sub> consumption kinetics of homoacetogenic bacteria is available in literature [177]. The available data show that  $\mu_{max}$  differs by one order of magnitude between strains (0.02-0.5 h<sup>-1</sup>) [4, 177]. This may or may not be lower than the  $\mu_{max}$  for SSRM 0.057-5.5 h<sup>-1</sup> [4, 38, 178] and methanogens 0.032-1.4 h<sup>-1</sup> [38, 178]. Krumholz et al. [176] showed that homoacetogens were not able to compete effectively for H<sub>2</sub> in the presence of SSRM in a subsurface sandstone ecosystem at 30 °C regardless of  $pH_2$ , and despite significant homoacetogenesis at excess H<sub>2</sub>. Findings by Berta et al. [4] for a groundwater sediment held under excess  $pH_2$  and 20 °C contrasts this as homoacetogenesis rates were up to 21 times higher than SSR.

Environmental conditions may be a crucial determinant for the competitiveness of homoacetogens, as low temperatures (~15 °C) [179, 180] and low pH values [62, 181] favor their growth over methanogens. Under excess  $pH_2$ , homoacetogenic strains with high  $\mu_{max}$  such as *Acetobacterium bakii* will outcompete methanogens [180]. The outstanding metabolic flexibility of homoacetogens for utilizing a vast variety of substrates may additionally explain why homoacetogens can compete with more specialized microorganisms like SSRM or methanogens [63, 180, 182].

As for the competitiveness of methanogens and SSRM, the  $H_2$  thresholds of methanogens may be comparable (1-15 nM) or higher (>15-95) than for sulfate reducers and significantly lower than for sulfur reducers (<<2500 nM; Table 1), indicating an advantage of sulfate reducers over methanogens and sulfur reducers in most non-engineered, low  $pH_2$  environments. In line with this, Lackner [183] recently reviewed that sulfate reducers outcompete methanogens for  $H_2$  in most studies. However, at excess  $H_2$ , methanogens and sulfate reducers would be expected to process equal shares of the in situ  $H_2$  pool [178]. Also, since concentrations of sulfate are much lower than bicarbonate in non-marine natural environments [38] (Table 2), the growth of sulfate reducers at excess  $H_2$  will be limited by the availability of their electron acceptor, making it possible for methanogens to compete [38]. As a general rule pH values below 7 favor the growth of methanogens over sulfate reducers [132]. Above pH 7.5, sulfate reducers grow faster than methanogens and would be expected to outcompete them [132].

Syntrophic relationships between different functional groups have been documented frequently (whereby the metabolic products of one group serve as substrates for the other). For example, SSRM and homoacetogens were shown to participate cooperatively in microbial induced corrosion of steel where SSRM grew on acetate produced by homoacetogenesis [68]. Substrate

provision by the co-culturing *Desulfovibrio vulgaris* enhanced growth of the dehalogen *Dehalococcoides ethenogenes* 195 by 24 % and caused three times higher dechlorination rates [184]. Syntrophy may also explain the detection of a combination of the SSRMs *Desulfovibrio* and the homoacetogens *Acetobacterium* in petroleum and subsurface CO<sub>2</sub> reservoirs [60, 185], and the presence of H<sub>2</sub>-producing heterotrophs along with methanogens in petroleum reservoirs where the latter rely on H<sub>2</sub>-transfer by the former [186].

#### **2.4 Microbial ecology in natural gas and petroleum reservoirs**

Recent years have seen a considerable effort in describing deep subsurface microbial communities, including those from gas and petroleum reservoirs. Isolated hydrogenotrophic microbes from these habitats are from the SSRM families *Hydrogenothermaceae*, *Sulfurospirillaceae*, *Rhodobacteraceae*, *Ectothiorhodospiraceae*, *Desulfomicrobiaceae*, *Peptococcaceae*, *Archaeoglobaceae*, *Desulforobacteraceae*, *Desulfobulbaceae*, *Desulfovibrionaceae*, *Syntrophobacteraceae* where the latter seven families are known with certainty to be capable of thiosulfate reduction [22, 54, 96, 97, 187-194], the *Eubacteriaceae* and *Sporomusaceae* families which host homoacetogenic strains [96, 190, 195], and the methanogen families *Methanosarcinaceae*, *Methanobacteriaceae*, *Methanomicrobiaceae*, *Methanopyraceae*, *Methanococcaceae*, *Methanocalculaceae* and *Methanosaetaceae* [97, 116, 190] in addition to uncultured microbial taxa [54, 188, 189, 194, 196]. Our collection of hydrogenotrophs (Fig. 1) lists many examples of the above microbial families, including the strain that holds the highest critical temperature for a methanogen, *Methanopyrus kandleri*. Sulphur reducing families that define the upper temperature limits for SSRM like *Thermoproteaceae* and *Pyrodictiaceae* were not reported. The cause for their absence may be a predominance of mesophilic and thermophilic sites but may also reflect a generally stronger growth of sulfate reducers over sulfur reducers in oil and gas reservoirs.

Ranchou-Peyrouse et al. [97] showed that the microbial community in 35 out of 36 subsurface wells from seven natural gas storage sites was dominated by sulfate reducers.

## **2.5 Effect of high hydrogen concentrations on the microbial metabolism and community structure**

A range of studies investigated the metabolism of methanogens at excess H<sub>2</sub> and ambient pressure, with unambiguous results. Conrad et al. [197] demonstrated that excess H<sub>2</sub> stimulated methanogenesis and growth rates in a paddy soil (species not specified). Opposed to this, results by Topcuoglu et al. [186] and Stewart et al. [198] suggest an inhibitory effect of high partial pressures of H<sub>2</sub>,  $p\text{H}_2$ , expressed as a ~10-fold drop in the growth yield (cells per mole CH<sub>4</sub>) of *Methanocaldococcus jannaschii* and a slight drop of ~0.1-0.7 h<sup>-1</sup> in the growth rate. Similar observations were made for *Methanothermobacter thermoautotrophicus* [199]. However, within the excess H<sub>2</sub> experiment, higher H<sub>2</sub> concentrations stimulated growth [186], suggesting a complex influence of  $p\text{H}_2$ . Methanogens seem to express a  $p\text{H}_2$ -dependent change in their ecological strategy, i.e. maximum growth rate vs. maximum growth yield, as a means to cope with different environmental conditions [186]. Indeed, *M. jannaschii* is capable of sensing subtle changes in dissolved H<sub>2</sub> concentration and restraining the energy-intensive growth of flagella to H<sub>2</sub>-limiting conditions whereas at excess H<sub>2</sub> cells are mostly flagella devoid [200].

Only few studies investigated microbial H<sub>2</sub> turnover at high  $p\text{H}_2$  of up to 1.5-24.8 MPa [4, 201, 202]. Methanogens (*M. jannaschii*) showed a strong inhibitory effect at high  $p\text{H}_2$  [201]. However, the authors added CO<sub>2</sub> at a pressure of at least 0.2 MPa to the hydrogen gas mixture which at  $p\text{CO}_2 > 0.1$  MPa can be toxic methanogens [167]. Hence it is not clear whether H<sub>2</sub> or CO<sub>2</sub> performed the toxic action. For homoacetogens and SSRM, the H<sub>2</sub> consumption was shown not to change in response to different  $p\text{H}_2$  of 0.1-3.5 MPa [4, 202], indicating neither

stimulation nor toxicity at different levels of excess H<sub>2</sub>. The comparison to limiting H<sub>2</sub> conditions was not made.

Apart from the microbial metabolism, the microbial community may also change in response to high  $p\text{H}_2$ . Given a perturbation by H<sub>2</sub> injection it can be anticipated that other types of microorganisms, e.g. the in hydrocarbon reservoirs, common fermenters [21, 95, 97, 187] will decrease in abundance while hydrogenotrophs will increase [7], in line with the Baas Becking principle [203]. An increase in hydrogenotrophs in response to H<sub>2</sub> addition was recently confirmed for soils, however H<sub>2</sub> consumption increased in only one of the investigated soils, suggesting a pronounced influence of the indigenous microbial community [204]. Bioreactor experiments support a decrease in microbial diversity in response to high  $p\text{H}_2$  as well [205, 206]. Puente-Sanchez et al. [207] were the first to report differences in the subsurface H<sub>2</sub>-consuming community in response to varying  $p\text{H}_2$  within the Iberian Pyrite Belt. Ranchou-Peyruse et al. [97] showed that town gas storage with more than 50 % H<sub>2</sub> changed the microbial community from a predominantly sulfate reducing community to a dominance of methanogens, and this balance was active even decades after injection stopped, possibly via H<sub>2</sub> trapping in the microporous system [97]. It was suspected that all sulfate was initially used up by SSRM following increased growth of methanogens [97].

### **3. Evaluating the potential hydrogen consumption in DOGFs**

#### **3.1 Calculation of the microbial growth**

We screened 42 DOGF in the North Sea and the Irish Sea and five H<sub>2</sub> storage test sites for temperature, salinity, pH and pressure data (Fig. 1, Table A.4). We discovered significant differences in the salinity for the DOGF reported by sources [208] and [172]. Because we relied on the solution compositions for the calculation of the potential microbial growth in the fields, which are available from [172], we chose to use the salinity data from the same source.

The environmental data from the DOGF and H<sub>2</sub> storage test sites were aligned with the constraints for growth of methanogens, homoacetogens and SSRM (Fig. 1-2) to select in which fields growth can be expected. For the few fields that fulfil the growth constraints of all investigated microorganisms, we calculated a first-order estimate of the microbial growth using the elemental cell composition as a proxy for the nutrient requirement [209, 210] (Text A.1).

Our calculations assumed that the supply of N and C are covered by diazotrophic and autotrophic growth, respectively. Requirements for trace elements were neglected in the calculation due to a lack of information on the relevant trace element contents in the reservoirs. Where a nutrient for a specific field was not available we used the average value from the fields given in Table 2. Any effect of the  $p\text{H}_2$  on microbial growth was neglected. We assumed that cells neither die nor are removed, and that nutrients are not replenished by inflow, remineralization from decaying biomass or mineral dissolution. Simultaneous growth by different microorganisms was not considered.

Percentages of nutrients in the cells (Text A.1) were converted to mass using a wet cell mass of  $1.77 \cdot 10^{-12}$  g for methanogens [211],  $3.2\text{-}6.2 \cdot 10^{-13}$  g for homoacetogens and  $7.81 \cdot 10^{-13}$  g for SSRM. The cell wet weight of homoacetogens was calculated by dividing the cell volume of  $1.62\text{-}3.14 \mu\text{m}^3$  for the subsurface mixotrophic homoacetogen *Acetobacterium psammolithicum* [176] with an assumed bacterial density of  $1 \cdot 10^{-12}$  g  $\mu\text{m}^{-3}$  [212]. The cell wet weight of SSRMs was calculated using a cell dry weight of  $3.125 \cdot 10^{-13}$  g for *Desulfovibrio desulfuricans* [213] and dividing this with a general bacterial dry weight to wet weight ratio of 0.4 [214]. Subsequently, the concentrations of C, H, O, Ca, K, Na, S, Mg, P and Fe in the DOGF (Table 2) were divided by the mass of the respective cell nutrients per microbial cell calculated above. This resulted in the maximum cell count within each microbial group,  $G$ , that could potentially

be created based on a single nutrient, where the lowest  $G$  indicated the limiting nutrient for cell growth. For an example of those calculations, see Text A.1.

### **3.2 Estimation of the cell-specific hydrogen consumption**

Hydrogen may be consumed by growing and resting microbial cells at rates of  $0.02-5.0 \cdot 10^5$   $\text{nM h}^{-1}$  for homoacetogens,  $0.02-5.8 \cdot 10^5$   $\text{nM h}^{-1}$  for methanogens and  $0.005-130 \cdot 10^5$   $\text{nM h}^{-1}$  for SSRM (Tables A.1-A.3), the latter considering sulfate concentrations in the range of  $0-2.3 \cdot 10^{-2}$  M in the DOGF (Table 2). In a few studies, the microbial  $\text{H}_2$  consumption was related to growth (Tables A.1-A.3), enabling the calculation of the  $\text{H}_2$  consumption per synthesized cell and the time for when the microbial cell count  $G$  would be reached (Text A.2).

### **3.3 Calculation of the hydrogen consumption in a hydrogen storage system**

We calculated the minimum  $\text{H}_2$  consumption for the DOGF Frigg and Hamilton by dividing the  $\text{H}_2$  consumption per synthesized cell with the microbial cell count. The calculation of the moles of  $\text{H}_2$  the in aquifer anticipated equal volumes of  $\text{H}_2$  and water and used the ideal gas law and the field size, temperature and pressure data in Table 2 and Table A.3. The percentage of  $\text{H}_2$  that was consumed as a function of growing and resting microbial cells was calculated by dividing the potential  $\text{H}_2$  consumption with the  $\text{H}_2$  concentration in the reservoir. Text A.3 shows our calculations for the Frigg reservoir and methanogens.

## **4. Results and discussion**

### **4.1 Characterization of the likelihood for growth in 42 DOGF**

Using the environmental limits constraining microbial growth on  $\text{H}_2$ , we analyzed the physicochemical parameters for 42 DOGF in the British and Norwegian North Sea and the Irish Sea and five  $\text{H}_2$  storage test sites (Fig. 1, Table A.4). Of the 47 fields, five fields have a temperature of  $122$  °C or higher and may be considered sterile with respect to  $\text{H}_2$ -consuming

microorganisms. Where long-term injection of cold sea water has been a practice, cooling of reservoirs is a likely scenario. Therefore, any H<sub>2</sub> storage operation in these fields will require a renewed measurement of the reservoir temperature. Thirty-two fields have a temperature >72 °C, implying that homoacetogenesis cannot take place. Twenty fields have a temperature ≥90 °C implying that homoacetogenesis and sulfate reduction cannot take place. Fourteen DOGF have a temperature >90 °C and <122 °C and pressures of 18.2-44 MPa where (piezophile) methanogens and SSRM will grow.

Of the fifteen sites with temperatures <72 °C where all investigated groups of microorganisms will grow, only six fields (Frigg, Hamilton, Veslefrikk, Ketzin, Lehen and Lobodice) fulfill the remaining pressure and salinity requirements for growth. Five fields, Lennox, North Morecambe and South Morecambe, Leman and Rhyl, have salinities ≥4.4 M where no cultivated microbial H<sub>2</sub>-oxidizing microorganisms can grow but not cultivable SSRM may still be active. This finding is supported by stable gas compositions at the similarly saline H<sub>2</sub>-storage test sites of the H2STORE project, Emsland and Altmark (Fig. 1, Table A.4), where low microbial populations of ~10<sup>2</sup> cells ml<sup>-1</sup> were present [215]. Hamilton North, Camelot and The V gas field complex with salinities of 2.9-5.0 M may permit the growth of SSRM and homoacetogens. The Viking field has temperature of 65-80 °C and a salinity of 3.8 M and so is likely to host only mesophilic SSRM, although pressures >30 MPa could become growth inhibiting [155]. The H<sub>2</sub>-storage test site Ketzin has similar salinity to the Viking field but a lower pressure (4.0 M NaCl, 35 °C, 6 MPa). Here SSRM were suspected to cause a 2-4 % decrease in H<sub>2</sub> and a reduction in the concentration of sulfate from 22 to 8\*10<sup>-3</sup> M [215].

#### **4.2 Microbial growth estimates for three low-temperature and low-salinity DOGF**

Our first order approach to calculating microbial growth, designed to give a first approximation to microbial numbers, only, yielded a maximum of 1\*10<sup>8</sup> methanogenic cells mL<sup>-1</sup>, 2\*10<sup>8</sup>



SSRM cells  $\text{mL}^{-1}$  or  $5 \cdot 10^8$  homoacetogenic cells  $\text{mL}^{-1}$  in the Veslefrikk reservoir. The Frigg reservoir a maximum of  $1 \cdot 10^8$  methanogenic cells  $\text{mL}^{-1}$ ,  $1 \cdot 10^8$  SSRM cells  $\text{mL}^{-1}$  or  $2 \cdot 10^8$  homoacetogenic cells  $\text{mL}^{-1}$ . The Hamilton reservoir could host a maximum of  $1 \cdot 10^7$  methanogenic cells  $\text{mL}^{-1}$ ,  $2 \cdot 10^7$  SRCM cells  $\text{mL}^{-1}$  or  $6 \cdot 10^7$  homoacetogenic cells  $\text{mL}^{-1}$ . These cell counts describe a maximum cell growth for each hydrogenotrophic group because simultaneous growth of hydrogenotrophs was not considered. The higher growth of homoacetogens over SSRM and methanogens results from a lower wet cell mass that causes a lower nutrient demand per cell (see Text A.1). Our calculations are in line with total cell concentrations of  $10^5$ - $10^{15}$  cells/  $\text{mL}^{-1}$  in oil reservoirs [216], and equal to or up to four order of magnitudes higher than cell counts from gas reservoirs ( $0.001$ - $1.2 \cdot 10^7$  cells  $\text{mL}^{-1}$ )[49, 96, 189].

Acknowledging that trace elements were not accounted for in our calculation, N and P are the first limiting nutrients in the reservoirs Frigg, Hamilton and Veslefrikk. However, this does not imply that microbial growth is N and P limited, as many microorganisms may use of ammonium (not measured) as N-source, and in the Hamilton reservoir the C:P ratio was between 59:1 and 158:1, whereas the limiting C:P ratio for microbial growth is in the range of 400:1 to 800:1 (reported for the SSRM *D. desulfuricans*) [217]. At moderately acidic pH values such as the pH of 5.8 in the Hamilton reservoir, P may further be continuously replenished by mineral buffering with apatite.

### **4.3 Hydrogen consumption in three low-temperature and low-salinity DOGF**

The  $\text{H}_2$  consumption in the Frigg reservoir by homoacetogens constitutes <0.01- 3.2 % of the  $\text{H}_2$  in the aquifer, <0.01- 1.3 % for methanogens and <0.01- 1.3 % for SSRM. In the Hamilton reservoir, the rates are <0.01- 2.0 %, <0.01- 2.3 % and <0.01- 0.5 % for homoacetogens, methanogens and SSRM, respectively. For actively growing cells these consumption rates may

be reached after only 0.1-19.1 days, which is the time it takes for the microorganisms to grow up to their maximum cell counts, based on the dissolved nutrient concentrations. Resting cells, i.e. cells that undergo no or only very little cell division, need 2.5-3.5 months (SSRM) or up to 3.6-6.6 years (methanogens) to reach the maximum cell count and consume the given percentage H<sub>2</sub>.

In a real aquifer system, nutrients are likely to at least partly be replenished by decaying cells, mineral weathering and inflowing brine, and cells will continue to consume H<sub>2</sub> beyond the time it takes to reach the maximum cell count (maintenance). As such our H<sub>2</sub> consumption estimates may be regarded as minima. On the other hand, considering that, with the exception of one study (Berta et al. [4]), our calculations employ laboratory H<sub>2</sub> consumption rates at optimal nutrient supply and optimal physicochemical conditions (Tables A.1-A.3), the H<sub>2</sub> consumption in the oligotrophic subsurface is likely overpredicted. Comparing the employed laboratory H<sub>2</sub> consumption rates to H<sub>2</sub> consumption rates by SSR and methanogenesis in oil and natural gas reservoirs of  $\sim 0.4\text{-}330\text{ nM h}^{-1}$  and  $0.02\text{-}1205\text{ nM h}^{-1}$ , respectively ( $\text{SO}_4^{2-}$ :  $8.3\text{-}805 \cdot 10^{-5}\text{ M}$ ;  $\text{HCO}_3^-$ :  $3.5\text{-}246 \cdot 10^{-4}\text{ M}$ ) [49, 187], shows that the field H<sub>2</sub> consumption by SSR is 1.5 times to five orders of magnitude lower, and 1.4 times to 7 orders of magnitude lower for methanogenesis. Within the operation and injection wells of a natural gas reservoir, H<sub>2</sub> consumption rates by SSR and methanogenesis were 2393 and 4475 nM h<sup>-1</sup>, respectively, [49], which falls within the lower range of the values reported from laboratory studies. Acknowledging the unknown but presumably low  $p\text{H}_2$  in above experiments, and that maintenance requirements were not included in our H<sub>2</sub>-consumption calculations, we expect the actual H<sub>2</sub> consumption in a H<sub>2</sub> storage system to lie within the higher range of our calculated values.

Our upper end results are in agreement with H<sub>2</sub> losses of ~3 % by methanogens and 2-4 % by sulfate reducers at the H<sub>2</sub> storage test sites in Lehen, Austria [27] and Ketzin, Germany [215], respectively. Reports from H<sub>2</sub>-rich town gas in Beynes, France, reports are contradictory ranging from no H<sub>2</sub> consumption during storage operations [218] to significant (unspecified) reductions of H<sub>2</sub> and CO<sub>2</sub> contents along with increases in CH<sub>4</sub> [219, 220]. A H<sub>2</sub> consumption of 17 % by methanogens at the Lobodice town gas storage site over a time span of seven months [218, 221] seems exceptional in the light of our calculations and the reported SSR and methanogenesis rates from the field. With a very low salinity of 0.03 M, temperatures of 20-45 °C, a pH of 6.7 and 4 MPa pressure, Lobodice is among the few sites which has highly favorable conditions for microbial growth considering *all* of these parameters (Table A.4). *The high H<sub>2</sub> consumption at Lobodice highlights the importance of environmental parameters for controlling microbial activity, as H<sub>2</sub> storage may face serious economic and technical problems if a site with growth-favoring conditions is selected.*

As mentioned, Berta et al. [4] measured high H<sub>2</sub> consumption rates under excess H<sub>2</sub> and oligotrophic conditions (P < 9.7\*10<sup>-7</sup> M; SO<sub>4</sub><sup>2-</sup> ≤ 9.5\*10<sup>-4</sup> M; DOC = 2.6\*10<sup>-4</sup> M), indicating that nutrient scarcity does not imply low H<sub>2</sub> consumption. A comparison to the nutrient concentrations in the DOGF reveals that many of them have a higher nutrient status (P = 0.002-0.452\*10<sup>-3</sup> M; SO<sub>4</sub><sup>2-</sup> = up to 23.1\*10<sup>-3</sup> M; organic acids = 1.2-8.1\*10<sup>-3</sup> M, Table 2), implying that H<sub>2</sub> consumption in DOGF under excess H<sub>2</sub> conditions may be even higher than reported in [4]. The experiment by Berta et al. [4] is further highly relevant because cells were at steady state, i.e. at the predominant growth stage in nature, but still consumed vast amounts of H<sub>2</sub>. Indeed the H<sub>2</sub> consumption of cells at steady state or resting may be just as high as or higher than for growing cells but growth is low or absent (Tables A.1-A.3).

#### **4.4 Knowledge gaps and future research**

More work is needed to predict the magnitude of microbial growth, H<sub>2</sub> consumption rates, and (not least) the mutual interaction of the microbial processes in DOGFs. The list of unknowns and uncertainties is long. To begin with are the poorly elucidated nutrient requirements of the microorganisms, especially in mixed cultures (e.g., [69]). Adding to this are the missing or incomplete datasets on the physical environment of certain reservoirs along with their gas phase and brine compositions, including chaotropy and kosmotropy characteristics. A better elucidation of the latter would allow the calculation of the dominating microbial processes via their free energies of the reaction. Combined with an analysis of the microbial community and metabolism this could give new insights into whether or not we can theoretically predict which microbial processes occur in DOGF and to which extend.

A further complication is the non-cultivability of many microorganisms in the deep subsurface, including DOGF [12, 32, 54, 97, 187]. Considering tiny culturabilities of  $\leq 0.1\%$  of the total viable cell count in many subsurface environments [32], any attempts to assign sterile habitats or quantify microbial H<sub>2</sub> consumption via cultivated microorganisms, only, are characterized by a significant uncertainty. In gas reservoirs, the percentage of cultured bacteria may be higher, ranging between 86-95% within each phylum [97]. Field-based metabolic activity measurements could circumvent any non-cultivability issues observed in laboratory experiments. Initially, however, DNA-based laboratory tests are recommended to obtain general cell numbers. The number of cultivable microbes may be maximized using a large array of modern cultivation techniques [222-227].

The lack of knowledge about the changes in microbial ecology as a response to increased H<sub>2</sub> concentrations beyond the level of functional groups is one of the major hurdles in our attempt to understand of the effect of high H<sub>2</sub> concentrations on the subsurface microbiology. Emerging evidence on the subject highlights species-specific responses to high  $p\text{H}_2$  [97, 205, 207], and

that H<sub>2</sub> injection may leave its fingerprint on the subsurface microbial community for decades [97]. Knowledge about the initial effect of a drastic increase in  $p\text{H}_2$  in the subsurface is lacking. Considering the pressure increase and the toxicity of high  $p\text{H}_2$  on methanogens [186, 199, 201], one possibility is that more EPS will be produced as a response to the perturbation with elevated H<sub>2</sub> pressures, as has been shown for other types of perturbation [18, 217, 228], with possible adverse effects on gas injectivity and withdrawal.

Future research should address the effect of high  $p\text{H}_2$  on the metabolisms of different functional groups in different geological settings and under changing nutritional supply and physicochemical conditions. Mixed culture studies at low and high  $p\text{H}_2$  can give insight into competitive and syntrophic relations under these conditions and reveal changes in the microbial community structure due to the perturbation with elevated H<sub>2</sub>. More base-line research includes determinations of the critical salinities and pressure tolerances that to date are missing for many cultivated strains, as well as the study of the brine compositional effects on the microbial community and metabolism. Future lab-based research should aim to employ chemostat studies that mimic the natural environment [17].

## **5. Conclusion**

In this work we presented the growth conditions of 518 cultivated strains from the three major groups of H<sub>2</sub>-oxidizing microorganisms and aligned those with physicochemical data from 42 DOGF in the British and Norwegian North Sea and the Irish Sea to predict where microbial growth can be expected in a future H<sub>2</sub> storage scenario. Our results can –with some certainty– exclude life in several high-temperature, i.e. deeper reservoirs. For low-salinity and low-temperature reservoirs our initial calculations indicate significant microbial growth and a small H<sub>2</sub> consumption, both of which may further increase during repeated storage cycles, giving replenishment of nutrients by mineral weathering, decaying microbial cells and inflowing

water. Hence, from the point of view of minimizing H<sub>2</sub> loss, clogging and corrosion, sites with more extreme conditions may be chosen over low-temperature and low-salinity reservoirs where the majority of microorganisms can proliferate. Yet, any storage operation will have to consider increased operational difficulties and costs with increased depth. Experimental investigations of subsurface life on H<sub>2</sub> are needed to verify our calculations and manifest whether H<sub>2</sub> consumption in low-temperature aquifers is a threat to H<sub>2</sub> storage. All sites of interest to H<sub>2</sub> storage should be carefully investigated and tested for microbial growth beforehand.

## ASSOCIATED CONTENT

**Appendix.** Figure A.1 shows the solubility of hydrogen as a function of temperature and pressure. Laboratories studies investigating homoacetogenesis, methanogenesis and SSR are listed in Tables A.1, A.2 and A.3, respectively. Table A.4 provides the reservoir conditions for 42 DOGF and five H<sub>2</sub> storage test sites. Text A.1 holds a discussion of the importance of other hydrogen oxidizing processes for hydrogen storage. A detailed calculation of the number SSRM cells that could grow based on the N content in the Frigg reservoir can be found in Text A.1. Text A.2 and A.3 hold an example calculation of the hydrogen consumption per synthesized cell and the calculation of the potential hydrogen consumption in a hydrogen storage system, respectively.

### **Data availability**

A dataset related to this article can be found at <http://dx.doi.org/10.17632/4dksb2x4zn.1>, an open-source online data repository hosted at Mendeley Data.

### **Author Contributions**

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

### **Declaration of interest**

The authors declare no competing financial interest.

### **ACKNOWLEDGEMENT**

This work was carried out for the HYSTORPOR project and was supported by the Engineering and Physical Science Research Council (EPSRC) [grant number EP/S027815/1]. I would like to thank Pete M. Higgins for excellent advice and discussions on how to estimate microbial numbers from few data.

### **LIST OF REFERENCES**

1. Kharel S, Shabani B. Hydrogen as a long-term large-scale energy storage solution to support renewables. *Energies* 2018;11(10):1-17.
2. Duan HX. The public perspective of carbon capture and storage for CO<sub>2</sub> emission reductions in China. *Energy Policy* 2010;38(9):5281-9.
3. Beckingham LE, Winningham L. Critical knowledge gaps for understanding water-rock-working phase interactions for compressed energy storage in porous formations. *ACS Sustain Chem Eng* 2020;8(1):2-11.
4. Berta M, Dethlefsen F, Ebert M, Schafer D, Dahmke A. Geochemical effects of millimolar hydrogen concentrations in groundwater: An experimental study in the context of subsurface hydrogen storage. *Environ Sci Technol* 2018;52(8):4937-49.
5. Heinemann N, Booth MG, Haszeldine RS, Wilkinson M, Scafidi J, Edlmann K. Hydrogen storage in porous geological formations - onshore play opportunities in the Midland Valley (Scotland, UK). *Int J Hydrogen Energy* 2018;43(45):20861-74.

6. Zivar D, Kumar S, Foroozesh J. Underground hydrogen storage: A comprehensive review. *Int J Hydrogen Energy* 2020;in press.
7. Gregory SP, Barnett MJ, Field LP, Milodowski AE. Subsurface microbial hydrogen cycling: Natural occurrence and implications for industry. *Microorganisms* 2019;7(53):1-27.
8. Fredrickson JK, McKinley JP, Bjornstad BN, Long PE, Ringelberg DB, White DC, et al. Pore-size constraints on the activity and survival of subsurface bacteria in a late cretaceous shale-sandstone sequence, northwestern New Mexico. *Geomicrobio J* 1997;14(3):183-202.
9. Colman DR, Poudel S, Stamps BW, Boyd ES, Spear JR. The deep, hot biosphere: Twenty-five years of retrospection. *PNAS Perspective* 2017;114(3):6895-903.
10. Lovley D, Chapelle FH. Deep subsurface microbial processes *Rev Geophys* 1995;33(3):365-81.
11. Hallbeck L, Pedersen K. Characterization of microbial processes in deep aquifers of the Fennoscandian Shield. *J Appl Geochem* 2008;23:1796-819.
12. Payler SJ, Biddle JF, Lollar BS, Fox-Powell MG, Edwards T, Ngwenya BT, et al. An ionic limit to life in the deep subsurface. *Front Microbiol* 2019;10:1-16.
13. Krumholz LR, McKinley JP, Ulrich GA, Suflita JM. Confined subsurface microbial communities in cretaceous rock. *Nature* 1997;386(6):64-6.
14. Methé BA, Nelson KE, Eisen JA, Paulsen IT, Nelson W, Heidelberg JF, et al. Genome of *Geobacter sulfurreducens*: Metal reduction in subsurface environments. *Science* 2003;302(5652):1967-9.
15. Aüllo T, Ranchou-Peyruse A, Ollivier B, Magot M. *Desulfotomaculum* spp. and related gram-positive sulfate-reducing bacteria in deep subsurface environments. *Front Microbiol* 2013;4:1-12.



16. Roh Y. Isolation and characterization of metal-reducing *Thermoanaerobacter* strains from deep subsurface environments of the Piceance Basin, Colorado. *Appl Environ Microbiol* 2002;68(2):6013-20.
17. Pedersen K. Microbial processes in radioactive waste disposal. Stockholm, Sweden; 2000.
18. Yin W, Wang Y, Liu L, He J. Biofilms: The microbial “protective clothing” in extreme environment. *Int J Mol Sci* 2019;20(3423):1-18.
19. Coombs P, Wagner D, Bateman K, Harrison H, Milodowski AE, Noy D, et al. The role of biofilms in subsurface transport processes *Q J Eng Geol* 2010;43:131-9.
20. Zgonnik V. The occurrence and geoscience of natural hydrogen: A comprehensive review. *Earth-Sci Rev* 2020;203(103140):1-50.
21. Pannekens M, Kroll L, Mueller H, Mbow FT, Meckenstock RU. Oil reservoirs, an exceptional habitat for microorganisms. *N Biotechnol* 2019;49:1-9.
22. Kleinitz W, Boehling E. Underground gas storage in porous media- operating experience with bacteria on gas quality 67th EAGE Conference and Exhibition SPE EUROPEC 13-16 June Madrid, Spain Society of Petroleum Engineers 2005 p. 1-6.
23. Dopffel N, Jansen S, Gerritse J. Microbial side effects of underground hydrogen storage - Knowledge gaps, risks and opportunities for successful implementation. *Int J Hydrogen Energy* 2021;46:8594-606.
24. Loto CA. Microbiological corrosion: mechanism, control and impact—a review. *Int J Adv Manuf Tech* 2017;92:4241-52.
25. Heinemann N, Alcalde J, Miocic J, Hangz SJT, Kallmeyer J, Ostertag-Henning, et al. Enabling large-scale hydrogen storage in porous media. *Energy Environ Sci* 2021.
26. Strobel G, Hagemann B, Huppertz TM, Ganzer L. Underground bio-methanation: Concept and potential. *Renew Sust Energ Rev* 2020;123(109747):1-11.

27. Bauer S. Underground Sun Storage. Final Report Vienna, Austria; 2017.
28. Taylor SW, Jaffe PR. Biofilm growth and the related changes in the physical properties of a porous medium 1. Experimental investigation. *Water Resour Res* 1990;26(9):2153-9.
29. Harris SH, Smith RL, Suflita JM. In situ hydrogen consumption kinetics as an indicator of subsurface microbial activity. *Fems Microbiol Ecol* 2007;60(2):220-8.
30. Hoehler TM, Barker Joergensen B. Microbial life under extreme energy limitation. *Nat Rev* 2013;11:83-94.
31. Maier RM, Pepper IL, Gerba CP. *Environmental Microbiology Second Edition* ed: Academic Press 2009.
32. Parkes RJ, Sass H. Deep sub-surface In: Schaechter M, editor. *Encyclopaedia of Microbiology* Elsevier Academic Press 2009.
33. Lovley D, Goodwin S. Hydrogen concentrations as an indicator of the predominant terminal electron-accepting reactions in aquatic sediments. *Geochim Cosmochim Acta* 1988;52:2993-3003.
34. Heimann A, Jakobsen R, Blodau C. Energetic constraints on H<sub>2</sub>-dependent terminal electron accepting processes in anoxic environments: A review of observations and model approaches. *Environ Sci Technol* 2010;44:24-33.
35. Appelo CAJ, Postma D. *Geochemistry, groundwater and pollution* second ed. Leiden A.A.Balkema Publishers 2007.
36. Loeffler FE, Tiedje JM, Sanford RA. Fraction of electrons consumed in electron acceptor reduction and hydrogen thresholds as indicators of halo-respiratory physiology. *Appl Environ Microbiol* 1999;65(9):4049-56.

37. Cord-Ruwisch R, Seitz HJ, Conrad R. The capacity of hydrogenotrophic anaerobic bacteria to compete for traces of hydrogen depends on the redox potential of the terminal electron acceptor. *Arch Microbiol* 1988;149:350-7.
38. Karadagli F, Rittmann BE. Kinetic characterization of *Methanobacterium bryantii* *M.o.H.* *Environ Sci Technol* 2005;39:4900-5.
39. Amend JP, Shock EL. Energetics of overall metabolic reactions of thermophilic and hyperthermophilic archaea and bacteria. *FEMS Microbiol Rev* 2001;25:175-243.
40. Machel HG. Bacterial and thermochemical sulfate reduction in diagenetic settings: old and new insights. *Sediment Geol* 2001;140:143-75.
41. Muyzer G, Stams AJM. The ecology and biotechnology of sulphate-reducing bacteria. *Nat Rev* 2008;6:441-54.
42. Findlay AJ. Microbial impact on polysulfide dynamics in the environment. *Fems Microbiol Lett* 2016;363:1-12.
43. Hedderich R, Klimmek O, Kroeger A, Dirmeier R, Keller M, Stetter KO. Anaerobic respiration with elemental sulfur and with disulfides. *FEMS Microbiol Rev* 1999;22:353-81.
44. Rickard D, Luther GW. Chemistry of iron sulfides. *Chem Rev* 2007;107:514-62.
45. Lovley D. Dissimilatory Fe(III) and Mn(IV) reduction. *Microbiol Rev* 1991;55(2):259-87.
46. Hermsdorf AW, Amano Y, Miyakawa K, Ise K, Suzuki Y, Ananhaman K, et al. Potential for microbial H<sub>2</sub> and metal transformations associated with novel bacteria and archaea in deep terrestrial subsurface sediments. *Nature* 2017;11:1915-29.
47. Javaherdashti R. *Microbially Influenced Corrosion . An Engineering Insight* second ed. Switzerland Springer; 2008.

48. Wiegel J, Hanel J, Aygen K. Chemolithoautotrophic thermophilic iron(III)-reducer. In: Ljungdahl LG, Adams MW, Barton LL, Ferry JG, Johnson MK, editors. Biochemistry and physiology of anaerobic bacteria. New York: Springer; 2003. p. 235-51.
49. Ivanova AE, Borzenkov IA, Tarasov AL, Milekhina EI, Belyaev SS. A microbiological study of an underground gas storage in the process of gas extraction. *Microbiology* 2007;76:461-8.
50. Slobodkin AI, Jeanthnon C, L'Haridon S, Nazina T, Miroshnichenko M, Bonch-Osmolokskaya EA. Dissimilatory reduction of Fe(III) by thermophilic bacteria and archaea in deep subsurface petroleum reservoirs of Western Siberia. *Curr Microbiol* 1999.
51. Kashefi K, Lovley D. Reduction of Fe(III), Mn(IV), and toxic metals at 100°C by *Pyrobaculum islandicum*. *Appl Environ Microbiol* 2000;66(3):1050-6.
52. Jiang Y, Zhang B, He C, Shi J, Borthwick AGL, Huang X. Synchronous microbial vanadium (V) reduction and denitrification in groundwater using hydrogen as the sole electron donor. *Water Resour Res* 2018;141:289-96.
53. Wisotzksy F, Eckert P. Sulfat-dominiertes BTEX Abbau im Grundwasser eines ehemaligen Gaswerks-standortes. *Grundwasser* 1997;2:11-20.
54. Ranchou-Peyruse A, Gasc C, Guignard M, Auello T, Sequidt D, Peyret P, et al. The sequence capture by hybridization: a new approach for revealing the potential of mono-aromatic hydrocarbons bioattenuation in a deep oligotrophic aquifer. *Microb Biotechnol* 2017;10(2):469-79.
55. Zettlitzer M, Moeller F, Morozova D, Lokay P, Würdemann H. Re-establishment of the proper injectivity of the CO<sub>2</sub>-injection well Ktzi 201 in Ketzin, Germany. *Int J Greenh Gas Control* 2010;4(6):952-9.

56. Bath A. Drilling fluid tracers – Review and update of industry experience and issues for RWMD site characterisation programme. Loughborough, UK: Nuclear Decommissioning Authority, Radioactive Waste Management Directorate; 2011.
57. Gittel A, Soerensen KB, Skovhus KI, Schramm A. Prokaryotic community structure and sulfate reducer activity in water from high-temperature oil reservoirs with and without nitrate treatment. *Appl Environ Microbiol* 2009;75(22):7086-96.
58. Laturnus F, Lauritsen FR, Groen C. Chloroform in a pristine aquifer system: Toward an evidence of biogenic origin. *Water Resour Res* 2000;36(10):2999-3009.
59. Hoehener P, Werner D, Balsiger C, Pasteris G. Worldwide occurrence and fate of chlorofluorocarbons in groundwater. *Crit Rev Environ Sci Technol* 2003;33(1):1-29.
60. Grabowski A, Nercessian O, Fayolle F, Blanchet D, Jeanthon C. Microbial diversity in production waters of a low-temperature biodegraded oil reservoir. *Fems Microbiol Ecol* 2005;54:427-43.
61. Liu F, Conrad R. Chemolithotrophic acetogenic H<sub>2</sub>/CO<sub>2</sub> utilization in Italian rice field soil. *Isme J* 2011;5:1526-39.
62. Kuesel K, Drake HL. Acetogens In: Thiel JRV, editor. *Encyclopedia of Geobiology*: Springer Science+Business Media B.V; 2011.
63. Schuchmann K, Mueller V. Energetics and Application of Heterotrophy in Acetogenic Bacteria. *Appl Environ Microbiol* 2016;82(14):4056-69.
64. Bengelsdorf FR, Beck MH, Erz C, Hoffmeister S, Karl MM, Riegler P, et al. Chapter four- Bacterial anaerobic synthesis gas (syngas) and CO<sub>2</sub> + H<sub>2</sub> fermentation. *Adv Appl Microbiol* 2018;103:143-221.
65. Esteve-Nunez A, Nunez C, Lovley DR. Preferential reduction of Fe(III) over fumarate by *Geobacter sulfurreducens*. *J Bacteriol* 2004;186(9):2897-9.

66. Eecke HCV, Akerman NH, Huber JA, Butterfield DA, Holden JF. Growth kinetics and energetics of a deep-sea hyperthermophilic methanogen under varying environmental conditions. *Environ Microbiol Rep* 2013;5(5):665-71.
67. Freitag TE, Prosser JI. Correlation of methane production and functional gene transcriptional activity in a peat soil. *Appl Environ Microbiol* 2009;75(21).
68. Usher K, Kaksonen A, Bouquet D, Cheng KY, Geste Y, Chapman PG, et al. The role of bacterial communities and carbon dioxide on the corrosion of steel. *Corros Sci* 2015;98:354-65.
69. Choong YY, Norli I, Abdullah AZ, Yhaya MF. Impacts of trace element supplementation on the performance of anaerobic digestion process: A critical review. *Bioresour Technol* 2016;209:369-79.
70. Pedersen K, Karlsson F. Investigations of subterranean microorganisms. Their importance for performance assessment of radioactive waste disposal. Swedish Nuclear Fuel and Waste Management Co; 1995.
71. Moench TT, Zeikus JG. Nutritional growth requirements for *Butyribacterium methylotrophicum* on single carbon substrates and glucose. *Curr Microbiol* 1983;9:151-4.
72. Magot M, Basso O, Tardy-Jacquenod C, Caumette P. *Desulfovibrio bastinii* sp. nov. and *Desulfovibrio gracilis* sp. nov., moderately halophilic, sulfate reducing bacteria isolated from deep subsurface oilfield water. *Int J Syst Evol Microbiol* 2004;54:1693-7.
73. Steinsbu BO, Thorseth IH, Nagakawa S, Inagaki F, Lever MA, Engelen B, et al. *Archaeoglobus sulfaticallidus* sp. nov., a thermophilic and facultatively lithoautotrophic sulfate-reducer isolated from black rust exposed to hot ridge flank crustal fluids. *Int J Syst Evol Microbiol* 2010;60:2745-52.

74. Casar CP, Kruger BR, Flynn TM, Masterson AL, Momper LM, Osburn MR. Mineral-hosted biofilm communities in the continental deep subsurface, Deep Mine Microbial Observatory, SD, USA. *Geobiology* 2020;18:508-22.
75. Samuels T, Bryce C, Landenmark H, Marie-Loudon C, Nicholson N, Stevens AH, et al. Microbial weathering of minerals and rocks in natural environments. In: Dontsova K, Balogh-Brunstad Z, Le Roux G, editors. *Biogeochemical cycles: Ecological drivers and environmental impact*: American Geophysical Union. John Wiley & Sons, Inc.; 2020.
76. Włodarczyk A, Lirski M, Fogtman A, Koblowska M, Bidzinski G, Matlakowska R. The oxidative metabolism of fossil hydrocarbons and sulfide minerals by the lithobiontic microbial community inhabiting deep subterrestrial kupferschiefer black shale. *Front Microbiol* 2018;9(972):1-14.
77. Napieralski S, Buss HL, Brantley SL, Lee S, Xu H, Roden EE. Microbial chemolithotrophy mediates oxidative weathering of granitic bedrock. *PNAS* 2019;116(52):26394-401.
78. Huang J, Sheng X-F, Xi J, Lin-Yan H, Huang Z, Wang Q, et al. Depth-related changes in community structure of culturable mineral weathering bacteria and in weathering patterns caused by them along two contrasting soil profiles. *Appl Environ Microbiol* 2014;80(1):29-42.
79. Alber BE. Autotrophic CO<sub>2</sub> metabolism. In: Schaechter M, editor. *Encyclopedia of Microbiology* Elsevier Academic Press; 2009.
80. Liu S, Suflita JM. H<sub>2</sub>-CO<sub>2</sub>-Dependent anaerobic O-demethylation activity in subsurface sediments and by an isolated bacterium. *Appl Environ Microbiol* 1993;59(5):1325-31.
81. Kotelnikova S, Pedersen K. Evidence for methanogenic archaea and homoacetogenic bacteria in deep granitic rock aquifers. *FEMS Microbiol Rev* 1997;20:339-49.

82. Londry KL, Jahnke LL, Des Marais DJ. Stable carbon isotope ratios of lipid biomarkers and biomass for sulfate-reducing bacteria grown with different substrates. Goldschmidt Conference 2001.
83. Camacho A. Sulfur bacteria In: Likens GE, editor. Encyclopedia of inland waters 1: Elsevier Science 2009.
84. Welsh DT, Bourges S, de Wit R, Herbert RA. Seasonal variations in nitrogen-fixation (acetylene reduction) and sulphate-reduction rates in the rhizosphere of *Zostera noltii*: Nitrogen fixation by sulphate-reducing bacteria Mar Biol 1996;125:619-28.
85. Whitman WB, Bowen TL, Boone DR. The methanogenic bacteria. In: Balows A, Truper HG, Dworkin M, Harder W, Schleifer K-H, editors. The Prokaryotes New York: Springer-Verlag; 1992. p. 719-67.
86. Drake HL. Acetogenesis London, United Kingdom Chapman & Hall; 2012.
87. Kapili BJ, Barnett SE, Buckley DH, Dekas AE. Evidence for phylogenetically and catabolically diverse active diazotrophs in deep-sea sediment Isme J 2020;14:971-83.
88. Herbert BN, Gilber PD, Stockdsle H, Watkinson RJ. Factors controlling the activity of sulphate-reducing bacteria in reservoirs during water injection. Society of Petroleum Engineers; 1985. Report No.: SPE-13978-MS Contract No.: SPE-13978-MS.
89. Chen Y, Cheng JJ, Creamer KS. Inhibition of anaerobic digestion process: A review. Bioresour Technol 2008;99:4044-64.
90. Vroblesky DA, Bradley PM, Chapelle FH. Influence of electron donor on the minimum sulfate concentration required for sulfate reduction in a petroleum hydrocarbon-contaminated aquifer. Environ Sci Technol 1996;30 1377-81.
91. Havig JR, Hamilton TL, Bachan A, Kump LR. Sulfur and carbon isotopic evidence for metabolic pathway evolution and a four-stepped Earth system progression across the Archean and Paleoproterozoic. Earth Sci Rev 2017;174:1-21.



92. Erb TJ, Kiefer P, Hattendorf B, Guenther D, Vorholt JA. GFAJ-1 is an arsenate-resistant, phosphate-dependent organism. *Science* 2012;337:467-71.
93. Taylor GT, Pirt SJ. Nutrition and factors limiting the growth of a methanogenic bacterium (*Methanobacterium thermoautotrophicum*) *Arch Microbiol* 1977;113:17-22.
94. van Houten RT, Yun SY, Lettinga G. Thermophilic sulphate and sulphite reduction in lab-scale gas-lift reactors using H<sub>2</sub> and CO<sub>2</sub> as energy and carbon source. *Biotechnol Bioeng* 1997;55:807-14.
95. Slobodkin AI, Slobodkina GB. Thermophilic prokaryotes from deep subterranean habitat. *Mikrobiologiya* 2014;83(3):255-70.
96. Basso O, Lascourreges JF, Le Borgne F, Le Goff C, Magot M. Characterization by culture and molecular analysis of the microbial diversity of a deep subsurface gas storage aquifer. *Res Microbiol* 2009;160:107-9.
97. Ranchou-Peyruse M, Auguet J-C, Maziere C, Restrepo-Ortiz CX, Guignard M, Dequidt D, et al. Geological gas-storage shapes deep life. *J Environ Biol* 2019;21(10):3953-64.
98. Magasanik B. Regulation of nitrogen utilization. In: FC N, Curtiss RI, Ingraham J, Lin E, Low K, Magasanik B, et al., editors. *Escherichia coli and Salmonella: cellular and molecular biology*. 2nd edition ed. Washington, DC: ASM Press; 1996. p. 1344–56.
99. Matos CR, Carneiro JF, Silva PP. Overview of large-scale underground energy storage technologies for integration of renewable energies and criteria for reservoir identification. *J Energy Storage* 2019;21:241-58.
100. Shi Z, Jessen K, Tsotsis TT. Impacts of the subsurface storage of natural gas and hydrogen mixtures *Int J Hydrogen Energy* 2020;45(15):f8757-73.

101. Hassanpouryouzband A, Joonaki E, Edlmann K, Heinemann N, Yang J. Thermodynamic and transport properties of hydrogen containing streams. *Sci Data* 2020;1(1):1-14.
102. Yen TF. *Microbial enhanced oil recovery: Principle and practice* Boca Raton, Florida: CRC Press; 1990.
103. Salwan R, Sharma V. *Physiological and biotechnological aspects of extremophiles*. London, United Kingdom Elsevier 2020.
104. Holden JF. *Extremophiles: Hot Environments* In: Schaechter M, editor. *Encyclopaedia of Microbiology* Elsevier Academic Press; 2009.
105. Takai K, Nakamura K, Toki T, Tsunogai U, Miyazaki M, Miyazaki J, et al. Cell proliferation at 122°C and isotopically heavy CH<sub>4</sub> production by a hyperthermophilic methanogen under high-pressure cultivation. *PNAS* 2008;105(31):10949-54.
106. Jaenicke R, Sterner R. Life at high temperatures. *Prokaryotes* 2006;2:167-209.
107. Hosh S, Lepcha K, Basak A, Mahanty AK. *Thermophiles and thermophilic hydrolases* In: Salwan R, Sharma V, editors. *Physiological and biotechnological aspects of extremophiles*. London, United Kingdom Elsevier Academic Press; 2020.
108. Miller JF, Nelson CM, Ludlow JM, Shah NN, Clark DS. High pressure-temperature bioreactor: assays of thermostable hydrogenase with fiber optics. *Biotechnol Bioeng* 1989;34:1015-21.
109. Pley U, Schipka J, Gambacorta A, Jannasch HW, Fricke H, Rachel R, et al. *Pyrodictium abyssi* sp. nov. represents a novel heterotrophic marine archaeal hyperthermophile growing at 110°C *Syst Appl Microbiol* 1991;14:245-53.
110. Basen M, Geiger I, Henke L, Mueller V. A genetic system for the thermophilic acetogenic bacterium *Thermoanaerobacter kivui*. *Appl Environ Microbiol* 2018;84(3):1-11.

111. Leigh JA, Mayer F, Wolfe RS. *Acetogenium kivui*, a new thermophilic hydrogen-oxidizing, acetogenic bacterium. Arch Microbiol 1981;129:275-80.
112. Balk M, Weijma J, Friedrich MW, Stams AJM. Methanol utilization by a novel thermophilic homoacetogenic bacterium, *Moorella mulderi* sp. nov., isolated from a bioreactor. Arch Microbiol 2003;179:315-20.
113. Oren A. Life at high salt concentrations In: Rosenberg E, editor. The Prokaryotes Prokaryotic Communities and Ecophysiology. Volume 4. Singapore: Springer; 2013.
114. Oren A. The bioenergetic basis for the decrease in metabolic diversity at increasing salt concentrations: implications for the functioning of salt lake ecosystems. Hydrobiologia 2001;466(1-3):61-72.
115. Oren. Thermodynamic limits to microbial life at high salt concentrations. J Environ Biol 2011;13(8):1908-23.
116. Ollivier B, Fardeau M-L, Cayol J-L, Magot M, Patel BKC, Prensier G, et al. *Methanocalculus halotolerans* gen. nov., sp. nov., isolated from an oil-producing well. Int J Syst Bacteriol 1998;48:821-8.
117. Zhilina TN, Zavarzina DG, Kevbrin VV, Kolganova TV. *Methanocalculus natronophilus* sp. nov., a new alkaliphilic hydrogenotrophic methanogenic archaeon from a soda lake, and proposal of the new family Methanocalculaceae. Microbiology 2013;82(6):698-706.
118. Ollivier B, Hatchikian G, Guezennec J, Garcia J-L. *Desulfohalobium retbaense* gen. nov. sp. nov. a halophilic sulfate-reducing bacterium from sediments of a hypersaline lake in Senegal. Int J Syst Bacteriol 1991;41(1):74-81.
119. Jakobsen RF, Kjeldsen KU, Ingvordsen K. *Desulfohalobium utahense* sp. nov., a moderately halophilic, sulfate-reducing bacterium isolated from Great Salt Lake. Int J Syst Evol Microbiol 2006;56:2063-9.

120. Krekeler D, Sigalevich P, Teske A, Cypionka H, Cohen Y. A sulfate-reducing bacterium from the oxic layer of a microbial mat from Solar Lake (Sinai), *Desulfovibrio oxycliniae* sp. nov. *Arch Microbiol* 1997;167:369-75.
121. Zhilina TN, Zavarzin GA, Detkova EN, Rainey FA. *Natroniella acetigena* gen. nov. sp. nov., an extremely haloalkaliphilic, homoacetic bacterium: a new member of *Haloanaerobiales*. *Curr Microbiol* 1996;32:320-6.
122. Zhilina TN, Zavarzin GA. Extremely halophilic, methylotrophic, anaerobic bacteria. *FEMS Microbiol Rev* 1990 87:315-22.
123. Porter D, Roychoudhury AN, Cowan D. Dissimilatory sulfate reduction in hypersaline coastal pans: Activity across a salinity gradient. *Geochim Cosmochim Acta* 2007;71:5102-16.
124. Borin S, Brusetti L, Mapelli F, D'Auria G, Brusa T, Marzorati M, et al. Sulfur cycling and methanogenesis primarily drive microbial colonization of the highly sulfidic Urania deep hypersaline basin. *PNAS* 2009;106(23):9151-6.
125. Foti M, Sorokin DY, Lomans B, Mussman M, Zacharova EE, Pimenov NV, et al. Diversity, activity, and abundance of sulfate-reducing bacteria in saline and hypersaline soda lakes. *Appl Environ Microbiol* 2007;73(2):2093–100.
126. Motamedi M, Karland O, Pedersen K. Survival of sulfate reducing bacteria at different water activities in compacted bentonite. *Fems Microbiol Lett* 1996;141(1):83-7.
127. Kushner DJ. *Microbial life in extreme environments*. London: Academic Press 1978.
128. Hallsworth JE, Yakimov MM, Golyshin PN, Gillion JLM, D'Auria G, de Lima ALves F, et al. Limits of life in MgCl<sub>2</sub>-containing environments: chaotropicity defines the window. *Environ Microbiol* 2007;9(3):801-13.

129. Steinle L, Knittel K, Felber N, Casalino C, de Lange G, Tessarolo C, et al. Life on the edge: active microbial communities in the Kryos MgCl<sub>2</sub>- brine basin at very low water activity. *Isme J* 2018;12:1414-26.
130. Cray JA, Russell JT, Timson DJ, Singhai RS, Hallsworth JE. A universal measure of chaotropicity and kosmotropicity. *Environ Microbiol* 2013;15(1):287-96.
131. Yuan H, Chen Y, Zhang H, Jiang S, Zhou Q, Gu G. Improved Bioproduction of Short-Chain Fatty Acids (SCFAs) from Excess Sludge under Alkaline Conditions. *Environ Sci Technol* 2006;40:2025-9.
132. O'Flatherty V, Mahony T, O'Kennedy R, Colleran E. Effect of pH on growth kinetics and sulphide toxicity of a range of methanogenic, syntrophic and sulphate-reducing bacteria. *Process Biochem* 1998;33(5):555-69.
133. Sorokin DY, Abbas B, Merkel AY, Riipstra EIC, Sinninghe Samste JS, Sukhacheva MV, et al. *Methanosalsum natronophilum* sp. nov., and *Methanocalculus alkaliphilus* sp. nov., haloalkaliphilic methanogens from hypersaline soda lakes. *Int J Syst Evol Microbiol* 2015;65:3739-45.
134. Stewart LC, Jung J-H, Kim Y-T, Kwon S-W, Park C-S, Holden JF. *Methanocaldococcus bathoardescens* sp. nov., a hyperthermophilic methanogen isolated from a volcanically active deep-sea hydrothermal vent. *Int J Syst Evol Microbiol* 2015;65:1280-3.
135. Takai K, Inoue A, Horikoshi K. *Methanothermococcus okinawensis* sp. nov., a thermophilic, methane-producing archaeon isolated from a Western Pacific deep-sea hydrothermal vent system. *Int J Syst Evol Microbiol* 2002;51:1089-95.
136. Ganzert L, Schirmack J, Alawi M, Mangelsdorf K, Sand W, Hillebrand-Voiculescu A, et al. *Methanosarcina spelaei* sp. nov., a methanogenic archaeon isolated from a floating biofilm of a subsurface sulphurous lake. *Int J Syst Evol Microbiol* 2014;64:3478–84.

137. Braeuer SL, Cadillo-Quiroz H, Ward RJ, Yavitt JB, Zinder SH. *Methanoregula boonei* gen. nov., sp. nov., an acidiphilic methanogen isolated from an acidic peat bog. *Int J Syst Evol Microbiol* 2011;61:45-52.
138. Zhilina TN, Zavarzin GA, Rainey FA, Pikuta EN, Osipov GA, Kostrikina NA. *Desulfonatronovibrio hydrogenovorans* gen. nov., sp. nov., an alkaliphilic, sulfate-reducing bacterium. *Int J Syst Bacteriol* 1997;47(1):144-9.
139. Sorokin DY, Muyzer G. *Desulfurispira natronophila* gen. nov. sp. nov.: an obligately anaerobic dissimilatory sulfur-reducing bacterium from soda lakes. *Extremophiles* 2010;14:349-55.
140. Nga DP, Ha DTC, Hien LT, Stan-Lotter H. *Desulfovibrio vietnamensis* sp.nov., a halophilic sulfate-reducing bacterium from Vietnamese oil fields. *Anaerobe* 1996;2:385-92.
141. Hallberg KB, Lindstroem EB. Characterization of *Thiobacillus caldus* sp. nov., a moderately thermophilic acidophile. *Microbiology* 1994;140:3451-1456.
142. Segerer A, Neuner A, Kristjansson JK, Stetter KO. *Acidianus infernus* gen. nov. sp. nov. and *Acidianus brierleyi* comb. nov.: facultatively aerobic, extremely acidophilic thermophilic sulfur-metabolizing archaeobacteria. *Int J Syst Bacteriol* 1986;36(4):559-64.
143. Segerer A, Trincone A, Gahrtz M, Stetter KO. *Stygiolobus azoricus* gen. nov., sp. nov. represents a novel genus of anaerobic, extremely thermoacidophilic archaeobacteria of the order *Sulfolobales*. *Int J Syst Bacteriol* 1991;41(4):495-501.
144. Fliermans CB, Brock TD. Ecology of sulfur-oxidizing bacteria in hot acid soils. *J Bacteriol* 1972;111(2):343-50.
145. Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F. *The Prokaryotes. Firmicutes and Tenericutes* fourth edition ed. Heidelberg: Springer Science and Business Media 2014.

146. Inokuma K, Nakashimada Y, Akahoshi T, Nishio N. Characterization of enzymes involved in the ethanol production of *Moorella* sp. HUC22-1. *Arch Microbiol* 2007;188:37-45.
147. Schnuerer A, Schink B, Svensson BH. *Clostridium ultunense* sp. nov., a mesophilic bacterium oxidizing acetate in syntrophic association with a hydrogenotrophic methanogenic bacterium. *Int J Syst Bacteriol* 1996;46(4):1145-52.
148. Hirano S, Masuda N. Characterization of NADP-Dependent 7,B-Hydroxysteroid Dehydrogenases from *Peptostreptococcus productus* and *Eubacterium aerofaciens*. *Appl Environ Microbiol* 1982;43(5):1057-63.
149. Zhilina TN, Zavarzina DG, Detkova EN, Patutina EO, Kuznetsov BB. *Fuchsiella ferrireducens* sp. nov., a novel haloalkaliphilic, lithoautotrophic homoacetogen capable of iron reduction, and emendation of the description of the genus *Fuchsiella*. *Int J Syst Evol Microbiol* 2015;85:2432-40.
150. Wiegel J, Braun M, Gottschalk G. *Clostridium thermoautotrophicum* species novum, a thermophile producing acetate from molecular hydrogen and carbon dioxide. *Curr Microbiol* 1981;5:255-60.
151. Tanner RS, Miller LM, Yang D. *Clostridium ljungdahlii* sp. nov., an acetogenic species in clostridial rRNA homology group I *Int J Syst Bacteriol* 1993;43(2):232-6.
152. Kaneuchi C, Benno Y, Mitsuoka T. *Clostridium coccooides*, a new species from the faeces of mice. *Int J Syst Bacteriol* 1976;26(4):482-6.
153. Kuesel K, Dorsch T, Acker G, Stackebrandt E, Drake HL. *Clostridium scatologenes* strain SL1 isolated as an acetogenic bacterium from acidic sediments. *Int J Syst Evol Microbiol* 2000;50:537-46.

154. Gößner AS, Picardal F, Tanner RS, Drake HL. Carbon metabolism of the moderately acid-tolerant acetogen *Clostridium drakai* isolated from peat. *Fems Microbiol Lett* 2008;287:236-42.
155. Abe F, Kato C, Horikoshi K. Pressure-regulated metabolism in microorganisms. *Trends Microbiol* 1999;7(11):447-53.
156. Jebbar M, Franzetti B, Girard E, Oger P. Microbial diversity and adaptation to high hydrostatic pressure in deep-sea hydrothermal vents prokaryotes. *Extremophiles* 2015;19:721-40.
157. Vikromvarasiri N, S. B, Pisutpaisal N. Comparative performance of *Halothiobacillus Neapolitanus* and *Paracoccus Pantotrophus* in sulphur oxidation. *Energy Procedia* 2015;79:885-9.
158. Cao J, Gayet N, Zeng X, Shao Z, Jebbar M, Alain K. *Pseudodesulfovibrio indicus* gen. nov., sp. nov., a piezophilic sulfate-reducing bacterium from the Indian Ocean and reclassification of four species of the genus *Desulfovibrio*. *Int J Syst Evol Microbiol* 2016;66:3904-11.
159. Takai K, Miyazaki M, Hirayama H, Nakagawa S, Querellou J, Godfroy A. Isolation and physiological characterization of two novel, piezophilic, thermophilic chemolithoautotrophs from a deep-sea hydrothermal vent chimney. *J Environ Biol* 2009;11(8):1983-97.
160. Fichtel K, Logemann J, Fichtel J, Rullkoetter J, Cypionka H, Engelen B. Temperature and pressure adaptation of a sulfate reducer from the deep subsurface. *Front Microbiol* 2015 6:1-13.
161. Kurr M, Huber R, Koenig H, Jannasch HW, Fricke H, Trincone A, et al. *Methanopyrus kandleri*, gen. and sp. nov. represents a novel group of hyperthermophilic methanogens, growing at 110 ° C\*. *Arch Microbiol* 1991;156:239-47.



162. Ntagia E, Chatzigiannidou I, Williamson AJ, Arends JBA, Rabaey K. Homoacetogenesis and microbial community composition are shaped by pH and total sulfide concentration. *Microb Biotechnol* 2020;13(4):1026-38.
163. Suleimenov OM, Seward TM. A spectrophotometric study of hydrogen sulphide ionisation in aqueous solutions to 350°C. *Geochim Cosmochim Acta* 1997;61(24):5187-98.
164. Kushkeyvych I, Dordevic D, Vitezova M. Toxicity of hydrogen sulfide toward sulfate-reducing bacteria *Desulfovibrio piger* Vib-7. *Arch Microbiol* 2019;201(3):389-97.
165. Winfrey MR, Zeikus JG. Effect of sulfate on carbon and electron flow during microbial methanogenesis in freshwater sediments. *Appl Environ Microbiol* 1977;33(2):275-81.
166. Choi E, Rim JM. Competition and inhibition of sulfate reducers and methane producers in anaerobic treatment. *Water Sci Technol* 1991;23:1259-64.
167. Dupraz S, Fabbri A, Joulian C, Dictor M-C, Battaglia-Brunet F, Menez B, et al. Impact of CO<sub>2</sub> concentration on autotrophic metabolisms and carbon fate in saline aquifers – A case study. *Geochim Cosmochim Acta* 2013;119:61-76.
168. Froestl JM, Seifritz C, Drake HL. Effect of nitrate on the autotrophic metabolism of the acetogens *Clostridium thermoautotrophicum* and *Clostridium thermoaceticum* *J Bacteriol* 1996;178(15):4597–603.
169. Wang H, Zhang Y, Angelidakis I. Ammonia inhibition on hydrogen enriched anaerobic digestion of manure under mesophilic and thermophilic conditions. *Water Resour Res* 2016;105:314-9.
170. Conrad R. Contribution of hydrogen to methane production and control of hydrogen concentrations in methanogenic soils and sediments. *Fems Microbiol Ecol* 1999;28:193-202.
171. Barth T, Riis M. Interactions between organic acid anions in formation waters and reservoir mineral phases. *Org Geochem* 1992;19(4-6):455-82.

172. Oil and gas field data from the North Sea [Internet]. Oil and Gas Authority 2020 [cited 19.5.2020]. Available from: <https://www.ogauthority.co.uk/data-centre/>.
173. Hoehler TM, Albert DB, Alperin MJ, Martens CS. Acetogenesis from CO<sub>2</sub> in an anoxic marine sediment. *Limnol Oceanogr* 1999;44(3):662-7.
174. Pedersen K. Microbial life in deep granitic rock. *Fems Microbiol Ecol* 1997;20:399-414.
175. Breznak JA. Acetogenesis from carbon dioxide in termite guts. In: H.L. D, editor. *Acetogenesis Chapman & Hall Microbiology Series (Physiology / Ecology / Molecular Biology / Biotechnology)*. Boston, MA: Springer; 1994.
176. Krumholz LR, Harris SH, Tay ST, Suflita JM. Characterization of two subsurface H<sub>2</sub>-utilizing bacteria, *Desulfomicrobium hypogeium* sp. nov. and *Acetobacterium psammolithicum* sp. nov., and their ecological role. *Appl Environ Microbiol* 1999;65(6):2300-6.
177. Phillips J. Extracellular electron uptake by acetogenic bacteria: Does H<sub>2</sub> consumption favor the H<sub>2</sub> evolution reaction on a cathode or metallic iron? *Front Microbiol* 2020;10:1-13.
178. Robinson JA, Tiedje JM. Competition between sulfate-reducing and methanogenic bacteria for H<sub>2</sub> under resting and growing conditions. *Arch Microbiol* 1984;137:26-32.
179. Fu B, Jin X, Conrad R, Liu H, Liu H. Competition between chemolithotrophic acetogenesis and hydrogenotrophic methanogenesis for exogenous H<sub>2</sub>/CO<sub>2</sub> in anaerobically digested sludge. *Front Microbiol* 2019;10:1-9.
180. Kotsyurbenko OR, Glagolev MV, Nozhevnikova AN, Conrad R. Competition between homoacetogenic bacteria and methanogenic archaea for hydrogen at low temperature. *Fems Microbiol Ecol* 2001;38:153-9.

181. Van Verseveld HW, Duine JA, editors. Proceedings of the 5th international symposium on microbial growth on C<sub>1</sub> compounds. International Symposium on microbial growth on C<sub>1</sub> compounds; 1986; University of Groningen, The Netherlands.
182. Lever MA. Acetogenesis in the energy-starved deep biosphere—a paradox? *Front Microbiol* 2012;2:1-18.
183. Lackner N, Wagner AO, Ilmer P. Effect of sulfate addition on carbon flow and microbial community composition during thermophilic digestion of cellulose. *Appl Microbiol Biotechnol* 2020;104:4605-15.
184. Men Y, Feil H, VerBerkmoes NC, Shah MB, Johnson DR, Lee PKH, et al. Sustainable syntrophic growth of *Dehalococcoides ethenogenes* strain 195 with *Desulfovibrio vulgaris* Hildenborough and *Methanobacterium congolense*: global transcriptomic and proteomic analyses. *Isme J* 2012;6:410-2.
185. Freedman AJE, BoonFei T, Thompson JR. Microbial potential for carbon and nutrient cycling in a geogenic supercritical carbon dioxide reservoir. *Environ Microbiol* 2017;19:2228-45.
186. Topcuoglu BD, Meydan C, Nguyen TB, Lang SQ, Holden JF. Growth kinetics, carbon isotope fractionation, and gene expression in the hyperthermophile *Methanocaldococcus jannaschii* during hydrogen-limited growth and interspecies hydrogen transfer. *Appl Environ Microbiol* 2019;85(9):1-14.
187. Bonch-Osmolokskaya EA, Miroshnichenko ML, Lebedinsky AV, Chernyh TN, Nazina TN, Ivoilov VS, et al. Radioisotopic, culture-based, and oligonucleotide microchip analyses of thermophilic microbial communities in a continental high-temperature petroleum reservoir. *Appl Environ Microbiol* 2003;69(10):6143–51.

188. Tian H, Gao P, Chen Z, Li Y, Li Y, Wang Y, et al. Compositions and abundances of sulfate-reducing and sulfur-oxidizing microorganisms in water-flooded petroleum reservoirs with different temperatures in China. *Front Microbiol* 2017;8(143):1-14.
189. Auello T, Berlendis S, Lascourreges JF, Dessort D, Duclerc D, Saint-Laurent S, et al. New bio-indicators for long term natural attenuation of monoaromatic compounds in deep terrestrial aquifers. *Front Microbiol* 2016;7(122):1-16.
190. Mori K, Tsurumaru H, Harayama S. Iron corrosion activity of anaerobic hydrogen-consuming microorganisms isolated from oil facilities. *J Biosci Bioeng* 2010;110(4):426-30.
191. Okpala GN, Chen C, Fida T, Voordouw G. Effect of thermophilic nitrate reduction on sulfide production in high temperature oil reservoir samples. *Front Microbiol* 2017;8:1-13.
192. Priha O, Nyyssoenen M, Bomberg M, Laitila A, Simell J, Kapanen A, et al. Application of denaturing high-performance liquid chromatography for monitoring sulfate-reducing bacteria in oil fields. 2013;79(17).
193. Stetter KO, Huber R, Bloechl E, Kurr M, Eden RD, Fielder M, et al. Hyperthermophilic archaea are thriving in deep North Sea and Alaskan oil reservoirs *Nature* 1993;365:743-5.
194. Kross B, Amann A, Thuens N, Stanjek H, Ostertag-Henning C, Krueger M, et al. Transport von Wasserstoff in Gesteinen unter Berücksichtigung abiotischer und mikrobieller Redoxreaktionen (H<sub>2</sub>\_ReacT). Aachen, Germany Bundesministerium fuer Bildung und Forschung 2020.
195. Balk M, Mehboob F, Gelder A, Riipstra WIC, Sinninghe Damste JS, Stams AJM. (Per)chlorate reduction by an acetogenic bacterium, *Sporomusa sp.*, isolated from an underground gas storage. *Appl Microbiol Biotechnol* 2010;88:595-603.

196. Berlendis S, Lascourreges JF, Schraauwers B, Sivadon P, Magot M. Anaerobic biodegradation of BTEX by original bacterial communities from an underground gas storage aquifer. *Environ Sci Technol* 2010;44:3621-8.
197. Conrad R, Schuetz H, Babbel M. Temperature limitation of hydrogen turnover and methanogenesis in anoxic paddy soil. *Fems Microbiol Ecol* 1987;45:281-9.
198. Stewart LC, Algar CK, Fortunato CS, Larson BI, Vallino JJ, Huber JA, et al. Fluid geochemistry, local hydrology, and metabolic activity define methanogen community size and composition in deep-sea hydrothermal vents. *Isme J* 2019;13(7):1711-21.
199. Enoki M, Shinzato N, Sato H, Nakamura K, Y. K. Comparative proteomic analysis of *Methanothermobacter themautotrophicus* DH in pure culture and in co-culture with a butyrate-oxidizing bacterium. *Plos One* 2011;6(8):1-10.
200. Mukhopadhyay B, Johnson EF, Wolfe RS. A novel pH<sub>2</sub> control on the expression of flagella in the hyperthermophilic strictly hydrogenotrophic methanarchaeon *Methanococcus jannaschii*. *PNAS* 2000;97(21):11522-7.
201. Miller JF, Shah NN, Nelson CM, Ludlow JM, Clark DS. Pressure and temperature effects on growth and methane production of the extreme thermophile *Methanococcus jannaschi*. *Appl Environ Microbiol* 1988;54(12):3039-42.
202. Schieche D, Murty MVS, Kermode RI, Bhattacharyya D. Biohydrogenation of fumarate using *Desulfovibrio desulfuricans*: Experimental results and kinetic rate modelling. *J Chem Technol Biotechnol* 1997;70(3):316-22.
203. Baas Becking LGM. Geobiologie of inleiding tot de milieukunde. Den Hague, the Netherlands: W.P. Van Stockum & Zoon; 1934.
204. Xu Y, Teng Y, Wang XB, Li R, Christie P. Exploring bacterial community structure and function associated with polychlorinated biphenyl biodegradation in two hydrogen-amended soils *Sci Total Environ* 2020;745(140839):1-12.

205. Braga Nan L, Trably E, Santa-Catalina G, Bernet N, Delgenes J-P, Escudie R. Biomethanation processes: new insights on the effect of a high H<sub>2</sub> partial pressure on microbial communities. *Biotechnol Biofuels* 2020;13(141):1-17.
206. Treu L, Kogias PG, de Diego-Diaz B, Campanaro S, Bassani I, Fernandez-Rodriguez J, et al. Two-year microbial adaptation during hydrogen-mediated biogas upgrading process in a serial reactor configuration. *Bioresour Technol* 2018;264(140-147).
207. Puente-Sánchez F, Arce-Rodríguez A, Oggerind M, García-Villadangosa M, Moreno-Paza M, Blanco Y, et al. Viable cyanobacteria in the deep continental subsurface. *PNAS* 2018;115(42):10702-7.
208. Gluyas JG, Hichens HM. The United Kingdom oil and gas fields commemorative millennium volume Gluyas JG, Hichens HM, editors: *Memoirs of the Geological Society of London*; 2003.
209. Zhang Y, Zhang Z, Suzuki K, Maekawa T. Uptake and mass balance of trace metals for methane producing bacteria. *Biomass Bioenerg* 2003;25:427–33.
210. Scherer P, Lippert H, Wolff G. Composition of the major elements and trace-elements of 10 methanogenic bacteria determined by inductively coupled plasma emission-spectrometry. *Biol Trace Elem Res* 1983;5(3):149-63.
211. Amid A, Mignard D, Wilkinson M. Seasonal storage of hydrogen in a depleted natural gas reservoir. *Int J Hydrogen Energy* 2016;41(12):5549-58.
212. Kettle H, Louis P, Holtrop G, Duncan SH, Flint HJ. Modelling the emergent dynamics and major metabolites of the human colonic microbiota. *Environ Microbiol* 2015;17(5):1615-30.
213. Littlewood D, Postgate JR. On the osmotic behaviour of *Desulphovibrio desulphuricans* *J Gen Microbiol* 1957;17:378-89.

214. Bratbak G, Dundas I. Bacterial dry matter content and biomass estimations. *Appl Environ Microbiol* 1984;48(4):755-7.
215. Würdemann H, Halm H, Lerm S, Kleyböcker A. Verbund-Forschungsvorhaben H2STORE: Untersuchung der geohydraulischen, mineralogischen, geochemischen und biogenen Wechselwirkungen bei der Untertage-Speicherung von H<sub>2</sub> in konvertierten Gaslagerstätten: Teilprojekt 4- Mikrobiologie: Abschlussbericht: Berichtszeitraum: 01.08.2012 bis 31.12.2015. Potsdam: Helmholtz-Zentrum Potsdam Deutsches GeoForschungsZentrum GFZ; 2016.
216. Nazina TN, Pavlova NK, Tatarkin YV, Shestakova NM, Babich TL, Sokolova DS, et al. Microorganisms of the carbonate petroleum reservoir 302 of the Romashkinskoe oilfield and their biotechnological potential. *Microbiology* 2013;82(2):190-200.
217. Okabe S. Rate and stoichiometry of sulfate reducing bacteria in suspended and biofilm cultures. Montana, USA: Montana State University; 1992.
218. Stolten D, Emonts B. Hydrogen science and engineering: materials, processes, systems, and technology, 2 volume set: Wiley-VCH; 2016.
219. Ebrahimiyehta A. Characterization of geochemical interactions and migration of hydrogen in sandstone sedimentary formations: application to geological storage. Université d'Orléans: Earth Sciences; 2017. Contract No.: ffNNT : 2017ORLE2016.
220. Panfilov M. Underground storage of hydrogen: In situ self-organisation and methane generation. *Transp Porous Media* 2010;85(3):841-65.
221. Smigan P, Greksak M, Kozankova J, Buzek F, Onderka V, Wolf I. Methanogenic bacteria as a key factor involved in changes of town gas stored in an underground reservoir. *Fems Microbiol Ecol* 1990;73(3):221-4.

222. Kato S, Yamagishi A, Daimon S, Kawasaki K, Tamaki H, Kitagawa W, et al. Isolation of previously uncultured slow-growing bacteria by using a simple modification in the preparation of agar media. *Appl Environ Microb* 2018;84(19):1-9.
223. Stevenson BS, Eichhorst SA, Wertz JT, Schmidt TM, Breznak JA. New strategies for cultivation and detection of previously uncultured microbes. *Appl Environ Microbiol* 2004;70(8):4748–55.
224. Henson M, Pitre D, Weckhorst J, Lanclos V, Webber A, Thrash J. Artificial seawater media facilitate cultivating members of the microbial majority from the Gulf of Mexico. *mSphere* 2016;1(2).
225. Zengler K, Toledo, Rappe M, Elkins J, Mathur EJ, Short JM, et al. Cultivating the uncultured. *PNAS* 2002;26(24):15681–6.
226. Berdy B, Spoering AL, Ling LL, Epstein SS. In situ cultivation of previously uncultivable microorganisms using the ichip. *Nat Protoc* 2017;12(10):2232-422.
227. Kaeberlein T, Lewis K, Epstein SS. Isolating uncultivable microorganisms in pure culture in a simulated natural environment. *Science* 2002 296:1127-9
228. Mitchell AC, Phillips AJ, Hiebert R, Gerlach R, Spangler LH, Cunningham AB. Biofilm enhanced geologic sequestration of supercritical CO<sub>2</sub>. *Int J Greenh Gas Control* 2009;3:90-9.