

# Improving the ionospheric state estimate during geomagnetic storm time through assimilation of neutral density data

EGU 2022 – G5.1 Ionosphere, thermosphere and space weather: monitoring and modelling.



*Isabel Fernandez-Gomez, Timothy Kodikara, Claudia Borries, Ehsan Forootan, Michael Schmidt and Mihail Codrescu.*



Knowledge for Tomorrow





# Can we improve electron density by assimilating neutral density during storm conditions?



- **Assimilated data:** Swarm – A TMD observations normalized to the common altitude of 400 km.
- **Period:** 16-19 March 2015 containing St. Patrick's Day storm
- Days are classified as quiet (16), main phase (17) and recovery (18)
- **State vector:** Updates the forcing parameters and the necessary quantities to calculate neutral density.
- **Assimilation window:** 10 minutes
- TMD uncertainty is 10%




# What happens on the thermosphere? Neutral density



## Swarm – B/C TMD along the orbit RMSE improvement


$$RMSE = \sqrt{\frac{\sum (Obs - Mod)^2}{N}}$$

$$IMP(\%) = \frac{(RMSE_r - RMSE_a)}{RMSE_r} 100$$






# What happens on the ionosphere? Electron density (400 km)



**Swarm – A / GRACE Ne along the orbit RMSE improvement**


$$RMSE = \sqrt{\frac{\sum(\text{Obs} - \text{Mod})^2}{N}}$$

$$IMP(\%) = \frac{(RMSE_r - RMSE_a)}{RMSE_r} 100$$



# What happens on the ionosphere? Global Electron density improvement

- **Electron density global improvement** at 400 km between analysis and reference with respect to the **B-Spline electron density model**.
- For the three days of the storm
- Lower RMSE → Better fit of the model to observations
- Improvement (%) of RMSE of the analysis and reference differences.
- Positive values are **areas of improvement** (red)
- The main area of improvement is around the **equatorial region** (-45, 45) deg latitude.
- The **effect depends on altitude**. The positive improvement decreases for altitudes higher than 600 km.



$$\text{IMP}(\%) = \frac{(\text{RMSE}_r - \text{RMSE}_a)}{\text{RMSE}_r} 100$$



# Summary

- **Assimilation of neutral density** measurements into a physics-based model **during storm conditions** is capable of **correct the thermosphere and the ionosphere** (with limitations).
  - Neutral density improves along the orbit of the non assimilated Swarm – B/C satellites up to 40%
  - Electron density difference maps (analysis – reference) show the effects of TMD DA
  - Electron density improvement along the orbit of Swarm-A and GRACE are 8% and 22% respectively.
  - The **global electron density improvement** map shows the areas affected by TMD assimilation.
  - The **largest improvement in the electron density** estimates takes place during the **recovery phase** (negative storm driven by composition changes)

