



DNA Double Strand Break Repair and Its Control by Nucleosome Remodeling

Leonhard Andreas Karl[†], Martina Peritore[†], Lorenzo Galanti[†] and Boris Pfander^{*‡}

Research Group DNA Replication and Genome Integrity, Max Planck Institute of Biochemistry, Martinsried, Germany

OPEN ACCESS

Edited by:

Matthias Altmeyer,
University of Zurich, Switzerland

Reviewed by:

Jessica Downs,
Institute of Cancer Research (ICR),
United Kingdom

Alessandro A. Sartori,
University of Zurich, Switzerland

*Correspondence:

Boris Pfander
bpfander@biochem.mpg.de

[†]These authors have contributed
equally to this work

[‡]Present address:

Boris Pfander,
German Aerospace Center (DLR),
Institute of Aerospace Medicine,
Cologne, Germany;
Medical Faculty,
CECAD, Institute of Genome Stability
in Ageing and Disease, University of
Cologne, Cologne, Germany

Specialty section:

This article was submitted to
Human and Medical Genomics,
a section of the journal
Frontiers in Genetics

Received: 24 November 2021

Accepted: 23 December 2021

Published: 12 January 2022

Citation:

Karl LA, Peritore M, Galanti L and
Pfander B (2022) DNA Double Strand
Break Repair and Its Control by
Nucleosome Remodeling.
Front. Genet. 12:821543.
doi: 10.3389/fgene.2021.821543


DNA double strand breaks (DSBs) are repaired in eukaryotes by one of several cellular mechanisms. The decision-making process controlling DSB repair takes place at the step of DNA end resection, the nucleolytic processing of DNA ends, which generates single-stranded DNA overhangs. Dependent on the length of the overhang, a corresponding DSB repair mechanism is engaged. Interestingly, nucleosomes—the fundamental unit of chromatin—influence the activity of resection nucleases and nucleosome remodelers have emerged as key regulators of DSB repair. Nucleosome remodelers share a common enzymatic mechanism, but for global genome organization specific remodelers have been shown to exert distinct activities. Specifically, different remodelers have been found to slide and evict, position or edit nucleosomes. It is an open question whether the same remodelers exert the same function also in the context of DSBs. Here, we will review recent advances in our understanding of nucleosome remodelers at DSBs: to what extent nucleosome sliding, eviction, positioning and editing can be observed at DSBs and how these activities affect the DSB repair decision.

Keywords: nucleosome remodeling, double strand break, DNA repair, DNA end resection, cell cycle, genome stability

INTRODUCTION

DNA double strand breaks are a highly toxic form of DNA damage, arising from intrinsic and extrinsic sources (Ciccio and Elledge, 2010). Eukaryotes are equipped with several mechanisms to repair DSBs, including non-homologous end joining (NHEJ), alternative end joining (alt-EJ), homologous recombination (HR) and single strand annealing (SSA) (Chang et al., 2017; Ranjha et al., 2018). Notably, these pathways do not only have different prerequisites (for example HR requiring a homologous donor sequence), but they also differ in the repair outcome and the potential to introduce genetic changes (such as mutations and chromosomal rearrangements). The cellular repair pathway decision is therefore critical for the survival of the affected cell or organism as well as for the stability of its genome (Symington and Gautier, 2011). Moreover, the fact that DSB repair is controlled by endogenous factors is a major limitation for genome editing strategies, which can nowadays involve efficient delivery of DSBs at the gene of interest, but often lead to a heterogeneous outcome of the genome editing reaction across cell populations.

The cellular DSB repair pathway decision is made at the step of DNA end resection, the nucleolytic processing of DSB ends (Symington and Gautier, 2011; Cejka, 2015; Daley et al., 2015; Symington, 2016; Bonetti et al., 2018). Resection involves endo- and exonucleolytic cleavage of DNA ends that reveals 3' single-stranded DNA overhangs. Notably, resection destroys the substrate for repair by NHEJ and increasing amounts of 3' single-stranded DNA (ssDNA) predisposes for repair



by different mechanisms (alt EJ < HR < SSA, Blier et al., 1993; Falzon et al., 1993; Ira et al., 2004). The enzymatic process of resection has been subject of excellent reviews in this issue and elsewhere (Symington and Gautier, 2011; Cejka and Symington, 2021; Elbakry and Löbrich, 2021; Sanchez et al., 2021). Here we focus on how resection and thereby the repair pathway decision is regulated by nucleosomes and nucleosome remodelers, enzymes that can evict, position and edit nucleosomes. For general reviews on how DNA damage triggers post-translational histone modifications, we refer to the following articles (Smeenk and van Attikum, 2013; Van and Santos, 2018).

Nucleosomes form obstacles to the resection nucleases (Figure 1). Initial short-range resection is carried out by the Mre11-complex (Mre11-C in the following, consisting of Mre11-Rad50-Xrs2 with the Sae2 activator in budding yeast, and analogously of MRE11-RAD50-NBS1 with CtIP in human) (Symington and Gautier, 2011; Cejka and Symington, 2021; Elbakry and Löbrich, 2021; Sanchez et al., 2021). Endonucleolytic cleavage by Mre11-C occurs preferentially within nucleosome-free linker DNA, suggesting that nucleosomal DNA is protected and/or that chromatin binding of Mre11-C is guided by nucleosomes (Mimitou et al., 2017; Wang et al., 2017). Moreover, the nucleases that carry out long-range resection are directly inhibited by the presence of nucleosomes: biochemical studies with yeast proteins have shown that the Exo1 exonuclease is unable to act on a nucleosome substrate and the combined helicase-endonuclease STR-Dna2 (Sgs1-Top3-Rmi1-Dna2) can only process nucleosomal DNA, if sufficient nucleosome-free DNA is present (Adkins et al., 2013). Therefore, nucleosomes are a barrier to the resection process and resection control factors are expected to modify the permeability of this barrier.

Nucleosome remodelers have received attention as regulators of DNA end resection and DSB repair pathway choice. These nucleosome remodelers are enzymes that in

ATP-dependent fashion catalyze the breakage of histone-DNA-contacts within the nucleosome and translocate the DNA relative to histone proteins (Clapier and Cairns, 2009; Clapier et al., 2017). All eukaryotes possess several nucleosome remodelers - often in the form of multi-protein complexes - which are grouped into several sub-families according to the conservation of their ATPase subunit (Flaus et al., 2006). Biochemical and structural data suggest that the overall enzymatic mechanism of DNA translocation and breakage of DNA-histone contacts is highly related (Clapier and Cairns, 2009; Clapier et al., 2017) (with the potential exception of Fun30/SMARCAD1, see below). Nonetheless, studies on gene transcription and general chromatin organization have revealed that specific remodelers appear to have specific enzymatic activities, by which they slide, evict, position or edit nucleosomes (Clapier and Cairns, 2009; Clapier et al., 2017). Here, we will investigate whether such distinct roles can also be found in the context of DNA double strand breaks and how these activities may affect DSB repair. A natural focus of this review will be the budding yeast system, where remodelers have been studied comprehensively also in the context of DSBs, but we will additionally address whether the picture emerging from these studies is conserved in higher eukaryotes.

RESECTION IS AFFECTED BY NUCLEOSOMES

In many eukaryotes, DNA end resection is carried out by three resection enzymes, Mre11-C, STR-Dna2 and Exo1, which act specifically at one of the two stages of the resection process (short-range resection/resection initiation and long-range resection/resection elongation, Figure 1). Notably, all three act by distinct molecular mechanisms and it is therefore unsurprising that nucleosomes have distinct effects on each of them. Mre11-C recognizes the DSB end either directly or through a DSB end-binding protein (most likely the end-binding factor Ku) and, after activation by Sae2/CtIP, induces a single-strand break on the 5'-strand (Sartori et al., 2007; Cannavo and Cejka, 2014; Anand et al., 2016; Deshpande et al., 2016; Reginato et al., 2017; Wang et al., 2017). From this point, bidirectional resection occurs: Mre11-C catalyzes 3'-5' exonucleolytic resection towards the break, while Dna2 and Exo1 exonucleases catalyze long-range resection with 5'-3' polarity into undamaged chromatin (Mimitou and Symington, 2008; Zhu et al., 2008; Cejka et al., 2010; Niu et al., 2010; Garcia et al., 2011; Shibata et al., 2014).

Preferential cleavage of linker DNA indicates that nucleosomal DNA may be refractory to endonucleolytic clipping by Mre11-C (Mimitou et al., 2017; Wang et al., 2017). However, nucleosomes per se are not a barrier to Mre11-C. Rather, it can slide or reach over nucleosomes (Myler et al., 2017; Wang et al., 2017). In cases where such bypass occurs, the nucleosome located between DSB and incision site could then potentially constitute a barrier to the 3'-5' exonuclease activity of Mre11-C. Given the dual

endo- and exonucleolytic activities of Mre11-C, this question has so far been difficult to address.

Long-range resection enzymes are even more strongly affected by the presence of nucleosomes. For example, *in vitro* studies have shown that Exo1 cannot resect through nucleosomes (Adkins et al., 2013), suggesting that additional activities are needed to overcome the chromatin barrier. Interestingly, changing nucleosome composition may be sufficient to allow Exo1-mediated resection. Incorporation of the H2A-variant H2A.Z decreases nucleosome stability and increases accessibility of nucleosomal DNA (Abbott et al., 2001; Zhang et al., 2005; Jin and Felsenfeld, 2007; Adkins et al., 2013; Watanabe et al., 2013; Lewis et al., 2021), which may allow Exo1 to bypass the nucleosomal barrier (Adkins et al., 2013).

In contrast to Exo1, the other long-range resection enzyme STR-Dna2 is in principle able to bypass nucleosomes. This may be due to a different enzymatic mechanism. While during long-range resection STR-Dna2 has the net effect of an exonuclease, STR-Dna2 utilizes the combined action of the Sgs1 helicase that unwinds DNA, followed by endonucleolytic cleavage of the emerging flap structure by Dna2 (Cejka et al., 2010; Niu et al., 2010). Apparently, the Sgs1 helicase motor is powerful enough to disrupt nucleosomes, allowing STR-Dna2 to resect nucleosomal DNA (Adkins et al., 2013). However, in order to carry out resection of nucleosomal DNA, STR-Dna2 will need as much as 300 bp of free DNA to be able to traverse through nucleosomes (Adkins et al., 2013). This distance is greater than the nucleosomal linker DNA-length and, consistently, STR-Dna2 is effectively inhibited by a nucleosomal array (Adkins et al., 2013).

Therefore, both long-range resection enzymes are blocked by chromatin and will require the activity of additional factors. One factor that could help to overcome the nucleosomal barrier is Mre11-C. Speculatively, Mre11-C could catalyze further endonucleolytic incisions downstream of the nucleosome from which long-range nucleases could (re-)initiate and thereby allow to bypass the nucleosome barrier. Currently, such an auxiliary role of Mre11-C in long-range resection lacks experimental support, but recent data suggest that short-range and long-range resection nucleases work in a coordinated fashion (Ceppi et al., 2020).

Alternatively, resection enzymes will need assistance by chromatin remodelers to get past nucleosomes and it is therefore important to consider how these enzymes may be able to modify the nucleosome barrier.

REMODELERS HAVE DISTINCT ROLES IN CHROMATIN ORGANIZATION

Eukaryotes express several nucleosome remodelers (Flaus et al., 2006) and chromatin immunoprecipitation (ChIP) and related techniques have localized several of them to DSBs (Bantele et al., 2017; Bennett and Peterson, 2015; Bennett et al., 2013; Bird et al., 2002; Chai et al., 2005; Chen et al., 2012; Costelloe et al., 2012; Downs et al., 2004; Eapen et al., 2012; Gnugnoli et al., 2021; Lademann et al., 2017; Morrison et al., 2004; Shim et al., 2005;

2007; Tsukuda et al., 2005; van Attikum et al., 2004; 2007). This raises the question, whether these remodelers have distinct functions at DSBs or whether they act redundantly.

Nucleosome remodelers are found to be either single protein enzymes or multi-protein complexes. Historically, four major sub-families of remodelers have been proposed (Clapier and Cairns, 2009), but phylogenetic analysis based on sequence conservation of the catalytic ATPase subunits showed the existence of additional sub-families (Flaus et al., 2006). Five sub-families are found throughout eukaryotes – ISWI, SWI/SNF, CHD1, INO80 and Fun30/ETL. In contrast, ALC1, CHD7 and Mi2/NURD sub-families are not found throughout eukaryotes, with ALC1 and CHD7 orthologues specifically found in metazoans (Tong et al., 1998; Xue et al., 1998; Zhang et al., 1998; Ma et al., 2008; Bouazoune and Kingston, 2012). **Table 1** summarizes the different remodeler sub-families with their putative catalytic activities and involvement in DSB repair.

It seems expedient to group remodelers not only by evolutionary conservation, but also by functional similarity (**Figure 2, Table 1**). *In vitro* and *in vivo* we can discriminate at least three activities of nucleosome remodelers: 1) sliding/eviction leads to movement of nucleosomes along DNA that can even result in the removal of the entire nucleosome (**Figure 2A**); 2) positioning involves movement of nucleosomes to form regularly spaced nucleosomal arrays (**Figure 2B**); 3) editing involves the exchange of histones (commonly H2A-H2B dimers) to alter the composition of nucleosomes (Clapier and Cairns, 2009; Clapier et al., 2017). Based on studies of genome-wide chromatin organization, we currently think that SWI/SNF sub-family complexes (SWI/SNF and RSC in yeast) act as major sliding/eviction enzymes, that ISWI and CHD1 sub-family remodelers as well as INO80-C act as positioning enzymes and that INO80 sub-family complexes (SWR1 and INO80 in yeast) catalyze editing (**Table 1**, Clapier and Cairns, 2009; Clapier et al., 2017). In the following, we will investigate whether nucleosome remodelers carry out the same activities at DSBs.

NUCLEOSOME EVICTION AND RESECTION ARE COUPLED

With nucleosomes forming a barrier to resection, nucleosome eviction is the most straight-forward solution to allow spreading of resection into chromatin (**Figure 3**). Indeed, nucleosomes are lost around DSBs in the region where resection occurs (Bantele and Pfander, 2019; Chen et al., 2008; Mimitou et al., 2017; Tsukuda et al., 2005; 2009; van Attikum et al., 2007). While it was proposed that nucleosomes may associate in some form with resected, single-stranded DNA to form single-stranded nucleosomes (Adkins et al., 2017; Huang et al., 2018), a dedicated study did not find evidence to support wide-spread association of nucleosomes with single-stranded DNA *in vivo* (Peritore et al., 2021). But how do nucleosomes become evicted and how do sliding/evicting nucleosome remodelers of the SWI/SNF sub-family facilitate this eviction (**Figure 3A**)? In budding yeast, the SWI/SNF and RSC complexes are specifically recruited to DSBs (Chai et al., 2005; Shim et al., 2005, 2007; Kent et al.,

TABLE 1 | Overview of nucleosome remodeler sub-families and their members.

Family	Sub-family	Putative activity	<i>S. cerevisiae</i>	<i>H. sapiens</i> orthologues	Function at DSBs
Snf2-like	SWI/SNF	Nucleosome sliding/eviction	SWI/SNF RSC	BAF PBAF	Delamarre et al. (2020), Hays et al. (2020), Hu et al. (2020), Kakarougkas et al. (2014), Kent et al. (2007), Lee et al. (2010), Meisenberg et al. (2019), Ogiwara et al. (2011), Peng et al. (2009), Peritore et al. (2021), Qi et al. (2015), Shim et al. (2005), Shim et al. (2007), Ui et al. (2014), Watanabe et al. (2014), Wiest et al. (2017)
	ISWI	Nucleosome positioning	lsw1a	ACF CHRAC	Casari et al. (2021), Delamarre et al. (2020), Helfricht et al. (2013), Lan et al. (2010), Nakamura et al. (2011), Pessina and Lowndes, (2014), Sánchez-Molina et al. (2011), Sheu et al. (2010), Smeenk et al. (2012), Toiber et al. (2013), Vidi et al. (2014), Xiao et al. (2009)
			lsw1b	NoRC RSF	
			lsw2	WICH NURF CERF	
	CHD-I	Nucleosome positioning	Chd1	CHD1, CHD2	Delamarre et al. (2020), Gnugnoli et al. (2021), Kari et al. (2016), Luijsterburg et al. (2016), Zhou et al. (2018)
	CHD-II	?	-	Mi-2/NuRD	Chou et al. (2010), Goodarzi et al. (2011), Larsen et al. (2010), Luijsterburg et al. (2012), Pan et al. (2012), Polo et al. (2010), Qi et al. (2016), Smeenk et al. (2010), Smith et al. (2018), Spruijt et al. (2016)
CHD-III	?	-	CHD6, CHD7, CHD8, CHD9	Rother et al. (2020)	
Swr1-like	ALC1	?	-	ALC1	Ahel et al. (2009), Blessing et al. (2020), Juhász et al. (2020), Sellou et al. (2016)
	INO80	Nucleosome editing	INO80	INO80	Adkins et al. (2013), Alatwi and Downs, (2015), Bennett et al. (2013), Brahma et al. (2017), Chen et al. (2012), Downs et al. (2004), Kalocsay et al. (2009), Lademann et al. (2017), Morillo-Huesca et al. (2010), Morrison et al. (2004), Oberbeckmann et al. (2021b), Papamichos-Chronakis et al. (2006), Tsukuda et al. (2005), van Attikum et al. (2004), van Attikum et al. (2007)
	SWR1	Nucleosome positioning	SWR1	SRCAP TRAPP/Tip60	Bantele et al. (2017), Chen et al. (2012), Costelloe et al. (2012), Densham et al. (2016), Eapen et al. (2012)
	Fun30/ETL	?	Fun30	SMARCAD1	

Nucleosome remodelers are grouped into two families based on conservation of the ATPase subunit: Snf2-like and Swr1-like. Both families have several sub-families.



Snf2-like: The SWI/SNF (switch/sucrose non-fermentable) sub-family consists of two members in budding yeast - SWI/SNF and RSC (remodels the structure of chromatin) – as well as in human – BAF and PBAF. For human BAF variant complexes can be found harbouring ATPase subunit paralogs (Mittal and Roberts, 2020). The ISWI (imitation switch) sub-family in yeast contains 3 active complexes – lsw1a, lsw1b, lsw2 - that combine 2 different catalytic subunits - lsw1 and lsw2 - with different sets of proteins. For humans the setup with 2 catalytic subunits is similar, but with a higher number of different complexes: ACF, CHRAC, NoRC, RSF, WICH, NURF, CERF (Aydin et al., 2014). The CHD (chromodomain helicase DNA-binding) sub-family has a single member in yeast – Chd1 - and 3 subfamilies with in total 9 members in human: CHD1-2, CHD3-5 – forming NuRD/Mi-2 complex and CHD6-9 (Marfella and Imbalzano, 2007). The ALC1 sub-family carries a macrodomain for poly(ADP-ribose)-binding instead of a chromodomain and is found in human (Ahel et al., 2009).

Swr1-like: The INO80 (inositol requiring) sub-family has two members in yeast: INO80 and SWR1. In humans again there is additional complexity of this sub-family with INO80, SRCAP and TRAPP/Tip60 complexes (Willhoft and Wigley, 2020). The Fun30/ETL sub-family contains Fun30 in yeast and SMARCAD1 in human (Bantele and Pfander, 2019). Even though nucleosome remodelers appear to follow a highly similar enzymatic mechanism, they appear to exhibit distinct activities in chromatin organization. These putative activities are given along studies showing possible functions at DNA double strand breaks.

2007; Liang et al., 2007; Bennett et al., 2013; Bennett and Peterson, 2015; Wiest et al., 2017), suggesting that they may act during DSB repair or signaling. To interrogate the function of SWI/SNF and RSC, deletion of non-essential subunits or conditional depletion of the essential catalytic subunits have been used. Interestingly, interfering with either SWI/SNF or RSC function induced a defect already in the association of Mre11-C with DSBs (Shim et al., 2007; Wiest et al., 2017), suggesting that these remodelers could act at an early stage of DSB repair. Notably, under single-mutant conditions, resection and DSB repair were found to be delayed or reduced, but not abolished. Recently, experimental conditions were established that allowed to simultaneously induce the degradation of the ATPase subunits of both SWI/SNF and RSC (Peritore et al., 2021). Under these double mutant conditions, we find that nucleosome eviction and resection are both blocked (Peritore et al., 2021). This indicates that 1) SWI/SNF and RSC are redundantly required for DNA end resection and that 2) resection and nucleosome eviction are intrinsically coupled. Altogether, these data are consistent with SWI/SNF and RSC complexes playing a major role as nucleosome evictors also

in the context of DSBs. The single mutant data (Shim et al., 2007; Wiest et al., 2017), which showed defects in recruitment of Mre11-C, suggest that both complexes play an early role and evict or move DSB-proximal nucleosomes to allow binding of Mre11-C as well as resection initiation. Whether SWI/SNF or RSC influence Mre11-C activity, endonucleolytic clipping in particular, remains to be tested.

These findings raise the question of how these nucleosome remodelers sense the presence of a DSB and become recruited to DSB-proximal chromatin at such an early stage. Notably, RSC and SWI/SNF localize to the proximity of DSBs independently of each other and follow different recruitment kinetics, suggesting that they recognize DSBs by different mechanisms. RSC is recruited to a DSB within 10 min and thereby precedes resection initiation (Chai et al., 2005). While its recruitment kinetics are therefore similar to those of Mre11-C (Shim et al., 2007), we currently do not understand which signal is being recognized by RSC. SWI/SNF in contrast shows significantly slower recruitment (Chai et al., 2005) that depends on nucleosome modifications. Specifically, histone



acetylation is recognized by SWI/SNF and appears to lead to its DSB recruitment, consistent with the presence of several acetylation-binding bromodomains in the SWI/SNF complex (Bennett and Peterson, 2015; Cheng et al., 2021). Notably, the histone acetyltransferase NuA4 is specifically recruited to DSBs and this recruitment was shown to depend on Mre11-C (Cheng et al., 2021). The fact that SWI/SNF was found to be required for recruitment of Mre11-C, but at the same time also dependent on Mre11-C activity (Shim et al., 2005, 2007; Wiest et al., 2017) is not

necessarily a contradiction, but could suggest the presence of a positive feedback loop that promotes resection initiation.

Biochemical data suggest that long-range resection should be particularly dependent on nucleosome eviction (Adkins et al., 2013). Consistently, SWI/SNF appears to stimulate long range resection (Wiest et al., 2017), but this has yet to be correlated with nucleosome eviction. Altogether, these data show that RSC and SWI/SNF complexes promote DNA end resection in budding yeast and likely do so by acting as nucleosome evictors. Detailed biochemical and cell biological analysis will however be needed to

pinpoint exactly at which steps of DNA end resection these nucleosome remodelers act. These studies also need to account for the fact that RSC and SWI/SNF may also influence the long-range chromatin response to DSBs (on the 10 kb–1 Mb range). Indeed, γ H2A - the long-range chromatin mark of DSBs - was found to be reduced in RSC mutants (Kent et al., 2007; Shim et al., 2007), but it is unclear whether this effect relates to nucleosome eviction.

Lastly, SWI/SNF is also required later during HR, as SWI/SNF mutants show defects in synapsis and strand invasion (Chai et al., 2005). It is currently unclear whether this is due to defects in resection, due to a second “late” role in HR or due to long-range chromatin changes on the broken chromosome.

Nucleosome eviction appears to be conserved in human remodeler complexes. Human BAF and PBAF complexes are recruited to sites of DSBs (Park et al., 2006; Hays et al., 2020). Moreover, they appear to promote resection, possibly by acting on the Mre11-C activator CtIP (Hays et al., 2020). This suggests an early role in resection and it will be interesting to investigate whether this function is linked to nucleosome eviction.

While SWI/SNF and RSC are the major players in nucleosome eviction, it could be possible that also other nucleosome remodelers evict nucleosomes during DSB repair and resection. In particular, the INO80 complex has been linked to the eviction of nucleosomes at sites of transcription and DSBs as well (Tsukuda et al., 2005; van Attikum et al., 2007; Qiu et al., 2020), but given several functions of INO80 during DSB repair (see below) this activity is particularly challenging to ascertain. Additionally, whatever this INO80 complex function is, it appears to act differently from SWI/SNF and RSC complexes (Peritore et al., 2021). In all, we therefore conclude that 1) histone eviction occurs at DSBs, that 2) it is critical for DSB resection and repair and that 3) it appears to be mediated by the major cellular eviction activities of the SWI/SNF sub-family complexes.

THE ROLE OF NUCLEOSOME POSITIONING AT DOUBLE STRAND BREAKS REMAINS TO BE DETERMINED

Nucleosomes are positioned in a non-random fashion throughout the genome. In particular, a specific organization is seen at sites of transcribed genes, where a nucleosome-free region marks or neighbors the transcription start site, followed by regularly spaced nucleosomal arrays (Yuan et al., 2005; Weiner et al., 2010; Baldi et al., 2020). Positioning remodelers are responsible for the characteristic spacing of nucleosomes within such nucleosome arrays (Baldi et al., 2020). To generate the specific spacing of nucleosomes within the array, positioning remodelers use intrinsic ruler mechanisms as well as sensing of DNA shapes (Yamada et al., 2011; Krietenstein et al., 2016; Oberbeckmann et al., 2021a; Oberbeckmann et al., 2021b). The generation of nucleosome arrays has been extensively studied in budding yeast, where a combination of *in vitro* and *in vivo* studies suggests that four remodelers – Chd1, ISW1a, ISW2 and INO80 – can specifically position nucleosomes to form nucleosome

arrays (Gkikopoulos et al., 2011; Krietenstein et al., 2016; Ocampo et al., 2016; Kubik et al., 2019; Oberbeckmann et al., 2021b). Importantly, these remodelers also sense the presence of barrier-factors bound at specific sites in the genome to which the array is aligned to or “phased” (Eaton et al., 2010; Li et al., 2015; Krietenstein et al., 2016; Kubik et al., 2018; Rossi et al., 2018). Typical barrier factors are DNA-binding factors, like the abundant general regulatory factors Abf1, Rap1 or Reb1 in budding yeast or genome organizing factors like CTCF in mammals or Phaser in flies (Fu et al., 2008; Wiechens et al., 2016; Baldi et al., 2018). Importantly, recent *in vitro* work suggests that also DSBs are sensed as a barrier-factor by nucleosome remodelers and guide the formation of nucleosome arrays (Oberbeckmann et al., 2021a).

The finding that regularly spaced nucleosome arrays can form around DSBs in *in vitro* systems raises two questions: do remodelers position nucleosomes to form arrays around DSBs also *in vivo* and would such arrays promote DNA end resection? Experimentally, nucleosome positioning is typically investigated using micrococcal nuclease (MNase), which cleaves preferentially non-nucleosomal DNA. Several studies that used MNase to investigate nucleosome localization around a single DSB showed eviction of DSB-proximal nucleosomes, but came to different conclusions as to whether DSB-distal nucleosomes would shift their position (Kent et al., 2007; Shim et al., 2007; Tsabar et al., 2016). While these results are seemingly contradictory, this may simply be due to the fact that results from a single DSB are difficult to interpret. For example the newly formed array can be indistinguishable from the initial nucleosome positions, if the DSB and initial barrier factor are located at the same position. To overcome these limitations, a recent study utilized the *PHO5* gene, with its well characterized nucleosomal array and found evidence for eviction of the break-proximal nucleosome as well as repositioning of further distal nucleosomes (Tripuraneni et al., 2021). Further studies will need to show whether repositioned nucleosomes are indeed aligned to the DSB and whether the DSB itself or DSB-associated proteins serve as barrier. Furthermore, studies need to identify, if arrays are generated by positioning remodelers Chd1, ISW1a, ISW2 or INO80.

Interestingly, several studies in both yeast and human cells, point towards a function of these specific remodelers in promoting homologous recombination (Lan et al., 2010; Nakamura et al., 2011; Smeenk et al., 2012; Toiber et al., 2013; Kari et al., 2016; Zhou et al., 2018; Rother et al., 2020; Casari et al., 2021; Gnugnoli et al., 2021). In particular, remodelers of ISWI, CHD1 and CHD7 sub-families appear to be recruited to sites of DNA damage and to stimulate resection (Smeenk et al., 2012; Toiber et al., 2013; Kari et al., 2016; Delamarre et al., 2020; Rother et al., 2020; Gnugnoli et al., 2021). The precise mechanism by which these remodelers promote resection and HR is however uncertain. Moreover, even if these remodelers established nucleosome arrays around DSBs, it is at this point entirely unclear whether such arrays will have a positive function in DSB repair or whether they are simply a consequence of the enzymatic mechanism of positioning remodelers (Baldi et al.,

2018). Therefore, despite first hints that nucleosome arrays could form in the proximity of DSBs, the role of positioning remodelers in DSB repair still needs to be determined.

NUCLEOSOME EDITING AND H2A.Z EXCHANGE GUIDE DOUBLE STRAND BREAK REPAIR

Nucleosome editing describes the activity of exchanging canonical histone subunits with non-canonical histone variants and vice versa (Das and Tyler, 2013; Venkatesh and Workman, 2015). Nucleosome remodelers can facilitate editing by catalyzing the exchange of histone dimers. In eukaryotes, several histone variants exist primarily for H2A and H3 (Draizen et al., 2016; Talbert and Henikoff, 2010; 2017; 2021). While the principal mechanism for the incorporation of H3 and its variants is via *de novo* assembly of nucleosomes, H2A and its variants can be incorporated into existing nucleosomes by H2A-H2B dimer exchange (reviewed in Luger et al., 2012). In this chapter, we will therefore concentrate on nucleosome editing of H2A. Nucleosome editing has been extensively studied in budding yeast, where two H2A variants exist: H2A (which includes features of H2A.X) and H2A.Z (Santisteban et al., 2000). The SWR1 complex catalyzes the incorporation of H2A.Z-H2B dimers (Krogan et al., 2003; Mizuguchi et al., 2004). Furthermore, the INO80 complex is thought to catalyze the reverse reaction, the exchange of H2A.Z-H2B with H2A-H2B dimers (Papamichos-Chronakis et al., 2011; Brahma et al., 2017). This model of INO80 function is based on the principal finding that deletion of the H2A.Z gene *HTZ1* genetically suppresses many phenotypes of mutants deficient in INO80 function (Lademann et al., 2017; Papamichos-Chronakis et al., 2006; 2011).

The SWR1 complex is the prototypical nucleosome editing remodeler: mechanistically, it is able to translocate short stretches of DNA with no changes in nucleosome position, which then allows H2A-H2B dimers to be exchanged for H2A.Z-H2B dimers (Wu et al., 2009; Luk et al., 2010; Ranjan et al., 2015; Willhoft et al., 2018; Singh et al., 2019). In budding yeast, the SWR1 complex incorporates H2A.Z into chromatin around DSBs, as indicated by 1) the recruitment of SWR1 to DSB sites (van Attikum et al., 2007; Morillo-Huesca et al., 2010) and 2) a transient increase in H2A.Z occupancy in the DSB-surrounding chromatin shortly after DSB induction (Kalocsay et al., 2009). A transiently increased incorporation of H2A.Z into DSB-proximal chromatin was observed also in human cells (Xu et al., 2012; Nishibuchi et al., 2014; Alatwi and Downs, 2015; Guroy-Yuzugullu et al., 2015). Compared to canonical nucleosomes, H2A.Z-containing nucleosomes are more labile (Abbott et al., 2001; Zhang et al., 2005; Jin and Felsenfeld, 2007) suggesting that their presence will promote DNA end resection. Consistently, yeast cells lacking H2A.Z show a pronounced resection defect (Kalocsay et al., 2009; Lademann et al., 2017). In contrast, the absence of SWR1 causes a much milder resection phenotype (van Attikum et al., 2007; Chen et al., 2012; Adkins et al., 2013). These data suggest that either 1) H2A.Z becomes incorporated at DSB sites by an SWR1-independent mechanism or that 2) H2A.Z-incorporation into DSB-surrounding chromatin is not a major regulator of resection and that H2A.Z regulates resection by means independent from its incorporation in DSB-surrounding chromatin.

If H2A.Z-incorporation into DSB-proximal chromatin promotes resection, there are two putative mechanisms by which it could do so. First, the aforementioned reduction of nucleosome stability may allow remodelers or even resection nucleases to bypass and evict H2A.Z-containing nucleosomes (Adkins et al., 2013). Second, H2A.Z could serve as binding platform for associated factors (Xu et al., 2012) as has been shown for nucleotide excision repair (Yu et al., 2013). Binding of factors to H2A.Z or SUMO-modified H2A.Z is for example thought to lead to relocalization of DSBs to the nuclear periphery (Nagai et al., 2008; Kalocsay et al., 2009; Oza et al., 2009; Horigome et al., 2014). Relocalization of DSBs is also observed in *Drosophila*, where heterochromatic DSBs are first brought to the periphery of the heterochromatic domain (Chiolo et al., 2011) and then to the nuclear pore complex (Ryu et al., 2015). Similarly, in mammalian cells DSB relocation to discrete clusters in the periphery of heterochromatin has been observed (Jakob et al., 2011; Tsouroula et al., 2016; Schrank et al., 2018), but a connection between DSB relocation and H2A.Z has not been shown so far. Therefore, nucleosome editing and H2A.Z incorporation are used to regulate DSB repair, but the underlying molecular mechanisms warrant further investigation.

The importance of nucleosome editing for DSB repair raises the question whether H2A.Z incorporation becomes reversed at some point. Indeed, studies in budding yeast have shown that the INO80 complex is not only recruited to DSBs (Downs et al., 2004; Morrison et al., 2004; van Attikum et al., 2004; Bennett et al., 2013), but that it also counteracts H2A.Z incorporation (Papamichos-Chronakis et al., 2011). Also in human cells H2A.Z is removed from chromatin surrounding DSB sites (Xu et al., 2012; Nishibuchi et al., 2014; Alatwi and Downs, 2015; Guroy-Yuzugullu et al., 2015; Clouaire et al., 2018). While INO80's role as nucleosome editing and H2A.Z removal enzyme was initially controversial (Papamichos-Chronakis et al., 2011; Watanabe et al., 2013; Jeronimo et al., 2015; Tramantano et al., 2016; Wang et al., 2016; Watanabe and Peterson, 2016), recent structural work showed that besides its nucleosome positioning activity, the INO80 complex may be able to catalyze translocation of short stretches of DNA without nucleosome sliding, consistent with histone dimer exchange activity (Ayala et al., 2018; Eustermann et al., 2018). This suggests that at DSBs INO80 may have at least two activities: 1) a nucleosome positioning activity (see above) and 2) a nucleosome editing activity (Papamichos-Chronakis et al., 2006; Alatwi and Downs, 2015; Brahma et al., 2017; Lademann et al., 2017). Consistent with INO80 antagonizing the SWR1 complex and removing H2A.Z from chromatin, mutants deficient in INO80 complex function accumulate H2A.Z around DSBs (Papamichos-Chronakis et al., 2006; Alatwi and Downs, 2015; Lademann et al., 2017). The dual remodeling activity of the INO80 complex complicates the interpretation of *ino80* mutant phenotypes. To overcome this issue, deletion of the H2A.Z gene *HTZ1* has been used, because it suppresses phenotypes arising from an H2A.Z removal defect. Using this approach, an H2A.Z removal function of the INO80 complex was found to promote the formation of the Rad51 nucleosome-protein filament downstream of resection

(Lademann et al., 2017). In contrast, a resection-promoting function of the INO80 complex was found to be independent of H2A.Z (Lademann et al., 2017) and therefore unrelated to nucleosome editing. Moreover, also in human cells, nucleosome editing by the INO80 complex is important for DSB repair and acts after DNA end resection (Alatwi and Downs, 2015). Taken together, a picture emerges whereby nucleosome editing and H2A.Z incorporation by the SWR1 complex is involved in regulation of DNA end resection in yeast, while generally and throughout eukaryotes H2A.Z removal in DSB-surrounding chromatin is important for DSB repair, but likely acts only after resection.

FUN30/SMARCAD1 PROMOTE RESECTION BY ANTAGONIZING RESECTION-INHIBITORY FACTORS

Fun30 (from budding yeast), ETL1 (from mouse) and SMARCAD1 (from human) are the prototypical members of a sub-family of nucleosome remodelers that is evolutionary conserved throughout eukaryotes (Clark et al., 1992; Adra et al., 2000; Flaus et al., 2006). Historically they have not been considered major nucleosome remodelers and their molecular mechanisms have not yet been entirely elucidated (Bantele and Pfander, 2019). Recently, a study by the Luger lab suggested that SMARCAD1 evicts and also assembles entire nucleosomes by a mechanism that involves unique contacts between remodeler and nucleosome (Markert et al., 2021). Work with yeast Fun30 suggests that it can slide nucleosomes and mediate histone dimer exchange (Awad et al., 2010).


A key function of yeast Fun30 and human SMARCAD1 appears to be the stimulation of long-range resection (Chen et al., 2012; Costelloe et al., 2012; Eapen et al., 2012). For example, in budding yeast cells lacking Fun30, long-range resection of a non-repairable DSB is 2-3-fold slower than in WT cells (Eapen et al., 2012; Bantele et al., 2017). Accordingly, *fun30* mutants scored similarly to mutants deficient in the long-range resection nucleases, when they were initially found in screens for resection-dependent repair of DSBs (Chen et al., 2012; Costelloe et al., 2012). Moreover, an evolutionary conserved pathway facilitates recruitment of Fun30 to sites of DNA end resection. This pathway requires the 9-1-1 complex as recruitment platform at the ssDNA-dsDNA junction and is activated during cell cycle phases (S-M phase), when also resection is activated (Chen et al., 2016; Bantele et al., 2017).

In contrast, Fun30 did not stimulate Exo1's ability to resect through a nucleosome in an *in vitro* system (Adkins et al., 2013). This finding raises the possibility that a crucial factor was missing from these reconstituted systems. Consistently, *fun30* mutant phenotypes can be suppressed by the additional depletion of the resection inhibitor Rad9 from yeast cells (Chen et al., 2012; Bantele et al., 2017). These data indicate a functional antagonism between Fun30 and Rad9. Notably, also in human cells SMARCAD1 acts as resection activator, while the Rad9 orthologue 53BP1 is a resection inhibitor (Lazzaro et al., 2008; Bunting et al., 2010; Bothmer et al., 2011; Costelloe et al., 2012; Densham et al., 2016), suggesting that the antagonism of both factors is conserved throughout eukaryotic

evolution (please see (Sanchez et al., 2021)) in this issue for a detailed review on the interaction between 53BP1 and BRCA1 in the DSB repair decision). Notably, Rad9, 53BP1, as well as the fission yeast orthologue Crb2 associate with chromatin and have all been shown to bind to nucleosomes, where they recognize specific histone modifications (Huyen et al., 2004; Nakamura et al., 2004; Sanders et al., 2004; Wysocki et al., 2005; Botuyan et al., 2006; Du et al., 2006; Toh et al., 2006; Grenon et al., 2007; Hammett et al., 2007; Fradet-Turcotte et al., 2013; Wilson et al., 2016; Hu et al., 2017; Kilic et al., 2019). Rad9 orthologues appear to recognize distinct histone marks, but in each case two or more histone marks are bound (reviewed in Marini et al., 2019; Panier and Boulton, 2014), suggesting that Rad9 orthologues are multivalent histone binders. We therefore hypothesize that both Fun30 and Rad9 influence DSB-surrounding chromatin in an antagonistic fashion and that Fun30 specifically acts on Rad9-bound nucleosomes (Bantele and Pfander, 2019).

In budding yeast cells lacking both Fun30 and Rad9, resection and nucleosome eviction are fully functional (Peritore et al., 2021), suggesting that Fun30 is not required to overcome the general nucleosome barrier and that it is not the essential nucleosome evictor at DSBs. Alternatively, Fun30 may rather catalyze the direct removal of Rad9 from nucleosomes (Figure 4A) or it may counteract Rad9 association with nucleosomes by catalyzing histone dimer exchange which may remove one or more binding site(s) for Rad9 (Figure 4B). Lastly, it is possible that Fun30 slides or even entirely evicts Rad9-bound nucleosomes (Figure 4C). Given that Rad9 and Fun30 antagonize each other on multiple levels, including also the competition for binding to the scaffold protein Dpb11 (Granata et al., 2010; Pfander and Diffley, 2011; Bantele et al., 2017), future biochemical and structural studies will be needed to reveal the mechanism by which Fun30 promotes DNA end resection.

Also human SMARCAD1 antagonizes 53BP1. Depletion of SMARCAD1 stabilizes 53BP1 around DSB sites (Densham et al., 2016). However, resection regulation in human cells is more complex compared to yeast as besides SMARCAD1 a second resection promoting factor exist, the BRCA1-BARD1 complex (reviewed in Densham and Morris, 2019; Sanchez et al., 2021). BRCA1-BARD1 form an E3 ubiquitin-ligase complex that mediates ubiquitylation of H2A (Kalb et al., 2014; Densham et al., 2016; Leung et al., 2017; Nakamura et al., 2019). BRCA1-BARD1 is likely to act upstream of SMARCAD1, as ubiquitin-modified H2A promotes SMARCAD1 binding to nucleosomes around DSBs (Densham et al., 2016). Therefore, SMARCAD1 function has to be seen in the context of post-translational histone modifications, which affect DSB-surrounding chromatin. DSB-localized SMARCAD1 may also become post-translationally modified itself, including phosphorylation by the ATM kinase and ubiquitylation by the RING1 ubiquitin ligase (Chakraborty et al., 2018), which appears to activate the pro-resection function of SMARCAD1. These factors need to be taken into consideration for biochemical studies that ultimately will allow to understand whether the Fun30/SMARCAD1 sub-family remodelers facilitate resection by nucleosomes sliding and eviction, positioning or editing and whether it acts on nucleosomes or rather on nucleosome-associated proteins.



CONCLUSION

In all, we think that previous studies collectively indicate that nucleosome remodelers may serve similar roles during DSB repair as during gene transcription with nucleosome eviction, editing and potentially even positioning taking place at DSBs. Knowledge of the specific activities of individual nucleosome remodelers and of their redundancies thereby offers the potential to get to grips with chromatin changes occurring at DSBs. Moreover, we think that studies of DSB resection and repair may be generally inspired by analogies to gene transcription. Both processes appear to be similarly affected by the presence of chromatin, with nucleosomes forming a dynamic barrier and nucleosome remodelers facilitating its bypass.

Importantly, while nucleosomes clearly form a barrier to the resection nucleases, nucleosome remodelers equip cells with multiple ways to overcome this barrier. In this review, we have outlined several putative mechanisms of how bypass may occur. These include eviction, sliding and editing of nucleosomes. While we are still only beginning to understand how the nucleosome barrier is overcome, a key future question will be which bypass mechanism is chosen in which cellular scenario. Importantly, the nucleosome barrier and its dynamic nature offers additional possibilities to regulate resection and DSB repair. Moreover, critical factors of the DSB repair decision, such as 53BP1 and BRCA1, are proteins that bind and modify nucleosomes. Therefore, we propose that convergence of

resection-regulatory pathways on nucleosomes is a central part of the cellular DSB repair decision.

AUTHOR CONTRIBUTIONS

LAK, MP, and LG contributed equally to writing of the manuscript. LAK, MP, LG and BP developed concepts presented here and wrote the manuscript.

FUNDING

This work was supported by the Max Planck Society and a grant by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)–Project-ID 213249687–SFB 1064 (all to BP).

ACKNOWLEDGMENTS

We thank Till Bartke, Sigurd Braun and Philip Korber as well as members of the Pfander lab for discussions. We apologize to those researchers, whose work we could not cite due to space restrictions.

REFERENCES

- Abbott, D. W., Ivanova, V. S., Wang, X., Bonner, W. M., and Ausió, J. (2001). Characterization of the Stability and Folding of H2A.Z Chromatin Particles. *J. Biol. Chem.* 276, 41945–41949. doi:10.1074/jbc.M108217200
- Adkins, N. L., Niu, H., Sung, P., and Peterson, C. L. (2013). Nucleosome Dynamics Regulates DNA Processing. *Nat. Struct. Mol. Biol.* 20, 836–842. doi:10.1038/nsmb.2585
- Adkins, N. L., Swygert, S. G., Kaur, P., Niu, H., Grigoryev, S. A., Sung, P., et al. (2017). Nucleosome-like, Single-Stranded DNA (ssDNA)-Histone Octamer Complexes and the Implication for DNA Double Strand Break Repair*. *J. Biol. Chem.* 292, 5271–5281. doi:10.1074/jbc.m117.776369
- Adra, C. N., Donato, J.-L., Badovinac, R., Syed, F., Kheraj, R., Cai, H., et al. (2000). SMARCAD1, a Novel Human Helicase Family-Defining Member Associated with Genetic Instability: Cloning, Expression, and Mapping to 4q22-Q23, a Band Rich in Breakpoints and Deletion Mutants Involved in Several Human Diseases. *Genomics* 69, 162–173. doi:10.1006/geno.2000.6281
- Ahel, D., Hořejší, Z., Wiechens, N., Polo, S. E., Garcia-Wilson, E., Ahel, I., et al. (2009). Poly(ADP-ribose)-dependent Regulation of DNA Repair by the Chromatin Remodeling Enzyme ALC1. *Science* 325, 1240–1243. doi:10.1126/science.1177321
- Alatwi, H. E., and Downs, J. A. (2015). Removal of H2A.Z by INO 80 Promotes Homologous Recombination. *EMBO Rep.* 16, 986–994. doi:10.15252/embr.201540330
- Anand, R., Ranjha, L., Cannavo, E., and Cejka, P. (2016). Phosphorylated CtIP Functions as a Co-factor of the MRE11-RAD50-NBS1 Endonuclease in DNA End Resection. *Mol. Cell* 64, 940–950. doi:10.1016/j.molcel.2016.10.017
- Awad, S., Ryan, D., Prochasson, P., Owen-Hughes, T., and Hassan, A. H. (2010). The Snf2 Homolog Fun30 Acts as a Homodimeric ATP-dependent Chromatin-Remodeling Enzyme. *J. Biol. Chem.* 285, 9477–9484. doi:10.1074/jbc.M109.082149
- Ayala, R., Willhoft, O., Aramayo, R. J., Wilkinson, M., McCormack, E. A., Ocloo, L., et al. (2018). Structure and Regulation of the Human INO80-Nucleosome Complex. *Nature* 556, 391–395. doi:10.1038/s41586-018-0021-6
- Aydin, Ö. Z., Vermeulen, W., and Lans, H. (2014). ISWI Chromatin Remodeling Complexes in the DNA Damage Response. *Cell Cycle* 13, 3016–3025. doi:10.4161/15384101.2014.956551
- Baldi, S., Jain, D. S., Harpprecht, L., Zabel, A., Scheibe, M., Butter, F., et al. (2018). Genome-wide Rules of Nucleosome Phasing in *Drosophila*. *Mol. Cell* 72, 661–672. e4. doi:10.1016/j.molcel.2018.09.032
- Baldi, S., Korber, P., and Becker, P. B. (2020). Beads on a String-Nucleosome Array Arrangements and Folding of the Chromatin Fiber. *Nat. Struct. Mol. Biol.* 27, 109–118. doi:10.1038/s41594-019-0368-x
- Bantele, S. C., Ferreira, P., Gritenaite, D., Boos, D., and Pfander, B. (2017). Targeting of the Fun30 Nucleosome Remodeller by the Dpb11 Scaffold Facilitates Cell Cycle-Regulated DNA End Resection. *eLife Sci.* 6, e21687. doi:10.7554/eLife.21687
- Bantele, S. C. S., and Pfander, B. (2019). Nucleosome Remodeling by Fun30SMARCAD1 in the DNA Damage Response. *Front. Mol. Biosci.* 6. doi:10.3389/fmolb.2019.00078
- Bennett, G., Papamichos-Chronakis, M., and Peterson, C. L. (2013). DNA Repair Choice Defines a Common Pathway for Recruitment of Chromatin Regulators. *Nat. Commun.* 4, 2084. doi:10.1038/ncomms3084
- Bennett, G., and Peterson, C. L. (2015). SWI/SNF Recruitment to a DNA Double-Strand Break by the NuA4 and Gcn5 Histone Acetyltransferases. *DNA Repair* 30, 38–45. doi:10.1016/j.dnarep.2015.03.006
- Bird, A. W., Yu, D. Y., Pray-Grant, M. G., Qiu, Q., Harmon, K. E., Megee, P. C., et al. (2002). Acetylation of Histone H4 by Esa1 Is Required for DNA Double-Strand Break Repair. *Nature* 419, 411–415. doi:10.1038/nature01035
- Blessing, C., Mandemaker, I. K., Gonzalez-Leal, C., Preisser, J., Schomburg, A., and Ladurner, A. G. (2020). The Oncogenic Helicase ALC1 Regulates PARP Inhibitor Potency by Trapping PARP2 at DNA Breaks. *Mol. Cell* 80, 862–875. e6. doi:10.1016/j.molcel.2020.10.009
- Blier, P. R., Griffith, A. J., Craft, J., and Hardin, J. A. (1993). Binding of Ku Protein to DNA. Measurement of Affinity for Ends and Demonstration of Binding to Nicks. *J. Biol. Chem.* 268, 7594–7601. doi:10.1016/s0021-9258(18)53216-6
- Bonetti, D., Colombo, C. V., Clerici, M., and Longhese, M. P. (2018). Processing of DNA Ends in the Maintenance of Genome Stability. *Front. Genet.* 9, 390. doi:10.3389/fgene.2018.00390
- Bothmer, A., Robbiani, D. F., Di Virgilio, M., Bunting, S. F., Klein, I. A., Feldhahn, N., et al. (2011). Regulation of DNA End Joining, Resection, and Immunoglobulin Class Switch Recombination by 53BP1. *Mol. Cell* 42, 319–329. doi:10.1016/j.molcel.2011.03.019
- Botuyan, M. V., Lee, J., Ward, I. M., Kim, J.-E., Thompson, J. R., Chen, J., et al. (2006). Structural Basis for the Methylation State-specific Recognition of Histone H4-K20 by 53BP1 and Crb2 in DNA Repair. *Cell* 127, 1361–1373. doi:10.1016/j.cell.2006.10.043
- Bouazoune, K., and Kingston, R. E. (2012). Chromatin Remodeling by the CHD7 Protein Is Impaired by Mutations that Cause Human Developmental Disorders. *Proc. Natl. Acad. Sci. U S A.* 109, 19238–19243. doi:10.1073/pnas.1213825109
- Brahma, S., Udagama, M. I., Kim, J., Hada, A., Bhardwaj, S. K., Hailu, S. G., et al. (2017). INO80 Exchanges H2A.Z for H2A by Translocating on DNA Proximal to Histone Dimers. *Nat. Commun.* 8, 15616. doi:10.1038/ncomms15616
- Bunting, S. F., Callén, E., Wong, N., Chen, H.-T., Polato, F., Gunn, A., et al. (2010). 53BP1 Inhibits Homologous Recombination in Brca1-Deficient Cells by Blocking Resection of DNA Breaks. *Cell* 141, 243–254. doi:10.1016/j.cell.2010.03.012
- Cannavo, E., and Cejka, P. (2014). Sae2 Promotes dsDNA Endonuclease Activity within Mre11-Rad50-Xrs2 to Resect DNA Breaks. *Nature* 514, 122–125. doi:10.1038/nature13771
- Casari, E., Gobbin, E., Gnugnoli, M., Mangiagalli, M., Clerici, M., and Longhese, M. P. (2021). Dpb4 Promotes Resection of DNA Double-Strand Breaks and Checkpoint Activation by Acting in Two Different Protein Complexes. *Nat. Commun.* 12, 4750. doi:10.1038/s41467-021-25090-9
- Cejka, P., Cannavo, E., Polaczek, P., Masuda-Sasa, T., Pokharel, S., Campbell, J. L., et al. (2010). DNA End Resection by Dna2-Sgs1-RPA and its Stimulation by Top3-Rmi1 and Mre11-Rad50-Xrs2. *Nature* 467, 112–116. doi:10.1038/nature09355
- Cejka, P. (2015). DNA End Resection: Nucleases Team up with the Right Partners to Initiate Homologous Recombination. *J. Biol. Chem.* 290, 22931–22938. doi:10.1074/jbc.r115.675942
- Cejka, P., and Symington, L. S. (2021). DNA End Resection: Mechanism and Control. *Annu. Rev. Genet.* 55, 285–307. doi:10.1146/annurev-genet-071719-020312
- Ceppi, I., Howard, S. M., Kasaciunaite, K., Pinto, C., Anand, R., Seidel, R., et al. (2020). CtIP Promotes the Motor Activity of DNA2 to Accelerate Long-Range DNA End Resection. *Proc. Natl. Acad. Sci. USA* 117, 8859–8869. doi:10.1073/pnas.2001165117
- Chai, B., Huang, J., Cairns, B. R., and Laurent, B. C. (2005). Distinct Roles for the RSC and Swi/Snf ATP-dependent Chromatin Remodelers in DNA Double-Strand Break Repair. *Genes Dev.* 19, 1656–1661. doi:10.1101/gad.1273105
- Chakraborty, S., Pandita, R. K., Hambarde, S., Mattoo, A. R., Charaka, V., Ahmed, K. M., et al. (2018). SMARCAD1 Phosphorylation and Ubiquitination Are Required for Resection during DNA Double-Strand Break Repair. *iScience* 2, 123–135. doi:10.1016/j.isci.2018.03.016
- Chang, H. H. Y., Pannunzio, N. R., Adachi, N., and Lieber, M. R. (2017). Non-homologous DNA End Joining and Alternative Pathways to Double-Strand Break Repair. *Nat. Rev. Mol. Cell Biol.* 18, 495–506. doi:10.1038/nrm.2017.48
- Chen, C.-C., Carson, J. J., Feser, J., Tamburini, B., Zabaronic, S., Linger, J., et al. (2008). Acetylated Lysine 56 on Histone H3 Drives Chromatin Assembly after Repair and Signals for the Completion of Repair. *Cell* 134, 231–243. doi:10.1016/j.cell.2008.06.035
- Chen, X., Cui, D., Papusha, A., Zhang, X., Chu, C.-D., Tang, J., et al. (2012). The Fun30 Nucleosome Remodeller Promotes Resection of DNA Double-Strand Break Ends. *Nature* 489, 576–580. doi:10.1038/nature11355
- Chen, X., Niu, H., Yu, Y., Wang, J., Zhu, S., Zhou, J., et al. (2016). Enrichment of Cdk1-Cyclins at DNA Double-Strand Breaks Stimulates Fun30 Phosphorylation and DNA End Resection. *Nucleic Acids Res.* 44, 2742–2753. doi:10.1093/nar/gkv1544
- Cheng, X., Côté, V., and Côté, J. (2021). NuA4 and SAGA Acetyltransferase Complexes Cooperate for Repair of DNA Breaks by Homologous Recombination. *Plos Genet.* 17, e1009459. doi:10.1371/journal.pgen.1009459
- Chiolo, I., Minoda, A., Colmenares, S. U., Polyzos, A., Costes, S. V., and Karpen, G. H. (2011). Double-Strand Breaks in Heterochromatin Move outside of a

- Dynamic HP1a Domain to Complete Recombinational Repair. *Cell* 144, 732–744. doi:10.1016/j.cell.2011.02.012
- Chou, D. M., Adamson, B., Dephoure, N. E., Tan, X., Nottke, A. C., Hurov, K. E., et al. (2010). A Chromatin Localization Screen Reveals Poly (ADP Ribose)-Regulated Recruitment of the Repressive Polycomb and NuRD Complexes to Sites of DNA Damage. *Proc. Natl. Acad. Sci.* 107, 18475–18480. doi:10.1073/pnas.1012946107
- Ciccio, A., and Elledge, S. J. (2010). The DNA Damage Response: Making it Safe to Play with Knives. *Mol. Cell* 40, 179–204. doi:10.1016/j.molcel.2010.09.019
- Clapier, C. R., and Cairns, B. R. (2009). The Biology of Chromatin Remodeling Complexes. *Annu. Rev. Biochem.* 78, 273–304. doi:10.1146/annurev.biochem.77.062706.153223
- Clapier, C. R., Iwasa, J., Cairns, B. R., and Peterson, C. L. (2017). Mechanisms of Action and Regulation of ATP-dependent Chromatin-Remodelling Complexes. *Nat. Rev. Mol. Cell Biol* 18, 407–422. doi:10.1038/nrm.2017.26
- Clark, M. W., Zhong, W. W., Keng, T., Storms, R. K., Barton, A., Kaback, D. B., et al. (1992). Identification of a *Saccharomyces cerevisiae* Homolog of the SNF2 Transcriptional Regulator in the DNA Sequence of an 8-6 Kb Region in the LTE1-CYS1 Interval on the Left Arm of Chromosome I. *Yeast* 8, 133–145. doi:10.1002/yea.320080208
- Clouaire, T., Rocher, V., Lashgari, A., Arnould, C., Aguirrebengoa, M., Biernacka, A., et al. (2018). Comprehensive Mapping of Histone Modifications at DNA Double-Strand Breaks Deciphers Repair Pathway Chromatin Signatures. *Mol. Cell* 72, 250–262.e6. doi:10.1016/j.molcel.2018.08.020
- Costelloe, T., Louge, R., Tomimatsu, N., Mukherjee, B., Martini, E., Khadaroo, B., et al. (2012). The Yeast Fun30 and Human SMARCAD1 Chromatin Remodelers Promote DNA End Resection. *Nature* 489, 581–584. doi:10.1038/nature11353
- Daley, J. M., Niu, H., Miller, A. S., and Sung, P. (2015). Biochemical Mechanism of DSB End Resection and its Regulation. *DNA Repair* 32, 66–74. doi:10.1016/j.dnarep.2015.04.015
- Das, C., and Tyler, J. K. (2012). Histone Exchange and Histone Modifications during Transcription and Aging. *Biochim. Biophys. Acta (Bba) - Gene Regul. Mech.* 1819, 332–342. doi:10.1016/j.bbagr.2011.08.001
- Delamarre, A., Barthe, A., de la Roche Saint-André, C., Luciano, P., Forey, R., Padioulet, I., et al. (2020). MRX Increases Chromatin Accessibility at Stalled Replication Forks to Promote Nascent DNA Resection and Cohesin Loading. *Mol. Cell* 77, 395–410.e3. doi:10.1016/j.molcel.2019.10.029
- Densham, R. M., Garvin, A. J., Stone, H. R., Strachan, J., Baldock, R. A., Daza-Martin, M., et al. (2016). Human BRCA1-BARD1 Ubiquitin Ligase Activity Counteracts Chromatin Barriers to DNA Resection. *Nat. Struct. Mol. Biol.* 23, 647–655. doi:10.1038/nsmb.3236
- Densham, R. M., and Morris, J. R. (2019). Moving Mountains-The BRCA1 Promotion of DNA Resection. *Front. Mol. Biosci.* 6. doi:10.3389/fmolb.2019.00079
- Deshpande, R. A., Lee, J.-H., Arora, S., and Paull, T. T. (2016). Nbs1 Converts the Human Mre11/Rad50 Nuclease Complex into an Endo/Exonuclease Machine Specific for Protein-DNA Adducts. *Mol. Cell* 64, 593–606. doi:10.1016/j.molcel.2016.10.010
- Downs, J. A., Allard, S., Jobin-Robitaille, O., Javaheri, A., Auger, A., Bouchard, N., et al. (2004). Binding of Chromatin-Modifying Activities to Phosphorylated Histone H2A at DNA Damage Sites. *Mol. Cell* 16, 979–990. doi:10.1016/j.molcel.2004.12.003
- Draizen, E. J., Shaytan, A. K., Mariño-Ramírez, L., Talbert, P. B., Landsman, D., and Panchenko, A. R. (2016). HistoneDB 2.0: a Histone Database with Variants-An Integrated Resource to Explore Histones and Their Variants. *Database* 2016, baw014. doi:10.1093/database/baw014
- Du, L.-L., Nakamura, T. M., and Russell, P. (2006). Histone Modification-dependent and -independent Pathways for Recruitment of Checkpoint Protein Crb2 to Double-Strand Breaks. *Genes Dev.* 20, 1583–1596. doi:10.1101/gad.1422606
- Eapen, V. V., Sugawara, N., Tsabar, M., Wu, W.-H., and Haber, J. E. (2012). The *Saccharomyces cerevisiae* Chromatin Remodeler Fun30 Regulates DNA End Resection and Checkpoint Deactivation. *Mol. Cell Biol* 32, 4727–4740. doi:10.1128/MCB.00566-12
- Eaton, M. L., Galani, K., Kang, S., Bell, S. P., and MacAlpine, D. M. (2010). Conserved Nucleosome Positioning Defines Replication Origins. *Genes Dev.* 24, 748–753. doi:10.1101/gad.1913210
- Elbakry, A., and Löbrich, M. (2021). Homologous Recombination Subpathways: A Tangle to Resolve. *Front. Genet.* 12, 723847. doi:10.3389/fgene.2021.723847
- Eustermann, S., Schall, K., Kostrewa, D., Lakomek, K., Strauss, M., Moldt, M., et al. (2018). Structural Basis for ATP-dependent Chromatin Remodelling by the INO80 Complex. *Nature* 556, 386–390. doi:10.1038/s41586-018-0029-y
- Falzon, M., Fewell, J. W., and Kuff, E. L. (1993). EBP-80, a Transcription Factor Closely Resembling the Human Autoantigen Ku, Recognizes Single- to Double-Strand Transitions in DNA. *J. Biol. Chem.* 268, 10546–10552. doi:10.1016/s0021-9258(18)82233-5
- Flaus, A., Martin, D. M. A., Barton, G. J., and Owen-Hughes, T. (2006). Identification of Multiple Distinct Snf2 Subfamilies with Conserved Structural Motifs. *Nucleic Acids Res.* 34, 2887–2905. doi:10.1093/nar/gkl295
- Fradet-Turcotte, A., Canny, M. D., Escobedo-Díaz, C., Orthwein, A., Leung, C. C. Y., Huang, H., et al. (2013). 53BP1 Is a Reader of the DNA-Damage-Induced H2A Lys 15 Ubiquitin Mark. *Nature* 499, 50–54. doi:10.1038/nature12318
- Fu, Y., Sinha, M., Peterson, C. L., and Weng, Z. (2008). The Insulator Binding Protein CTCF Positions 20 Nucleosomes Around its Binding Sites across the Human Genome. *PLoS Genet.* 4, e1000138. doi:10.1371/journal.pgen.1000138
- Garcia, V., Phelps, S. E. L., Gray, S., and Neale, M. J. (2011). Bidirectional Resection of DNA Double-Strand Breaks by Mre11 and Exo1. *Nature* 479, 241–244. doi:10.1038/nature10515
- Gkikopoulos, T., Schofield, P., Singh, V., Pinskaya, M., Mellor, J., Smolle, M., et al. (2011). A Role for Snf2-Related Nucleosome-Spacing Enzymes in Genome-wide Nucleosome Organization. *Science* 333, 1758–1760. doi:10.1126/science.1206097
- Gnugnoli, M., Casari, E., and Longhese, M. P. (2021). The Chromatin Remodeler Chd1 Supports MRX and Exo1 Functions in Resection of DNA Double-Strand Breaks. *PLoS Genet.* 17, e1009807. doi:10.1371/journal.pgen.1009807
- Goodarzi, A. A., Kurka, T., and Jeggo, P. A. (2011). KAP-1 Phosphorylation Regulates CHD3 Nucleosome Remodeling during the DNA Double-Strand Break Response. *Nat. Struct. Mol. Biol.* 18, 831–839. doi:10.1038/nsmb.2077
- Granata, M., Lazzaro, F., Novarina, D., Panigada, D., Puddu, F., Abreu, C. M., et al. (2010). Dynamics of Rad9 Chromatin Binding and Checkpoint Function Are Mediated by its Dimerization and Are Cell Cycle-Regulated by CDK1 Activity. *PLoS Genet.* 6, e1001047. doi:10.1371/journal.pgen.1001047
- Grenon, M., Costelloe, T., Jimeno, S., O'Shaughnessy, A., FitzGerald, J., Zgheib, O., et al. (2007). Docking onto Chromatin via the *Saccharomyces cerevisiae* Rad9 Tudor Domain. *Yeast* 24, 105–119. doi:10.1002/yea.1441
- Gursoy-Yuzugullu, O., Ayrapetov, M. K., and Price, B. D. (2015). Histone chaperone Anp32e removes H2A.Z from DNA double-strand breaks and promotes nucleosome reorganization and DNA repair. *Proc. Natl. Acad. Sci. USA* 112, 7507–7512. doi:10.1073/pnas.1504868112
- Hammet, A., Magill, C., Heierhorst, J., and Jackson, S. P. (2007). Rad9 BRCT Domain Interaction with Phosphorylated H2AX Regulates the G1 Checkpoint in Budding Yeast. *EMBO Rep.* 8, 851–857. doi:10.1038/sj.embor.7401036
- Hays, E., Nettleton, E., Carter, C., Morales, M., Vo, L., Passo, M., et al. (2020). The SWI/SNF ATPase BRG1 Stimulates DNA End Resection and Homologous Recombination by Reducing Nucleosome Density at DNA Double Strand Breaks and by Promoting the Recruitment of the CtIP Nuclease. *Cell Cycle* 19, 3096–3114. doi:10.1080/15384101.2020.1831256
- Helfricht, A., Wiegant, W., Thijssen, P., Vertegaal, A., Luijsterburg, M., and van Attikum, H. (2013). Remodeling and Spacing Factor 1 (RSF1) Deposits Centromere Proteins at DNA Double-Strand Breaks to Promote Non-homologous End-Joining. *Cell Cycle* 12, 3070–3082. doi:10.4161/cc.26033
- Horigome, C., Oma, Y., Konishi, T., Schmid, R., Marcomini, I., Hauer, M. H., et al. (2014). SWR1 and INO80 Chromatin Remodelers Contribute to DNA Double-Strand Break Perinuclear Anchorage Site Choice. *Mol. Cell* 55, 626–639. doi:10.1016/j.molcel.2014.06.027
- Hu, K., Li, Y., Wu, W., Xie, L., Yan, H., Cai, Y., et al. (2020). ATM-Dependent Recruitment of BRD7 Is Required for Transcriptional Repression and DNA Repair at DNA Breaks Flanking Transcriptional Active Regions. *Adv. Sci.* 7, 2000157. doi:10.1002/adv.20200157
- Hu, Q., Botuyan, M. V., Cui, G., Zhao, D., and Mer, G. (2017). Mechanisms of Ubiquitin-Nucleosome Recognition and Regulation of 53BP1 Chromatin Recruitment by RNF168/169 and RAD18. *Mol. Cell* 66, 473–487.e9. doi:10.1016/j.molcel.2017.04.009
- Huang, T.-H., Fowler, F., Chen, C.-C., Shen, Z.-J., Sleckman, B., and Tyler, J. K. (2018). The Histone Chaperones ASF1 and CAF-1 Promote MMS22L-TONSL-

- Mediated Rad51 Loading onto ssDNA during Homologous Recombination in Human Cells. *Mol. Cell* 69, 879–892.e5. doi:10.1016/j.molcel.2018.01.031
- Huyen, Y., Zgheib, O., DiTullio Jr, R. A., Jr, Gorgoulis, V. G., Zacharatos, P., Petty, T. J., et al. (2004). Methylated Lysine 79 of Histone H3 Targets 53BP1 to DNA Double-Strand Breaks. *Nature* 432, 406–411. doi:10.1038/nature03114
- Ira, G., Pelliccioli, A., Balijja, A., Wang, X., Fiorani, S., Carotenuto, W., et al. (2004). DNA End Resection, Homologous Recombination and DNA Damage Checkpoint Activation Require CDK1. *Nature* 431, 1011–1017. doi:10.1038/nature02964
- Jakob, B., Splinter, J., Conrad, S., Voss, K.-O., Zink, D., Durante, M., et al. (2011). DNA Double-Strand Breaks in Heterochromatin Elicit Fast Repair Protein Recruitment, Histone H2AX Phosphorylation and Relocation to Euchromatin. *Nucleic Acids Res.* 39, 6489–6499. doi:10.1093/nar/gkr230
- Jeronimo, C., Watanabe, S., Kaplan, C. D., Peterson, C. L., and Robert, F. (2015). The Histone Chaperones FACT and Spt6 Restrict H2A.Z from Intragenic Locations. *Mol. Cell* 58, 1113–1123. doi:10.1016/j.molcel.2015.03.030
- Jin, C., and Felsenfeld, G. (2007). Nucleosome Stability Mediated by Histone Variants H3.3 and H2A.Z. *Genes Dev.* 21, 1519–1529. doi:10.1101/gad.1547707
- Juhász, S., Smith, R., Schauer, T., Speckhardt, D., Mamar, H., Zentout, S., et al. (2020). The Chromatin Remodeler ALC1 Underlies Resistance to PARP Inhibitor Treatment. *Sci. Adv.* 6, eabb8626. doi:10.1126/sciadv.abb8626
- Kakarougkas, A., Ismail, A., Chambers, A. L., Riballo, E., Herbert, A. D., Künzel, J., et al. (2014). Requirement for PBAF in Transcriptional Repression and Repair at DNA Breaks in Actively Transcribed Regions of Chromatin. *Mol. Cell* 55, 723–732. doi:10.1016/j.molcel.2014.06.028
- Kalb, R., Mallery, D. L., Larkin, C., Huang, J. T. J., and Hiom, K. (2014). BRCA1 Is a Histone-h2a-specific Ubiquitin Ligase. *Cel Rep.* 8, 999–1005. doi:10.1016/j.celrep.2014.07.025
- Kalocsay, M., Hiller, N. J., and Jentsch, S. (2009). Chromosome-wide Rad51 Spreading and SUMO-H2A.Z-dependent Chromosome Fixation in Response to a Persistent DNA Double-Strand Break. *Mol. Cell* 33, 335–343. doi:10.1016/j.molcel.2009.01.016
- Kari, V., Mansour, W. Y., Raul, S. K., Baumgart, S. J., Mund, A., Grade, M., et al. (2016). Loss of CHD1 Causes DNA Repair Defects and Enhances Prostate Cancer Therapeutic Responsiveness. *EMBO Rep.* 17, 1609–1623. doi:10.15252/embr.201642352
- Kent, N. A., Chambers, A. L., and Downs, J. A. (2007). Dual Chromatin Remodeling Roles for RSC during DNA Double Strand Break Induction and Repair at the Yeast MAT Locus. *J. Biol. Chem.* 282, 27693–27701. doi:10.1074/jbc.m704707200
- Kilic, S., Lezaja, A., Gatti, M., Bianco, E., Michelena, J., Imhof, R., et al. (2019). Phase Separation of 53 BP 1 Determines Liquid-like Behavior of DNA Repair Compartments. *Embo J.* 38, e101379. doi:10.15252/embj.2018101379
- Krietenstein, N., Wal, M., Watanabe, S., Park, B., Peterson, C. L., Pugh, B. F., et al. (2016). Genomic Nucleosome Organization Reconstituted with Pure Proteins. *Cell* 167, 709–721.e12. doi:10.1016/j.cell.2016.09.045
- Krogan, N. J., Keogh, M.-C., Datta, N., Sawa, C., Ryan, O. W., Ding, H., et al. (2003). A Snf2 Family ATPase Complex Required for Recruitment of the Histone H2A Variant Htz1. *Mol. Cell* 12, 1565–1576. doi:10.1016/s1097-2765(03)00497-0
- Kubik, S., Bruzzone, M. J., Challal, D., Dreos, R., Mattarocci, S., Bucher, P., et al. (2019). Opposing Chromatin Remodelers Control Transcription Initiation Frequency and Start Site Selection. *Nat. Struct. Mol. Biol.* 26, 744–754. doi:10.1038/s41594-019-0273-3
- Kubik, S., O'Duibhir, E., de Jonge, W. J., Mattarocci, S., Albert, B., Falcone, J.-L., et al. (2018). Sequence-Directed Action of RSC Remodeler and General Regulatory Factors Modulates +1 Nucleosome Position to Facilitate Transcription. *Mol. Cell* 71, 89–102.e5. doi:10.1016/j.molcel.2018.05.030
- Lademann, C. A., Renkawitz, J., Pfander, B., and Jentsch, S. (2017). The INO80 Complex Removes H2A.Z to Promote Presynaptic Filament Formation during Homologous Recombination. *Cel Rep.* 19, 1294–1303. doi:10.1016/j.celrep.2017.04.051
- Lan, L., Ui, A., Nakajima, S., Hatakeyama, K., Hoshi, M., Watanabe, R., et al. (2010). The ACF1 Complex Is Required for DNA Double-Strand Break Repair in Human Cells. *Mol. Cell* 40, 976–987. doi:10.1016/j.molcel.2010.12.003
- Larsen, D. H., Poinssignon, C., Gudjonsson, T., Dinant, C., Payne, M. R., Hari, F. J., et al. (2010). The Chromatin-Remodeling Factor CHD4 Coordinates Signaling and Repair after DNA Damage. *J. Cel Biol* 190, 731–740. doi:10.1083/jcb.200912135
- Lazzaro, F., Sapountzi, V., Granata, M., Pelliccioli, A., Vaze, M., Haber, J. E., et al. (2008). Histone Methyltransferase Dot1 and Rad9 Inhibit Single-Stranded DNA Accumulation at DSBs and Uncapped Telomeres. *Embo J.* 27, 1502–1512. doi:10.1038/emboj.2008.81
- Lee, H.-S., Park, J.-H., Kim, S.-J., Kwon, S.-J., and Kwon, J. (2010). A Cooperative Activation Loop Among SWI/SNF, γ -H2AX and H3 Acetylation for DNA Double-Strand Break Repair. *Embo J.* 29, 1434–1445. doi:10.1038/emboj.2010.27
- Leung, J. W. C., Makhharashvili, N., Agarwal, P., Chiu, L.-Y., Pourpre, R., Cammarata, M. B., et al. (2017). ZMYM3 Regulates BRCA1 Localization at Damaged Chromatin to Promote DNA Repair. *Genes Dev.* 31, 260–274. doi:10.1101/gad.292516.116
- Lewis, T. S., Sokolova, V., Jung, H., Ng, H., and Tan, D. (2021). Structural Basis of Chromatin Regulation by Histone Variant H2A.Z. *Nucleic Acids Res.* 49, 11379–11391. doi:10.1093/nar/gkab907
- Li, M., Hada, A., Sen, P., Olufemi, L., Hall, M. A., Smith, B. Y., et al. (2015). Dynamic Regulation of Transcription Factors by Nucleosome Remodeling. *eLife Sci.* 4, e06249. doi:10.7554/elife.06249
- Liang, B., Qiu, J., Ratnakumar, K., and Laurent, B. C. (2007). RSC Functions as an Early Double-Strand-Break Sensor in the Cell's Response to DNA Damage. *Curr. Biol.* 17, 1432–1437. doi:10.1016/j.cub.2007.07.035
- Luger, K., Dechassa, M. L., and Tremethick, D. J. (2012). New Insights into Nucleosome and Chromatin Structure: an Ordered State or a Disordered Affair. *Nat. Rev. Mol. Cel Biol* 13, 436–447. doi:10.1038/nrm3382
- Luijsterburg, M. S., Acs, K., Ackermann, L., Wiegant, W. W., Bekker-Jensen, S., Larsen, D. H., et al. (2012). A New Non-catalytic Role for Ubiquitin Ligase RNF8 in Unfolding Higher-Order Chromatin Structure. *EMBO J.* 31, 2511–2527. doi:10.1038/emboj.2012.104
- Luijsterburg, M. S., de Krijger, I., Wiegant, W. W., Shah, R. G., Smeenk, G., de Groot, A. J. L., et al. (2016). PARP1 Links CHD2-Mediated Chromatin Expansion and H3.3 Deposition to DNA Repair by Non-homologous End-Joining. *Mol. Cell* 61, 547–562. doi:10.1016/j.molcel.2016.01.019
- Luk, E., Ranjan, A., FitzGerald, P. C., Mizuguchi, G., Huang, Y., Wei, D., et al. (2010). Stepwise Histone Replacement by SWR1 Requires Dual Activation with Histone H2A.Z and Canonical Nucleosome. *Cell* 143, 725–736. doi:10.1016/j.cell.2010.10.019
- Ma, N. F., Hu, L., Fung, J. M., Xie, D., Zheng, B. J., Chen, L., et al. (2008). Isolation and Characterization of a Novel Oncogene, Amplified in Liver Cancer 1, within a Commonly Amplified Region at 1q21 in Hepatocellular Carcinoma. *Hepatology* 47, 503–510. doi:10.1002/hep.22072
- Marfella, C. G. A., and Imbalzano, A. N. (2007). The Chd Family of Chromatin Remodelers. *Mutat. Research/Fundamental Mol. Mech. Mutagenesis* 618, 30–40. doi:10.1016/j.mrfmmm.2006.07.012
- Marini, F., Rawal, C. C., Liberi, G., and Pelliccioli, A. (2019). Regulation of DNA Double Strand Breaks Processing: Focus on Barriers. *Front. Mol. Biosci.* 6. doi:10.3389/fmolb.2019.00055
- Markert, J., Zhou, K., and Luger, K. (2021). SMARCAD1 is an ATP-Dependent Histone Octamer Exchange Factor with De Novo Nucleosome Assembly Activity. *Sci. Adv.* 7. doi:10.1126/sciadv.abk2380
- Meisenberg, C., Pinder, S. I., Hopkins, S. R., Wooller, S. K., Benstead-Hume, G., Pearl, F. M. G., et al. (2019). Repression of Transcription at DNA Breaks Requires Cohesin throughout Interphase and Prevents Genome Instability. *Mol. Cell* 73, 212–223.e7. doi:10.1016/j.molcel.2018.11.001
- Mimitou, E. P., and Symington, L. S. (2008). Sae2, Exo1 and Sgs1 Collaborate in DNA Double-Strand Break Processing. *Nature* 455, 770–774. doi:10.1038/nature07312
- Mimitou, E. P., Yamada, S., and Keeney, S. (2017). A Global View of Meiotic Double-Strand Break End Resection. *Science* 355, 40–45. doi:10.1126/science.aak9704
- Mittal, P., and Roberts, C. W. M. (2020). The SWI/SNF Complex in Cancer - Biology, Biomarkers and Therapy. *Nat. Rev. Clin. Oncol.* 17, 435–448. doi:10.1038/s41571-020-0357-3
- Mizuguchi, G., Shen, X., Landry, J., Wu, W.-H., Sen, S., and Wu, C. (2004). ATP-driven Exchange of Histone H2AZ Variant Catalyzed by SWR1 Chromatin Remodeling Complex. *Science* 303, 343–348. doi:10.1126/science.1090701

- Morillo-Huesca, M., Clemente-Ruiz, M., Andújar, E., and Prado, F. (2010). The SWR1 Histone Replacement Complex Causes Genetic Instability and Genome-wide Transcription Misregulation in the Absence of H2A.Z. *PLoS One* 5, e12143. doi:10.1371/journal.pone.0012143
- Morrison, A. J., Highland, J., Krogan, N. J., Arbel-Eden, A., Greenblatt, J. F., Haber, J. E., et al. (2004). INO80 and γ -H2AX Interaction Links ATP-dependent Chromatin Remodeling to DNA Damage Repair. *Cell* 119, 767–775. doi:10.1016/j.cell.2004.11.037
- Myler, L. R., Gallardo, I. F., Soniat, M. M., Deshpande, R. A., Gonzalez, X. B., Kim, Y., et al. (2017). Single-Molecule Imaging Reveals How Mre11-Rad50-Nbs1 Initiates DNA Break Repair. *Mol. Cell* 67, 891–898.e4. doi:10.1016/j.molcel.2017.08.002
- Nagai, S., Dubrana, K., Tsai-Pflugfelder, M., Davidson, M. B., Roberts, T. M., Brown, G. W., et al. (2008). Functional Targeting of DNA Damage to a Nuclear Pore-Associated SUMO-dependent Ubiquitin Ligase. *Science* 322, 597–602. doi:10.1126/science.1162790
- Nakamura, K., Kato, A., Kobayashi, J., Yanagihara, H., Sakamoto, S., Oliveira, D. V. N. P., et al. (2011). Regulation of Homologous Recombination by RNF20-dependent H2B Ubiquitination. *Mol. Cell* 41, 515–528. doi:10.1016/j.molcel.2011.02.002
- Nakamura, K., Saredi, G., Becker, J. R., Foster, B. M., Nguyen, N. V., Beyer, T. E., et al. (2019). H4K20me0 Recognition by BRCA1-BARD1 Directs Homologous Recombination to Sister Chromatids. *Nat. Cell Biol* 21, 311–318. doi:10.1038/s41556-019-0282-9
- Nakamura, T. M., Du, L.-L., Redon, C., and Russell, P. (2004). Histone H2A Phosphorylation Controls Crb2 Recruitment at DNA Breaks, Maintains Checkpoint Arrest, and Influences DNA Repair in Fission Yeast. *Mol. Cell Biol* 24, 6215–6230. doi:10.1128/MCB.24.14.6215-6230.2004
- Nishibuchi, I., Suzuki, H., Kinomura, A., Sun, J., Liu, N.-A., Horikoshi, Y., et al. (2014). Reorganization of Damaged Chromatin by the Exchange of Histone Variant H2A.Z-2. *Int. J. Radiat. Oncology*Biophysics* 89, 736–744. doi:10.1016/j.ijrobp.2014.03.031
- Niu, H., Chung, W.-H., Zhu, Z., Kwon, Y., Zhao, W., Chi, P., et al. (2010). Mechanism of the ATP-dependent DNA End-Resection Machinery from *Saccharomyces cerevisiae*. *Nature* 467, 108–111. doi:10.1038/nature09318
- Oberbeckmann, E., Krietenstein, N., Niebauer, V., Wang, Y., Schall, K., Moldt, M., et al. (2021a). Genome Information Processing by the INO80 Chromatin Remodeler Positions Nucleosomes. *Nat. Commun.* 12, 3231. doi:10.1038/s41467-021-23016-z
- Oberbeckmann, E., Niebauer, V., Watanabe, S., Farnung, L., Moldt, M., Schmid, A., et al. (2021b). Ruler Elements in Chromatin Remodelers Set Nucleosome Array Spacing and Phasing. *Nat. Commun.* 12, 3232. doi:10.1038/s41467-021-23015-0
- Ocampo, J., Chereji, R. V., Eriksson, P. R., and Clark, D. J. (2016). The ISW1 and CHD1 ATP-dependent Chromatin Remodelers Compete to Set Nucleosome Spacing *In Vivo*. *Nucleic Acids Res.* 44, 4625–4635. doi:10.1093/nar/gkw068
- Ogiwara, H., Ui, A., Otsuka, A., Satoh, H., Yokomi, I., Nakajima, S., et al. (2011). Histone Acetylation by CBP and P300 at Double-Strand Break Sites Facilitates SWI/SNF Chromatin Remodeling and the Recruitment of Non-homologous End Joining Factors. *Oncogene* 30, 2135–2146. doi:10.1038/onc.2010.592
- Oza, P., Jaspersen, S. L., Miele, A., Dekker, J., and Peterson, C. L. (2009). Mechanisms that Regulate Localization of a DNA Double-Strand Break to the Nuclear Periphery. *Genes Dev.* 23, 912–927. doi:10.1101/gad.1782209
- Pan, M.-R., Hsieh, H.-J., Dai, H., Hung, W.-C., Li, K., Peng, G., et al. (2012). Chromodomain Helicase DNA-Binding Protein 4 (CHD4) Regulates Homologous Recombination DNA Repair, and its Deficiency Sensitizes Cells to poly(ADP-Ribose) Polymerase (PARP) Inhibitor Treatment. *J. Biol. Chem.* 287, 6764–6772. doi:10.1074/jbc.M111.287037
- Panier, S., and Boulton, S. J. (2014). Double-strand Break Repair: 53BP1 Comes into Focus. *Nat. Rev. Mol. Cell Biol* 15, 7–18. doi:10.1038/nrm3719
- Papamichos-Chronakis, M., Krebs, J. E., and Peterson, C. L. (2006). Interplay between Ino80 and Swr1 Chromatin Remodeling Enzymes Regulates Cell Cycle Checkpoint Adaptation in Response to DNA Damage. *Genes Dev.* 20, 2437–2449. doi:10.1101/gad.1440206
- Papamichos-Chronakis, M., Watanabe, S., Rando, O. J., and Peterson, C. L. (2011). Global Regulation of H2A.Z Localization by the INO80 Chromatin-Remodeling Enzyme Is Essential for Genome Integrity. *Cell* 144, 200–213. doi:10.1016/j.cell.2010.12.021
- Park, J.-H., Park, E.-J., Lee, H.-S., Kim, S. J., Hur, S.-K., Imbalzano, A. N., et al. (2006). Mammalian SWI/SNF Complexes Facilitate DNA Double-Strand Break Repair by Promoting γ -H2AX Induction. *Embo J.* 25, 3986–3997. doi:10.1038/sj.emboj.7601291
- Peng, G., Yim, E.-K., Dai, H., Jackson, A. P., Burgt, I. V. D., Pan, M.-R., et al. (2009). BRIT1/MCPH1 Links Chromatin Remodelling to DNA Damage Response. *Nat. Cell Biol* 11, 865–872. doi:10.1038/ncb1895
- Peritore, M., Reuswig, K.-U., Bantele, S. C. S., Straub, T., and Pfander, B. (2021). Strand-specific ChIP-Seq at DNA Breaks Distinguishes ssDNA versus dsDNA Binding and Refutes Single-Stranded Nucleosomes. *Mol. Cell* 81, 1841–1853.e4. doi:10.1016/j.molcel.2021.02.005
- Pessina, F., and Lowndes, N. F. (2014). The RSF1 Histone-Remodelling Factor Facilitates DNA Double-Strand Break Repair by Recruiting Centromeric and Fanconi Anaemia Proteins. *Plos Biol.* 12, e1001856. doi:10.1371/journal.pbio.1001856
- Pfander, B., and Diffley, J. F. X. (2011). Dpb11 Coordinates Mec1 Kinase Activation with Cell Cycle-Regulated Rad9 Recruitment. *EMBO J.* 30, 4897–4907. doi:10.1038/emboj.2011.345
- Polo, S. E., Kaidi, A., Baskcomb, L., Galanty, Y., and Jackson, S. P. (2010). Regulation of DNA-Damage Responses and Cell-Cycle Progression by the Chromatin Remodelling Factor CHD4. *Embo J.* 29, 3130–3139. doi:10.1038/emboj.2010.188
- Qi, W., Chen, H., Xiao, T., Wang, R., Li, T., Han, L., et al. (2016). Acetyltransferase P300 Collaborates with Chromodomain Helicase DNA-Binding Protein 4 (CHD4) to Facilitate DNA Double-Strand Break Repair. *Mutagenesis* 31, 193–203. doi:10.1093/mutage/gev075
- Qi, W., Wang, R., Chen, H., Wang, X., Xiao, T., Boldogh, I., et al. (2015). BRG1 Promotes DNA Double-Strand Break Repair by Facilitating the Replacement of RPA with RAD51. *J. Cell Sci.* 128, 317–330. doi:10.1242/jcs.159103
- Qiu, H., Biernat, E., Govind, C. K., Rawal, Y., Chereji, R. V., Clark, D. J., et al. (2020). Chromatin Remodeler Ino80C Acts Independently of H2A.Z to Evict Promoter Nucleosomes and Stimulate Transcription of Highly Expressed Genes in Yeast. *Nucleic Acids Res.* 48, 8408–8430. doi:10.1093/nar/gkaa571
- Ranjana, A., Wang, F., Mizuguchi, G., Wei, D., Huang, Y., and Wu, C. (2015). H2A Histone-fold and DNA Elements in Nucleosome Activate SWR1-Mediated H2A.Z Replacement in Budding Yeast. *eLife Sci.* 4, e06845. doi:10.7554/eLife.06845
- Ranjha, L., Howard, S. M., and Cejka, P. (2018). Main Steps in DNA Double-Strand Break Repair: an Introduction to Homologous Recombination and Related Processes. *Chromosoma* 127, 187–214. doi:10.1007/s00412-017-0658-1
- Reginato, G., Cannavo, E., and Cejka, P. (2017). Physiological Protein Blocks Direct the Mre11-Rad50-Xrs2 and Sae2 Nuclease Complex to Initiate DNA End Resection. *Genes Dev.* 31, 2325–2330. doi:10.1101/gad.308254.117
- Rossi, M. J., Lai, W. K. M., and Pugh, B. F. (2018). Genome-wide Determinants of Sequence-specific DNA Binding of General Regulatory Factors. *Genome Res.* 28, 497–508. doi:10.1101/gr.229518.117
- Rother, M. B., Pellegrino, S., Smith, R., Gatti, M., Meisenberg, C., Wiegant, W. W., et al. (2020). CHD7 and 53BP1 Regulate Distinct Pathways for the Re-ligation of DNA Double-Strand Breaks. *Nat. Commun.* 11, 1–19. doi:10.1038/s41467-020-19502-5
- Ryu, T., Spatola, B., Delabaere, L., Bowlin, K., Hopp, H., Kunitake, R., et al. (2015). Heterochromatic Breaks Move to the Nuclear Periphery to Continue Recombinational Repair. *Nat. Cell Biol* 17, 1401–1411. doi:10.1038/ncb3258
- Sanchez, A., Lee, D., Kim, D. I., and Miller, K. M. (2021). Making Connections: Integrative Signaling Mechanisms Coordinate DNA Break Repair in Chromatin. *Front. Genet.* 12. doi:10.3389/fgene.2021.747734
- Sánchez-Molina, S., Mortusewicz, O., Bieber, B., Auer, S., Eckey, M., Leonhardt, H., et al. (2011). Role for hACF1 in the G2/M Damage Checkpoint. *Nucleic Acids Res.* 39, 8445–8456. doi:10.1093/nar/gkr435
- Sanders, S. L., Portoso, M., Mata, J., Bähler, J., Allshire, R. C., and Kouzarides, T. (2004). Methylation of Histone H4 Lysine 20 Controls Recruitment of Crb2 to Sites of DNA Damage. *Cell* 119, 603–614. doi:10.1016/j.cell.2004.11.009
- Santisteban, M. S., Kalashnikova, T., and Smith, M. M. (2000). Histone H2A.Z Regulates Transcription and Is Partially Redundant with Nucleosome Remodeling Complexes. *Cell* 103, 411–422. doi:10.1016/S0092-8674(00)00133-1

- Sartori, A. A., Lukas, C., Coates, J., Mistrik, M., Fu, S., Bartek, J., et al. (2007). Human CtIP Promotes DNA End Resection. *Nature* 450, 509–514. doi:10.1038/nature06337
- Schrank, B. R., Aparicio, T., Li, Y., Chang, W., Chait, B. T., Gundersen, G. G., et al. (2018). Nuclear ARP2/3 Drives DNA Break Clustering for Homology-Directed Repair. *Nature* 559, 61–66. doi:10.1038/s41586-018-0237-5
- Sellou, H., Lebeaupin, T., Chapuis, C., Smith, R., Hegele, A., Singh, H. R., et al. (2016). The poly(ADP-ribose)-dependent Chromatin Remodeler Alc1 Induces Local Chromatin Relaxation upon DNA Damage. *MBoC* 27, 3791–3799. doi:10.1091/mbc.E16-05-0269
- Sheu, J. J.-C., Guan, B., Choi, J.-H., Lin, A., Lee, C.-H., Hsiao, Y.-T., et al. (2010). Rsf-1, a Chromatin Remodeling Protein, Induces DNA Damage and Promotes Genomic Instability. *J. Biol. Chem.* 285, 38260–38269. doi:10.1074/jbc.M110.138735
- Shibata, A., Moiani, D., Arvai, A. S., Perry, J., Harding, S. M., Genois, M.-M., et al. (2014). DNA Double-Strand Break Repair Pathway Choice Is Directed by Distinct MRE11 Nuclease Activities. *Mol. Cell* 53, 7–18. doi:10.1016/j.molcel.2013.11.003
- Shim, E. Y., Hong, S. J., Oum, J.-H., Yanez, Y., Zhang, Y., and Lee, S. E. (2007). RSC Mobilizes Nucleosomes to Improve Accessibility of Repair Machinery to the Damaged Chromatin. *Mol. Cell Biol* 27, 1602–1613. doi:10.1128/mcb.01956-06
- Shim, E. Y., Ma, J.-L., Oum, J.-H., Yanez, Y., and Lee, S. E. (2005). The Yeast Chromatin Remodeler RSC Complex Facilitates End Joining Repair of DNA Double-Strand Breaks. *Mol. Cell Biol* 25, 3934–3944. doi:10.1128/mcb.25.10.3934-3944.2005
- Singh, R. K., Fan, J., Gioacchini, N., Watanabe, S., Bilsel, O., and Peterson, C. L. (2019). Transient Kinetic Analysis of SWR1C-Catalyzed H2A.Z Deposition Unravels the Impact of Nucleosome Dynamics and the Asymmetry of Histone Exchange. *Cel Rep.* 27, 374–386. e4. doi:10.1016/j.celrep.2019.03.035
- Smeenk, G., and van Attikum, H. (2013). The Chromatin Response to DNA Breaks: Leaving a Mark on Genome Integrity. *Annu. Rev. Biochem.* 82, 55–80. doi:10.1146/annurev-biochem-061809-174504
- Smeenk, G., Wiegant, W. W., Marteiijn, J. A., Luijsterburg, M. S., Sroczynski, N., Costelloe, T., et al. (2012). Poly(ADP-ribosylation) Links the Chromatin Remodeler SMARCA5/SNF2H to RNF168-dependent DNA Damage Signaling. *J. Cell Sci.* 126, 889–903. doi:10.1242/jcs.109413
- Smeenk, G., Wiegant, W. W., Vrolijk, H., Solari, A. P., Pastink, A., and van Attikum, H. (2010). The NuRD Chromatin-Remodeling Complex Regulates Signaling and Repair of DNA Damage. *J. Cell Biol* 190, 741–749. doi:10.1083/jcb.201001048
- Smith, R., Sellou, H., Chapuis, C., Huet, S., and Timinszky, G. (2018). CHD3 and CHD4 Recruitment and Chromatin Remodeling Activity at DNA Breaks Is Promoted by Early poly(ADP-ribose)-dependent Chromatin Relaxation. *Nucleic Acids Res.* 46, 6087–6098. doi:10.1093/nar/gky334
- Spruijt, C. G., Luijsterburg, M. S., Menafrá, R., Lindeboom, R. G. H., Jansen, P. W. T. C., Edupuganti, R. R., et al. (2016). ZMYND8 Co-localizes with NuRD on Target Genes and Regulates Poly(ADP-ribose)-dependent Recruitment of GATAD2A/NuRD to Sites of DNA Damage. *Cel Rep.* 17, 783–798. doi:10.1016/j.celrep.2016.09.037
- Symington, L. S., and Gautier, J. (2011). Double-strand Break End Resection and Repair Pathway Choice. *Annu. Rev. Genet.* 45, 247–271. doi:10.1146/annurev-genet-110410-132435
- Symington, L. S. (2016). Mechanism and Regulation of DNA End Resection in Eukaryotes. *Crit. Rev. Biochem. Mol. Biol.* 51, 195–212. doi:10.3109/10409238.2016.1172552
- Talbert, P. B., and Henikoff, S. (2010). Histone Variants - Ancient Wrap Artists of the Epigenome. *Nat. Rev. Mol. Cell Biol* 11, 264–275. doi:10.1038/nrm2861
- Talbert, P. B., and Henikoff, S. (2021). Histone Variants at a Glance. *J. Cell Sci.* 134, jcs244749. doi:10.1242/jcs.244749
- Talbert, P. B., and Henikoff, S. (2017). Histone Variants on the Move: Substrates for Chromatin Dynamics. *Nat. Rev. Mol. Cell Biol* 18, 115–126. doi:10.1038/nrm.2016.148
- Toh, G. W.-L., O'Shaughnessy, A. M., Jimeno, S., Dobbie, I. M., Grenon, M., Maffini, S., et al. (2006). Histone H2A Phosphorylation and H3 Methylation Are Required for a Novel Rad9 DSB Repair Function Following Checkpoint Activation. *DNA Repair* 5, 693–703. doi:10.1016/j.dnarep.2006.03.005
- Toiber, D., Erdel, F., Bouazoune, K., Silberman, D. M., Zhong, L., Mulligan, P., et al. (2013). SIRT6 Recruits SNF2H to DNA Break Sites, Preventing Genomic Instability through Chromatin Remodeling. *Mol. Cell* 51, 454–468. doi:10.1016/j.molcel.2013.06.018
- Tong, J. K., Hassig, C. A., Schnitzler, G. R., Kingston, R. E., and Schreiber, S. L. (1998). Chromatin Deacetylation by an ATP-dependent Nucleosome Remodelling Complex. *Nature* 395, 917–921. doi:10.1038/27699
- Tramantano, M., Sun, L., Au, C., Labuz, D., Liu, Z., Chou, M., et al. (2016). Constitutive Turnover of Histone H2A.Z at Yeast Promoters Requires the Preinitiation Complex. *eLife Sci.* 5, e14243. doi:10.7554/eLife.14243
- Tripuraneni, V., Memisoglu, G., MacAlpine, H. K., Tran, T. Q., Zhu, W., Hartemink, A. J., et al. (2021). Local Nucleosome Dynamics and Eviction Following a Double-Strand Break Are Reversible by NHEJ-Mediated Repair in the Absence of DNA Replication. *Genome Res.* 31, 775–788. doi:10.1101/gr.271155.120
- Tsabar, M., Hicks, W. M., Tsaponina, O., and Haber, J. E. (2016). Re-establishment of Nucleosome Occupancy during Double-Strand Break Repair in Budding Yeast. *DNA Repair* 47, 21–29. doi:10.1016/j.dnarep.2016.09.005
- Tsouroula, K., Furst, A., Rogier, M., Heyer, V., Maglott-Roth, A., Ferrand, A., et al. (2016). Temporal and Spatial Uncoupling of DNA Double Strand Break Repair Pathways within Mammalian Heterochromatin. *Mol. Cell* 63, 293–305. doi:10.1016/j.molcel.2016.06.002
- Tsukuda, T., Fleming, A. B., Nickoloff, J. A., and Osley, M. A. (2005). Chromatin Remodelling at a DNA Double-Strand Break Site in *Saccharomyces cerevisiae*. *Nature* 438, 379–383. doi:10.1038/nature04148
- Tsukuda, T., Lo, Y.-C., Krishna, S., Sterk, R., Osley, M. A., and Nickoloff, J. A. (2009). INO80-dependent Chromatin Remodeling Regulates Early and Late Stages of Mitotic Homologous Recombination. *DNA Repair* 8, 360–369. doi:10.1016/j.dnarep.2008.11.014
- Ui, A., Ogiwara, H., Nakajima, S., Kanno, S., Watanabe, R., Harata, M., et al. (2014). Possible Involvement of LKB1-AMPK Signaling in Non-homologous End Joining. *Oncogene* 33, 1640–1648. doi:10.1038/onc.2013.125
- van Attikum, H., Fritsch, O., and Gasser, S. M. (2007). Distinct Roles for SWR1 and INO80 Chromatin Remodeling Complexes at Chromosomal Double-Strand Breaks. *Embo J.* 26, 4113–4125. doi:10.1038/sj.emboj.7601835
- van Attikum, H., Fritsch, O., Hohn, B., and Gasser, S. M. (2004). Recruitment of the INO80 Complex by H2A Phosphorylation Links ATP-dependent Chromatin Remodeling with DNA Double-Strand Break Repair. *Cell* 119, 777–788. doi:10.1016/j.cell.2004.11.033
- Van, H. T., and Santos, M. A. (2018). Histone Modifications and the DNA Double-Strand Break Response. *Cell Cycle* 17, 2399–2410. doi:10.1080/15384101.2018.1542899
- Venkatesh, S., and Workman, J. L. (2015). Histone Exchange, Chromatin Structure and the Regulation of Transcription. *Nat. Rev. Mol. Cell Biol* 16, 178–189. doi:10.1038/nrm3941
- Vidi, P.-A., Liu, J., Salles, D., Jayaraman, S., Dorfman, G., Gray, M., et al. (2014). NuMA Promotes Homologous Recombination Repair by Regulating the Accumulation of the ISWI ATPase SNF2h at DNA Breaks. *Nucleic Acids Res.* 42, 6365–6379. doi:10.1093/nar/gku296
- Wang, F., Ranjan, A., Wei, D., and Wu, C. (2016). Comment on "A Histone Acetylation Switch Regulates H2A.Z Deposition by the SWR-C Remodeling Enzyme". *Science* 353, 358. doi:10.1126/science.aad5921
- Wang, W., Daley, J. M., Kwon, Y., Krasner, D. S., and Sung, P. (2017). Plasticity of the Mre11-Rad50-Xrs2-Sae2 Nuclease Ensemble in the Processing of DNA-Bound Obstacles. *Genes Dev.* 31, 2331–2336. doi:10.1101/gad.307900.117
- Watanabe, R., Ui, A., Kanno, S.-I., Ogiwara, H., Nagase, T., Kohno, T., et al. (2014). SWI/SNF Factors Required for Cellular Resistance to DNA Damage Include ARID1A and ARID1B and Show Interdependent Protein Stability. *Cancer Res.* 74, 2465–2475. doi:10.1158/0008-5472.CAN-13-3608
- Watanabe, S., and Peterson, C. L. (2016). Response to Comment on "A Histone Acetylation Switch Regulates H2A.Z Deposition by the SWR-C Remodeling Enzyme". *Science* 353, 358. doi:10.1126/science.aad6398
- Watanabe, S., Radman-Livaja, M., Rando, O. J., and Peterson, C. L. (2013). A Histone Acetylation Switch Regulates H2A.Z Deposition by the SWR-C Remodeling Enzyme. *Science* 340, 195–199. doi:10.1126/science.1229758

- Weiner, A., Hughes, A., Yassour, M., Rando, O. J., and Friedman, N. (2010). High-resolution Nucleosome Mapping Reveals Transcription-dependent Promoter Packaging. *Genome Res.* 20, 90–100. doi:10.1101/gr.098509.109
- Wiechens, N., Singh, V., Gkikopoulos, T., Schofield, P., Rocha, S., and Owen-Hughes, T. (2016). The Chromatin Remodelling Enzymes SNF2H and SNF2L Position Nucleosomes Adjacent to CTCF and Other Transcription Factors. *Plos Genet.* 12, e1005940. doi:10.1371/journal.pgen.1005940
- Wiest, N. E., Houghtaling, S., Sanchez, J. C., Tomkinson, A. E., and Osley, M. A. (2017). The SWI/SNF ATP-dependent Nucleosome Remodeler Promotes Resection Initiation at a DNA Double-Strand Break in Yeast. *Nucleic Acids Res.* 45, 5887–5900. doi:10.1093/nar/gkx221
- Willhoft, O., Ghoneim, M., Lin, C.-L., Chua, E. Y. D., Wilkinson, M., Chaban, Y., et al. (2018). Structure and Dynamics of the Yeast SWR1-Nucleosome Complex. *Science* 362, eaat7716. doi:10.1126/science.aat7716
- Willhoft, O., and Wigley, D. B. (2020). INO80 and SWR1 Complexes: the Non-identical Twins of Chromatin Remodelling. *Curr. Opin. Struct. Biol.* 61, 50–58. doi:10.1016/j.sbi.2019.09.002
- Wilson, M. D., Benlekber, S., Fradet-Turcotte, A., Sherker, A., Julien, J.-P., McEwan, A., et al. (2016). The Structural Basis of Modified Nucleosome Recognition by 53BP1. *Nature* 536, 100–103. doi:10.1038/nature18951
- Wu, W.-H., Wu, C.-H., Ladurner, A., Mizuguchi, G., Wei, D., Xiao, H., et al. (2009). N Terminus of Swr1 Binds to Histone H2AZ and Provides a Platform for Subunit Assembly in the Chromatin Remodeling Complex. *J. Biol. Chem.* 284, 6200–6207. doi:10.1074/jbc.M808830200
- Wysocki, R., Javaheri, A., Allard, S., Sha, F., Côté, J., and Kron, S. J. (2005). Role of Dot1-dependent Histone H3 Methylation in G1 and S Phase DNA Damage Checkpoint Functions of Rad9. *Mol. Cell Biol.* 25, 8430–8443. doi:10.1128/MCB.25.19.8430-8443.2005
- Xiao, A., Li, H., Shechter, D., Ahn, S. H., Fabrizio, L. A., Erdjument-Bromage, H., et al. (2009). WSTF Regulates the H2A.X DNA Damage Response via a Novel Tyrosine Kinase Activity. *Nature* 457, 57–62. doi:10.1038/nature07668
- Xu, Y., Ayrapetov, M. K., Xu, C., Gursoy-Yuzugullu, O., Hu, Y., and Price, B. D. (2012). Histone H2A.Z Controls a Critical Chromatin Remodeling Step Required for DNA Double-Strand Break Repair. *Mol. Cell* 48, 723–733. doi:10.1016/j.molcel.2012.09.026
- Xue, Y., Wong, J., Moreno, G. T., Young, M. K., Côté, J., and Wang, W. (1998). NURD, a Novel Complex with Both ATP-dependent Chromatin-Remodeling and Histone Deacetylase Activities. *Mol. Cell* 2, 851–861. doi:10.1016/s1097-2765(00)80299-3
- Yamada, K., Frouws, T. D., Angst, B., Fitzgerald, D. J., DeLuca, C., Schimmele, K., et al. (2011). Structure and Mechanism of the Chromatin Remodelling Factor ISW1a. *Nature* 472, 448–453. doi:10.1038/nature09947
- Yu, Y., Deng, Y., Reed, S. H., Millar, C. B., and Waters, R. (2013). Histone Variant Htz1 Promotes Histone H3 Acetylation to Enhance Nucleotide Excision Repair in Htz1 Nucleosomes. *Nucleic Acids Res.* 41, 9006–9019. doi:10.1093/nar/gkt688
- Yuan, G.-C., Liu, Y.-J., Dion, M. F., Slack, M. D., Wu, L. F., Altschuler, S. J., et al. (2005). Genome-Scale Identification of Nucleosome Positions in *S. cerevisiae*. *Science* 309, 626–630. doi:10.1126/science.1112178
- Zhang, H., Roberts, D. N., and Cairns, B. R. (2005). Genome-Wide Dynamics of Htz1, a Histone H2A Variant that Poises Repressed/Basal Promoters for Activation through Histone Loss. *Cell* 123, 219–231. doi:10.1016/j.cell.2005.08.036
- Zhang, Y., LeRoy, G., Seelig, H.-P., Lane, W. S., and Reinberg, D. (1998). The Dermatomyositis-specific Autoantigen Mi2 Is a Component of a Complex Containing Histone Deacetylase and Nucleosome Remodeling Activities. *Cell* 95, 279–289. doi:10.1016/s0092-8674(00)81758-4
- Zhou, J., Li, J., Serafim, R. B., Ketchum, S., Ferreira, C. G., Liu, J. C., et al. (2018). Human CHD1 Is Required for Early DNA-Damage Signaling and Is Uniquely Regulated by its N Terminus. *Nucleic Acids Res.* 46, 3891–3905. doi:10.1093/nar/gky128
- Zhu, Z., Chung, W.-H., Shim, E. Y., Lee, S. E., and Ira, G. (2008). Sgs1 Helicase and Two Nucleases Dna2 and Exo1 Resect DNA Double-Strand Break Ends. *Cell* 134, 981–994. doi:10.1016/j.cell.2008.08.037

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Karl, Peritore, Galanti and Pfander. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.