



OPEN

Enhancement of astaxanthin accumulation using black light in *Coelastrum* and *Monoraphidium* isolated from Malaysia

Marshila Kaha^{1,5}, Koji Iwamoto^{1,5}✉, Nurul Ashyikin Yahya¹, Noraiza Suhaimi¹, Norio Sugiura¹, Hirofumi Hara¹, Nor'Azizi Othman², Zuriati Zakaria¹ & Kengo Suzuki^{3,4}

Microalgae are important microorganisms which produce potentially valuable compounds. Astaxanthin, a group of xanthophyll carotenoids, is one of the most powerful antioxidants mainly found in microalgae, yeasts, and crustaceans. Environmental stresses such as intense light, drought, high salinity, nutrient depletion, and high temperature can induce the accumulation of astaxanthin. Thus, this research aims to investigate the effect of black light, also known as long-wave ultraviolet radiation or UV-A, as a stressor on the accumulation of astaxanthin as well as to screen the antioxidant property in two tropical green algal strains isolated from Malaysia, *Coelastrum* sp. and *Monoraphidium* sp. SP03. *Monoraphidium* sp. SP03 showed a higher growth rate (0.66 day⁻¹) compared to that of *Coelastrum* sp. (0.22 day⁻¹). *Coelastrum* sp. showed significantly higher accumulation of astaxanthin in black light (0.999 g mL culture⁻¹) compared to that in control condition (0.185 g mL⁻¹). Similarly, *Monoraphidium* sp. SP03 showed higher astaxanthin content in black light (0.476 g mL culture⁻¹) compared to that in control condition (0.363 g mL culture⁻¹). *Coelastrum* sp. showed higher scavenging activity (30.19%) when cultured in black light condition, indicating a correlation between the antioxidant activity and accumulation of astaxanthin. In this study, black light was shown to possess great potential to enhance the production of astaxanthin in microalgae.

In recent years, there has been a surge in attention to extract bioactive compounds from natural resources due to their efficiency in the treatment of various diseases. Microalgae, which are oxygen generating photosynthetic microorganisms, are gaining great interest from biotechnological viewpoint. Microalgae can be classified as prokaryotic and eukaryotic organisms that are mostly found in the aquatic environment. These organisms possess unique characteristics of higher plants (oxygenic photosynthesis) combined with biotechnological attributes such as fast growth rates, ease of cultivation and the ability to accumulate primary and secondary metabolites^{1,2}. These useful features led to the selection of microalgae in various applications such as biofuels, food, aquaculture, cosmetics, nutrition and pharmaceuticals.

Depending on the microalgae species, various valuable secondary metabolites can be extracted from its biomass, including carotenoids, polysaccharides, fatty acids and vitamins. Among all the secondary metabolites identified from microalgae, carotenoids have received particular attention due to their importance in therapeutic value, including antioxidant, neuroprotective, anti-angiogenic and anti-inflammatory activities^{1,2}. Carotenoids comprise a huge group of natural compounds that are found predominantly in green plants, algae and some bacteria which belong to photosynthetic organisms³. It also can be found in non-photosynthetic microorganism such as fungi and yeast. Carotenoids are a class of terpenoid pigments consisting of carotenes (non-oxygenated molecule) and xanthophylls (oxygenated molecule) which can be distinguished based on their chemical structure⁴.

Among the carotenoids, astaxanthin, a group of xanthophyll carotenoids is recognized as one of the most powerful antioxidants found in nature that has been noted to surpass those of β -carotene and α -tocopherol (vitamin E)⁵. It is a red fat-soluble pigment found mainly in microalgae, yeast, and some crustaceans and gained

¹Department of Chemical and Environmental Engineering, Malaysia-Japan International Institute of Technology, Universiti Teknologi Malaysia, Jalan Sultan Yahya Petra, 54100 Kuala Lumpur, Malaysia. ²Department of Mechanical Precision Engineering, Malaysia-Japan International Institute of Technology, Universiti Teknologi Malaysia, Jalan Sultan Yahya Petra, 54100 Kuala Lumpur, Malaysia. ³Euglena Co., Ltd, Tokyo 108-0014, Japan. ⁴Microalgae Production Control Technology Laboratory, RIKEN, Yokohama, Kanagawa 230-0045, Japan. ⁵These authors contributed equally: Marshila Kaha and Koji Iwamoto. ✉email: k.iwamoto@utm.my




Figure 1. Light microscopic image with $\times 40$ magnification. (A) *Coelastrum* sp., (B) *Monoraphidium* sp. SP3. Scale bar: 20 μm .

a significant interest from pharmaceutical and nutraceutical industries due to the presence of two hydroxyl (OH) and two keto groups in the side chains of astaxanthin⁶. These applications include protection from ultraviolet (UV) light oxidation owing to astaxanthin's anti-inflammatory and anti-aging properties⁷. Antioxidants are compounds that can neutralize the reactive oxygen species (ROS) and other free radicals that generated by various metabolic processes and environmental stresses such as smoke, air pollution and UV damage.

Production of astaxanthin can be carried out by two methods, namely biological synthesis and chemical synthesis⁸. Biological synthesis includes the production of natural astaxanthin from potential organisms such as microalgae, yeast and some crustaceans as a by-product, whereas chemical synthesis includes the production of synthetic astaxanthin from petrochemicals. Synthetic astaxanthin was not approved by the US Food and Drug Administration (FDA) for human consumption due to its low bioavailability and health issues⁹. Thus, production of natural astaxanthin is gaining interest as it is shown to have greater stability and more biological functions than synthetic astaxanthin.

In microalgae, astaxanthin is accumulated in the cytosolic lipid bodies (LBs)¹⁰ and acts as a defense chemical that reacts to environmental stresses such as nutrient starvation, high light intensity, high salinity, and high temperature. Light availability is one of the most important factors which control the photosynthesis process and production of carotenoids in the photosynthetic membrane vesicles¹¹. In this study, black light was used to enhance the astaxanthin production in microalgae. Black light, also referred to as UV-A light, has a purple colour when it is turned on. UV light is classified into three wavelengths, UV-A (320–400 nm), UV-B (280–320 nm) and UV-C (200–280 nm)¹². As the world is experiencing global warming and ozone-layer depletion¹³, UV radiation issue has become worse as it affects various organisms including humans and microalgae. UV radiation is harmful as it generates reactive oxygen species such as hydrogen peroxide (H_2O_2), superoxide anion (O_2^-) and hydroxyl radical ($\cdot\text{OH}$) which possess a strong oxidation ability¹⁴. However, UV radiation is considered to be a stressor for many physiological processes such as triggering the production of secondary carotenoids to protect the cells from damage¹⁵. Therefore, in this study, black light was used to increase the accumulation of astaxanthin in tropical microalgae isolated from Malaysia.

Results

Isolation of tropical microalgae and growth analysis. Algal strains used in this study were isolated from brackish water environment at Kuala Selangor Nature Park, Malaysia (3.361057, 101.244325). The pH of the collected water was 7.2 and the temperature at the sampling site was in a range of 27–29 °C. Prescreening was done by observing the microalgae under a microscope to monitor the growth and ability of the isolated species to accumulate carotenoids, which was indicated by the colour change (yellowish-orange, data not shown). From this screening, two out of ten isolated strains were chosen for further analysis. Their morphological characteristics are shown in Fig. 1. The alga in Fig. 1A was considered *Coelastrum* sp. because it showed typical morphology, i.e. the spherical cells are arranged in pherical, about 30-celled coemobia¹⁶. The alga in Fig. 1B seemed to be *Monoraphidium* sp. from the morphology, a single cell with curved, sigmoid, or spiral, and gradually tapered towards the apex¹⁷. Strain SP03 was identified as a *Monoraphidium* representatives as it established a phyletic




Figure 2. Algal growth in control and black light condition. (A) *Coelastrum* sp., (B) *Monoraphidium* sp. SP03; white square: control condition (continuous illumination of white fluorescence lamps); black square: black light condition (1 h illumination of black light with continuous illumination of white fluorescence lamps per day). Error bars correspond to the standard deviation of triplicate cultures.

lineage associated with *Monoraphidium* strains in the 18S rRNA gene-based phylogenetic analysis (MW507790) (Fig. S1).

Figure 2 shows the growth profiles of *Coelastrum* sp. and *Monoraphidium* sp. SP03 cultured in control (without black light) and black light conditions. In this study, the isolated strains were exposed to black light at different initiation times since the time to reach stationary phase of growth varied between different strains. *Coelastrum* sp. and *Monoraphidium* sp. SP03 were exposed to black light at day 13 and 9, respectively. *Monoraphidium* sp. SP03 culture exhibited higher growth rate (0.66 day^{-1}) compared to that of *Coelastrum* sp. culture (0.22 day^{-1}) both under control and black light conditions as shown in Fig. 2B.

Astaxanthin production in both strains cultured in black light condition. Figure 3A shows the typical chromatogram and retention time for standard astaxanthin at 8.499 min ($\log k' = 1.35$, Fig. S2), whereas Fig. 3B–E show the chromatograms of astaxanthin extracted from *Coelastrum* sp. and *Monoraphidium* sp. SP03 cultured in black light and control conditions after 30 days of cultivation. Both strains showed similar retention time and $\log k'$ to that of standard astaxanthin in either condition (Fig. 3, Fig. S2) with the absorbance maxima at 480 nm (personal communication from Dr. Tharek, A¹⁸, Fig. S3) and 474 nm¹⁹, respectively. The $\log k'$ s of contaminated carotenoids in *Coelastrum* sp. and *Monoraphidium* sp. SP03 in both control and black light conditions were shown in Fig. S2. Data of chromatogram obtained after 15 days of cultivation was not shown in this study. Figure 4 shows the production of astaxanthin in *Coelastrum* sp. and *Monoraphidium* sp. SP03 after 15 and 30 days of cultivation. After 15 days, *Coelastrum* sp. cultured in black light and control conditions produced astaxanthin with the concentrations of 0.178 $\mu\text{g}/\text{mL}$ culture and 0.139 $\mu\text{g}/\text{mL}$ culture, respectively (Fig. 4A). Interestingly, when cultured in black light condition, the astaxanthin production in *Coelastrum* sp. increased remarkably (5.4-fold per volume culture) compared to that in control condition. The concentrations of astaxanthin in black light and control conditions were 0.999 $\mu\text{g}/\text{mL}$ culture and 0.185 $\mu\text{g}/\text{mL}$ culture, respectively. This difference was significant as verified statistically by using *t*-test ($p < 0.05$). In *Monoraphidium* sp. SP03, no production of astaxanthin was observed in either control or black light condition after 15 days of cultivation (Fig. 4B). After 30 days of culture, the production of astaxanthin was significantly increased (0.23-fold per volume culture) by black light compared to those cultivated in control condition. The concentrations of astaxanthin in black light and control conditions were 0.476 $\mu\text{g}/\text{mL}$ culture and 0.363 $\mu\text{g}/\text{mL}$ culture, respectively.

Antioxidant profile of astaxanthin extracted from *Coelastrum* and *Monoraphidium*. Scavenging activity is the ability of an antioxidant to inhibit oxidation. Thus, the higher the scavenging activity, higher the ability of the antioxidant to scavenge the free radicals. Both *Coelastrum* sp. and *Monoraphidium* sp. SP03 cultured in black light condition showed higher radical scavenging activity compared to those cultured in control condition (Fig. 5). In this study, correlation analysis was conducted to test the relationship between the scavenging activity and the amount of astaxanthin produced. The results indicated that both *Coelastrum* sp. (Fig. 5A) and *Monoraphidium* sp. SP03 (Fig. 5B) showed a correlation between the production of astaxanthin




Figure 3. HPLC chromatogram of astaxanthin extracted after 30 days of cultivation. (A) astaxanthin standard, (B) *Coelastrum* sp. in control condition, (C) *Coelastrum* sp. in black light condition, (D) *Monoraphidium* sp. SP03 in control condition (E) *Monoraphidium* sp. SP03 in black light condition. Control condition: continuous illumination of white fluorescense lamps); Black light condition: 1 h illumination of black light with continuous illumination of white fluorescense lamps per day.




Figure 4. Astaxanthin production after 15 and 30 days of cultivation under black light and control conditions. (A) *Coelastrum* sp., (B) *Monoraphidium* sp. SP03. ND, not detected; white bar: black light condition; grey bar: control condition. Error bars correspond to the standard deviation of triplicate cultures. The asterisk shows that the p-value is less than 0.05.

and scavenging activity with r-values 0.75 and 0.99, respectively. The scavenging activity of astaxanthin extracted from *Coelastrum* cells was approximately 30.19% per volume culture, which was higher compared to that of *Monoraphidium* sp. SP03. This was consistent with the higher astaxanthin production by *Coelastrum* sp.




Figure 5. Astaxanthin production and scavenging activity in (A) *Coelastrum* sp. and (B) *Monoraphidium* sp. SP03 after 30 days of culture. Grey bar: astaxanthin content ($\mu\text{g mL culture}^{-1}$); dashed line: scavenging activity. The asterisk shows that the p-value is less than 0.05.

Discussion

In the present study, *Coelastrum* sp. that belong to the family of *Scenedesmeceae* and *Monoraphidium* sp. SP03 that comes from *Selenastraceae* family were successfully isolated from brackish water in Malaysia. The isolation of tropical strain would be the promising approach for the adaptation to local changing environment and can provide more productive culture. There were only few reports on the production of astaxanthin by these strains. A report by Liu et al.²⁰ showed that *Coelastrum* sp. HA-1 produced 1.6-fold of astaxanthin as compared to control condition when induced with linoleic. Similarly, *Coelastrum* sp. TISTR 9501RE showed increase in carotenoid biosynthesis in nutrient-depleted condition²¹. The carotenoid production by *Coelastrum* cf. *pseudomicroporum* was also reported using urban wastewater²². More recently, astaxanthin production by *Coelastrum* sp. was shown to be enhanced by methyl viologen as the reactive oxygen species enhancer¹⁸. In case of *Monoraphidium* strain, previous study improved the photoautotrophic culture condition and outdoor cultivation method for increasing the astaxanthin content in *Monoraphidium* GK12^{23,24}. In the present study, black light was introduced to increase the accumulation of astaxanthin in *Coelastrum* sp. and *Monoraphidium* sp. SP03. These strains were subjected to a stress condition induced by the black light at their initial stationary phase of growth. Carotenogenesis has been reported to be enhanced when microalgae are subjected to unfavourable or stress conditions such as nutrient depletion, intense irradiation, high salinity or high temperature which usually occur when the algal cells reach the stationary phase²⁵.

Astaxanthin was extracted from *Coelastrum* sp. and *Monoraphidium* sp. SP03 at day 15 and 30 and compared. Based on the results shown in Fig. 4, both strains showed higher astaxanthin production on day 30 compared to that on day 15. Both strains showed higher astaxanthin content in black light condition compared to those cultured in control condition. However, only *Coelastrum* sp. showed a significant difference in the amount of astaxanthin produced between the control and black light conditions, as verified by using *t*-test analysis ($p < 0.05$). On the other hand, *Monoraphidium* sp. SP03 cultured in control and black light conditions showed no statistically significant difference ($p > 0.05$).

Light intensity is one of the factors that influence the carotenogenesis in microalgae²⁶. Photosynthesis can proceed at both low and high light intensities. Under low light intensity, fewer oxygen radicals are generated, whereas under high light intensity, surplus energy activates more oxygen molecules, since the cells are unable to utilize all energy that is generated by photosynthesis. The excess energy leads to the formation of astaxanthin²⁷. Thus, the early culture stage of the strains resulted in low accumulation and no detection of astaxanthin due to exposure of the cultures to low light intensity (Fig. 3).

The black light, referred to as UV-A radiation, has a purple color when it is turned on. The term UV radiation is used to describe the UV wavelength region ranging from 200 to 400 nm and is classified into three categories, UVC (200–280 nm), UVB (280–320 nm) and UVA (320–400 nm)¹². UV radiation exerts both positive and negative effects on microalgae cells²⁸. For instance, UV radiation was known to cause DNA damage in the cells. Photodamage and conformational change caused by absorption of UV radiation by nucleic acids and proteins directly impair the vital metabolic functions such as DNA replication, translation, and transcription²⁹. UV-A irradiation has been shown to inhibit photosynthesis³⁰. Thus, it needs to be counteracted or avoided. Despite the inactivation of photosynthesis, the irradiation drives the accumulation of secondary metabolites³¹ and induced oxidative stress that resulted in the accumulation of carotenoids³².

At present, the effect of black light radiation on the accumulation of astaxanthin has not been reported. There were only a few reports on the effect of UV-A radiation on the production of carotenoids in microalgae. Salguero et al. studied the effect of adding UV-A radiation with different intensities to photosynthetically active radiation in marine microalga *Dunaliella bardawil*³³. They observed that exposure to UV-A enhanced the production of total carotenoids, mainly β -carotene, lutein, and zeaxanthin, as well as the cell growth and the photosynthetic efficiency of *D. bardawil*. Higher β -carotene accumulation was observed in *D. bardawil* when UV-A radiation was coupled with nitrogen starvation³⁴. The findings of the present study are consistent with the results of the previous studies which showed that UV-A could be one of the stressors that can enhance the accumulation of carotenoids in the algal cells.

Carotenoids show significant absorption in the UV-A spectrum, and this can be the logic behind the correlation between the higher production of astaxanthin (this study) as well as the higher production of β -carotene (previous studies) in microalgae and UV-A radiation (Figs. 3, 4). Astaxanthin has been known as an excellent antioxidant in numerous applications, for example, as an additive in feed industries, cosmetics as well as pharmaceuticals³⁵. It has been reported that antioxidants play a role in preventing certain tumors as well as ameliorating degenerative and cardiovascular diseases³⁶.

In this study, DPPH assay was performed as a fast screening method to analyze the antioxidant profile of astaxanthin extracted from *Coelastrum* sp. and *Monoraphidium* sp. SP03 after 30 days of culture. DPPH assay is a simple and easy method to perform antioxidant assays. In addition, DPPH is a stable radical compound that does not disintegrate in methanol, ethanol and water. Thus, in this study, ethanol was used as a solvent to dilute DPPH. DPPH assay has been successfully used to analyze the antioxidant properties of edible seed oils, wheat grains, and herbs³⁷.

In this study, both *Coelastrum* sp. and *Monoraphidium* sp. SP03 showed a correlation between the production of astaxanthin and scavenging activity. *Coelastrum* sp. showed 30.19% scavenging activity that correlated with the higher amount of extracted astaxanthin. Generally, carotenoids in microalgae act as excellent antioxidants by antagonizing oxidative stress and nitrogen stress through neutralizing the excess of free radicals and ROS. In addition, strong antioxidants are commonly found in the carotenoids, especially from microalgae, due to their unique molecular structure, which consists of an additional oxygenic functional group (carbonyl, hydroxyl and epoxy) and several conjugated double bonds. Apart from astaxanthin, other carotenoids such as β -carotene, lutein, α -tocopherol, canthaxanthin, and fucoxanthin were also reported in microalgae due to their ability to scavenge free radicals. Nonetheless, the antioxidant properties of these carotenoids are not as strong as that of astaxanthin.

Astaxanthin is chemically similar to β -carotene but more powerful in terms of singlet oxygen quenching mechanism (antioxidant activity) compared to other members of carotenoids such as β -carotene, lutein, and lycopene³⁸. The antioxidant activity of astaxanthin was tenfold higher than that of β -carotene and 500-fold higher than that of α -tocopherol³⁹. A number of studies have validated the superior antioxidant activity of astaxanthin. In particular, the potent antioxidant properties of astaxanthin, zeaxanthin, lutein, β -carotene, and lycopene were studied in specific physio-chemical interactions with membranes⁴⁰. Their result showed that astaxanthin could suppress lipid peroxidation and maintain the membrane integrity, whereas lutein and β -carotene could not prevent the distortion of the membrane structure and production of high levels of lipid hydroperoxides. To summarize, astaxanthin plays an important role in scavenging the ROS produced by exposure to black light. In addition, *Coelastrum* sp. showed astaxanthin accumulation following exposure to black light. To clarify the effect of black light on the production of astaxanthin in algae, further study would be required such as the comparative analysis of astaxanthin production among *Coelastrum* sp., *Haematococcus*, and other algal strains with black light.

Materials and methods

Isolation and culture conditions. Water samples were collected from a freshwater environment at Kuala Selangor Nature Park, Selangor Malaysia (3.361057, 101.244325) by using 25 μ m plankton net from a known distance. The pH of water samples was measured by using a compact pH meter (B-71X, Horiba Scientific, Japan), while the temperature of water samples and surroundings were measured by Thermo recorder TR-71wf (T&D, Japan). Microalgae cells from the water samples were isolated using traditional isolation technique by picking up the single cells using a Pasteur pipette⁴¹. This technique included the use of a Pasteur pipette attached to a latex tube and the isolation process was performed under a microscope (Eclipse TS100, Nikon, Japan). A drop of sterilized AF-6 medium⁴² was placed on a glass slide and into the microplate. A single cell of microalgae from the collected water samples was transferred to the AF-6 medium on the glass slide. The isolated cells were washed several times (2–3 times) on the washing plate before being transferred to the 96-well microplates containing AF-6 medium. The microplates were incubated at (28 ± 1) °C. Throughout this process, sterilized equipments and media were used to avoid contamination and to get unialgal cultures.

Isolated culture (5 mL) was transferred to 250 mL Erlenmeyer flask containing 100 mL of medium (5% v/v inoculation). The Erlenmeyer flask was placed on an orbital shaker (Orbitron, InforsHT, Switzerland) with the speed of 120 rpm at (28 ± 1) °C under continuous illumination by white fluorescence lamps (18 W/865, TL-D, Philips, Nederland) at 35 ± 5 μ mol photon $m^{-2} s^{-1}$. After the cultures reached stationary phase, the light intensity was increased to 165 ± 5 μ mol photon $m^{-2} s^{-1}$ with continuous illumination. All inoculations were prepared in triplicate.

Exposure to black light. To investigate the effect of black light on the production of astaxanthin, triplicate culture of each strain was prepared with 5% v/v inoculation. These triplicate cultures were maintained in identical conditions until they reached the early stationary phase. The cultures were subjected to continuous illumina-

nation ($165 \pm 5 \mu\text{mol photon m}^{-2} \text{ s}^{-1}$) and black light (365 nm, 18 W, TL-D, Philips, Nederland). The triplicate cultures were exposed to black light for 1 h every day.

Growth measurement. The optical density at 750 nm was measured every day by using a colorimeter (Spectronic 20A, Shimadzu, Japan). The specific growth rate was calculated using the Eq. (1)⁴³

$$\text{Specific growth rate: } \mu = (\ln\text{OD}_{t_2} - \ln\text{OD}_{t_1}) / (t_2 - t_1) \quad (1)$$

where OD_{t_2} = Highest absorbance (OD_{750}) at exponential phase, OD_{t_1} = Lowest absorbance (OD_{750}) at exponential phase, t_2 = time (day) at OD_{t_2} , t_1 = time (day) at OD_{t_1} .

Phylogenetic analysis. Plant DNA Preparation Kit (Jena Bioscience, Germany) was employed to extract the genomic DNA with slight modification of the manufacturer's protocol. Two eukaryotic universal primers to amplify the specific region of nuclear-encoded small subunit ribosomal RNA gene (18S rDNA), SR1 (forward, 5'-TACCTGGTTGATCCTGCCAG-3') and SR12 (reverse, 5'-CCTTCCGCAGGTTACCTAC-3') were used⁴⁴. PCR mastermixes were prepared by using Promega Kit (Promega, USA). The PCR reaction conditions were as described; 30 cycles of initial denaturation at 94 °C for 2 min, denaturation at 98 °C for 10 s, annealing at 53 °C for 30 s, and extension at 72 °C for 2 min. Amplified DNA was sequenced by First Base Laboratory Sdn. Bhd. (Kuala Lumpur, Malaysia), and the sequences were analysed by BLAST algorithm available on National Center for Biotechnology Information (NCBI) website. Genbank database sequences having 90–99% similarity with the analyzed sequence were aligned by using MEGA X software. The phylogenetic tree was constructed using Maximum likelihood method with bootstrap test (1000 replicates).

Extraction of astaxanthin. Astaxanthin from the isolated microalgae was extracted by the acid-acetone method as described previously⁴⁵. The samples (10 mL) were centrifuged, and the harvested biomass was lyophilized and used for the extraction process. The biomass was treated with 1 mL of 1 N HCl (Sigma-Aldrich, USA) at 70 °C for 45 min. The sample was cooled at room temperature (25 °C) and centrifuged (10,000g) for 10 min. Then, the treated sample was washed twice with deionized water (1 mL) and suspended in 1 mL of acetone (Sigma-Aldrich) prior to incubation for 1 h in an orbital shaker (200 rpm). The sample was centrifuged (10,000g) for 5 min. The supernatant was filtered through 0.45 μm membrane filters (Whatman, Germany) and was injected into vials for high-performance liquid chromatography (HPLC) analysis.

Quantification of astaxanthin content. The astaxanthin content was analyzed by HPLC system (1220 Infinity LC, Agilent Technologies, USA) with a diode-array detector (DAD, threshold 0.001 mAU). The chromatographic separation was achieved on Eclipse Plus C18 column (5 μm , 4.6 \times 250 mm, Agilent Technologies). The extracts (10 μL) were injected onto the column with 0.8 mL/min flow rate. The optical density at 475 nm detected by DAD with 4 nm of slit width and 4 nm band width was used for quantifying astaxanthin. The mobile phase consisted of solvent A (acetone HPLC grade) (Sigma-Aldrich) and solvent B (methanol: H_2O , 9:1) (Sigma-Aldrich) with the following 30 min gradient: 0 min (80% solvent B, 20% solvent A), 10 min (50% solvent A, 50% solvent B), and 20 min (20% solvent B, 80% solvent A). Standard astaxanthin (Sigma-Aldrich) was used to compare the concentrations of astaxanthin in isolated algal samples. The peaks were identified from the retention time and $\log k'$ ⁴⁶.

Antioxidant activity assay. The DPPH assay was performed as described previously⁴⁷. The antioxidant activity was assessed colorimetrically at 517 nm. The reagent 1, 1'-diphenyl-2-picrylhydrazyl (DPPH) (Sigma-Aldrich) (2.4 mg) was mixed with ethanol (100 mL) and stored at 20 °C in dark until further use. Then, sample (0.2 mL) was added to ethanol (2 mL), incubated for 20 min, and assessed as a diluted sample. For the blank, distilled water (0.2 mL) was added to ethanol (2 mL) and incubated for 20 min. Both dilutions were stored at -20 °C in dark until further use. Then, the diluted sample (1.5 mL) was mixed with DPPH solution (0.06 mM, 1.5 mL). This reaction was incubated for 30 min in dark at room temperature. This step was repeated for the blank solution. Finally, the solution was analyzed spectrophotometrically at 517 nm. The percentage of radical scavenging activity was calculated using the Eq. (2).

$$\text{Radical scavenging activity (\%)} = \frac{\text{Abs of blank} - \text{Abs of sample}}{\text{Abs of blank}} \times 100. \quad (2)$$

Statistical analysis. All the assays used in this research were conducted in triplicates and the results were expressed as mean values. A two-tailed paired t-test was performed to check whether the utilization of black light could affect the production of astaxanthin by the microalgae. The *p*-value of less than 0.05 was used to indicate statistically significant differences⁴⁸. In addition, correlation analysis was also conducted to check the linear relationship between the antioxidant profile and astaxanthin production in isolated species. Pearson's correlation coefficient (r-value) was used to measure the correlation between the two variables, where the correlation coefficient value of 0.7 to 1.0 indicated a strong positive linear relationship. Microsoft Excel 2010 equipped with the Analysis Toolpak was used as a statistical tool.

Data availability

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Received: 20 April 2020; Accepted: 19 May 2021

Published online: 03 June 2021

References

1. Gateau, H., Solymosi, K., Marchand, J. & Schoefs, B. Carotenoids of microalgae used in food industry and medicine. *Mini Rev. Med. Chem.* **17**, 1140–1172 (2017).
2. Mimouni, V. *et al.* The potential of microalgae for the production of bioactive molecules of pharmaceutical interest. *Curr. Pharm. Biotechnol.* **13**, 2733–2750 (2012).
3. Bijttebier, S. *et al.* Ultra high performance liquid chromatography versus high performance liquid chromatography: Stationary phase selectivity for generic carotenoid screening. *J. Chromatogr. A* **1332**, 46–56 (2014).
4. Ahmed, F. *et al.* Profiling of carotenoids and antioxidant capacity of microalgae from subtropical coastal and brackish waters. *Food Chem.* **165**, 300–306 (2014).
5. Higuera-Ciajara, I., Félix-Valenzuela, L. & Goycoolea, F. M. Astaxanthin: A review of its chemistry and applications. *Crit. Rev. Food Sci. Nutr.* **46**, 185–196 (2006).
6. Britton, G. Chapter 10: Functions of intact carotenoids. In *Carotenoids. Volume 4: Natural Functions* (eds Britton, G. *et al.*) 189–211 (Birkhauser, Basel, 2004).
7. Yang, Y., Kim, B. & Lee, J. Astaxanthin structure, metabolism, and health benefits. *J. Hum. Nutr. Food Sci.* **1**, 1003. <https://doi.org/10.3390/md12010128> (2013).
8. Haque, F., Dutta, A., Thimmanagari, M. & Chiang, Y. W. Intensified green production of astaxanthin from *Haematococcus pluvialis*. *Food Bioprod. Process.* **99**, 1–11 (2016).
9. Solymosi, K., Latruffe, N., Morant-Manceau, A. & Schoefs, B. Food colour additives of nature in *Colour Additives for Foods and Beverages* (eds Scotter, M.J.) 1–34, (Woodhead Publishing, 2015).
10. Pick, U., Zarka, A., Boussiba, S. & Davidi, L. A hypothesis about the origin of carotenoid lipid droplets in the green algae *Dunaliella* and *Haematococcus*. *Planta* **249**, 31–47 (2019).
11. Lemoine, Y. & Schoefs, B. Secondary ketocarotenoid astaxanthin biosynthesis in algae: a multifunctional response to stress. *Photosynth. Res.* **106**, 155–177 (2010).
12. Mavarakis, E., Miyamura, Y., Bowen, M. P. & Correa, G. Light, including ultraviolet. *J. Autoimmun.* **34**, 247–257 (2010).
13. Rozanov, E. Preface: Ozone evolution in the past and future. *Atmosphere* **11**, 709 (2020).
14. de Jager, T. L., Cockrell, A. E. & Du Plessis, S. S. Ultraviolet light induced generation of reactive oxygen species. *Adv. Exp. Med. Biol.* **996**, 15–23 (2017).
15. Thoma, F., Somborn-Schulz, A., Schlehner, D., Keuter, V. & Deerberg, G. Effects of light on secondary metabolites in selected leafy greens: A review. *Front. Plant Sci.* **11**, 497 (2020).
16. Comas, A. & Krienitz, L. Comparative LM- and SEM-studies on *Coelastrum* (Chlorophyta, Chlorococcales) under culture conditions. *Algol. Stud.* **87**, 87–98 (1997).
17. Ramos, G. J. P., Bicudo, C. E. D., Neto, A. G. & Moura, C. W. D. *Monoraphidium* and *Ankistrodesmus* (Chlorophyceae, Chlorophyta) from Pantanal dos Marimbus, Chapada Diamantina, Bahia State, Brazil. *Hoehnea* **39**, 421–434 (2012).
18. Tharek, A. *et al.* Enhanced astaxanthin production by oxidative stress using methyl viologen as a reactive oxygen species (ROS) reagent in green microalgae *Coelastrum* sp. *Indones. J. Biotechnol.* **25**, 95–101 (2020).
19. Mulders, K. J. M. *et al.* Nitrogen-depleted *Chlorella zofingiensis* produces astaxanthin, ketolutein and their fatty acid esters: a carotenoid metabolism study. *J. Appl. Phycol.* **27**, 125–140 (2014).
20. Liu, Z. *et al.* Enhancement of linoleic acid content stimulates astaxanthin esterification in *Coelastrum* sp. *Bioresour. Technol.* **300**, 122649 (2020).
21. Rauytanapanit, M. *et al.* Nutrient deprivation-associated changes in green microalga *Coelastrum* sp. TISTR9501RE enhanced potent antioxidant carotenoids. *Mar. Drugs.* **17**, 328 (2019).
22. Úbeda, B., Gálvez, J. Á., Michel, M. & Bartual, A. Microalgae cultivation in urban wastewater: *Coelastrum* cf. pseudomicroporum as a novel carotenoid source and a potential microalgae harvesting tool. *Bioresour. Technol.* **228**, 210–217 (2017).
23. Fujii, K., Nakajima, H. & Anno, Y. Potential of *Monoraphidium* sp. GK12. *J. Chem. Technol. Biotechnol.* **83**, 1578–1584 (2008).
24. Fujii, K. *et al.* Potential use of the astaxanthin-producing microalga, *Monoraphidium* sp. GK12, as a functional aquafeed for prawns. *J. Appl. Phycol.* **22**, 363–369 (2010).
25. Minhas, A. K., Hodgson, P., Barrow, C. J. & Adhholeya, A. A review on the assessment of stress conditions for simultaneous production of microalgal lipids and carotenoids. *Front. Microbiol.* **7**, 546; <https://doi.org/10.3389/fmicb.2016.00546> (2016).
26. Přibyl, P., Pilný, J., Cepák, V. & Kaštánek, P. The role of light and nitrogen in growth and carotenoid accumulation in *Scenedesmus* sp. *Algal Res.* **16**, 69–75 (2016).
27. Wu, K. *et al.* High irradiance compensated with CO₂ enhances the efficiency of *Haematococcus lacustris* growth. *Biotechnol. Rep.* **26**, e00444. <https://doi.org/10.1016/j.btre.2020.e00444> (2020).
28. Wulff, A., Roleda, M., Zacher, K. & Wiencke, C. UV radiation effects on pigments, photosynthetic efficiency and DNA of an Antarctic marine benthic diatom community. *Aquat. Biol.* **3**, 167–177 (2008).
29. Buma, A. G. J., Boelen, P. & Jeffrey, W. A. UVR-induced DNA damage in aquatic organism in *Comprehensive Series in Photochemistry and Photobiology 1, UV Effects in Aquatic Organism and Ecosystem* (eds Helbling, E. W. & Zagarese, H.) 291–327, (Royal Society of Chemistry, 2003).
30. Larkum, A. W. D. & Wood, W. F. The effect of UV-B radiation on photosynthesis and respiration of phytoplankton, benthic macroalgae and seagrasses. *Photosyn. Res.* **36**, 17–23 (1993).
31. Holzinger, A. & Lutz, C. Algae and UV-irradiation: Effects on ultrastructure and related metabolic functions. *Micron* **37**, 190–207 (2006).
32. Jahnke, L. S. Massive carotenoid accumulation in *Dunaliella bardawil* induced by ultraviolet: A radiation. *J. Photochem. Photobiol. B Biol.* **48**, 68–74 (1999).
33. Salguero, A. *et al.* UV-A mediated induction of carotenoid accumulation in *Dunaliella bardawil* with retention of cell viability. *Appl. Microbiol. Biotech.* **66**, 506–511 (2005).
34. Mogedas, B., Salguero, A., Casal, C. & Vilchez, C. UV-A promotes long-term carotenoid production of *Dunaliella* in photobioreactors with retention of cell viability in *Communicating Current Research and Educational Topics and Trends in Applied Microbiology* (ed. Méndez-Vilas, A.) 348–355 (Formatex, 2007).
35. Kim, H. C. *et al.* Treating high-strength saline piggy wastewater using the heterotrophic cultivation of *Acutodesmus obliquus*. *Biochem. Eng. J.* **110**, 51–58 (2016).
36. Abe, K., Hattori, H. & Hirano, M. Accumulation and antioxidant activity of secondary carotenoids in the aerial microalga *Coelastrum striolata* var. *multistriata*. *Food Chem.* **100**, 656–661 (2007).
37. Cheng, Z., Moore, J. & Yu, L. High-throughput relative DPPH radical scavenging capacity assay. *J. Agric. Food Chem.* **54**, 7429–7436 (2006).
38. Edge, R. & Truscott, T. G. Singlet oxygen and free radical reactions of retinoids and carotenoids—a review. *Antioxidants* **7**, 5. <https://doi.org/10.3390/antiox7010005> (2018).

39. Ceron, M. C. *et al.* Antioxidant activity of *Haematococcus pluvialis* cells grown in continuous culture as a function of their carotenoid and fatty acid content. *Appl. Microbiol. Biotechnol.* **74**, 1112–1119 (2007).
40. McNulty, H. P., Byun, J., Lockwood, S. F., Jacob, R. F. & Mason, R. P. Differential effects of carotenoids on lipid peroxidation due to membrane: X-ray diffraction analysis. *Biochem. Biophys. Acta.* **1768**, 167–174 (2007).
41. Throndsen, J. Special method-micromanipulators in *Handbook of Phycological Methods: Culture Methods and Growth Measurement* (ed. Stein J. R.) 139–144 (Cambridge University Press, 1973).
42. Kawachi, M. *et al.* in *MCC-NIES List of Strains* 9th edn Microalgae, endangered macroalgae and protists (Microbial Culture Collection at the National Institute for Environmental Studies, Tsukuba, Japan, 2013).
43. Li, T., Zheng, Y., Yu, L. & Chen, S. Mixotrophic cultivation of a *Chlorella sorokiniana* strain for enhanced biomass and lipid production. *Biomass Bioenergy* **66**, 204–213 (2014).
44. Sekimoto, S. *et al.* A multigene phylogeny of *Olpidium* and its implications for early fungal evolution. *BMC Evol. Biol.* **11**, 331. <https://doi.org/10.1186/1471-2148-11-331> (2011).
45. Sarada, R., Vidhyavathi, R., Usha, D. & Ravishankar, G. A. An efficient method for extraction of astaxanthin from green alga *Haematococcus pluvialis*. *J. Agric. Food Chem.* **54**, 7585–7588 (2006).
46. Darko, E., Schoefs, B. & Lemoine, Y. Improved liquid chromatographic method for the analysis of photosynthesis pigments of higher plants. *J. Chromatogr. A.* **876**, 111–116 (2000).
47. Marinova, G. & Batchvarov, V. Evaluation of the methods for determination of the free radical scavenging activity by DPPH. *Bulg. J. Agric. Sci.* **17**, 11–24 (2011).
48. Damiani, M. C., Popovich, C. A., Constenla, D. & Leonardi, P. I. Lipid analysis in *Haematococcus pluvialis* to assess its potential use as a biodiesel feedstock. *Bioresour. Technol.* **101**, 3801–3807 (2010).

Acknowledgements

We are grateful to the Algal Biomass and Metabolic & Molecular Biological Engineering Laboratory of Malaysia-Japan International Institute of Technology for providing the materials and equipments throughout this research. We also thank Editage (www.editage.com) for English language editing. This study was supported by the International Grant of University of Tsukuba (R.K130000.7343.4B335).

Author contributions

K.I., H.H. and Z.Z. conceived and designed the experiments; M.K., N.A.Y. and A.S. performed the experiment; K.S. and K.I. analyzed the data; N.O., Z.Z. and N.S. contributed reagents/materials/analytical equipment; M.K., K.I. and K.S. prepared and revised the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-91128-z>.

Correspondence and requests for materials should be addressed to K.I.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021