

Fibrations and stable generalized complex structures

Gil R. Cavalcanti and Ralph L. Klaasse

ABSTRACT

A generalized complex structure is called stable if its defining anticanonical section vanishes transversally, on a codimension-two submanifold. Alternatively, it is a zero elliptic residue symplectic structure in the elliptic tangent bundle associated to this submanifold. We develop Gompf–Thurston symplectic techniques adapted to Lie algebroids, and use these to construct stable generalized complex structures out of log-symplectic structures. In particular we introduce the notion of a boundary Lefschetz fibration for this purpose and describe how they can be obtained from genus one Lefschetz fibrations over the disc.

Contents

1. Introduction	1242
2. Stable generalized complex and log-symplectic structures	1245
3. Lie algebroids and Lie algebroid symplectic structures	1251
4. Constructing Lie algebroid symplectic structures	1259
5. Boundary maps and boundary Lefschetz fibrations	1263
6. Constructing boundary Lefschetz fibrations	1271
7. Constructing stable generalized complex structures	1273
8. Examples and applications	1276
References	1279

1. Introduction

Generalized complex structures [16, 17, 20] capture aspects of both symplectic and complex geometry. While slightly misleading, a generalized complex structure can be seen as a Poisson structure together with a suitably compatible complex structure normal to the (singular) symplectic leaves. In general the symplectic leaves may have varying dimension, leading to the notion of type change, where complex and symplectic behavior is mixed.

In this paper we are interested in a class of generalized complex manifolds for which the type need not be constantly equal to zero, but differs from zero only in the mildest way possible. These are called stable generalized complex structures [9, 14], and are those generalized complex structures whose defining anticanonical section vanishes transversally. Whenever a generalized complex structure is of type 0, it is isomorphic after a B -field transformation to a symplectic structure. In this sense a stable generalized complex structure is a symplectic-like structure that fails to be symplectic on at most a codimension-two submanifold.

In [7, 8, 15, 32, 33], many examples of stable generalized complex structures on four-manifolds were constructed, in particular on manifolds without a symplectic nor complex structure. A thorough study of stable generalized complex manifolds was initiated in [9].

Received 1 September 2017; revised 7 June 2018; published online 28 September 2018.

2010 *Mathematics Subject Classification* 53D18, 53D17, 53D05 (primary).

This research was supported by VIDI grant number 639.032.221 from NWO, the Netherlands Organisation for Scientific Research.

© 2018 The Author(s). The Proceedings of the London Mathematical Society is copyright © London Mathematical Society. This is an open access article under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Stable generalized complex structures can be seen as the generalized geometric analogue of a log-symplectic structure, which is a type of mildly degenerate Poisson structure that has recently received a lot of attention [6, 10, 18, 19, 25, 26].

In [9] it was noted that stable generalized complex structures can alternatively be viewed as symplectic forms on a Lie algebroid, the elliptic tangent bundle, constructed out of the anticanonical section and its zero set (they are further of zero elliptic residue: see Section 2 for details). This symplectic viewpoint allows one to apply symplectic techniques to the study of stable generalized complex structures. In this paper we focus on constructing stable generalized complex structures on total spaces of fibration-like maps.

More precisely, in this paper we extend Gompf–Thurston symplectic techniques to arbitrary Lie algebroids. We introduce Lie algebroid fibrations and Lie algebroid Lefschetz fibrations, and give criteria when these can be equipped with compatible Lie algebroid symplectic structures. We discuss morphisms between elliptic and log-tangent bundles, which are Lie algebroids whose symplectic structures describe stable generalized complex structures and log-symplectic structures respectively. We then introduce a class of maps called boundary maps. These induce morphisms between the elliptic and log tangent bundles, and we use them to construct stable generalized complex structures out of log-symplectic structures.

The following is the main result about Lie algebroid Lefschetz fibrations and Lie algebroid symplectic structures. A more precise version exists (Theorem 4.8), as well as an analogous result for Lie algebroid fibrations (Theorem 4.5).

THEOREM 1.1. *Let $(\varphi, f): \mathcal{A}_X^4 \rightarrow \mathcal{A}_\Sigma^2$ be a Lie algebroid Lefschetz fibration with connected fibers. Assume that \mathcal{A}_Σ admits a symplectic structure and there exists a closed \mathcal{A}_X -two-form η such that $\eta|_{\ker \varphi}$ is nondegenerate. Then X admits an \mathcal{A}_X -symplectic structure.*

This result can be used as a tool to obtain geometric structures that admit an interpretation as \mathcal{A} -symplectic structures using Lefschetz-type maps. What is most crucial for being able to apply it is understanding how to obtain the above ‘fiberwise symplectic’ form η .

A specific type of boundary map we call a boundary Lefschetz fibration is shown to induce a Lie algebroid Lefschetz fibration between the elliptic and log-tangent bundle. These are essentially genus one Lefschetz fibrations over surfaces, except that generic fibers can collapse to circles over a codimension-one submanifold, which is often the boundary of the surface. This concept extends and formalizes the generalized Lefschetz fibrations of [8]. Using this correspondence we prove the following result (Theorem 7.1) which motivated this work.

THEOREM 1.2. *Let $f: (X^4, D^2) \rightarrow (\Sigma^2, Z^1)$ be a boundary Lefschetz fibration with D coorientable such that (Σ, Z) carries a log-symplectic structure. Assume $[F] \neq 0 \in H_2(X \setminus D; \mathbb{R})$ for the generic fiber F . Then (X, D) admits a stable generalized complex structure.*

A similar result is true in arbitrary dimension, using boundary fibrations instead (Theorem 7.2). This result validates the definition of a boundary Lefschetz fibration as being the type of fibration-like map which should be linked with stable generalized complex structures. This extends the original relationship between Lefschetz fibrations and symplectic structures due to Gompf [13], that between broken Lefschetz fibrations and near-symplectic structures [1, 3], and between achiral Lefschetz fibrations and log-symplectic structures [6, 10].

To produce concrete examples, one has to construct boundary Lefschetz fibrations on explicit four-manifolds. This can be done using genus one Lefschetz fibrations over punctured surfaces with boundary monodromy given by powers of Dehn twists using a process we call completion (see Corollary 6.6). In particular, we have the following corollary (Theorem 8.1).

THEOREM 1.3. *Let $f: X^4 \rightarrow \mathbb{D}^2$ be a genus one Lefschetz fibration over the disc whose monodromy around the boundary is a power of a Dehn twist. Then all possible completions $\tilde{f}: (\tilde{X}, D) \rightarrow (\mathbb{D}^2, \partial\mathbb{D}^2)$ admit a stable generalized complex structure.*

Together with Stefan Behrens we have classified boundary Lefschetz fibrations over the disc \mathbb{D}^2 degenerating over its boundary, establishing the following.

THEOREM 1.4 [4]. *Let $f: (X^4, D^2) \rightarrow (\mathbb{D}^2, \partial\mathbb{D}^2)$ be a relatively minimal boundary Lefschetz fibration. Then X is diffeomorphic to one of the following:*

- (1) $S^1 \times S^3$;
- (2) $\#m(S^2 \times S^2)$, including S^4 for $m = 0$;
- (3) $\#m\mathbb{C}P^2 \#n\mathbb{C}P^2$ with $m > n \geq 0$.

In all cases the generic fiber is nontrivial in $H_2(X \setminus D; \mathbb{R})$. In case (1), D is coorientable, while in cases (2) and (3), D is coorientable if and only if m is odd.

Here relative minimality refers to the nonexistence of spheres with self-intersection -1 in the fibers, nor of those vanishing cycles which are isotopic to the cycle specifying the boundary monodromy of the boundary Lefschetz fibration (see [4, Definition 3.3]). This result can be combined with Theorem 1.2 to equip the manifolds $S^1 \times S^3 \#n\overline{\mathbb{C}P^2}$, $\#(2m+1)\mathbb{C}P^2 \#n\overline{\mathbb{C}P^2}$ and $\#(2m+1)S^2 \times S^2$ with a stable generalized complex structure whose type-change locus has a single component. In particular, these manifolds admit a generalized complex structure with connected type-change locus. Further, they provide a complete list of four-manifolds whose stable generalized complex structures are obtained from boundary Lefschetz fibrations over the disc degenerating over its boundary. Note that in the list of Theorem 1.4, the submanifold $D \subseteq X$ is coorientable if and only if X admits an almost-complex structure, as is required for the existence of a generalized complex structure.

All manifolds in this paper will be compact, unless specifically stated otherwise. Note however that they are not necessarily closed. Throughout we will identify de Rham cohomology with singular cohomology with \mathbb{R} -coefficients.

Organization of the paper

In Section 2 we recall the definition of a stable generalized complex structure and introduce the language of divisors. We further define Poisson structures of divisor-type and recall the definition of a log-symplectic structure.

In Section 3 we recall the definition of a Lie algebroid, and introduce the two relevant Lie algebroids (the elliptic and log tangent bundle) constructed out of divisors. We further define Lie algebroid fibrations and Lie algebroid Lefschetz fibrations, and discuss Lie algebroid symplectic structures and their Poisson counterparts.

In Section 4 we discuss the Lie algebroid version of the Gompf–Thurston argument to construct Lie algebroid symplectic structures (Theorem 4.3). Moreover, we prove results for Lie algebroid fibrations (Theorem 4.5) and Lie algebroid Lefschetz fibrations (Theorem 4.8).

In Section 5 we define the normal Hessian of a map and introduce boundary maps. We define boundary fibrations and boundary Lefschetz fibrations and prove several normal form results for boundary maps (Proposition 5.8, Proposition 5.10, and Proposition 5.15).

In Section 6 we introduce a standard boundary fibration (Proposition 6.2) and use this to obtain boundary Lefschetz fibrations out of genus one Lefschetz fibrations over a punctured surface using monodromy data (Corollary 6.6) via a completion process.

In Section 7 we then prove our main results: Theorem 7.1, that four-dimensional boundary Lefschetz fibrations give rise to stable generalized complex structures; and Theorem 7.2,

that the same holds for boundary fibrations. Moreover, we show compatibility with existing fibrations over T^2 and S^1 can be achieved in the compact case (Corollary 7.6).

In Section 8 we give examples of stable generalized complex manifolds constructed using our methods. In particular, we recover the examples $m\mathbb{C}P^2 \# n\overline{\mathbb{C}P}^2$ for m odd from [8].

2. Stable generalized complex and log-symplectic structures

In this section we recall the notion of a stable generalized complex structure as defined in [9], and that of log-symplectic structures [19]. Moreover, we introduce the language of divisors to study them. For a more comprehensive account, see also [21]. We start by recalling the definition of a generalized complex structure, for which a general reference is [17].

2.1. Generalized complex structures

Let X be a $2n$ -dimensional manifold equipped with a closed three-form $H \in \Omega_{\text{cl}}^3(X)$. Recall that the *double tangent bundle* $\mathbb{T}X := TX \oplus T^*X$ is a Courant algebroid whose anchor is the projection $p: \mathbb{T}X \rightarrow TX$. It carries a natural pairing $\langle V + \xi, W + \eta \rangle = \frac{1}{2}(\eta(V) + \xi(W))$ of split signature and an H -twisted Courant bracket $[[V + \xi, W + \eta]]_H = [V, W] + \mathcal{L}_V \eta - \iota_W d\xi + \iota_V \iota_W H$ for $V, W \in \Gamma(TX)$ and $\xi, \eta \in \Gamma(T^*X)$. Two-forms $B \in \Omega^2(X)$ act via B -field (or gauge) transformations $e^B: \mathbb{T}X \rightarrow \mathbb{T}X$ given by $e^B: V + \xi \mapsto V + \xi + \iota_V B$. This takes $(\mathbb{T}X, H)$ to $(\mathbb{T}X, H + dB)$, leading to the *Severa class* $[H] \in H^3(X; \mathbb{R})$ determining $\mathbb{T}X$ up to Courant isomorphism. The Courant automorphisms of $\mathbb{T}X$ are generated by the diffeomorphisms and closed B -field transformations.

DEFINITION 2.1. A *generalized complex structure* on (X, H) is a complex structure \mathcal{J} on $\mathbb{T}X$ that is orthogonal with respect to $\langle \cdot, \cdot \rangle$, and whose $+i$ -eigenbundle is involutive under $[[\cdot, \cdot]]_H$.

There is an alternative definition of a generalized complex structure using spinors. To state it, recall that sections $v = V + \xi \in \Gamma(\mathbb{T}X)$ of the double tangent bundle act on differential forms via Clifford multiplication, given by $v \cdot \rho = \iota_V \rho + \xi \wedge \rho$ for $\rho \in \Omega^\bullet(X)$.

DEFINITION 2.2. A *generalized complex structure* on (X, H) is given by a complex line bundle $K_{\mathcal{J}} \subset \wedge^\bullet T_{\mathbb{C}}^*X$ pointwise generated by a differential form $\rho = e^{B+i\omega} \wedge \Omega$ with Ω a decomposable complex k -form, satisfying $\Omega \wedge \overline{\Omega} \wedge \omega^{n-k} \neq 0$, and such that $d\rho + H \wedge \rho = v \cdot \rho$ for any local section $\rho \in \Gamma(K_{\mathcal{J}})$ and some $v \in \Gamma(\mathbb{T}X)$.

Both definitions are related using that $K_{\mathcal{J}} = \text{Ann}(E_{\mathcal{J}})$ is the annihilator under the Clifford action of $E_{\mathcal{J}}$, the $+i$ -eigenbundle of \mathcal{J} . The bundle $K_{\mathcal{J}}$ is called the *canonical bundle* of \mathcal{J} . For later use, we further introduce the analogue of a Calabi–Yau manifold in generalized geometry. Denote by $d^H = d + H \wedge$ the H -twisted de Rham differential.

DEFINITION 2.3. A generalized complex structure \mathcal{J} on (X, H) is *generalized Calabi–Yau* if its canonical bundle $K_{\mathcal{J}}$ is determined by a global nowhere vanishing d^H -closed form.

EXAMPLE 2.4. The following provide examples of generalized complex structures on $(X, 0)$.

- Let ω be a symplectic structure on X . Then $K_{\mathcal{J}_\omega} := \langle e^{i\omega} \rangle$ defines a generalized complex structure \mathcal{J}_ω .
- Let J be a complex structure on X with canonical bundle $K_J = \wedge^{n,0} T^*X$. Then $K_{\mathcal{J}_J} := K_J$ defines a generalized complex structure \mathcal{J}_J .
- Let $P \in \Gamma(\wedge^{2,0} TX)$ a holomorphic Poisson structure with respect to a complex structure J . Then $K_{\mathcal{J}_{P,J}} := e^P K_J$ defines a generalized complex structure $\mathcal{J}_{P,J}$.

The automorphisms \mathcal{J}_ω , \mathcal{J}_J and $\mathcal{J}_{P,J}$ are given by, with $\pi = \text{Im}(P)$:

$$\mathcal{J}_\omega = \begin{pmatrix} 0 & -\omega^{-1} \\ \omega & 0 \end{pmatrix}, \quad \mathcal{J}_J = \begin{pmatrix} -J & 0 \\ 0 & J^* \end{pmatrix}, \quad \mathcal{J}_{P,J} = \begin{pmatrix} -J & \pi \\ 0 & J^* \end{pmatrix}.$$

We next introduce the type of a generalized complex structure \mathcal{J} , which colloquially provides a measure for how many complex directions there are. The type is an integer-valued upper semicontinuous function on X whose parity is locally constant.

DEFINITION 2.5. Let \mathcal{J} be a generalized complex structure on X . The *type* of \mathcal{J} is a map $\text{type}(\mathcal{J}): X \rightarrow \mathbb{Z}$ whose value at a point $x \in X$ is the integer k above, the degree of Ω . The *type change locus* $D_{\mathcal{J}}$ of \mathcal{J} is the subset of X where $\text{type}(\mathcal{J})$ is not locally constant.

The complement of $D_{\mathcal{J}}$ is an open dense set where the type is minimal. Using the type, generalized complex structures are seen to interpolate between symplectic and complex structures. At points where $\text{type}(\mathcal{J}) = 0$, the generalized complex structure is equivalent to a symplectic structure, in that it is equivalent to the generalized complex structure \mathcal{J}_ω of a symplectic structure under a B -field transformation.

Any generalized complex structure \mathcal{J} determines a Poisson structure $\pi_{\mathcal{J}}$ as the composition $\pi_{\mathcal{J}}^\# := p \circ \mathcal{J}|_{T^*X}$. The type of \mathcal{J} is related to the rank of $\pi_{\mathcal{J}}$ through the formula $\text{rank}(\pi_{\mathcal{J}}) = 2n - 2 \text{type}(\mathcal{J})$. Using $\pi_{\mathcal{J}}$ one can view a generalized complex structure \mathcal{J} as a foliation on X with symplectic leaves, and a suitably compatible complex structure transverse to the leaves.

2.2. Stable generalized complex structures

Generalized complex structures which are stable were introduced in [9, 14]. Their defining property is a natural condition and since stable generalized complex structures are not far from being symplectic, one can use symplectic techniques to study them.

Let \mathcal{J} be a generalized complex structure on (X, H) . The anticanonical bundle $K_{\mathcal{J}}^*$ has a section $s \in \Gamma(K_{\mathcal{J}}^*)$, given by $s(\rho) := \rho_0$ for $\rho \in \Gamma(K_{\mathcal{J}})$, with ρ_0 the degree-zero part of ρ .

DEFINITION 2.6. A generalized complex structure \mathcal{J} on (X, H) is *stable* if s is transverse to the zero section in $K_{\mathcal{J}}^*$. The set $D_{\mathcal{J}} := s^{-1}(0)$ is a codimension-two smooth submanifold of X called the *anticanonical divisor* of \mathcal{J} .

Outside of $D_{\mathcal{J}}$, the section s is nonvanishing hence the type of \mathcal{J} is equal to zero, while over $D_{\mathcal{J}}$ it is equal to two. Consequently, stable generalized complex structures can be seen as generalized complex structures which are close to being symplectic.

EXAMPLE 2.7. Consider $(\mathbb{C}^2, 0)$ with holomorphic Poisson structure $\pi = z\partial_z \wedge \partial_w$. This gives a stable generalized complex structure with $K_{\mathcal{J}} = \langle z + dz \wedge dw \rangle$ and $D_{\mathcal{J}} = \{z = 0\}$.

By Example 2.4, any holomorphic Poisson structure (J, P) defines a generalized complex structure $\mathcal{J}_{P,J}$ on $(X, 0)$ with $K_{\mathcal{J}_{P,J}} = e^P K_J$. Thus $K_{\mathcal{J}_{P,J}}$ is locally generated by $e^P \Omega$, with Ω a local trivialization of K_J . This immediately proves the following.

PROPOSITION 2.8 [9, Example 2.11]. *Let $(X, 0, J, P)$ be a holomorphic Poisson structure on a complex $2m$ -dimensional manifold. Then $\mathcal{J}_{P,J}$ is a stable generalized complex structure if and only if the Pfaffian $\wedge^m P$ is transverse to the zero section in $\Gamma(\wedge^{2m,0} TX)$.*

Any stable generalized complex structure \mathcal{J} is locally equivalent around points in $D_{\mathcal{J}}$ to $\langle e^{i\omega_0}(z + dz \wedge dw) \rangle$ on $\mathbb{C}^2 \times \mathbb{R}^{2n-4}$, with ω_0 the standard symplectic form on \mathbb{R}^{2n-4} , and

$\langle z + dz \wedge dw \rangle$ the structure on \mathbb{C}^2 of Example 2.7 with $D_{\mathcal{J}} = \{z = 0\}$, see [9, Section 3.4]. On $D_{\mathcal{J}}$ there is an induced type-1 generalized Calabi–Yau structure [9, Theorem 2.13].

Any compact type-1 generalized Calabi–Yau manifold, such as $D_{\mathcal{J}}$, fibers over the torus T^2 [2]. Moreover, the semilocal form of a stable generalized complex structure around its type change locus is given by its *linearization* along $D_{\mathcal{J}}$, which is the stable generalized complex structure naturally present on the normal bundle to this type-1 generalized Calabi–Yau manifold. We will not elaborate on this further here and instead refer to [2, 9].

The anticanonical bundle $K_{\mathcal{J}}^*$ of a stable generalized complex manifold together with its natural section s are a particular example of a *divisor*, which we will introduce shortly. As the theory of divisors permeates much of this work, we now turn to developing this concept.

2.3. Real and complex divisors

In this section we introduce real and complex divisors, which are extensions to the smooth setting of the notion of a divisor used in complex geometry. We further discuss the relation between the ideals they define, and define morphisms between divisors. See also [9, 22].

DEFINITION 2.9. Let X be a manifold. A *divisor* on X is a pair (U, σ) where $U \rightarrow X$ is a real/complex line bundle and $\sigma \in \Gamma(U)$ is a section whose zero set is nowhere dense.

In this paper we will mostly focus on real divisors. As such, we will often drop the prefix ‘real’, while explicitly stating when divisors are instead complex. Examples of divisors will be discussed in upcoming subsections.

Given a divisor (U, σ) , viewing $\sigma: \Gamma(U^*) \rightarrow C^\infty(X)$ specifies a locally principal ideal $I_\sigma := \sigma(\Gamma(U^*))$ which is locally generated by a function with nowhere dense zero set. Letting α be a local trivialization of U^* , we have $\alpha(\sigma) = g$ for some local function g . Then locally $I_\sigma = \langle \alpha(\sigma) \rangle = \langle g \rangle$. Conversely, out of any such ideal we can construct a divisor, which recovers the ideal via this evaluation process. This extends the correspondence between divisors and holomorphic line bundles in complex geometry.

PROPOSITION 2.10 [22]. Let I be a locally principal ideal on X generated by functions with nowhere dense zero set. Then there exists a divisor (U_I, σ) on X such that $I_\sigma = I$.

Proof. Let $\{U_\alpha\}$ be an open cover of X and $f_\alpha \in I(U_\alpha)$ be generators. Then on $U_\alpha \cap U_\beta$ we have $f_\alpha = g_{\alpha\beta} f_\beta$ with $g_{\alpha\beta} \in C^\infty(U_\alpha \cap U_\beta)$. Since $f_\alpha = g_{\alpha\beta} g_{\beta\alpha} f_\alpha$, and f_α has nowhere dense support, we see that each $g_{\alpha\beta}$ is a nonvanishing function and $g_{\beta\alpha} = g_{\alpha\beta}^{-1}$. Similarly the identity $f_\alpha = g_{\alpha\beta} g_{\beta\gamma} g_{\gamma\alpha} f_\alpha$ on $U_\alpha \cap U_\beta \cap U_\gamma$ implies that the functions $g_{\alpha\beta}$ satisfy the cocycle condition. We conclude that $\{(U_{\alpha\beta}, g_{\alpha\beta})\}$ defines a line bundle U_I on X . Further, setting $\sigma|_{U_\alpha} = f_\alpha$ on U_α specifies a section σ of this bundle with the desired properties. \square

The section σ constructed in the proof of the above proposition is unique up to multiplication by a smooth nonvanishing function. We next define morphisms between divisors in terms of the ideals they give rise to. Denote by $f^*I \subseteq C^\infty(X)$ the ideal generated by the pullback of an ideal $I \subseteq C^\infty(Y)$ along a map $f: X \rightarrow Y$.

DEFINITION 2.11. Let (U_X, σ_X) be a divisor on X and (U_Y, σ_Y) a divisor on Y . A map $f: X \rightarrow Y$ is a *morphism of divisors* if $f^*I_{\sigma_Y} = I_{\sigma_X}$.

Equivalently, one can require that $(U_X, \sigma_X) = (f^*U_Y, gf^*\sigma_Y)$ for some $g \in C^\infty(X; \mathbb{R}^*)$. Two divisors are *diffeomorphic* (denoted using \cong) if there exists a morphism of divisors between them which is in addition a diffeomorphism. Two divisors on a fixed manifold X are *isomorphic* (denoted using $=$) if the identity map on X is a morphism of divisors.

2.3.1. *Log divisors.* The simplest type of divisor is that of a log divisor. These are divisors whose zero set will be a smooth codimension-one submanifold, or *hypersurface*.

DEFINITION 2.12. A *log divisor* is a divisor (L, s) whose zero set $Z_s = s^{-1}(0)$ is a smooth hypersurface along which s is transverse to the zero section.

It follows from the definition that the first Stiefel–Whitney class of L is the \mathbb{Z}_2 -Poincaré dual of Z . Further, the intrinsic derivative $ds|_Z: NZ \rightarrow L|_Z$ is an isomorphism.

There is also a complex analogue of a (real) log divisor.

DEFINITION 2.13. A *complex log divisor* is a complex divisor (U, σ) whose zero set $D_\sigma = \sigma^{-1}(0)$ is a smooth codimension-two submanifold with σ transverse to the zero section.

Any complex log divisor (U, σ) has a complex conjugate $(\bar{U}, \bar{\sigma})$ with the same zero locus.

EXAMPLE 2.14. By definition the anticanonical bundle $K_{\mathcal{J}}^*$ of a stable generalized complex structure together with its natural section s is an example of a complex log divisor.

Let (L, s) be a log divisor. The associated ideal $I_Z := I_s$ is exactly the vanishing ideal of the hypersurface Z . Any hypersurface naturally gives rise to a unique log divisor.

PROPOSITION 2.15. *Let X be a manifold and $Z \subset X$ a hypersurface. Then Z carries a unique log divisor structure, that is, there exists a unique log divisor (L, s) with $s^{-1}(0) = Z$.*

Proof. Apply Proposition 2.10 to the vanishing ideal I_Z , giving a divisor $(U_{I_Z}, \sigma) =: (L, s)$. The section s vanishes transversely along Z as it is equal to a local defining function for Z in any trivializing open U_α of L containing Z . We conclude that (L, s) is a log divisor. \square

Because of this result we often identify a log divisor with the associated submanifold Z .

2.3.2. *Elliptic divisors.* More directly relevant to our study of stable generalized complex structures is the notion of an elliptic divisor.

DEFINITION 2.16. An *elliptic divisor* is a divisor (R, q) whose zero set $D_q = q^{-1}(0)$ is a smooth codimension-two critical submanifold of q along which its normal Hessian is definite.

We denote an elliptic divisor by $|D|$. The normal Hessian $\text{Hess}(q) \in \Gamma(D; \text{Sym}^2 N^*D \otimes R)$ of q is the leading term of its Taylor expansion. There is more information available in the elliptic divisor than just the zero set of q . The ideal $I_{|D|} := I_q$ is called an *elliptic ideal*, and both R and q (up to a nonzero smooth function) can be recovered from $I_{|D|}$ by Proposition 2.10.

EXAMPLE 2.17. Let $D = (U, \sigma)$ be a complex log divisor. Then $((U \otimes \bar{U})_{\mathbb{R}}, \sigma \otimes \bar{\sigma})$ is an elliptic divisor $|D|$. Using Proposition 2.20 below and the factorization $x^2 + y^2 = (x + iy)(x - iy) = w\bar{w}$, any elliptic divisor with coorientable zero set arises from a complex log divisor in this way, with (U, σ) being determined up to diffeomorphism by the choice of coorientation.

Note that I_D is not the vanishing ideal of D , but instead is locally generated by an even index Morse–Bott function in coordinates normal to D , as we now explain.

DEFINITION 2.18. Let $g \in C^\infty(X)$ be given. A compact connected submanifold $S \subset X$ is a *nondegenerate critical submanifold* of g if $S \subset \text{Crit}(g)$ and $\ker \text{Hess}(g) = T_p S$ for all $p \in S$. If $\text{Crit}(g)$ consists of nondegenerate critical submanifolds, then g is a *Morse–Bott function*.

Let $g \in C^\infty(X)$ and take S a nondegenerate critical submanifold of g . Consider the exact sequence $0 \rightarrow TS \rightarrow TX|_S \rightarrow NS \rightarrow 0$. For $p \in S$ we have $\text{Hess}(g)(p) \in \text{Sym}^2 T_p^* X$, and this vanishes when restricted to $T_p S$. But then $\text{Hess}(g)(p) \in \text{Sym}^2 N_p^* S$, giving a nondegenerate bilinear form $Q_g \in \Gamma(S; \text{Sym}^2 N^* S)$. The semi-global version of the Morse–Bott lemma says that g is diffeomorphic to this quadratic approximation in a tubular neighborhood of S .

LEMMA 2.19 [29, Proposition 2.6.2]. *Let $g \in C^\infty(X)$ be a Morse–Bott function and S a nondegenerate critical submanifold of g . Then there exists a neighborhood U of the zero section $S \subset NS$ and an open embedding $\Phi: U \rightarrow X$ such that $\Phi|_S = \text{id}_S$ and $\Phi^* g = Q_g$.*

Let $|D| = (R, q)$ be an elliptic divisor. Then R is orientable by q as it is a trivialization away from a codimension-two submanifold. Consequently R is always a trivial line bundle and if one were to orient R using q , the normal Hessian of q along D is positive definite. In other words, let α be a trivialization of R^* . Then $g := \alpha(q) \in C^\infty(X)$ is a function with $g^{-1}(0) = D$ and $\text{Hess}(g) = \alpha(\text{Hess}(q))$. Moreover, D is a nondegenerate critical submanifold of g , and g is locally Morse–Bott around D . As D is codimension two, $X \setminus D$ is connected, so that the sign of g on $X \setminus D$ is fixed. Replace α by $-\alpha$ if necessary so that this sign is positive, and then q and α induce compatible orientations. Call such a trivialization α compatible with q . For compatible trivializations we have $g \geq 0$ so that $\text{Hess}(g)$ is positive definite. As a consequence of Lemma 2.19 we obtain the following.

PROPOSITION 2.20. *Let $|D| = (R, q)$ be an elliptic divisor and α a compatible trivialization. Then there exists a neighborhood U of the zero section $D \subset ND$ and an open embedding $\Phi: U \rightarrow X$ such that $\Phi|_D = \text{id}_D$ and $(\Phi^* \alpha)(\Phi^* q) = Q_g$, where $g = \alpha(q) \in C^\infty(X)$.*

Given $p \in D$ we can locally trivialize the bundles R and ND , so that using Proposition 2.20 the section q can be written locally as $q(x_1, \dots, x_n) = \pm(x_1^2 + x_2^2)$ in normal bundle coordinates such that $ND = \langle \partial_{x_1}, \partial_{x_2} \rangle$. Consequently, the elliptic ideal $I_{|D|}$ is locally generated by r^2 , where $r^2 = x_1^2 + x_2^2$ is the squared radial distance from D inside ND .

REMARK 2.21. While Proposition 2.15 shows that hypersurfaces carry a unique log divisor structure, the same is not true for codimension-two submanifolds and elliptic divisors. A simple example is provided by $X = \mathbb{R}^2$ with $D = \{(0, 0)\}$ and coordinates (x, y) . Equip D with the elliptic ideals $I = \langle x^2 + y^2 \rangle$ and $I' = \langle x^2 + 2y^2 \rangle$. As these ideals are distinct, they supply D with two nonisomorphic yet diffeomorphic elliptic divisor structures.

2.4. Divisors and geometric structures

Divisors provide a convenient way to define and study specific classes of Poisson structures. We first describe the general way in which divisors relate to Poisson geometry. Let X be a $2n$ -dimensional manifold and π a Poisson structure on X . The Pfaffian $\wedge^n \pi$ of π is a section of the real line bundle $\wedge^{2n} TX$.

DEFINITION 2.22. A Poisson structure π is of *divisor-type* if $(\wedge^{2n} TX, \wedge^n \pi)$ is a divisor.

Note that this condition is equivalent to asking that π is generically symplectic. However, it is useful to encode using a divisor exactly what type of degeneracy π has. Poisson structures of divisor-type are more thoroughly explored in [21]. In this paper we will restrict our attention to the two types associated to log and elliptic divisors.

2.4.1. *Log-Poisson and log-symplectic structures.* In this section we discuss Poisson structures built out of log divisors. We refer to [10, 19, 21] for a more comprehensive treatment.

DEFINITION 2.23. Let X be a $2n$ -dimensional manifold. A *log-Poisson structure* is a Poisson structure π on X that is of log divisor-type.

While the name ‘log-Poisson structure’ is most consistent with other Poisson structures built out of divisors, these Poisson structures also carry the name of b -Poisson, b -symplectic, and *log-symplectic structures* [6, 18, 19, 25, 26, 28]. The latter two names emphasize the ability to view log-Poisson structures as suitably degenerate symplectic forms, see Proposition 3.33.

Let (X, Z) be a *log pair*, that is, a manifold X together with a log divisor Z . We say the pair (X, Z) *admits a log-Poisson structure* if there exists a log-Poisson structure π on X such that $Z_{\wedge^n \pi} = Z$. We will also denote $Z_{\wedge^n \pi}$ as Z_π .

The rank of a log-Poisson structure π is equal to $2n$ on $X \setminus Z$, and $2n - 2$ on Z . By the Weinstein splitting theorem [35], any log-Poisson structure π on X^{2n} is locally equivalent around points in Z_π to $x\partial_x \wedge \partial_y + \omega_0^{-1}$ on $\mathbb{R}^2 \times \mathbb{R}^{2n-2}$, where ω_0 is the standard symplectic structure, and $Z_\pi = \{x = 0\}$ (see [19, 28]). On Z_π , a log-symplectic structure π induces an equivalence class of *cosymplectic structures*, which is a pair (α, β) of closed one- and two-forms on Z such that $\alpha \wedge \beta^{n-1} \neq 0$. Here the equivalence relation is generated by addition of two-forms $df \wedge \alpha$, with $f \in C^\infty(Z)$. The one-form α is canonically determined by π , while the two-form β requires a choice of defining function for Z_π (better: a distance function $|z|$ on NZ_π from its zero section). In particular, Z_π carries a codimension-one symplectic foliation.

Any compact cosymplectic manifold, such as Z_π , fibers over S^1 [18, 23, 26, 30]. As can be read in [6, 26], a log-symplectic structure π is said to be *proper* if the induced foliation on Z_π is given by a submersion onto S^1 . This is determined by whether the cohomology class $[\alpha]$ of the one-form above is a real multiple of an integral class. Any log-symplectic structure can be perturbed slightly to become proper. The following innocuous result will be important later in Section 5. Recall that a hypersurface Z is *separating* if $[Z] = 0 \in H_{2n-1}(X; \mathbb{Z}_2)$.

PROPOSITION 2.24. *Let (X, Z) be an orientable log-Poisson pair. Then Z is separating.*

Proof. As X is orientable, $\wedge^{2n}TX$ is trivial. The section $\wedge^n \pi$ has transverse zeros, hence its zero set Z is Poincaré dual to the first Stiefel–Whitney class of the trivial line bundle. \square

Any trivialization $\sigma \in \Gamma(\wedge^{2n}TX)$ produces a global defining function h for Z via $\wedge^n \pi = h\sigma$.

2.4.2. *Elliptic Poisson and stable generalized complex structures.* We can also construct Poisson structures out of elliptic divisors, obtaining the notion of an elliptic Poisson structure. These are called Poisson structures of elliptic log-symplectic type in [9].

DEFINITION 2.25 [9, Definition 3.3]. Let X be a $2n$ -dimensional manifold. An *elliptic Poisson structure* is a Poisson structure π on X that is of elliptic divisor-type.

Let $(X, |D|)$ be an *elliptic pair*, that is, a manifold equipped with an elliptic divisor $|D|$. An elliptic pair $(X, |D|)$ *admits an elliptic Poisson structure* if there exists an elliptic Poisson structure π on X such that $|D_{\wedge^n \pi}| = |D|$. When there is no elliptic divisor structure on D , we say that (X, D) admits an elliptic Poisson structure if there exists some elliptic divisor structure $|D|$ on D such that $(X, |D|)$ admits an elliptic Poisson structure.

The underlying Poisson structure of a stable generalized complex structure \mathcal{J} can be shown to belong to this class, which in fact characterizes when \mathcal{J} is stable.

PROPOSITION 2.26 [9, Theorem 3.7]. *Let (X, H, \mathcal{J}) be a generalized complex manifold. Then \mathcal{J} is a stable generalized complex structure if and only if $\pi_{\mathcal{J}}$ is an elliptic Poisson structure.*

In Section 3.4 we will see that this proposition can be pushed further. Namely, gauge equivalence classes of stable generalized complex structures \mathcal{J} on X are in one-to-one correspondence with certain types of elliptic Poisson structures π via the map $\mathcal{J} \mapsto \pi_{\mathcal{J}}$ (see Theorem 3.32). Moreover, the closed three-form H required to state integrability of \mathcal{J} is determined by π .

An elliptic pair $(X, |D|)$ admits a stable generalized complex structure if there exists a closed three-form $H \in \Omega_{cl}^3(X)$ and a stable generalized complex structure \mathcal{J} on (X, H) such that $|D_{\mathcal{J}}| = |D|$. When no elliptic divisor structure is specified on D , we say that (X, D) admits a stable generalized complex structure if there exists some elliptic divisor structure $|D|$ on D such that $(X, |D|)$ admits a stable generalized complex structure.

3. Lie algebroids and Lie algebroid symplectic structures

In this section we deal with some aspects of the general theory of symplectic forms in a Lie algebroid. For interesting applications, the Lie algebroid should be chosen so that such symplectic forms describe a certain type of geometric structure. The main examples relevant to this paper are the elliptic tangent bundle and the log-tangent bundle, describing stable generalized complex structures and log-symplectic structures respectively. More examples of Lie algebroids with interesting symplectic geometry can be found in [21].

DEFINITION 3.1. A Lie algebroid is a vector bundle $\mathcal{A} \rightarrow X$ together with a Lie bracket $[\cdot, \cdot]_{\mathcal{A}}$ on $\Gamma(\mathcal{A})$ and a vector bundle map $\rho_{\mathcal{A}}: \mathcal{A} \rightarrow TX$ called the anchor, satisfying the Leibniz rule $[v, fw]_{\mathcal{A}} = f[v, w]_{\mathcal{A}} + \mathcal{L}_{\rho_{\mathcal{A}}(v)}f \cdot w$ for all $v, w \in \Gamma(\mathcal{A})$ and $f \in C^\infty(X)$.

We think of Lie algebroids as generalizations of the tangent bundle TX , chosen such that geometric constructions done using \mathcal{A} are more suitable to the situation at hand. This should be kept in mind especially when we define Lie algebroid fibrations in Section 3.3, as we will replace many usual notions by their Lie algebroid counterparts. The Lie algebroids we consider will be isomorphic via their anchor to the tangent bundle outside of a submanifold of positive codimension. We introduce the following subset to keep track of this.

DEFINITION 3.2. Let $\mathcal{A} \rightarrow X$ be a Lie algebroid with anchor $\rho_{\mathcal{A}}: \mathcal{A} \rightarrow TX$. The isomorphism locus of \mathcal{A} is the open set $X_{\mathcal{A}} \subseteq X$ where $\rho_{\mathcal{A}}$ is an isomorphism.

Any Lie algebroid gives rise to a graded algebra $\Omega^\bullet(\mathcal{A}) = \Gamma(\wedge^\bullet \mathcal{A}^*)$ of differential \mathcal{A} -forms. This algebra comes equipped with a differential $d_{\mathcal{A}}$ squaring to zero, constructed using the bracket $[\cdot, \cdot]_{\mathcal{A}}$ by means of the Koszul formula. This defines a cohomology theory for \mathcal{A} .

DEFINITION 3.3. The Lie algebroid cohomology of $\mathcal{A} \rightarrow X$ is $H_{\mathcal{A}}^k(X) := H^k(\Omega^\bullet(\mathcal{A}), d_{\mathcal{A}})$.

For notational convenience we denote by $\Omega_{\mathcal{A}}^k(U)$, for an open subset $U \subset X$, the set of \mathcal{A} - k -forms defined on U , so that $\Omega_{\mathcal{A}}^k(X) = \Omega^k(\mathcal{A})$. The inclusion $i: X_{\mathcal{A}} \hookrightarrow X$ of the isomorphism locus induces a bijection $\rho_{\mathcal{A}}^*: \Omega^k(X_{\mathcal{A}}) \rightarrow \Omega_{\mathcal{A}}^k(X_{\mathcal{A}})$.

3.1. The log-tangent bundle and elliptic tangent bundle

In this section we discuss the Lie algebroids that are of primary concern to us. These are the log-tangent bundle and the elliptic tangent bundle, constructed out of log divisors and

elliptic divisors respectively. Both are examples of *ideal Lie algebroids* [21], which we now introduce.

Let X be an n -dimensional manifold and let I be an ideal sheaf, and denote by \mathcal{V}_X the sheaf of vector fields on X . Let $\mathcal{V}_X(I) := \{v \in \mathcal{V}_X \mid \mathcal{L}_v I \subset I\}$ be the sheaf of derivations preserving I . This is a subsheaf of Lie algebras of \mathcal{V}_X . If $\mathcal{V}_X(I)$ is in addition locally finitely generated projective, it gives rise to a Lie algebroid \mathcal{A}_I with $\Gamma(\mathcal{A}_I) = \mathcal{V}_X(I)$ by the Serre–Swan theorem.

DEFINITION 3.4. The Lie algebroid \mathcal{A}_I is called the *ideal Lie algebroid* associated to I .

As $\mathcal{V}_X(I)$ is a submodule of $\Gamma(TX)$, the anchor of any ideal Lie algebroid is the natural inclusion on sections. However, as vector bundle map the anchor of \mathcal{A}_I need not be an isomorphism. More precisely, the isomorphism locus $X_{\mathcal{A}_I}$ of \mathcal{A}_I is the complement of the support $\text{supp}(C^\infty(X)/I)$ of the quotient sheaf, or equivalently, the complement of $\text{supp}(\mathcal{V}_X/\mathcal{V}_X(I))$.

3.1.1. *The log-tangent bundle.* Let $Z = (L, s)$ be a log divisor on X with associated ideal $I_Z = s(\Gamma(L^*))$. This ideal is exactly the vanishing ideal of Z , and $\mathcal{V}_X(I_Z)$ is the locally free sheaf of vector fields tangent to Z [27]. In local adapted coordinates (z, x_2, \dots, x_n) around $Z = \{z = 0\}$ with $I_z = \langle z \rangle$, one has $\mathcal{V}_X(I_Z) = \langle z\partial_z, \partial_{x_2}, \dots, \partial_{x_n} \rangle$.

DEFINITION 3.5. The *log-tangent bundle* $TX(-\log Z) \rightarrow X$ is the ideal Lie algebroid on X with $\Gamma(TX(-\log Z)) = \mathcal{V}_X(I_Z)$.

It is immediate that the isomorphism locus of $TX(-\log Z)$ is given by $X \setminus Z$. In analogy with the holomorphic case, denote $\Omega^k(\log Z) = \Omega^k(TX(-\log Z))$.

REMARK 3.6. The log-tangent bundle is also called the *b-tangent bundle* [27] and is then denoted by bTX . We use the name log-tangent bundle as it shows the similarities with the elliptic tangent bundle defined below, and its notation allows us to keep track of Z .

The log-tangent bundle admits a residue map. The restriction of $TX(-\log Z)$ to Z surjects onto TZ via the anchor map ρ , giving the following exact sequence

$$0 \rightarrow \mathbb{L}_Z \rightarrow TX(-\log Z)|_Z \xrightarrow{\rho} TZ \rightarrow 0,$$

where $\mathbb{L}_Z \rightarrow Z$ is the kernel of ρ . Dualizing gives a projection map $\text{Res}_Z : \Omega^k(\log Z) \rightarrow \Omega^{k-1}(Z)$, which fits in the residue sequence

$$0 \rightarrow \Omega^\bullet(X) \xrightarrow{\rho^*} \Omega^\bullet(\log Z) \xrightarrow{\text{Res}_Z} \Omega^{\bullet-1}(Z) \rightarrow 0.$$

In terms of the local coordinate system above, with $\Gamma(TX(-\log Z)) = \langle z\partial_z, \partial_{x_2}, \dots, \partial_{x_n} \rangle$, a given log k -form $\alpha \in \Omega^k(\log Z)$ can be expressed as

$$\alpha = d \log z \wedge \alpha_0 + \alpha_1,$$

with α_i smooth forms. The inclusion $j_Z : Z \hookrightarrow X$ gives $\text{Res}_Z(\alpha) = j_Z^* \alpha_0$. The following result referred to as the Mazzeo–Melrose theorem shows the above sequence splits, and identifies the Lie algebroid cohomology of the log-tangent bundle $TX(-\log Z)$ in terms of X and Z .

THEOREM 3.7 [27]. *Let (X, Z) be a log pair. Then $H^k(TX(-\log Z)) \cong H^k(X) \oplus H^{k-1}(Z)$.*

3.1.2. *The elliptic tangent bundle.* Similarly, let $(X, |D|)$ be an elliptic pair with associated ideal $I_{|D|} = q(\Gamma(R^*))$. Note this is not the vanishing ideal of D . The associated submodule $\mathcal{V}_X(I_{|D|})$ defines a sheaf of locally constant rank [9], locally generated in appropriate polar coordinates $(r, \theta, x_3, \dots, x_n)$ around $D = \{r = 0\}$ such that $I_{|D|} = \langle r^2 \rangle$ by $\langle r\partial_r, \partial_\theta, \partial_{x_3}, \dots, \partial_{x_n} \rangle$.

DEFINITION 3.8. The *elliptic tangent bundle* $TX(-\log |D|) \rightarrow X$ is the ideal Lie algebroid on X with $\Gamma(TX(-\log |D|)) = \mathcal{V}_X(I_{|D|})$.

The isomorphism locus of $TX(-\log |D|)$ is given by $X \setminus D$. As for the log-tangent bundle, we denote $\Omega^k(\log |D|) = \Omega^k(TX(-\log |D|))$.

REMARK 3.9. There is another ideal Lie algebroid defined in [9], namely the *complex log-tangent bundle* $TX(-\log D)$ (there called the *logarithmic tangent bundle*). These are associated to complex log divisors $D = (U, \sigma)$. This Lie algebroid is a generalization of the log-tangent bundle one can define on complex manifolds equipped with a divisor. We will not directly use this Lie algebroid in this paper, hence we will not elaborate on its properties.

The elliptic tangent bundle admits several residue maps [9], three of which we will now describe. The *elliptic residue* Res_q comes from considering the restriction of $TX(-\log |D|)$ to D , which fits in an exact sequence

$$0 \rightarrow \mathbb{R} \oplus \mathfrak{k} \rightarrow TX(-\log |D|)|_D \rightarrow TD \rightarrow 0, \tag{3.1}$$

with \mathbb{R} generated by the Euler vector field of ND , and $\mathfrak{k} \cong \wedge^2 N^*D \otimes R$. Choosing a coorientation for D , that is, a trivialization of ND , also trivializes \mathfrak{k} . Dualizing the above sequence we obtain a projection map $\text{Res}_q: \Omega^k(\log |D|) \rightarrow \Omega^{k-2}(D, \mathfrak{k}^*)$, with kernel $\Omega_0^*(\log |D|)$ the subcomplex of $\Omega^*(\log |D|)$ of zero elliptic residue forms.

Denote by S^1ND the circle bundle associated to ND . The *radial residue* Res_r arises from quotienting (3.1) by the Euler vector field of ND , giving the extension

$$0 \rightarrow \mathfrak{k} \rightarrow \text{At}(S^1ND) \rightarrow TD \rightarrow 0,$$

where $\text{At}(S^1ND)$ is the associated Atiyah algebroid of S^1ND . Noting that $TX(-\log |D|)|_D$ is a trivial extension of $\text{At}(S^1ND)$, the elliptic residue factors through the radial residue map $\text{Res}_r: \Omega^k(\log |D|) \rightarrow \Gamma(D, \wedge^{k-1} \text{At}(S^1ND)^*)$. When the elliptic residue vanishes, the radial residue naturally maps to $\Omega^{k-1}(D)$ without needing a coorientation. Finally, there is also a θ -residue $\text{Res}_\theta: \Omega_0^k(\log |D|) \rightarrow \Omega^{k-1}(D)$, which we will only define for forms with zero elliptic residue. We provide a description of these residue maps in local coordinates. In the adapted coordinate system around D as above, where $\Gamma(TX(-\log |D|)) = \langle r\partial_r, \partial_\theta, \partial_{x_3}, \dots, \partial_{x_n} \rangle$, a given elliptic k -form $\alpha \in \Omega^k(\log |D|)$ can be locally written as

$$\alpha = d \log r \wedge d\theta \wedge \alpha_0 + d \log r \wedge \alpha_1 + d\theta \wedge \alpha_2 + \alpha_3,$$

where each α_i is a smooth form. Using the inclusion $j_D: D \hookrightarrow X$ we have $\text{Res}_q(\alpha) = j_D^* \alpha_0$, $\text{Res}_r(\alpha) = (d\theta \wedge \alpha_0 + \alpha_1)|_D$, and we set $\text{Res}_\theta(\alpha) := j_D^* \alpha_2$. Moreover, we see that using these coordinates we have $\text{Res}_q(\alpha) = \iota_{\partial_\theta} \text{Res}_r(\alpha)$.

As for the log-tangent bundle, the Lie algebroid cohomology of the elliptic tangent bundle as well as its zero elliptic residue version can be expressed in terms of X and D .

THEOREM 3.10 [9, Theorems 1.8, 1.12]. *Let $(X, |D|)$ be an elliptic pair. Then one has $H^k(\log |D|) \cong H^k(X \setminus D) \oplus H^{k-1}(S^1ND)$. Moreover, $H_0^k(\log |D|) \cong H^k(X \setminus D) \oplus H^{k-1}(D)$.*

The above isomorphisms are induced by the maps (i^*, Res_r) with $i: X \setminus D \hookrightarrow X$ the inclusion of the divisor complement, noting Res_r naturally maps to $\Omega^{k-1}(D)$ when the elliptic residue vanishes (in that case, $\text{Res}_r(\alpha) = \alpha_1|_D$ in the above local coordinates). For later purposes we describe how the radial residue of representatives of classes in $H_0^k(\log |D|)$ can be controlled. Let $\Omega_{0,0}^k(\log |D|) \subseteq \Omega_0^k(\log |D|)$ be defined as the kernel $\ker(\text{Res}_r: \Omega_0^k(\log |D|) \rightarrow \Omega^{k-1}(D))$.

LEMMA 3.11. *Let $(X, |D|)$ be an elliptic pair. The short exact sequences of complexes of sheaves*

$$0 \rightarrow \Omega_{0,0}^\bullet(\log |D|) \rightarrow \Omega_0^\bullet(\log |D|) \xrightarrow{\text{Res}_r} \Omega^{\bullet-1}(D) \rightarrow 0 \tag{3.2}$$

induces short exact sequences in cohomology, that is,

$$0 \rightarrow H_{0,0}^k(\log |D|) \rightarrow H_0^k(\log |D|) \xrightarrow{\text{Res}_r} H^{k-1}(D) \rightarrow 0. \tag{3.3}$$

Moreover, $H_{0,0}^k(\log |D|) \cong \ker(\text{Res}_r : H_0^k(\log |D|) \rightarrow H^{k-1}(D)) \cong H^k(X \setminus D)$.

Proof. To show that the sequence in equation (3.2) is exact, we only have to show that Res_r surjects onto $\Omega^{\bullet-1}(D)$. This is indeed the case, as will now argue. Fix a choice of appropriate tubular neighborhood embedding $p: U \subseteq ND \rightarrow D$ for D provided by Proposition 2.20 (in which $TX(-\log |D|)$ is trivialized as in Section 3.1.2). Moreover, choose a bump function $\psi: \mathbb{R}_{\geq 0} \rightarrow [0, 1]$ for which $\psi \equiv 1$ near 0, and let r be a radial distance function from D . Given $\alpha \in \Omega^{k-1}(D)$, note that the form $p^*(\alpha) \wedge d(\psi(r) \log r) \in \Omega_0^k(\log |D|)$ maps onto α via Res_r .

To show there are induced short exact sequences in cohomology, we have to show that the induced map $H_0^k(\log |D|) \rightarrow H^{k-1}(D)$ is surjective. This follows from Theorem 3.10.

Finally, the isomorphism $H_{0,0}^k(\log |D|) \cong \ker(\text{Res}_r : H_0^k(\log |D|) \rightarrow H^{k-1}(D))$ follows immediately from exactness of the sequence in equation (3.3). \square

3.2. Lie algebroid morphisms

In this section we discuss morphisms between Lie algebroids, focusing specifically on those between the elliptic and log-tangent bundle.

DEFINITION 3.12. Let $\mathcal{A}, \mathcal{A}' \rightarrow X$ be Lie algebroids over the same base X . A *Lie algebroid morphism* is a vector bundle map $(\varphi, f): \mathcal{A} \rightarrow \mathcal{A}'$ such that $df \circ \rho_{\mathcal{A}} = \rho_{\mathcal{A}'} \circ \varphi$ and $\varphi[v, w]_{\mathcal{A}} = [\varphi(v), \varphi(w)]_{\mathcal{A}'}$ for all $v, w \in \Gamma(\mathcal{A})$.

The above definition does not immediately generalize to varying base, as a vector bundle map does not give a map on the space of sections, as is required in order to state the bracket condition. There is a description of bracket compatibility in terms of the pullback bundle $f^*\mathcal{A}'$, but we will instead use the following equivalent definition in terms of their duals.

Let $\mathcal{A} \rightarrow X$ and $\mathcal{A}' \rightarrow X'$ be Lie algebroids. Vector bundle maps $(\varphi, f): \mathcal{A} \rightarrow \mathcal{A}'$ are in one-to-one correspondence with algebra morphisms $\varphi^*: \Omega^\bullet(\mathcal{A}') \rightarrow \Omega^\bullet(\mathcal{A})$. Using this we can phrase the conditions that φ preserves anchors and brackets in terms of φ^* .

DEFINITION 3.13. A vector bundle map $(\varphi, f): \mathcal{A} \rightarrow \mathcal{A}'$ is a *Lie algebroid morphism* if $\varphi^*: \Omega^\bullet(\mathcal{A}') \rightarrow \Omega^\bullet(\mathcal{A})$ is a chain map, that is $d_{\mathcal{A}'} \circ \varphi^* = \varphi^* \circ d_{\mathcal{A}}$.

A Lie algebroid morphism $(\varphi, f): \mathcal{A} \rightarrow \mathcal{A}'$ is in particular a morphism of anchored vector bundles, that is, satisfies $df \circ \rho_{\mathcal{A}} = \rho_{\mathcal{A}'} \circ \varphi$. Moreover, a Lie algebroid morphism restricts pointwise to a linear map $\varphi: \ker \rho_{\mathcal{A}} \rightarrow \ker \rho_{\mathcal{A}'}$ between kernels of the respective anchor maps. Furthermore, $\rho_{\mathcal{A}}$ gives an isomorphism between $\ker \varphi$ and $\ker df$ when in the isomorphism locus. For the Lie algebroids we will consider, smooth maps of the underlying manifolds give rise to Lie algebroid morphisms, as long as they intertwine the anchor maps. This is true in general for anchored vector bundle morphisms between Lie algebroids with dense isomorphism loci.

PROPOSITION 3.14. *Let $\mathcal{A}_X \rightarrow X$ and $\mathcal{A}_Y \rightarrow Y$ be Lie algebroids such that $X_{\mathcal{A}_X}$ is dense. Suppose that $(\varphi, f): \mathcal{A}_X \rightarrow \mathcal{A}_Y$ is an anchored bundle morphism and $f^{-1}(Y_{\mathcal{A}_Y}) = X_{\mathcal{A}_X}$. Then $(\varphi, f): \mathcal{A}_X \rightarrow \mathcal{A}_Y$ is a Lie algebroid morphism.*

Proof. As φ is a vector bundle morphism, φ^* is an algebra morphism. In the isomorphism loci, φ must equal df , and (df, f) is a Lie algebroid morphism between TX and TY , that is, f^* is a chain map. By density of $X_{\mathcal{A}_X}$, the map φ^* is a chain map everywhere. \square

Consequently, for such Lie algebroids one can determine whether there is a Lie algebroid morphism (φ, f) by checking that f^* extends to a map φ^* on forms. This in turn will follow by the universal property of the exterior algebra if it holds on generators, so it suffices to check that f^* extends to $\varphi^*: \Omega^1(\mathcal{A}_Y) \rightarrow \Omega^1(\mathcal{A}_X)$. We will now perform such a check to obtain a Lie algebroid morphism between the elliptic and log tangent bundles.

3.2.1. *Morphisms between elliptic and log-tangent bundles.* Let $(X, |D|)$ be an elliptic pair and (Y, Z) a log pair, with associated ideals $I_{|D|} \subset C^\infty(X)$ and $I_Z \subset C^\infty(Y)$. A morphism of the corresponding divisors gives rise to a Lie algebroid morphism.

PROPOSITION 3.15. *Let $f: (X, |D|) \rightarrow (Y, Z)$ be a morphism of divisors. Then df induces a Lie algebroid morphism $(\varphi, f): TX(-\log |D|) \rightarrow TY(-\log Z)$ such that $\varphi = df$ on sections.*

Proof. It is immediate that $f^{-1}(Z) = D$. Note that the isomorphism loci of the elliptic tangent bundle and the log-tangent bundle are dense. By Proposition 3.14 it thus suffices to show that df induces a vector bundle morphism, which in turn is equivalent to showing that f^* extends to an algebra morphism from $\Omega^\bullet(Y; \log Z)$ to $\Omega^\bullet(X; \log |D|)$. Let $x \in D$ and denote $y = f(x) \in Z$. Consider suitable tubular neighborhood coordinates $(r, \theta, x_3, \dots, x_n)$ in a neighborhood U of x such that $U \cap D = \{r = 0\}$ with $I_{|D|} = \langle r^2 \rangle$, and (z, y_2, \dots, y_m) in a neighborhood V of y such that $V \cap Z = \{z = 0\}$ and $I_Z = \langle z \rangle$. In these coordinates we have $\Omega^1(U; -\log |D|) = \langle d \log r, d\theta, dx_3, \dots, dx_n \rangle$ and $\Omega^1(V; -\log Z) = \langle d \log z, dy_2, \dots, dy_m \rangle$.

The Lie algebroid one-forms dy_i can be pulled back using f^* as these are smooth. Moreover, the smooth one-forms inject into $\Omega^1(U; -\log D)$ using the anchor. We are left with checking that $d \log z$ is pulled back to a Lie algebroid one-form. As f is a morphism of divisors we have $f^*I_Z = I_{|D|}$, so that $f^*(z) = gr^2$ for g a smooth nonvanishing function on U . Consequently,

$$f^*d \log z = d \log f^*(z) = d \log (gr^2) = d \log |g| + 2d \log r \in \Omega^1(U; -\log |D|).$$

We conclude that df induces a Lie algebroid morphism as desired. \square

REMARK 3.16. Lie algebroid morphisms between log-tangent bundles arise out of so-called *b-maps*, which are maps $f: (X, Z_X) \rightarrow (Y, Z_Y)$ between log pairs such that $f^{-1}(Z_Y) = Z_X$ and f is transverse to Z_Y . Alternatively, they are exactly the corresponding morphisms of divisors, that is, those maps satisfying $f^*I_{Z_Y} = I_{Z_X}$ [21]. Similarly, morphisms between elliptic tangent bundles arise from those maps $f: (X, |D_X|) \rightarrow (Y, |D_Y|)$ between elliptic pairs for which $f^{-1}(D_Y) = D_X$ and f is transverse to D_Y . Again, these can be alternatively described as maps satisfying $f^*I_{|D_Y|} = I_{|D_X|}$.

3.2.2. *Lie algebroid morphisms and residue maps.* A Lie algebroid morphism from the elliptic tangent bundle $TX(-\log |D|)$ to the log tangent bundle $TY(-\log Z)$ intertwines the residue maps that were discussed in Section 3.1.

PROPOSITION 3.17. *Let $(\varphi, f): TX(-\log |D|) \rightarrow TY(-\log Z)$ be a Lie algebroid morphism. Then $\text{Res}_q \circ \varphi^* = 0$. Moreover, $(\text{Res}_r + \text{Res}_\theta) \circ \varphi^* = f^* \circ \text{Res}_Z$.*

To prove the above proposition, we briefly elaborate on how the residue maps come about. Given a short exact sequence $\mathcal{S}: 0 \rightarrow E \rightarrow W \rightarrow V \rightarrow 0$ of vector spaces, there is an associated dual sequence $\mathcal{S}^*: 0 \rightarrow V^* \rightarrow W^* \rightarrow E^* \rightarrow 0$. For a given $k \in \mathbb{N}$, by taking k th exterior powers we obtain a filtration of spaces $\mathcal{F}^i := \{\rho \in \wedge^k W^* \mid \iota_x \rho = 0 \text{ for all } x \in \wedge^i E\}$, for $i = 0, \dots, k + 1$. These spaces satisfy

$$\mathcal{F}^0 = 0, \quad \mathcal{F}^1 = \wedge^k V^*, \quad \mathcal{F}^i \subset \mathcal{F}^{i+1}, \quad \text{and} \quad \mathcal{F}^{i+1}/\mathcal{F}^i \cong \wedge^{k-i} V^* \otimes \wedge^i E^*.$$

Setting $\ell := \dim E$, we have $\mathcal{F}^{\ell+1} = \wedge^k W^*$. This allows for the following definition.

DEFINITION 3.18. The residue of $\rho \in \wedge^k W^*$ is $\text{Res}(\rho) = [\rho] \in \mathcal{F}^{\ell+1}/\mathcal{F}^\ell \cong \wedge^{k-\ell} V^* \otimes \wedge^\ell E^*$.

Upon a choice of trivialization of $\wedge^\ell E^*$, that is, a choice of orientation for E , one can view the residue $\text{Res}(\rho)$ as an element of $\wedge^{k-\ell} V^*$. One can further consider lower residue maps $\text{Res}_{-m}: \wedge^k W^* \rightarrow \mathcal{F}^{\ell+1}/\mathcal{F}^{\ell-m}$ for $m > 0$. These are always defined but have a better description for forms $\rho \in \wedge^k W^*$ whose higher residues vanish, because in this case we have

$$\text{Res}_{-m}(\rho) \in \mathcal{F}^{\ell-m+1}/\mathcal{F}^{\ell-m} \cong \wedge^{k-\ell+m} V^* \otimes \wedge^{\ell-m} E^*.$$

Given a map of short exact sequences $\Psi: \tilde{\mathcal{S}} \rightarrow \mathcal{S}$ with dual map $\Psi^*: \mathcal{S}^* \rightarrow \tilde{\mathcal{S}}^*$, there is a corresponding map of filtrations $\Psi^*: \mathcal{F}^i \rightarrow \tilde{\mathcal{F}}^i$. Setting $\tilde{\ell} := \dim \tilde{E}$, we have the following.

LEMMA 3.19. In the above setting, assume $\tilde{\ell} > \ell$. Then $\widetilde{\text{Res}}(\Psi^* \rho) = 0$ for all $\rho \in \wedge^k W^*$.

Proof. We have $\rho \in \mathcal{F}^{\ell+1}$ so that $\Psi^* \rho \in \tilde{\mathcal{F}}^{\ell+1}$. As $\tilde{\ell} > \ell$, we have $\tilde{\mathcal{F}}^{\ell+1} \subset \tilde{\mathcal{F}}^{\tilde{\ell}} \subset \tilde{\mathcal{F}}^{\tilde{\ell}+1}$, so that $\widetilde{\text{Res}}(\Psi^* \rho) = [\Psi^* \rho] \in \tilde{\mathcal{F}}^{\tilde{\ell}+1}/\tilde{\mathcal{F}}^{\tilde{\ell}}$ vanishes as desired. \square

Assuming $\tilde{\ell} > \ell$, all lower residues automatically vanish by degree reasons until considering $\Psi^* \rho \in \tilde{\mathcal{F}}^{\ell+1}$. Hence the first possibly nonzero residue is

$$\widetilde{\text{Res}}_{\ell-\tilde{\ell}}(\Psi^* \rho) = [\Psi^* \rho] \in \tilde{\mathcal{F}}^{\ell+1}/\tilde{\mathcal{F}}^{\tilde{\ell}} \cong \wedge^{k-\ell} \tilde{V}^* \otimes \wedge^\ell \tilde{E}^*.$$

Similarly to Lemma 3.19 we obtain the following.

LEMMA 3.20. In the above setting, assuming $\tilde{\ell} \geq \ell$, we have $\Psi^* \circ \text{Res} = \widetilde{\text{Res}}_{\ell-\tilde{\ell}} \circ \Psi^*$.

With this understood, we can turn to proving Proposition 3.17.

Proof of Proposition 3.17. By assumption we have $f^{-1}(Z) = D$ so that $df: TD \rightarrow TZ$. Restricting $TX(-\log|D|)$ to D and $TY(-\log Z)$ to Z gives the following commutative diagram.

$$\begin{array}{ccccccc} \tilde{\mathcal{S}}: 0 & \longrightarrow & \mathbb{R} \oplus \mathfrak{k} & \longrightarrow & TX(-\log|D|)|_D & \xrightarrow{\rho_X} & TD \longrightarrow 0 \\ & & \varphi \downarrow & & \varphi \downarrow & & \downarrow df \\ \mathcal{S}: 0 & \longrightarrow & \mathbb{L}_Z & \longrightarrow & TY(-\log Z)|_Z & \xrightarrow{\rho_Y} & TZ \longrightarrow 0 \end{array}$$

Consequently, we obtain a map $\varphi^*: \mathcal{S}^* \rightarrow \tilde{\mathcal{S}}^*$ between dual sequences, and also between spaces of sections. Using the notation preceding this proof we have $E = \mathbb{L}_Z$ so that $\ell = \dim(E) = 1$, and $\tilde{E} = \mathbb{R} \oplus \mathfrak{k}$ so that $\tilde{\ell} = \dim(\tilde{E}) = 2$. Recall that \mathbb{L}_Z carries a canonical trivialization. Given a form $\alpha \in \Omega^k(\log Z)$, we can identify $\text{Res}(\alpha) \in \Gamma(Z; \wedge^{k-1} T^* Z \otimes \mathbb{L}_Z^*)$ with $\text{Res}_Z(\alpha) \in \Omega^{k-1}(Z)$. Similarly, a choice of coorientation for ND trivializes $\mathfrak{k} = \wedge^2 N^* D \otimes R$. Given $\beta \in \Omega^k(\log|D|)$ this identifies $\widetilde{\text{Res}}(\beta) \in \Gamma(D; \wedge^{k-2} T^* D \otimes \mathfrak{k}^*)$ with $\text{Res}_q(\beta) \in \Omega^{k-2}(D)$, using that $\wedge^2(\mathbb{R} \oplus \mathfrak{k}) \cong$

†. Moreover, for β with $\text{Res}_q(\beta) = 0$, the radial residue $\text{Res}_r(\beta) \in \Omega^{k-1}(D)$ together with the θ -residue $\text{Res}_\theta(\beta) \in \Omega^{k-1}(D)$ is identified with $\widetilde{\text{Res}}_{-1}(\beta) \in \Omega^{k-1}(D; \mathbb{R} \oplus \mathfrak{t})$. As $2 = \tilde{\ell} \geq \ell = 1$ with $\ell - \tilde{\ell} = -1$ we obtain immediately from Lemma 3.19 and Lemma 3.20 that $\text{Res}_q(\varphi^*\alpha) = 0$ and that $(\text{Res}_r + \text{Res}_\theta)(\varphi^*\alpha) = f^*(\text{Res}_Z(\alpha))$. \square

REMARK 3.21. The proof of Proposition 3.17 shows that the residue maps are also intertwined for morphisms between log-tangent bundles, and between elliptic tangent bundles.

3.3. Lie algebroid Lefschetz fibrations

In this section we introduce the appropriate notions of fibration and Lefschetz fibration in the context of Lie algebroid morphisms.

DEFINITION 3.22. A Lie algebroid fibration is a Lie algebroid morphism $(\varphi, f): \mathcal{A} \rightarrow \mathcal{A}'$ for which $\varphi: \mathcal{A} \rightarrow f^*\mathcal{A}'$ is surjective. Equivalently, φ should be fiberwise surjective.

Note that if $f: X \rightarrow Y$ is a fibration, then $(df, f): TX \rightarrow TY$ is a Lie algebroid fibration.

REMARK 3.23. Our notion of a Lie algebroid fibration differs from the one used by other authors, notably Mackenzie [24]. Our Lie algebroid fibrations are not required to cover a surjective submersion. In other words, only φ should be fiberwise surjective, not both φ and df . This is in line with viewing \mathcal{A} as the replacement of TX .

We next introduce Lie algebroid Lefschetz fibrations, which are a simultaneous generalization of Lefschetz fibrations, as well as of Lie algebroid fibrations. We first recall the notion of a Lefschetz fibration. See [13] for more information.

DEFINITION 3.24. A Lefschetz fibration is a proper map $f: X^{2n} \rightarrow \Sigma^2$ between oriented connected manifolds which is injective on critical points and such that for each critical point $x \in X$ there exist orientation preserving complex coordinate charts centered at x and $f(x)$ in which f takes the form $f: \mathbb{C}^n \rightarrow \mathbb{C}$, $f(z_1, \dots, z_n) = z_1^2 + \dots + z_n^2$.

REMARK 3.25. The requirement that Lefschetz fibrations are injective on critical points is convenient but not essential. If it is not satisfied it can be ensured by a small perturbation.

Next we essentially separate two types of singular behavior, namely that of the anchors of the Lie algebroids, and that of the morphism between them. In the isomorphism locus of the Lie algebroid, the condition of being a Lie algebroid fibration is just that it be a fibration. We can weaken this condition here, and only here, to allow for Lefschetz-type singularities.

DEFINITION 3.26. A Lie algebroid Lefschetz fibration $(\varphi, f): \mathcal{A}_X^{2n} \rightarrow \mathcal{A}_\Sigma^2$ is a Lie algebroid morphism where \mathcal{A}_X has dense isomorphism locus and $f^{-1}(\Sigma_{\mathcal{A}_\Sigma}) = X_{\mathcal{A}_X}$, for which there is a discrete set $\Delta \subset X_{\mathcal{A}_X}$ with $f(\Delta) \subset \Sigma_{\mathcal{A}_\Sigma}$ and

- $f|_{X_{\mathcal{A}_X}}: X_{\mathcal{A}_X} \rightarrow \Sigma_{\mathcal{A}_\Sigma}$ is a Lefschetz fibration with $\text{Crit}(f|_{X_{\mathcal{A}_X}}) = \Delta$;
- $(\varphi, f): X \setminus f^{-1}(f(\Delta)) \rightarrow \Sigma \setminus f(\Delta)$ is a Lie algebroid fibration.

REMARK 3.27. Note that there are no Lefschetz singularities outside of $X_{\mathcal{A}_X}$. Moreover, whenever Δ is nonempty, both $X_{\mathcal{A}_X}$ and $\Sigma_{\mathcal{A}_\Sigma}$ are nonempty, and hence $\dim(X) = 2n$ and $\dim(\Sigma) = 2$.

In Section 4, we will use Lie algebroid Lefschetz fibrations whose generic fibers in $X_{\mathcal{A}_X}$ are connected. Unlike for usual Lefschetz fibrations [13], for Lie algebroid Lefschetz fibrations there is in general no exact sequence in homotopy by which we can ensure connected fibers.

3.4. Lie algebroid symplectic structures

In this section we discuss symplectic structures for a Lie algebroid. We start by defining Poisson structures for Lie algebroids, as the symplectic forms we consider in this paper will be obtained through their bivector counterparts. Throughout, let $\mathcal{A} \rightarrow X$ be a Lie algebroid. The bracket $[\cdot, \cdot]_{\mathcal{A}}$ extends in the natural way to an \mathcal{A} -Schouten bracket on $\Gamma(\wedge^{\bullet} \mathcal{A})$, again denoted by $[\cdot, \cdot]_{\mathcal{A}}$.

DEFINITION 3.28. An \mathcal{A} -Poisson structure is an \mathcal{A} -bivector $\pi_{\mathcal{A}} \in \Gamma(\wedge^2 \mathcal{A})$ with $[\pi_{\mathcal{A}}, \pi_{\mathcal{A}}]_{\mathcal{A}} = 0$.

DEFINITION 3.29. A Poisson structure π on X is said to be of \mathcal{A} -type for a Lie algebroid $\rho_{\mathcal{A}}: \mathcal{A} \rightarrow X$ if there exists an \mathcal{A} -bivector $\pi_{\mathcal{A}} \in \Gamma(\wedge^2 \mathcal{A})$ such that $\rho_{\mathcal{A}}(\pi_{\mathcal{A}}) = \pi$.

In the above situation we call $\pi_{\mathcal{A}}$ an \mathcal{A} -lift of π . Such a lift need not be unique if it exists (consider the zero Poisson structure and any \mathcal{A} with zero anchor), but it is unique if $X_{\mathcal{A}}$ is dense, as will be the case for us. Moreover, when $X_{\mathcal{A}}$ is dense, any \mathcal{A} -lift of a Poisson structure is automatically \mathcal{A} -Poisson. Note that we have $\pi^{\sharp} = \rho_{\mathcal{A}} \circ \pi_{\mathcal{A}}^{\sharp} \circ \rho_{\mathcal{A}}^*$ as maps.

DEFINITION 3.30. An \mathcal{A} -symplectic structure is a closed and nondegenerate \mathcal{A} -two-form $\omega_{\mathcal{A}} \in \Omega_{\mathcal{A}}^2(X)$. Denote the space of \mathcal{A} -symplectic forms by $\text{Symp}(\mathcal{A})$.

An \mathcal{A} -Poisson structure $\pi_{\mathcal{A}}$ is called *nondegenerate* if $\pi_{\mathcal{A}}^{\sharp}$ is an isomorphism. As when $\mathcal{A} = TX$, for any Lie algebroid of even rank there is a bijection between \mathcal{A} -symplectic forms and nondegenerate \mathcal{A} -Poisson structures. Namely, given an \mathcal{A} -symplectic structure $\omega_{\mathcal{A}}$, nondegeneracy implies we can invert the map $\omega_{\mathcal{A}}^b: \mathcal{A} \rightarrow \mathcal{A}^*$ to $(\omega_{\mathcal{A}}^b)^{-1} = \pi_{\mathcal{A}}^{\sharp}: \mathcal{A}^* \rightarrow \mathcal{A}$ for an \mathcal{A} -Poisson structure $\pi_{\mathcal{A}}$. The conditions $d_{\mathcal{A}}\omega_{\mathcal{A}} = 0$ and $[\pi_{\mathcal{A}}, \pi_{\mathcal{A}}]_{\mathcal{A}} = 0$ are equivalent.

Given an \mathcal{A} -symplectic structure $\omega_{\mathcal{A}}$, we call $\pi = \rho_{\mathcal{A}}(\pi_{\mathcal{A}})$ the *dual bivector* to $\omega_{\mathcal{A}}$. We say π is of *nondegenerate \mathcal{A} -type* if it admits a nondegenerate \mathcal{A} -lift $\pi_{\mathcal{A}}$. In this paper we will focus on nondegenerate \mathcal{A} -Poisson structures, as we wish to use symplectic techniques.

The elliptic Poisson structures from Section 2.4 are in one-to-one correspondence with symplectic forms in the associated elliptic tangent bundle, that is, *elliptic symplectic structures*.

PROPOSITION 3.31 [9, Lemma 3.4]. A Poisson structure π on X^{2n} is elliptic if and only if it is of nondegenerate \mathcal{A} -type, for $\mathcal{A} = TX(-\log |D|)$ with elliptic divisor $|D| = (\wedge^{2n} TX, \wedge^n \pi)$.

We can now state the extension of Proposition 2.26, giving a characterization of stable generalized complex structure purely in terms of elliptic symplectic structures.

THEOREM 3.32 [9, Theorem 3.7]. Let X be a compact manifold. There is a bijection $(\mathcal{J}, H) \rightarrow (\pi_{\mathcal{J}}^{-1}, \mathfrak{o})$ between stable generalized complex structures up to gauge equivalence and elliptic symplectic structures with vanishing elliptic residue and cooriented degeneracy locus.

The associated closed three-form H in the definition of a generalized complex structure can be determined via $[H] = \text{Res}_r([\omega_{\mathcal{A}}]) \wedge \text{PD}_X[D]$, where $\omega_{\mathcal{A}}$ is the Lie algebroid symplectic structure for $\mathcal{A} = TX(-\log |D_{\mathcal{J}}|)$ whose dual bivector is $\pi_{\mathcal{J}}$, and $D = (\wedge^n \pi_{\mathcal{J}})^{-1}(0)$. The Poincaré dual of D requires a choice of coorientation.

Similarly to Proposition 3.31, there is a bijective correspondence between log-Poisson structures, and log-symplectic structures. This was used in [10] to construct log-Poisson structures.

PROPOSITION 3.33 [19, Proposition 20]. *A Poisson structure π on X^{2n} is log if and only if it is of nondegenerate \mathcal{A} -type, where $\mathcal{A} = TX(-\log Z)$ with log divisor $Z = (\wedge^{2n}TX, \wedge^n\pi)$.*

In this correspondence, the one-form α discussed in Section 2.4.1 is obtained from π as $\alpha = \text{Res}_Z(\omega)$, and thus is canonically determined by ω (or π). Finally, we note that a Lie algebroid $\mathcal{A} \rightarrow X$ of rank 2 admits an \mathcal{A} -symplectic structure if and only if \mathcal{A} is orientable, that is, when $w_1(\mathcal{A}) = 0$. Namely, any nonvanishing section of $\wedge^2\mathcal{A}$ is a nondegenerate \mathcal{A} -Poisson structure by dimension reasons. For later use we record the following, also noted in [10].

LEMMA 3.34. *Let Σ^2 be a compact oriented surface. Then $(\Sigma, \partial\Sigma)$ admits a log-symplectic structure. For a hypersurface Z , the log pair (Σ, Z) admits a log-symplectic structure if and only if $[Z] = 0 \in H_1(\Sigma; \mathbb{Z}_2)$.*

Proof. By the previous discussion, the pair (Σ, Z) carries a log-symplectic structure if and only if $T\Sigma(-\log Z)$ is orientable. Note that $w_1(T\Sigma(-\log Z)) = w_1(\Sigma) + w_1(L)$ via the bundle isomorphism $T\Sigma(-\log Z) \oplus L \cong T\Sigma \oplus \mathbb{R}$ [21], where $L = L_{I_Z}$ in the notation of Proposition 2.10. As an orientable manifold with boundary has orientable boundary, the result follows, noting that $w_1(L) = \text{PD}_{\mathbb{Z}_2}(Z)$, the Poincaré dual of $[Z]$ with \mathbb{Z}_2 -coefficients. \square

4. Constructing Lie algebroid symplectic structures

In this section we consider the Thurston argument [31] for constructing symplectic structures extended by Gompf [12], and adapt it to the context of Lie algebroid symplectic forms. The case of the log-tangent bundle can be found in [10]. The guiding principle is to combine suitably symplectic-type structures from the base of a fibration-like map with a form that is symplectic on the tangent spaces of the fibers of that map. In the Lie algebroid case one uses Lie algebroid morphisms $(\varphi, f): \mathcal{A}_X \rightarrow \mathcal{A}_Y$. Special attention is required because $\rho_{\mathcal{A}_X}: \ker \varphi \rightarrow \ker df$ need not be an isomorphism (nor injective or surjective), hence one should interpret the tangent space to the fibers suitably.

We will use Lie algebroid almost-complex structures as certificates for nondegeneracy of forms, by using the notion of tameness. Let $\mathcal{A}_X \rightarrow X$ be a Lie algebroid and $\omega \in \text{Symp}(\mathcal{A}_X)$.

DEFINITION 4.1. An \mathcal{A}_X -almost-complex structure is a vector bundle complex structure J for \mathcal{A}_X . An \mathcal{A}_X -almost-complex structure J is ω -tame if $\omega(v, Jv) > 0$ for all $v \in \Gamma(\mathcal{A}_X)$. Given $(\varphi, f): \mathcal{A}_X \rightarrow \mathcal{A}_Y$ a Lie algebroid morphism and $\omega_Y \in \text{Symp}(\mathcal{A}_Y)$, J is (ω_Y, φ) -tame if $(\varphi_x^*\omega_Y)(v, Jv) > 0$ for all $v \in \mathcal{A}_{X,x} \setminus \ker \varphi_x$ and $x \in X$.

As usual, the space of taming \mathcal{A}_X -almost-complex structures for ω is convex and nonempty and is denoted by $\mathcal{J}(\omega)$. Note that any $\omega \in \Omega_{\mathcal{A}_X}^2(X)$ taming an \mathcal{A}_X -almost-complex structure J is necessarily nondegenerate. Hence if ω is a closed \mathcal{A}_X -two-form taming some J , then ω is \mathcal{A}_X -symplectic and J induces the same \mathcal{A}_X -orientation as ω . Note moreover that if J is (ω_Y, φ) -tame, then $\ker \varphi$ is a J -complex subspace of \mathcal{A}_X . Indeed, if $v \in \ker \varphi$ and $Jv \notin \ker \varphi$, we would have $0 = \varphi_x^*\omega_Y(v, Jv) = \varphi_x^*\omega_Y(Jv, J(Jv)) > 0$, which is a contradiction.

PROPOSITION 4.2. *The taming condition is open, that is, it is preserved under sufficiently small perturbations of ω and J , and of varying the point in X .*

Proof. The taming condition $\omega(v, Jv) > 0$ for the pair (ω, J) holds provided it holds for all $v \in \Sigma\mathcal{A}_X \subset \mathcal{A}_X$, the unit sphere bundle with respect to some preassigned metric, as X is compact. As $\Sigma\mathcal{A}_X$ is also compact, the continuous map $\tilde{\omega}: \Sigma\mathcal{A}_X \rightarrow \mathbb{R}$ given by $\tilde{\omega}(v) := \omega(v, Jv)$ for $v \in \Sigma\mathcal{A}_X$ is bounded from below by a positive constant on $\Sigma\mathcal{A}_X$. But then $\tilde{\omega}$ will remain positive under small perturbations of ω and J . Similarly the condition of ω taming J on $\ker \varphi$ is open. Consider $x \in X$ so that $\omega(v, Jv) > 0$ for all $v \in \ker \varphi_x$. As $\tilde{\omega}$ is continuous and $\Sigma\mathcal{A}_X$ is compact, there exists a neighborhood U of $\ker \varphi_x \cap \Sigma\mathcal{A}_X$ in $\Sigma\mathcal{A}_X$ on which $\tilde{\omega}$ is positive. Points $x' \in X$ close to x will then have $\ker \varphi_{x'} \subset U$ because $\ker \varphi$ is closed. \square

We will not use the associated notion of compatibility, where J further leaves ω invariant, as we use almost-complex structures as auxiliary structures to show nondegeneracy, and make use of the openness of this condition. For this reason, all almost-complex structures will only be required to be continuous as this avoids arguments to ensure smoothness.

The following is the Lie algebroid version of [12, Theorem 3.1] and is our main tool to construct \mathcal{A}_X -symplectic structures; see [10, Theorem 3.4] for the log-tangent version.

THEOREM 4.3. *Let $(\varphi, f): (X, \mathcal{A}_X) \rightarrow (Y, \mathcal{A}_Y)$ be a Lie algebroid morphism between compact connected manifolds, J an \mathcal{A}_X -almost-complex structure, ω_Y an \mathcal{A}_Y -symplectic form and η a closed \mathcal{A}_X -two-form. Assume that*

- (i) J is (ω_Y, φ) -tame;
- (ii) η tames $J|_{\ker \varphi_x}$ for all $x \in X$.

Then $\omega_X := \varphi^\omega_Y + t\eta$ is an \mathcal{A}_X -symplectic structure on X for $t > 0$ sufficiently small.*

The proof of this result is modeled on that by Gompf of [12, Theorem 3.1].

Proof. Let $t > 0$ and consider the form $\omega_t := \varphi^*\omega_Y + t\eta$. We show J is ω_t -tame for t small enough. By Proposition 4.2 it is enough to show that there exists a $t_0 > 0$ so that $\omega_t(v, Jv) > 0$ for every $t \in (0, t_0)$ and v in the unit sphere bundle $\Sigma\mathcal{A}_X \subset \mathcal{A}_X$ with respect to some metric. For $v \in \mathcal{A}_X$ we have $\omega_t(v, Jv) = \varphi^*\omega_Y(v, Jv) + t\eta(v, Jv)$. As J is (ω_Y, φ) -tame, the first term is positive for $v \in \mathcal{A}_X \setminus \ker \varphi$ and is zero otherwise. The second term $\eta(v, Jv)$ is positive on $\ker \varphi$ because $J|_{\ker \varphi}$ is η -tame, hence is also positive for all v in some neighborhood U of $\ker \varphi \cap \Sigma\mathcal{A}_X$ in \mathcal{A}_X by openness of the taming condition. We conclude that $\omega_t(v, Jv) > 0$ for all $t > 0$ when $v \in U$. The function $v \mapsto \eta(v, Jv)$ is bounded on the compact set $\Sigma\mathcal{A}_X \setminus U$. Furthermore, $\varphi^*\omega_Y(v, Jv)$ is also bounded from below there by a positive constant, as it is positive away from $\ker \varphi$, and thus also away from $\ker \varphi \cap \Sigma\mathcal{A}_X \subset U$. But then $\omega_t(v, Jv) > 0$ for all $0 < t < t_0$ for t_0 sufficiently small, so that ω_t is \mathcal{A}_X -symplectic for t small enough. \square

Given a map $f: X \rightarrow Y$ we denote by $F_y = f^{-1}(y)$ for $y \in Y$ the level set, or fiber, of f over y . In order to meaningfully apply Theorem 4.3 we must be able to construct closed \mathcal{A}_X -forms η as in (ii) of the statement. Note first that it suffices to have such closed forms in neighborhoods of fibers, all lying in the same global Lie algebroid cohomology class.

PROPOSITION 4.4. *Assume that there exists a class $c \in H^2_{\mathcal{A}_X}(X; \mathbb{R})$ and that for each $y \in Y$, F_y has a neighborhood W_y with a closed \mathcal{A}_X -two-form $\eta_y \in \Omega^2_{\mathcal{A}_X}(W_y)$ such that $[\eta_y] = c|_{W_y} \in H^2_{\mathcal{A}_X}(W_y; \mathbb{R})$, and η_y tames $J|_{\ker \varphi_x}$ for all $x \in W_y$. Then there exists a closed \mathcal{A}_X -two-form η such that $[\eta] = c$ and η tames $J|_{\ker \varphi_x}$ for all $x \in X$.*

Proof. Let $\xi \in \Omega^2_{\mathcal{A}_X}(X)$ be closed and such that $[\xi] = c$. Then for each $y \in Y$ we have $[\eta_y] = c|_{W_y} = [\xi]|_{W_y}$, so on W_y we have $\eta_y = \xi + d_{\mathcal{A}_X}\alpha_y$ for some $\alpha_y \in \Omega^1_{\mathcal{A}_X}(W_y)$. As each $X \setminus W_y$ and hence $f(X \setminus W_y)$ is compact, each $y \in Y$ has a neighborhood disjoint from $f(X \setminus W_y)$.

Cover Y by a finite amount of such open sets U_i so that each $f^{-1}(U_i)$ is contained in some W_{y_i} . Let $\{\psi_i\}$ be a partition of unity subordinate to the cover $\{U_i\}$ of Y , so that $\{\psi_i \circ f\}$ is a partition of unity of X . Define an \mathcal{A}_X -two-form $\eta \in \Omega^2_{\mathcal{A}_X}(X)$ on X via

$$\eta := \xi + d_{\mathcal{A}_X} \left(\sum_i (\psi_i \circ f) \alpha_{y_i} \right) = \xi + \sum_i (\psi_i \circ f) d_{\mathcal{A}_X} \alpha_{y_i} + \sum_i \rho_{\mathcal{A}_X}^* (d\psi_i \circ df) \wedge \alpha_{y_i}.$$

Then $d_{\mathcal{A}_X} \eta = 0$ and $[\eta] = c$. As $\rho_{\mathcal{A}_X}^* (d\psi_i \circ df) = d\psi_i \circ df \circ \rho_{\mathcal{A}_X} = dg \circ \rho_{\mathcal{A}_Y} \circ \varphi$ using that φ is a Lie algebroid morphism, the last of the above three terms vanishes when applied to a pair of vectors in $\ker \varphi_x$ for any $x \in X$, so on each $\ker \varphi_x$ we have

$$\eta = \xi + \sum_i (\psi_i \circ f) d_{\mathcal{A}_X} \alpha_{y_i} = \sum_i (\psi_i \circ f) (\xi + d_{\mathcal{A}_X} \alpha_{y_i}) = \sum_i (\psi_i \circ f) \eta_{y_i}.$$

From the above we see that on $\ker \varphi$, the \mathcal{A}_X -form η is a convex combination of J -taming \mathcal{A}_X -forms, so $J|_{\ker \varphi}$ is η -tame. □

One can further look for (local or global) closed two-forms $\tilde{\eta} \in \Omega^2(X)$ so that $\eta = \rho_{\mathcal{A}_X}^* \tilde{\eta}$ satisfies hypothesis (ii) of Theorem 4.3. When using such \mathcal{A}_X -forms which are pullbacks of regular forms, the behavior of the map $\rho_{\mathcal{A}_X} : \ker \varphi_x \rightarrow \ker df_x$ is important. In Section 7 we will see an example where there cannot be an η of the form $\eta = \rho_{\mathcal{A}_X}^* \tilde{\eta}$ making $\ker \varphi$ symplectic. This is unlike the situation for log-symplectic forms as studied in [10], where the anchor of the log-tangent bundle provides a pointwise isomorphism between $\ker \varphi$ and $\ker df$.

Using Theorem 4.3 we can prove the Lie algebroid version of Thurston’s result for symplectic fiber bundles with two-dimensional fibers [31], adapting the proof by Gompf in [11].

THEOREM 4.5. *Let $(\varphi, f) : (X, \mathcal{A}_X^{2n}) \rightarrow (Y, \mathcal{A}_Y^{2n-2})$ be a Lie algebroid fibration between compact connected manifolds. Assume that Y is \mathcal{A}_Y -symplectic and there exists a closed \mathcal{A}_X -two-form η which is nondegenerate on $\ker \varphi$. Then $\omega_X := \varphi^* \omega_Y + t\eta$ for $\omega_Y \in \text{Symp}(\mathcal{A}_Y)$ is an \mathcal{A}_X -symplectic structure on X for $t > 0$ sufficiently small.*

Proof. Let $\omega_Y \in \text{Symp}(\mathcal{A}_Y)$ and choose $J_Y \in J(\omega_Y)$. Fix the orientation for $\ker \varphi$ so that η is positive. Let g be a metric on \mathcal{A}_X and let $H \subset \mathcal{A}_X$ be the subbundle of orthogonal complements to $\ker \varphi$, so that $\varphi : H \rightarrow \mathcal{A}_Y$ is a fiberwise isomorphism. Define an \mathcal{A}_X -almost-complex structure J by letting $J|_H = \varphi^* J_Y$, and on $\ker \varphi$, use the metric and define J by $\frac{\pi}{2}$ -counterclockwise rotation, demanding φ is orientation preserving via the fiber-first convention. This determines J uniquely on \mathcal{A}_X by linearity. Moreover, J is (ω_Y, φ) -tame as $\varphi^* \omega_Y(v, Jv) = \omega_Y(\varphi v, J_Y \varphi v) > 0$, for all $v \in \mathcal{A}_X \setminus \ker \varphi \cong H$. Further, η tames J on $\ker \varphi$ as J is compatible with the orientation on $\ker \varphi$ determined by η . By Theorem 4.3 we obtain an \mathcal{A}_X -symplectic structure. □

REMARK 4.6. One can combine Theorem 4.5 with Proposition 4.4 to obtain a statement requiring only the existence of local forms η_y governed by a global cohomology class.

Sadly for general Lie algebroid morphisms there is no direct analogue of homological essentialness of generic fibers to replace the hypothesis of the existence of η as in Theorem 4.3 and Theorem 4.5. In other words, we cannot replace the condition on the existence of η by demanding that the generic fiber F satisfies $[F] \neq 0 \in H_2(X; \mathbb{R})$. This is a well-known necessary condition when constructing symplectic-like structures out of Lefschetz-type fibrations using Gompf–Thurston techniques. The reason for the lack of such an analogue is again the behavior of the map $\rho_{\mathcal{A}_X} : \ker \varphi \rightarrow \ker df$. While the codomain can be seen as the tangent space to the fiber at regular points, the domain cannot. For Lie algebroid submersions, surjectivity is only

demanded of φ , not of df , and there is no duality pairing between homology and Lie algebroid cohomology in general.

REMARK 4.7. In the special cases of the log-tangent bundle and the elliptic tangent bundle one can state such an analogue (see Theorems 7.1 and 7.2, and [10]), precisely because we understand their respective Lie algebroid cohomologies (see Theorem 3.7 and Theorem 3.10).

Recall that we defined a Lie algebroid Lefschetz fibration to be a Lie algebroid fibration away from the set of Lefschetz singularities, which must be a subset of the isomorphism locus of \mathcal{A}_X . Note that $X_{\mathcal{A}_X}$ is either empty or of codimension zero, so that we can restrict \mathcal{A}_X to $X_{\mathcal{A}_X}$ and again obtain a Lie algebroid. In this way the inclusion $i: X_{\mathcal{A}_X} \hookrightarrow X$ covers a Lie algebroid morphism ι , and we obtain a restriction map in cohomology $\iota^*: H_{\mathcal{A}_X}(X) \rightarrow H_{\mathcal{A}_X}(X_{\mathcal{A}_X})$. The anchor gives a bijection $\rho_{\mathcal{A}_X}^*: \Omega^\bullet(X_{\mathcal{A}_X}) \rightarrow \Omega_{\mathcal{A}_X}^\bullet(X_{\mathcal{A}_X})$ and hence an isomorphism $\rho_{\mathcal{A}_X}^*: H_{\text{dR}}(X_{\mathcal{A}_X}) \rightarrow H_{\mathcal{A}_X}(X_{\mathcal{A}_X})$ in cohomology over $X_{\mathcal{A}_X}$.

Assuming $\text{rank}(\mathcal{A}_X) = \text{rank}(\mathcal{A}_Y) + 2$, the second hypothesis in Theorem 4.3 demands the existence of a class $c \in H_{\mathcal{A}_X}(X)$ such that the de Rham class $(\rho_{\mathcal{A}_X}^*)^{-1}\iota^*(c)$ evaluates nonzero on each generic fiber. Even if such a class exists, a similar statement must hold over $X \setminus X_{\mathcal{A}_X}$. Hence, we need the existence of a two-form $\eta \in \Omega_{\text{cl}}^2(X; \mathcal{A}_X)$ such that $(\rho_{\mathcal{A}_X}^*)^{-1}\iota^*[\eta]$ evaluates nonzero on the fibers, and $\eta|_{\ker \varphi}$ is nowhere zero over $X \setminus X_{\mathcal{A}_X}$.

The notion of a Lie algebroid Lefschetz fibration is such that the hypotheses of Theorem 4.3 are still satisfied, when in dimension 4.

THEOREM 4.8. *Let $(\varphi, f): \mathcal{A}_X^4 \rightarrow \mathcal{A}_\Sigma^2$ be a Lie algebroid Lefschetz fibration with connected fibers between compact connected manifolds. Assume that \mathcal{A}_Σ admits a symplectic structure $\omega_{\mathcal{A}_\Sigma}$ and there exists $\eta \in \Omega_{\text{cl}}^2(\mathcal{A}_X)$ such that $(\rho_{\mathcal{A}_X}^*)^{-1}\iota^*[\eta]$ evaluates nonzero on the fibers, and $\eta|_{\ker \varphi}$ is nondegenerate over $X \setminus X_{\mathcal{A}_X}$. Then, for $t > 0$ sufficiently small, X admits an \mathcal{A}_X -symplectic structure that is equal to $\varphi^*\omega_\Sigma + t\eta$ in a neighborhood of $X \setminus X_{\mathcal{A}_X}$.*

The proof of this result is similar to [10, Theorem 4.7], which in turn is based on Gompf’s proof of [13, Theorem 10.2.18] using almost-complex structures. We will use Proposition 4.4 to glue together the given form η with a suitable adaptation in the isomorphism locus of \mathcal{A}_X . Recall that by assumption we assume that Lefschetz fibrations are injective on critical points. Given a critical value $y \in \Sigma$, we write $x = f^{-1}(y)$ for the unique critical point mapping to y .

REMARK 4.9. Note that if $\mathcal{A}_X = TX$, then $X_{\mathcal{A}_X} = X$ and we are demanding that the generic fiber is homologically essential, recovering the result by Gompf [13, Theorem 10.2.18].

Proof. If $X_{\mathcal{A}_X}$ is empty, $\eta|_{\ker \varphi}$ is nondegenerate everywhere and φ is a Lie algebroid fibration, so that the result follows from Theorem 4.5. If not, then X is four dimensional. Denote $\xi := (\rho_{\mathcal{A}_X}^*)^{-1}\iota^*\eta \in \Omega^2(X_{\mathcal{A}_X})$ and $c = [\xi] \in H_{\text{dR}}^2(X_{\mathcal{A}_X})$. Note that ξ orients the generic fibers F , as these are two dimensional. Recall that $\Delta \subset X_{\mathcal{A}_X}$ is the set of Lefschetz singularities, and let $\Delta' := f(\Delta)$ be the set of singular values of $f|_{X_{\mathcal{A}_X}}$.

Use the proof of Theorem 4.5 to obtain a (φ, ω_Σ) -tame almost-complex structure J on \mathcal{A}_X over $X \setminus f^{-1}(\Delta')$ compatible with the orientation on $\ker \varphi$, noting that $f: X \setminus f^{-1}(\Delta') \rightarrow \Sigma \setminus \Delta'$ is a Lie algebroid fibration. As Δ' is contained in $\Sigma_{\mathcal{A}_\Sigma}$, let $V \subseteq \Sigma$ be the disjoint union of open balls V_y disjoint from $\Sigma \setminus \Sigma_{\mathcal{A}_\Sigma}$ and centered at each point $y \in \Delta'$. Let $W := f^{-1}(V) \subset X$ be the union of the neighborhoods $W_y := f^{-1}(V_y)$ of singular fibers F_y . Let $C \subset X$ be the disjoint union of open balls $C_y \subseteq \Sigma_{\mathcal{A}_\Sigma}$ centered at each critical point $f^{-1}(y) = x$ for all $y \in \Delta'$ so that on each ball f takes on the local form in Definition 3.24. Possibly shrink C so that $\overline{C_y} \subset W_y$. The local description of f gives an almost-complex structure on C with fibers being holomorphic, and we glue this to the existing almost-complex structure J on $X \setminus C$ (this is

possible by holomorphicity of the fibers in the coordinates on C , see [12, Lemma 3.2; 13, Exercise 8.1.6]). This gives a global (ω_Σ, φ) -tame \mathcal{A}_X -almost-complex structure J .

As $\eta|_{\ker \varphi}$ is nondegenerate over $X \setminus X_{\mathcal{A}_X}$, the same is true in a neighborhood S around $X \setminus X_{\mathcal{A}_X}$ disjoint from Δ . Let $y \in Y \setminus f(S)$ be given. If $y \notin \Delta'$, let $D_y \subset Y \setminus f(S)$ disjoint from Δ' be a disc containing y , fully contained in a trivializing neighborhood of f around y . Define $W_y := f^{-1}(D_y) \cong D_y \times F_y$, with projection map $p: W_y \rightarrow D_y$. Let η'_y be an area form on F_y compatible with the preimage orientation. Define $\eta_y := \lambda_y \rho_X^* p^* \eta'_y$, where $\lambda_y \in \mathbb{R}$ is chosen such that $\langle [F], c \rangle = \langle [F], \eta_y \rangle$. As $H_2(W_y; \mathbb{R})$ is generated by $[F_y]$, it follows that $[\eta_y] = c|_{W_y} \in H^2(W_y; \mathbb{R})$. But then η_y tames J on $\ker \varphi \cong \ker df$ for all $x \in W_y$, as the restriction of $(\rho_X^*)^{-1} \eta_y = \lambda_y p^* \eta'_y$ is an area form for that fiber.

If $y \in \Delta'$, the singular fiber F_y either is indecomposable or consists of two irreducible components F_y^\pm which satisfy $[F_y^+] \cdot [F_y^-] = 1$ and $[F_y^\pm]^2 = -1$, see [13, p. 289]. In the latter case, note that $0 < \lambda = \langle c, [F] \rangle = \langle c, [F_y] \rangle = \langle c, [F_y^+] \rangle + \langle c, [F_y^-] \rangle$. If either term is nonpositive assume without loss of generality that $\langle c, [F_y^-] \rangle = r \leq 0$. Define $c' := c + (\frac{\lambda}{2} - r)c_y^+$, where $c_y^+ \in H^2(X; \mathbb{R})$ is a class dual to $[F_y^+]$. As $[F_y] \cdot [F_y^\pm] = 0$ we then have $\langle c', [F] \rangle = \langle c, [F] \rangle > 0$, and furthermore

$$\langle c', [F_y^+] \rangle = \langle c, [F_y^+] \rangle - \left(\frac{\lambda}{2} - r \right) = (\lambda - r) - \frac{\lambda}{2} + r = \frac{\lambda}{2} > 0 \quad \text{and} \quad \langle c', [F_y^-] \rangle = \frac{\lambda}{2} > 0.$$

Moreover, as different fibers do not intersect, we have $c|_{W_{y'}} = c'|_{W_{y'}}$ for $y' \neq y$. After finitely many repetitions, at most once for each $y \in \Delta'$, one obtains a class, again denoted by c , pairing positively with every fiber component (see [13, Exercise 10.2.19]).

Return to $y \in \Delta'$ and let σ be the standard symplectic form on C_y given locally in real coordinates by $\sigma = dx_1 \wedge dy_1 + dx_2 \wedge dy_2$, where $z_i = x_i + iy_i$. As all fibers F'_y in C_y are holomorphic, $\rho_X^* \sigma|_{F'_y \cap C_y}$ tames J for all $y' \in f(C_y)$, so that $\rho_X^* \sigma$ tames J on C_y . Let σ_y be an extension of σ to F_y as a positive area form with total area $\langle \sigma_y, [F_y] \rangle$ equal to $\langle c, [F_y] \rangle$. Let $p: W_y \rightarrow F_y$ be a retraction and let $\psi: C_y \rightarrow [0, 1]$ be a smooth radial function so that $\psi \equiv 0$ in a neighborhood of $x = f^{-1}(y) \in \Delta$ and $\psi \equiv 1$ in a neighborhood of ∂C_y , which is smoothly extended to W_y by being identically 1 outside C_y . On the ball C_y , the form σ is exact, say equal to $\sigma = d\alpha$ for $\alpha \in \Omega^1(C_y)$.

Define a two-form η'_y on W_y by $\eta'_y := p^*(\psi \sigma_y) + d((1 - \psi)\alpha)$, which is closed as $\psi \sigma_y$ is a closed area form on F_y . Near x we have $\psi \equiv 0$ so that $\eta'_y = d\alpha = \sigma$. Set $\eta_y := \rho_X^* \eta'_y$. Then there $\eta_y = \rho_X^* \sigma$ tames J , hence in particular tames $J|_{\ker \varphi}$. Similarly, σ_y is an area form on $F_y \setminus \{x\}$ for the orientation given by J under the isomorphism by ρ_X . But then $\rho_X^* \sigma_y$ tames J on $\ker \varphi_y \cong \ker df = TF_y$ on F_y , so that the same holds for η_y as this condition is convex. By openness of the taming condition, shrinking V_y and hence W_y and possibly C_y we can ensure that η_y tames $J|_{\ker \varphi}$ on W_y . Finally, note that $[\eta'_y] = c|_{W_y} \in H^2(W_y; \mathbb{R})$ by construction. For points $y \in f(S)$, take the Lie algebroid two-form $\eta_y := \eta$ on the neighborhood $W_y := S$ of F_y .

This gives the required neighborhoods W_y and forms η_y for all $y \in Y$ to apply Proposition 4.4 and we obtain a Lie algebroid closed two-form again denoted by η such that η tames J on $\ker \varphi_x$ for all $x \in X$. By Theorem 4.3 we obtain an \mathcal{A}_X -symplectic structure on X . \square

REMARK 4.10. When both \mathcal{A}_X and \mathcal{A}_Σ are log-tangent bundles, Theorem 4.8 recovers [10, Theorem 4.7], upon combination with the fact that $\ker \varphi$ and $\ker df$ are pointwise isomorphic.

5. Boundary maps and boundary Lefschetz fibrations

In this section we introduce the notion of a boundary map, which is a map degenerating suitably on a submanifold. When this submanifold has codimension two these will supply us with morphisms from elliptic to log divisors, and hence with Lie algebroid morphisms from the respective elliptic to log-tangent bundles by Proposition 3.15. After this we define the notion

of a boundary Lefschetz fibration, which is a Lefschetz-type fibration that can be interpreted as a Lie algebroid Lefschetz fibration between these Lie algebroids.

5.1. *The normal Hessian*

Let (X, D) be a pair, that is, $D \subseteq X$ is a submanifold of X .

DEFINITION 5.1. A map of pairs $f: (X, D) \rightarrow (Y, Z)$ is a map $f: X \rightarrow Y$ such that $f(D) \subseteq Z$. A strong map of pairs is a map of pairs $f: (X, D) \rightarrow (Y, Z)$ such that $f^{-1}(Z) = D$.

Given a map of pairs $f: (X, D) \rightarrow (Y, Z)$, note that $df(TD) \subseteq TZ$, so that df induces a map $\nu(df): ND \rightarrow NZ$ between normal bundles. When f is a strong map of pairs and f is transverse to Z , the map $\nu(df)$ is a fiberwise isomorphism.

Consider a map of pairs $f: (X, D) \rightarrow (Y, Z)$ such that $\nu(df): ND \rightarrow NZ$ is the zero map. Equivalently, one can assume that $\text{im } df \subset TZ$. Let z_1, \dots, z_ℓ be local defining functions for Z and consider their pullbacks $h_i := f^*(z_i)$ for $i = 1, \dots, \ell$. As $f(D) \subseteq Z$, the functions h_i vanish on D . Moreover, because $\nu(df)$ is the zero map, the derivatives dh_i vanish on D as well. Consequently, we can consider their Hessians $H(h_i): \text{Sym}^2(TX) \rightarrow \mathbb{R}$, which descend to maps $H(h_i): \text{Sym}^2(ND) \rightarrow \mathbb{R}$. As the differentials dz_i span N^*Z , these combine to give a map $H^\nu(f): \text{Sym}^2(ND) \rightarrow f^*(NZ)$, which one checks to be invariantly defined.

DEFINITION 5.2. Let $f: (X, D) \rightarrow (Y, Z)$ such that $\text{im } df \subset TZ$. The normal Hessian of f along D is the map $H^\nu(f): \text{Sym}^2(ND) \rightarrow f^*(NZ)$ over D .

When $\text{codim}(Z) = 1$, the normal Hessian $H^\nu(f)$ can be viewed as the matrix of second partial derivatives of the coordinate function of f normal to Z in directions normal to D .

5.2. *Boundary maps*

Let $f: (X, D) \rightarrow (Y, Z)$ be a strong map of pairs with Z a hypersurface and $\text{codim}(D) \geq 2$. Then f cannot be transverse to Z , as then $f^{-1}(Z) = D$ would be of codimension one. As Z is of codimension one, f being transverse to Z is equivalent to $\nu(df): ND \rightarrow NZ$ being nonzero, hence in this case $\nu(df): ND \rightarrow NZ$ is the zero map. Equivalently we have $\text{im } df \subset TZ$, so that the normal Hessian of f along D is well defined.

DEFINITION 5.3. Let $f: (X, D) \rightarrow (Y, Z)$ be a strong map of pairs, Z a hypersurface and $\text{codim}(D) \geq 2$. Then f is a boundary map if its normal Hessian $H^\nu(f)$ is definite along D .

As Z is a hypersurface, NZ is one dimensional. Because of this, $H^\nu(f)$ being definite makes sense, as locally it is a map $H^\nu(f): \text{Sym}^2(\mathbb{R}^d) \rightarrow \mathbb{R}$ where $d = \text{codim}(D)$. The choice of the name will become clearer after establishing some properties (see Remark 5.20). We will also call f a codimension- k boundary map if $\text{codim}(D) = k$, and sometimes implicitly assume that $\text{codim}(D) = 2$ when $\text{dim}(X) = 4$. Indeed, the main reason for introducing the notion of a boundary map comes from Proposition 5.6, where $\text{codim}(D) = 2$.

REMARK 5.4. Let $f: (X, D) \rightarrow (Y, Z)$ be a codimension- k boundary map and further let $f'': (X', D') \rightarrow (X, D)$ and $f': (Y, Z) \rightarrow (Y', Z')$ strong maps of pairs with f' and f'' transverse to Z' and D . Then $f'' \circ f \circ f': (X', D') \rightarrow (Y', Z')$ is a codimension- k boundary map.

REMARK 5.5. Assuming that $\text{codim}(D) \geq 2$ is only done to ensure that $\nu(df)$ is the zero map, as is required for the definition of the normal Hessian. If $\text{codim}(D) = 1$ yet this condition holds, it makes sense to talk about codimension-one boundary maps.

Codimension-two boundary maps naturally give rise to morphisms of divisors between elliptic and log divisors respectively, and hence to Lie algebroid morphisms by Proposition 3.15.

PROPOSITION 5.6. *Let $f: (X, D) \rightarrow (Y, Z)$ be a map of pairs with Z a hypersurface and $\text{codim}(D) = 2$. Then f is a boundary map if and only if $I_{|D|} := f^*I_Z$ is an elliptic ideal and f a morphism of divisors.*

In other words, a codimension-two boundary map uniquely specifies a compatible elliptic divisor structure on D .

Proof. Assume that f is a boundary map and consider $I_{|D|} = f^*I_Z$. Let z be a local defining function for Z so that locally $I_Z = \langle z \rangle$, and hence $I_{|D|} = \langle f^*(z) \rangle$. As f is a boundary map, $H^\nu(f)$ is definite, so that f^*z specifies the germ of a definite Morse–Bott function around D . By the discussion above Proposition 2.20 we see that $I_{|D|}$ is an elliptic ideal specifying an elliptic divisor structure on D by Proposition 2.10. The map f is a morphism of divisors by construction. Alternatively, one shows that for (L, s) the log divisor determined by Z , the pair $(R, q) := (f^*L, f^*s)$ is an elliptic divisor with $D_q = D$. The converse is similar, using again that f has definite normal Hessian if and only if $f^*(z)$ is locally Morse–Bott of even index around D , where z is a local defining function for Z . \square

COROLLARY 5.7. *Let $f: (X, D) \rightarrow (Y, Z)$ be a codimension-two boundary map. Then df induces a Lie algebroid morphism $(\varphi, f): TX(-\log |D|) \rightarrow TY(-\log Z)$ for the divisor structures of Proposition 5.6.*

The pointwise conclusion of the Morse–Bott lemma, Lemma 2.19, provides a local form for boundary maps around points in D .

PROPOSITION 5.8. *Let $f: (X^n, D^k) \rightarrow (Y^m, Z^{m-1})$ be a boundary map and $x \in D$. Then around x and $f(x) \in Z$ there exist coordinates (x_1, \dots, x_n) and (z, y_2, \dots, y_m) for which $\{x_1 = \dots = x_{n-k} = 0\} = D$ and $\{z = 0\} = Z$ such that for some map $g: \mathbb{R}^n \rightarrow \mathbb{R}^{k-1}$ we have $f(x_1, \dots, x_n) = (x_1^2 + \dots + x_{n-k}^2, g(x_1, \dots, x_n))$.*

Proof. Let $x \in D$ and z be a local defining function for Z around $f(x) \in Z$. As f is a boundary map, the proof of Proposition 5.6 shows that $f^*(z)$ is a local Morse–Bott function of index zero around x , after possibly replacing z by $-z$. By Lemma 2.19, after trivializing ND , there exist coordinates (x_1, \dots, x_n) of X around x with $\{x_1 = \dots = x_{n-k} = 0\} = D$ such that $f^*(z)(x_1, \dots, x_n) = x_1^2 + \dots + x_{n-k}^2$. Complete z to a coordinate system (z, y_2, \dots, y_m) of Y around $f(x)$. Then in these coordinates for some map $g: \mathbb{R}^n \rightarrow \mathbb{R}^{k-1}$ we have

$$f(x_1, \dots, x_n) = (f^*(z)(x_1, \dots, x_n), g(x_1, \dots, x_n)) = (x_1^2 + \dots + x_{n-k}^2, g(x_1, \dots, x_n)). \quad \square$$

Using either Proposition 5.8 or the proof Proposition 5.6, the Lie algebroid morphisms $(\varphi, f): TX(-\log |D|) \rightarrow TY(-\log Z)$ obtained from boundary maps $f: (X, D) \rightarrow (Y, Z)$ using Corollary 5.7 have the following extra property: for any k -form $\alpha \in \Omega^k(\log Z)$, we have $\text{Res}_\theta(\varphi^*\alpha) = 0$, because $\text{Res}_\theta(d \log f^*(z)) = 0$ for any local defining function z for Z .

5.3. Fibrating boundary maps

We next introduce specific boundary maps by demanding submersiveness on D .

DEFINITION 5.9. A *fibrating boundary map* is a boundary map $f: (X, D) \rightarrow (Y, Z)$ such that $f|_D: D \rightarrow Z$ is submersive.

Note that it is not required that D surjects onto Z . For fibrating boundary maps we can improve upon Proposition 5.8, obtaining again a local form around points in D .

PROPOSITION 5.10. *Let $f: (X^n, D^k) \rightarrow (Y^m, Z^{m-1})$ be a fibrating boundary map and z be a (local) defining function for Z . Then there exist coordinates (x_1, \dots, x_n) and $(\pm z, y_2, \dots, y_m)$ with $\{x_1 = \dots = x_{n-k} = 0\} = D$ such that $f(x_1, \dots, x_n) = (x_1^2 + \dots + x_{n-k}^2, x_{n-m+1}, \dots, x_n)$.*

In other words, we can simultaneously put both the components of f normal and tangent to D in standard form, and obtain a commuting diagram of normal bundles near points $x \in D$ and $f(x) \in Z$, viewing f as giving a map $f: ND \rightarrow NZ$ using a tubular neighborhood:

$$\begin{array}{ccc} ND & \xrightarrow{f} & NZ \\ \text{pr}_D \downarrow & & \downarrow \text{pr}_Z \\ D & \xrightarrow{f|_D} & Z \end{array}$$

In fact, the condition that f is a fibrating boundary map is equivalent to the existence of coordinates as in Proposition 5.10 around each point $x \in D$. Call a finite collection of functions an *independent set* at p if their differentials are linearly independent at p . By the implicit function theorem, an independent set can be completed to a coordinate system in a neighborhood of p . Independence is preserved under pulling back along a submersion.

Proof. Choose a tubular neighborhood embedding $\Phi: NZ \rightarrow V \subset Y$ for $\text{pr}_Z: NZ \rightarrow Z$ and let z be a local defining function for Z on an open subset $V' \subset V$ of $f(x)$. Let $U := f^{-1}(V') \subset X$. Choose coordinate functions $y_2, \dots, y_m: V' \rightarrow \mathbb{R}$ for Z . Then $\{y_2, \dots, y_m\}$ forms an independent set on Z , and because pr_Z is a submersion, the same is true for $\{z, \text{pr}_Z^*(y_2), \dots, \text{pr}_Z^*(y_m)\}$ on Y . By Proposition 5.8, after possibly shrinking U and V' and changing z to $-z$, there are tubular neighborhood coordinates (x_1, \dots, x_n) such that $f^*(z) = x_1^2 + \dots + x_{n-k}^2$. Consider $\{x_1, \dots, x_{n-k}, f^* \text{pr}_Z^*(y_j)\}$, which is an independent set on X , using submersiveness of $f|_D$. Complete this to a coordinate system $\{x_1, \dots, x_{n-m+1}, f^* \text{pr}_Z^*(y_j)\}$ on X , and relabel to (x_1, \dots, x_n) . Using these coordinates on X and the coordinates $(z, \text{pr}_Z^*(y_j))$ on Y , the map f is given by $f(x_1, \dots, x_n) = (x_1^2 + \dots + x_{n-k}^2, x_{n-m+1}, \dots, x_n)$ as desired. \square

The normal form result of Proposition 5.10 immediately implies the following.

COROLLARY 5.11. *Let $f: (X, D) \rightarrow (Y, Z)$ be a fibrating boundary map. Then f is submersive in a punctured neighborhood around D .*

Consequently, fibrating boundary maps have well-defined fibers of dimension $\dim(X) - \dim(Y)$ near D , and of dimension $\dim(X) - \dim(Y) - \text{codim}(D) + 1$ on D . In particular, when f is a fibrating boundary map, D will be a fiber bundle over certain components of Z .

REMARK 5.12. An alternative way of viewing the proof of Corollary 5.11 when $\text{codim}(D) = 2$ uses Proposition 5.6. Namely, let $f: (X, D) \rightarrow (Y, Z)$ give rise to a Lie algebroid morphism $(\varphi, f): TX(-\log |D|) \rightarrow TY(-\log Z)$. While $df: ND \rightarrow NZ$ is the zero map, $\varphi|_D$ is surjective because f is fibrating. This is an open condition, so that φ is surjective in a neighborhood around D . On $X \setminus D$, the Lie algebroid $TX(-\log |D|)$ is isomorphic to TX , hence f is submersive there. We see that fibrating boundary maps lead to Lie algebroid morphisms which are fibrations near D .

We proceed to obtain similar normal bundle commutativity around components of D whose image is coorientable. To prove this, we use a result by Bursztyn–Lima–Meinrenken [5] on normal bundle embeddings, which we now describe. Let $M \subseteq X$ be a submanifold and NM its normal bundle. Denote by $\mathcal{E}_M \in \mathfrak{X}(NM)$ the associated Euler vector field and given $V \in \mathfrak{X}(X)$ tangent to M , let $\nu(V) \in \mathfrak{X}(NM)$ be its linear approximation.

DEFINITION 5.13. Let $M \subseteq X$ be a submanifold. A vector field $V \in \mathfrak{X}(X)$ is *Euler-like* along M if it is complete, and satisfies $V|_M = 0$ and $\nu(V) = \mathcal{E}_M$.

A *strong tubular neighborhood embedding* for $M \subseteq X$ is an embedding $\Phi: NM \rightarrow X$ taking the zero section of NM to M , and such that the linear approximation $\nu(\Phi): \nu(NM, M) \rightarrow \nu(X, M)$ is the identity map. The following proposition says that Euler-like vector fields give rise to strong tubular neighborhood embeddings.

PROPOSITION 5.14 [5, Proposition 2.6]. Let $M \subseteq X$ be a submanifold and $V \in \mathfrak{X}(X)$ Euler-like along M . Then there exists a unique strong tubular neighborhood embedding $\Phi: NM \rightarrow X$ such that $\Phi_*(\mathcal{E}_M) = V$.

We use this result to construct compatible tubular neighborhood embeddings for fibrating boundary maps around components of D whose image is coorientable.

PROPOSITION 5.15. Let $f: (X, D) \rightarrow (Y, Z)$ be a fibrating boundary map and $D_j \subseteq D$ a connected component such that $f(D_j) =: Z_j \subset Z$ is coorientable. Then there exist a defining function z for Z_j and tubular neighborhood embeddings $\Phi_{D_j}: \tilde{U} \rightarrow U \subset X$ for D_j and $\Phi_{Z_j}: \tilde{V} \rightarrow V \subset Y$ for Z_j such that $\Phi_{D_j}^* f^*(z) = Q_{f^*(z)} \in \Gamma(D_j; \text{Sym}^2 N^* D_j)$ and the following diagram commutes:

$$\begin{array}{ccccccc}
 \tilde{U} & \xrightarrow{\Phi_{D_j}} & U & \xrightarrow{f} & V & \xleftarrow{\Phi_{Z_j}} & \tilde{V} \\
 \text{pr}_{D_j} \downarrow & & \downarrow & & \downarrow & & \downarrow \text{pr}_{Z_j} \\
 D_j & \xrightarrow{\text{id}} & D_j & \xrightarrow{f|_D} & Z_j & \xleftarrow{\text{id}} & Z_j
 \end{array}$$

Proof. Let $z: V \rightarrow \mathbb{R}$ be a defining function for Z_j and let $U' \subset f^{-1}(V)$ be the connected component containing D_j . Using Lemma 2.19 applied to $f^*(z)$, shrink U' so that $U' = \Phi_{D_j}(\tilde{U}')$ for some tubular neighborhood embedding $\Phi_{D_j}: \tilde{U}' \rightarrow U'$ of D_j . Choose a tubular neighborhood embedding $\Phi_{Z_j}: \tilde{V} \rightarrow V$ for Z_j . For $x \in D_j$, use Proposition 5.10 (possibly changing z to $-z$) to obtain an open $U_x \subset U'$ containing x and coordinates (x_1, \dots, x_n) so that $f^*(z) = x_1^2 + \dots + x_{n-k}^2$. Note that U' is connected so that $f^*(z)$ has a fixed sign. Consider $U := \cup_{x \in D_j} U_x \subset U'$ and extract a finite subcover $\{U_\alpha\}_{\alpha \in I}$. On each set U_α , define the vector field $\mathcal{E}_\alpha := x_1 \partial_{x_1} + \dots + x_{n-k} \partial_{x_{n-k}}$. It satisfies $\mathcal{L}_{\mathcal{E}_\alpha} f^*(z) = 2f^*(z)$, and $f_* \Phi_{\alpha*} \mathcal{E}_\alpha = z \partial_z = \mathcal{E}_{Z_j}$. Let $\{\psi_\alpha\}_{\alpha \in I}$ be a partition of unity subordinate to $\{U_\alpha\}_{\alpha \in I}$ and define $\mathcal{E} := \sum_{\alpha \in I} \psi_\alpha \mathcal{E}_\alpha$ on U' . Then $\mathcal{E}|_{D_j} = \sum_{\alpha \in I} \psi_\alpha \mathcal{E}_\alpha|_{D_j} = \sum_{\alpha \in I} \psi_\alpha \cdot 0 = 0$, and $\nu(\mathcal{E}) = \sum_{\alpha \in I} \psi_\alpha \nu(\mathcal{E}_\alpha) = \sum_{\alpha \in I} \psi_\alpha \mathcal{E}_{D_j} = \mathcal{E}_{D_j}$. Ensure that \mathcal{E} is complete by multiplying by a bump function and shrinking U' , so that $\mathcal{E} \in \mathfrak{X}(U)$ is Euler-like along D_j . Moreover, we now have that

$$\mathcal{L}_{\mathcal{E}} f^*(z) = \sum_{\alpha \in I} \mathcal{L}_{\psi_\alpha \mathcal{E}_\alpha} f^*(z) = \sum_{\alpha \in I} \psi_\alpha \mathcal{L}_{\mathcal{E}_\alpha} f^*(z) = \sum_{\alpha \in I} \psi_\alpha 2f^*(z) = 2f^*(z),$$

and similarly

$$f_*\mathcal{E} = \sum_{\alpha \in I} f_*(\psi_\alpha \mathcal{E}_\alpha) = \sum_{\alpha \in I} \psi_\alpha \mathcal{E}_Z = \mathcal{E}_Z.$$

Use Proposition 5.14 and possibly shrink U to obtain a tubular neighborhood embedding $\Phi'_{D_j} : \tilde{U} \rightarrow U$ such that $\Phi_*\mathcal{E}_{D_j} = \mathcal{E}$, $f_*\mathcal{E} = \mathcal{E}_Z$, and $\mathcal{L}_\mathcal{E}f^*(z) = 2f^*(z)$. This embedding satisfies all desired properties. Moreover, upon pulling back $f^*(z)$ to \tilde{U} we have

$$\Phi'_{D_j} f^*(z) = Q_{f^*(z)} \in \Gamma(D_j; \text{Sym}^2(N^*D_j)),$$

as $g := \Phi'_{D_j} f^*(z)$ is smooth and satisfies $\mathcal{L}_\mathcal{E}g = 2g$. □

As a consequence of the above proposition, when Z is coorientable this result implies a normal form for f around any point in Z and its entire inverse image, as then a semi-global defining function for Z can be used. We can summarize the above discussion as follows.

COROLLARY 5.16. *Let $f : (X^{2n}, D^{2n-2}) \rightarrow (\Sigma^2, Z^1)$ be a fibrating boundary map for which Z is coorientable. Then there are*

- neighborhoods U of D and V of Z and diffeomorphisms between these sets and neighborhoods of the zero sections of the corresponding normal bundles, $\Phi_D : U \rightarrow ND$ and $\Phi_Z : V \rightarrow \mathbb{R} \times Z$, and
- a bundle metric g on ND ,

such that the following diagram commutes, where $\text{pr}_D : ND \rightarrow D$ is the bundle projection:

$$\begin{array}{ccc} U & \xrightarrow{f} & V \\ \Phi_D \downarrow & & \downarrow \Phi_Z \\ ND & \xrightarrow{(\|\cdot\|_g^2, f|_D \circ \text{pr}_D)} & \mathbb{R} \times Z \end{array}$$

We can now obtain topological information about the generic fibers of f near D . The following result is the main reason for wanting compatible tubular neighborhood embeddings.

PROPOSITION 5.17. *Let $f : (X^n, D^k) \rightarrow (Y^m, Z^{m-1})$ be a fibrating boundary map with Z coorientable. Denote the fiber of $f|_D : D \rightarrow Z$ by F_D^{k-m+1} . Then the fiber F^{n-m} of f near D is an S^{n-k-1} -sphere bundle over F_D .*

Proof. Apply Corollary 5.16 using a semi-global defining function for Z , setting $\tilde{U} = \Phi_D(U)$ and similarly for \tilde{V} . Let $y \in \tilde{V} \setminus Z$. As f is submersive on $\tilde{U} \setminus D$, consider $F_y = \Phi_D \circ f^{-1} \circ \Phi_Z^{-1}(y)$, which is of dimension $n - m$. Consider $\text{pr}_Z(y) \in Z$ and its $(k - m + 1)$ -dimensional fiber $F_{D, \text{pr}_Z(y)} = f|_D^{-1}(\text{pr}_Z(y))$. Because the tubular neighborhood embeddings are compatible with f , we have $F_y = \text{pr}_D^{-1}(F_{D, \text{pr}_Z(y)})$, noting that pr_D is submersive. As a point in $\tilde{V} \subset NZ$ is given by a point in Z together with a distance, the fiber of $\text{pr}_D : F_y \rightarrow F_{D, \text{pr}_Z(y)}$ is given by a sphere, consisting of all points with fixed distance above the corresponding point in D . □

COROLLARY 5.18. *Let $f : (X^4, D^2) \rightarrow (Y^2, Z^1)$ be a codimension-two fibrating boundary map with Z coorientable. Then the connected components of the generic fibers of f near D are tori or Klein bottles. If either X is orientable or D is coorientable, only tori can occur.*

Proof. This follows from Proposition 5.17 using $n = 4$ and $k = m = 2$ by a dimension count. The generic fibers F of f near D satisfy $\dim(F) = 4 - 2 = 2$, and are S^1 -bundles over fibers of $f|_D$, which in turn are one dimensional. Thus F is a union of tori and Klein bottles.

Whether the fibers are tori or Klein bottles is determined by their orientability. Assuming the fibers of the fibrating boundary map near D are connected, let $D_j \subseteq D$ be a connected component and $y \in Z_j = f(D_j)$. Consider the pairing $\langle w_1(ND_j), [f^{-1}(y)] \rangle$, which determines whether the fiber F is the trivial S^1 -bundle over F_D or not. We see that if D_j is coorientable, then $w_1(ND_j) = 0$ and the fiber F near D_j is a torus. If X is orientable, then F is orientable, as there is a splitting $TX|_F \cong TF \oplus f^*TY|_F$ and the latter factor is trivial on F . This means that F inherits an orientation from X , so that the fiber F must again be a torus. \square

Recall that when Z is separating it admits a global defining function $z \in C^\infty(Y)$. In this case, any boundary map f to (Y, Z) must map onto one component of Y with respect to Z . More precisely, given such a generator $z \in I_Z$, consider $Y_+ := z^{-1}([0, \infty)) \subset Y$, a manifold with boundary given by $Z = z^{-1}(0)$. Then we have the following.

PROPOSITION 5.19. *Let $f: (X, D) \rightarrow (Y, Z)$ be a boundary map with X connected, and assume that Z is separating. Then there exists a global defining function z for Z so that $f(X) \subset Y_+$, and f defines a boundary map $f: (X, D) \rightarrow (Y', \partial Y')$, where Y' consists of the connected component of Y_+ which contains points in the image of the map f .*

Proof. Let z be a defining function for Z . Then $f^*(z)$ is globally defined on X , with zero set $D = f^{-1}(Z)$. As D has codimension at least two in X , its complement $X \setminus D$ is connected, so that $f^*(z)$ has fixed sign on $X \setminus D$. Consequently, by changing z to $-z$ if necessary, the function $f^*(z)$ is nonnegative. But then for all points $x \in X$, $z(f(x)) = f^*(z)(x) \geq 0$, so that $f(X) \subset Y_+$. Moreover, f is still a boundary map when restricting its codomain. \square

REMARK 5.20. The previous proposition explains the name ‘boundary map’, as the defining condition specifies the behavior of f normal to Z , the boundary of its restricted codomain.

5.4. Boundary (Lefschetz) fibrations

We introduce further submersiveness assumptions.

DEFINITION 5.21. A *boundary fibration* is a fibrating boundary map $f: (X, D) \rightarrow (Y, Z)$ such that $f|_{X \setminus D}: X \setminus D \rightarrow Y \setminus Z$ is a fibration.

DEFINITION 5.22. A *boundary Lefschetz fibration* is a fibrating boundary map $f: (X^{2n}, D) \rightarrow (\Sigma^2, Z)$ such that $f|_{X \setminus D}: X \setminus D \rightarrow \Sigma \setminus Z$ is a Lefschetz fibration.

The following is immediate from Corollary 5.7 together with Remark 5.12.

COROLLARY 5.23. *Let $f: (X, D) \rightarrow (Y, Z)$ be a codimension-two boundary (Lefschetz) fibration. Then f induces a Lie algebroid (Lefschetz) fibration $(\varphi, f): TX(-\log|D|) \rightarrow TY(-\log Z)$.*

The statement that f induces a Lie algebroid morphism is to be interpreted as in Proposition 3.15, namely that there is a Lie algebroid morphism (φ, f) such that $\varphi = df$ on sections. Moreover, the elliptic divisor structure on D is the one induced from f and Z . By adapting the usual argument, we can ensure that boundary (Lefschetz) fibrations have connected fibers.

PROPOSITION 5.24. *Let $f: (X, D) \rightarrow (Y, Z)$ be a codimension-two boundary (Lefschetz) fibration, and suppose that Z is separating. Define $(Y', \partial Y')$ as in Proposition 5.19. Then there exists a cover $g: \tilde{Y}' \rightarrow Y'$ of Y' and a boundary (Lefschetz) fibration $\tilde{f}: (X, D) \rightarrow (\tilde{Y}', \partial \tilde{Y}')$ with connected generic fibers such that $f = g \circ \tilde{f}$.*

Proof. As f is a global boundary map, by Proposition 5.19 we obtain a boundary map $f: (X, D) \rightarrow (Y', \partial Y')$ which immediately is also a boundary (Lefschetz) fibration. By definition, $f: X \setminus D \rightarrow Y' \setminus \partial Y'$ is a (Lefschetz) fibration. Consequently, denoting its generic fiber by F , there is a sequence in homotopy $\pi_1(F) \rightarrow \pi_1(X \setminus D) \rightarrow \pi_1(Y' \setminus \partial Y') \rightarrow \pi_0(F) \rightarrow 0$ [13, Proposition 8.1.9]. Applying Van Kampen’s theorem for each connected component of D shows that $\pi_1(X \setminus D)$ surjects onto $\pi_1(X)$ via the natural inclusion. Note that we have $\pi_1(Y' \setminus \partial Y') \cong \pi_1(Y')$. We obtain the following commutative diagram.

$$\begin{array}{ccccccc} \pi_1(F) & \longrightarrow & \pi_1(X \setminus D) & \longrightarrow & \pi_1(Y' \setminus \partial Y') & \longrightarrow & \pi_0(F) \longrightarrow 0 \\ & & \downarrow & & \cong \downarrow & & \\ & & \pi_1(X) & \longrightarrow & \pi_1(Y') & & \end{array}$$

The generic fiber F of $f: X \setminus D \rightarrow Y' \setminus \partial Y'$ is compact, hence $\pi_0(F)$ is finite. But then $f_*(\pi_1(X \setminus D)) \subset \pi_1(Y' \setminus \partial Y')$ is a subgroup of finite index. Making use of the isomorphism $\pi_1(Y' \setminus \partial Y') \cong \pi_1(Y')$, let G denote the corresponding finite index subgroup inside $\pi_1(Y')$, and let $g: \tilde{Y}' \rightarrow Y'$ be the associated cover. Then $f: X \rightarrow Y'$ lifts to $\tilde{f}: X \rightarrow \tilde{Y}'$ if and only if $f_*(\pi_1(X)) \subset G$, but this is an equality because the map $\pi_1(X \setminus D) \rightarrow \pi_1(X)$ surjects. For \tilde{f} we have that its restriction $\tilde{f}: X \setminus D \rightarrow \tilde{Y}' \setminus \partial \tilde{Y}'$ induces a surjection $\tilde{f}_*: \pi_1(X \setminus D) \rightarrow \pi_1(\tilde{Y}' \setminus \partial \tilde{Y}')$. But then $\pi_0(\tilde{F})$ is trivial, so that the generic fibers \tilde{F} of \tilde{f} are connected. \square

By the above we can replace a boundary Lefschetz fibration on (X, D) with Z separating by one for which the generic fibers in $X \setminus D$ are connected. We next study the fibrating case, concluding that the fibers of $f|_D: D \rightarrow Z$ are connected if those of f near D are.

PROPOSITION 5.25. *Let $f: (X, D) \rightarrow (Y, Z)$ be a fibrating boundary map whose generic fibers near D are connected, and suppose that Z is separating. Then the fibers of $f|_D: D \rightarrow Z$ are connected.*

Proof. Using Proposition 5.19, replace (Y, Z) by $(Y', \partial Y')$, where Y' is connected, and let $V \subset Y'$ be a connected open neighborhood of a point $y \in Z'$ so that $V \setminus V \cap Z'$ is connected, and $f|_{f^{-1}(V) \setminus D}$ is a fibration by Corollary 5.11. Let F be the generic fiber of f near D and denote $F_y := f^{-1}(y) = f|_D^{-1}(y)$. Let $F_{y,i}$ be the connected components of F_y , and let U_i be disjoint opens around $F_{y,i}$ in X . Note that for each i , the set $f(U_i) \subset Y'$ contains y in its interior. In fact, it follows from the normal form for f around D that if $x \in F_y$ and U is a neighborhood of x , then $f(U)$ contains y in its interior.

Choose a curve $\gamma: [0, 1] \rightarrow Y'$ such that $\gamma([0, 1)) \in V \setminus Z'$, $f(1) = y$ and γ is transverse to Z' . Then $M := f^{-1}(\gamma([0, 1])) \subset f^{-1}(V)$ is a compact submanifold of X with boundary equal to $f^{-1}(0)$. Choose a sequence of points $(y_n)_{n \in \mathbb{N}} \subseteq \gamma([0, 1])$ converging to y such that $y_n \neq y$ for all $n \in \mathbb{N}$. For each n , consider the set $W_n := F_{y_n} \setminus \bigsqcup_i (U_i \cap F_{y_n})$. As each intersection $U_i \cap M$ is open in M , the sets $U_i \cap F_{y_n}$ are disjoint opens in F_{y_n} .

Assume that F_y is not connected. Then for n large enough, there will be at least two indices i_1 and i_2 for which $U_{i_j} \cap F_{y_n} \neq \emptyset$. This is because $f(U_{i_1}) \cap f(U_{i_2})$ has y in its interior. As F_{y_n} is connected, we conclude that $W_n \neq \emptyset$ for n large enough, as F_{y_n} cannot be covered by disjoint opens. For each such n , let $x_n \in W_n$ be some element. By compactness of M , the

sequence $\{x_n\}_{n \in \mathbb{N}}$ has a convergent subsequence, so that after relabeling we have $x_n \rightarrow x$ for some $x \in M$. By definition $x \in M \setminus (\bigcup_i U_i)$, so that $x \notin U_i$ for all i , hence $x \notin F_y$. However, by continuity we have $f(x_n) = y_n \rightarrow y = f(x)$, which is a contradiction. We conclude that F_y is connected, so that the fibers of $f|_D: D \rightarrow Z'$ are connected. \square

6. Constructing boundary Lefschetz fibrations

In this section we discuss how to obtain four-dimensional boundary Lefschetz fibrations. In particular, we construct them out of genus one Lefschetz fibrations $f: X^4 \rightarrow \Sigma^2$ by surgery, replacing neighborhoods of fibers of points $x \in \Sigma$ by a certain standard boundary map.

Given $n \in \mathbb{Z}$, let $p: L_n \rightarrow T^2$ be the complex line bundle over T^2 with first Chern class equal to $n \in H^2(T^2; \mathbb{Z})$. Choose a Hermitian metric on L_n , which provides a fiberwise radial distance $r: \text{Tot}(L_n) \rightarrow \mathbb{R}_{\geq 0}$. Further, let $\text{pr}: T^2 \rightarrow S^1$ be the projection onto the first factor.

DEFINITION 6.1. The total space $\text{Tot}(L_n)$ together with the map $f: \text{Tot}(L_n) \rightarrow S^1 \times \mathbb{R}_{\geq 0}$ given by $f(x) := (\text{pr}(p(x)), r^2(x))$ for $x \in \text{Tot}(L_n)$, is called the *standard n -model*.

The first thing to note is that f as defined above is an example of a boundary fibration.

PROPOSITION 6.2. The map $f: (\text{Tot}(L_n), T^2) \rightarrow (S^1 \times \mathbb{R}_{\geq 0}, S^1 \times \{0\})$ is a boundary fibration, where T^2 is identified with the zero section in $\text{Tot}(L_n)$.

Proof. Certainly f is a strong map of pairs, and the codimension of $D = T^2$ inside $\text{Tot}(L_n)$ is (at least) two. The normal Hessian $H^\nu(f)$ of f along D is given by the quadratic form with constant matrix given by $2 \cdot \text{Id}$ in an appropriate orthonormal frame normal to D given by (x, y) with $x^2 + y^2 = r^2$, which is clearly definite. We conclude that f is a boundary map. Both the bundle projection $p: L_n \rightarrow T^2$ and the projection $\text{pr}: T^2 \rightarrow S^1$ are submersive, so that $f|_D: T^2 \rightarrow S^1 \times \{0\}$ is submersive, making f a fibrating boundary map. It is immediate that f is submersive when $r \neq 0$, that is, on $\text{Tot}(L_n) \setminus T^2$, so that f is a boundary fibration. \square

Let $f: X \rightarrow Y$ be a smooth map that is a fibration in a neighborhood of an embedded oriented circle $\gamma \subset Y$. Recall that the *monodromy* of f around γ is the element in the mapping class group of the fiber F corresponding to the mapping torus $f|_{f^{-1}(\gamma)}: f^{-1}(\gamma) \rightarrow \gamma \cong S^1$.

For the standard n -model, let $\varepsilon > 0$ be small and let $\gamma := S^1 \times \{\varepsilon\} \subset S^1 \times \mathbb{R}_{\geq 0}$ with its standard orientation. We compute the monodromy of $f: \text{Tot}(L_n) \rightarrow S^1 \times \mathbb{R}_{\geq 0}$ around γ .

PROPOSITION 6.3. The monodromy of f around γ is the $-n$ th power of a Dehn twist.

Proof. Let $M := f^{-1}(\gamma)$ be the T^2 -bundle $f: M \rightarrow \gamma$. To compute the monodromy of f around γ , note that using $p: M \rightarrow T^2$ we can view M as the principal ε -sphere bundle of L_n . The Hermitian metric gives rise to a connection $i\theta \in \Omega^1(M; \mathbb{R})$. As $L_n \rightarrow T^2$ has Euler class equal to n , its curvature equals $\frac{d\theta}{2\pi i} = nda \wedge db$, where da and db are generators of $H^1(T^2)$. Recall now that circle bundles are classified by their Euler class due to the Gysin sequence. Consider $M' := \mathbb{R}^3/\Gamma$, with $\Gamma = \langle \alpha_1, \alpha_2, \alpha_3 \rangle$ the integral lattice generated by the following group elements acting on \mathbb{R}^3 :

$$\alpha_1(x, y, z) = (x, y + 1, z), \quad \alpha_2(x, y, z) = (x, y, z + 1), \quad \alpha_3(x, y, z) = (x + 1, y, z - ny).$$

The projection $\text{pr}_{12}: \mathbb{R}^3 \rightarrow \mathbb{R}^2$ given by $(x, y, z) \mapsto (x, y)$ descends to a map $g: \mathbb{R}^3/\Gamma \rightarrow \mathbb{R}^2/\mathbb{Z}^2$. This is an S^1 -bundle over T^2 , with invariant one-forms $e_1 = dx$ and $e_2 = dy$, and connection one-form $e_3 = dz + nxdy$. As $de_3 = ne_1 \wedge e_2$, we conclude that \mathbb{R}^3/Γ has Euler class equal to n , so that $\mathbb{R}^3/\Gamma \cong M$, as circle bundles over T^2 . Note that the map $\text{pr}_1: \mathbb{R}^3 \rightarrow \mathbb{R}$ given by

$(x, y, z) \mapsto x$ also descends and exhibits M' as a T^2 -bundle over $\mathbb{R}/\mathbb{Z} \cong S^1$. To obtain a T^2 -bundle over S^1 out of $p: M \rightarrow T^2$, the choice of any projection $T^2 \rightarrow S^1$ gives isomorphic bundles. Hence we can assume that the projection pr_1 makes the following diagram commute.

$$\begin{array}{ccccc} M & \xrightarrow{p} & T^2 & \xrightarrow{\text{pr}} & S^1 \\ \cong \downarrow & & \cong \downarrow & & \cong \downarrow \\ M' & \xrightarrow{\text{pr}_{12}} & T^2 & \xrightarrow{\text{pr}_1} & S^1 \end{array}$$

We can thus compute the monodromy of f around γ by considering $\text{pr}_1: M' \rightarrow S^1$. Due to our concrete description it is immediate that

$$M' = T^2 \times S^1 / (x, 0) \sim (f(x), 1), \quad f \in \text{Aut}(T^2) \text{ with } f_* = \begin{pmatrix} 1 & -n \\ 0 & 1 \end{pmatrix}: H_2(T^2) \rightarrow H_2(T^2).$$

We conclude that M has monodromy equal to the $-n$ th power of a Dehn twist as desired. \square

We next describe the surgery process whereby we replace suitable neighborhoods of points by the above standard models.

DEFINITION 6.4. A *punctured surface* is an open surface Σ obtained from a closed surface by removing a finite number of discs.

A punctured surface naturally has a compact closure $\bar{\Sigma}$ by adding the circle boundaries of the removed discs. This closure comes with a natural isomorphism $j: \Sigma \xrightarrow{\cong} \bar{\Sigma} \setminus \partial\bar{\Sigma}$.

PROPOSITION 6.5. Let $f: X^4 \rightarrow \Sigma^2$ be a proper map over a punctured surface such that f is a genus one fibration in a neighborhood of $\partial\Sigma$. Assume that the monodromy of f around every boundary component is a power of a Dehn twist. Then there exists a compact elliptic pair $(\tilde{X}, |D|)$ with $i: X \xrightarrow{\cong} \tilde{X} \setminus D$ and a fibrating boundary map $\tilde{f}: (\tilde{X}, |D|) \rightarrow (\bar{\Sigma}, \partial\bar{\Sigma})$ such that the following diagram commutes:

$$\begin{array}{ccc} X & \xleftarrow{i} & (\tilde{X}, |D|) \\ f \downarrow & & \downarrow \tilde{f} \\ \Sigma & \xleftarrow{j} & (\bar{\Sigma}, \partial\bar{\Sigma}) \end{array}$$

In the above situation, we say $f: X \rightarrow \Sigma$ can be completed to $\tilde{f}: (\tilde{X}, |D|) \rightarrow (\bar{\Sigma}, \partial\bar{\Sigma})$. These completions are unique when the monodromy is not trivial.

Proof. At each boundary end E_i of Σ , glue in the standard n -model $\text{Tot}(L_{n_i})$, where $n_i \in \mathbb{Z}$ is such that $\text{Mon}(\gamma_i) = \delta^{n_i}$, with $\gamma_i = \partial E_i$. Here δ denotes a Dehn twist. As the monodromies agree, the fibrations are isomorphic, hence can be glued together to a new fibration.

It only remains to argue that if the monodromy is nontrivial, the completion is unique. Since the completion of each end of X is obtained by gluing L_n to the end of X , by identifying M , the ε -sphere bundle of L_n , with the preimage of a loop around a boundary component of Σ , we see that any other completion can be obtained by precomposing the gluing map by a diffeomorphism of M . However, due to a result of Waldhausen [34, Theorem 5.5] any such diffeomorphism is isotopic to a fiber-preserving diffeomorphism (covering a diffeomorphism of the base) and therefore extends to the ε -disc bundle of L_n . Hence the completion of the end is unique up to diffeomorphism. \square

COROLLARY 6.6. *Let $f: X^4 \rightarrow \Sigma^2$ be a genus one Lefschetz fibration over a punctured surface. Assume that the monodromy of f around every boundary component is a power of a Dehn twist. Then f can be completed to a boundary Lefschetz fibration $\tilde{f}: (\tilde{X}, |D|) \rightarrow (\tilde{\Sigma}, \partial\tilde{\Sigma})$.*

Proof. The Lefschetz singularities lie in the interior of Σ so that f is a genus one fibration near $\partial\Sigma$. After completing using Proposition 6.5, \tilde{f} is a fibrating boundary map and a genus one Lefschetz fibration in the interior, so that it is a boundary Lefschetz fibration. \square

Consequently, given a genus one Lefschetz fibration on X , removing several discs and their inverse image and then completing as in Corollary 6.6 gives a boundary Lefschetz fibration on \tilde{X} . Homological essentialness of generic fibers is preserved by the completion process.

PROPOSITION 6.7. *Let $\tilde{f}: X^4 \rightarrow \Sigma^2$ be a genus one Lefschetz fibration with boundary Lefschetz completion $\tilde{f}: (\tilde{X}, |D|) \rightarrow (\tilde{\Sigma}, \partial\tilde{\Sigma})$. Then the generic fibers of f are homologically essential if and only if those of \tilde{f} are.*

A codimension-two boundary Lefschetz fibration on a four-manifold X has singular fibers equal to the Euler characteristic of X , as is true for genus one Lefschetz fibrations.

PROPOSITION 6.8. *Let $f: (X^4, D) \rightarrow (\Sigma^2, Z)$ be a codimension-two boundary Lefschetz fibration. Then we have $\chi(X) = \mu$, with μ the number of singular fibers of $f|_{X \setminus D}$.*

Proof. Given two open sets $U, V \subseteq X$ we have $\chi(U \cup V) = \chi(U) + \chi(V) - \chi(U \cap V)$. Let $U := X \setminus D$ and take $V \subseteq ND$ to be a tubular neighborhood of D . Then V is homotopy equivalent to D , which is a union of tori, so that $\chi(V) = \chi(D) = 0$. Moreover, $U \cap V$ is deformation equivalent to S^1ND , which is a principal circle bundle over D , hence $\chi(U \cap V) = 0$ as well. Recall that if a manifold M admits a genus g Lefschetz fibration over a surface of genus h , it satisfies $\chi(M) = (2 - 2h)(2 - 2g) + \mu$, where μ is the number of singular fibers. Applying this to $U = X \setminus D$ with $g = 1$ we obtain that $\chi(X) = \chi(X \setminus D) = \mu$. \square

7. Constructing stable generalized complex structures

In this section we combine the results from previous sections to construct stable generalized complex structures using codimension-two boundary fibrations with two-dimensional fibers, and from codimension-two boundary Lefschetz fibrations in dimension 4. For notational convenience, we call a boundary map $f: (X^{2n}, D) \rightarrow (Y^{2n-2}, Z)$ *homologically essential* if the (generic) two-dimensional fibers F (near D) specify nonzero homology classes $[F] \neq 0 \in H_2(X \setminus D; \mathbb{R})$. We prove the following results stated in the introduction.

THEOREM 7.1. *Let $f: (X^4, D^2) \rightarrow (\Sigma^2, Z^1)$ be a homologically essential boundary Lefschetz fibration between compact oriented manifolds with D coorientable and $[Z] = 0 \in H_1(\Sigma; \mathbb{Z}_2)$. Then (X, D) admits a stable generalized complex structure.*

THEOREM 7.2. *Let $f: (X^{2n}, D^{2n-2}) \rightarrow (Y^{2n-2}, Z^{2n-3})$ be a homologically essential boundary fibration between compact oriented manifolds with D coorientable such that (Y, Z) admits a log-symplectic structure. Then (X, D) admits a stable generalized complex structure.*

In the remainder of this section, any mention of an elliptic divisor structure on (X, D) , or the Lie algebroid $TX(-\log |D|)$, will refer to the structure induced by a codimension-two boundary map via Proposition 5.6. Similarly, we will without further mention have codimension-two

boundary maps induce Lie algebroid morphisms between the appropriate elliptic and log-tangent bundles using Corollary 5.23. Before proving the above two results, we first establish that we can create a suitable closed Lie algebroid two-form on X .

PROPOSITION 7.3. *Let $f: (X^{2n}, D^{2n-2}) \rightarrow (Y^{2n-2}, Z^{2n-3})$ be a homologically essential fibrating boundary map with $f(D)$ coorientable. Then there exists a closed two-form $\eta \in \Omega_{0,0}^2(\log |D|)$, that is, $\text{Res}_q(\eta) = \text{Res}_r(\eta) = 0$, such that $\eta|_{\ker \varphi}$ is nondegenerate near D .*

Proof. Note that f is submersive with fibers F around D by Corollary 5.11. As $[F] \neq 0$, there exists $c \in H^2(X \setminus D)$ such that $\langle c, [F] \rangle = 1$. We will first construct local Lie algebroid two-forms around D as in Proposition 4.4. By Lemma 3.11, the cohomology $H^2(X \setminus D)$ equals $H_{0,0}^2(\log |D|)$, the cohomology of the subcomplex of zero radial (and elliptic) residue forms. Thus there exists a closed two-form $\xi \in \Omega_{0,0}^2(\log |D|)$ satisfying $[\xi] = c$.

Continuing, by Proposition 5.15 there exist open neighborhoods U and V around D and $f(D)$ and a defining function z for $f(D)$ on V such that f and $f|_D$ commute with the tubular neighborhood projections pr_D and pr_Z , and such that $f^*(z) = Q_{f^*(z)}$. Let $y \in f(D)$ and let $V' \subseteq V$ be a small open disc containing y on which NZ is trivial, and set $U' := f^{-1}(V')$. Then as in Proposition 5.10 we have $f(re^{i\theta}, x, y) = (r^2, x)$ using polar coordinates in normal directions. Moreover, in these coordinates we have $\ker \varphi = \langle \partial_\theta, \partial_y \rangle$. Let $\{V_i\}$ be a finite covering extracted from such open sets and let $U_i = f^{-1}(V_i)$, which together cover a neighborhood of D . Set $U_0 = X \setminus D$. As $f: U_i \setminus D \rightarrow V_i \setminus Z$ is a T^2 -bundle it is necessarily trivial, so that $H^2(U_i \setminus D) = H^2(T^2)$, with $H^1(T^2)$ generated by θ_i and γ_i say.

Define $\eta_0 = \xi|_{X \setminus D}$ and $\eta_i \in \Omega_{0,0}^2(U_i; \log |D|)$ for $i \geq 1$ via $\eta_i = \lambda_i \theta_i \wedge \gamma_i$, where $\lambda_i \in \mathbb{R}$ is chosen such that $\int_F \eta_i = \langle c, [F] \rangle$. Then $[\eta_i] = c|_{U_i} = [\xi|_{U_i}] \in H_{0,0}^2(U_i; \log |D|)$, so that by Lemma 3.11 there exist one-forms $\alpha_i \in \Omega_{0,0}^1(U_i; \log |D|)$ such that $\eta_i = \xi|_{U_i} + d\alpha_i$. As in Proposition 4.4, define a closed Lie algebroid two-form $\eta := \xi + d(\sum_i (\psi_i \circ f)\alpha_i)$ using a partition of unity $\{\psi_i\}$ subordinate to $\{V_i\}$. Note that $\text{Res}_q(\eta) = \text{Res}_r(\eta) = 0$. Finally, near D the form η is nondegenerate on $\ker \varphi$ as there it is given by the convex combination of forms $\eta = \sum_i (\psi \circ f)\eta_i$, where each η_i is nondegenerate on $\ker \varphi = \langle \partial_\theta, \partial_y \rangle$ by construction. \square

We can now prove Theorem 7.1 and Theorem 7.2 simultaneously, using Theorem 4.5 or Theorem 4.8 respectively.

Proof of Theorem 7.1 and Theorem 7.2. Let ω_Y be a log-symplectic structure on (Y, Z) . As Y is oriented, by Proposition 3.33 and Proposition 2.24 we know that Z is separating, and in particular coorientable. Using Proposition 5.19, replace (Y, Z) by $(Y', \partial Y')$, and then using Proposition 5.24, lift to a cover $g: \tilde{Y}' \rightarrow Y'$ so that $\tilde{f}: (X, D) \rightarrow (\tilde{Y}', \partial \tilde{Y}')$ has connected fibers. It is immediate that $g^*\omega_Y$ defines a log-symplectic structure on $(\tilde{Y}', \partial \tilde{Y}')$, and moreover that \tilde{f} is a boundary Lefschetz fibration. The generic fibers of $\tilde{f}: X \rightarrow \tilde{Y}'$ are either all homologically essential, or none are. Let $y \in Y' \setminus \partial Y'$ be a regular point and let $y_1, \dots, y_n \in g^{-1}(y) \subset \tilde{Y}'$ be its inverse images under the covering, with F_{y_1}, \dots, F_{y_n} their fibers under \tilde{f} . Then for some choice of signs we have $\pm[F_{y_1}] \pm \dots \pm [F_{y_n}] = [F]$. However, we have $[F] \neq 0$ for the generic fiber F by assumption, so there exists $i \in \{1, \dots, n\}$ such that $[F_{y_i}] \neq 0$. But then $[F_{y_i}] \neq 0$ for all $i \in \{1, \dots, n\}$, so \tilde{f} is homologically essential.

Apply Proposition 7.3 to the boundary (Lefschetz) fibration \tilde{f} . This yields a global closed Lie algebroid two-form $\eta \in \Omega_{\text{cl}}^2(X, \log |D|)$ such that $\eta|_{\ker \varphi}$ is nondegenerate near D (and globally so if f has no Lefschetz singularities). Moreover, its cohomology class $c = [\eta]$ pairs nontrivially with (generic) fibers in $X \setminus D$. Now apply either Theorem 4.5 or Theorem 4.8, possibly changing η to a form η' nondegenerate on $\ker \varphi$ everywhere, which agrees with η in a neighborhood of D . We obtain an elliptic symplectic structure $\omega_t = \varphi^*\omega_Y + t\eta'$ on $(X, |D|)$ for $t > 0$ small. As

$\text{Res}_q(\eta) = 0$ and η' agrees with η near D , we have $\text{Res}_q(\eta') = 0$ as well. Using Proposition 3.17 we see that $\text{Res}_q(\omega_t) = \text{Res}_q(\varphi^*\omega_Y) + t\text{Res}_q(\eta') = 0$. But then the elliptic symplectic structure ω_t has zero elliptic residue. By Theorem 3.32, we conclude that ω_t for $t > 0$ small determines a stable generalized complex structure on $(X, |D|)$. \square

The stable generalized complex structure on (X, D) constructed in the proof of Theorem 7.1 and Theorem 7.2 arises as an elliptic log-symplectic forms ω_X with zero elliptic residue through the correspondence of Theorem 3.32. As mentioned below Theorem 3.32, the three-form H required for integrability of the generalized complex structure is given by $[H] = \text{Res}_r([\omega_X]) \cup \text{PD}_X[D]$. The two-form η' introduced during the proof satisfies $\text{Res}_r(\eta') = 0$, because this holds for η . Using Proposition 3.17, together with the fact that $\omega_X = \varphi^*\omega_Y + t\eta'$ for some $t > 0$, we see that $\text{Res}_r(\omega_X) = \text{Res}_r(\varphi^*\omega_Y) + t\text{Res}_r(\eta') = f^*\text{Res}_Z(\omega_Y)$.

REMARK 7.4. Given a defining function z for Z we can locally write $\omega_Y = d \log z \wedge \alpha + \beta$, with (α, β) the induced cosymplectic structure on Z by the choice of z , and $\alpha = \text{Res}_Z(\omega_Y)$. Then $\varphi^*\omega_Y = d \log r \wedge f^*(\alpha) + f^*(\beta)$, and $\text{Res}_r(\varphi^*\omega_Y) = f^*(\text{Res}_Z(\omega_Y)) = f^*(\alpha)$.

7.1. *Fibering over T^2 and S^1*

Recall from Section 2 that in the compact case, for a stable generalized pair (X, D) and a log-Poisson pair (Y, Z) , the manifolds D and Z fiber over T^2 and S^1 respectively. After perturbing each structure, this can be achieved using one-forms on D and Z induced by the geometric structures (that is, they can be made proper). For a stable generalized complex structure \mathcal{J} on X , these one-forms are $(\text{Res}_r, \text{Res}_\theta)(\omega_X) \in \Omega^1(D) \times \Omega^1(D)$, where ω_X is the elliptic symplectic structure obtained from \mathcal{J} using Theorem 3.32, see [2, 9]. In the log-Poisson case, the one-form used instead is $\text{Res}_Z(\omega_Y)$ (see [6, 18, 19, 25, 30]), with ω_Y the log-symplectic structure induced by the log-Poisson structure using Proposition 3.33.

REMARK 7.5. If the log-symplectic structure is proper we improve upon Remark 7.4. Indeed, when Z fibers over S^1 using $\alpha := \text{Res}_Z(\omega_Y)$, the form α is Poincaré dual on Z to the fiber F_p of the induced fibration $p_Z: Z \rightarrow S^1$. In this case we obtain that $[H] = f^*(\text{PD}_Z[F_p]) \cup \text{PD}_X[D]$.

The stable generalized complex structures we construct out of boundary (Lefschetz) fibrations $f: (X, D) \rightarrow (Y, Z)$ are such that D fibers over Z . It is natural to ask whether this can be made compatible with the above fibrations $p_D: D \rightarrow T^2$ and $p_Z: Z \rightarrow S^1$. Note that $(\text{Res}_r, \text{Res}_\theta)(\omega_X) = (f^*\text{Res}_Z(\omega_Y), t\text{Res}_\theta(\eta))$ as $\omega_X = \varphi^*\omega_Y + t\eta$, and $\text{Res}_\theta(\varphi^*\omega_Y) = 0$ by the discussion below Proposition 5.8. This immediately shows we can make the fibrations compatible, as we can choose η such that $t\text{Res}_\theta(\eta)$ and $f^*\text{Res}_Z(\omega_Y)$ determine a fibration, instead of a just foliation. As in Section 6, let $p: T^2 \rightarrow S^1$ be projection onto the first factor.

COROLLARY 7.6. *Under the assumptions of Theorem 7.1 or Theorem 7.2, given a log-symplectic structure ω_Y on (Y, Z) , the elliptic two-form ω_X can be chosen such that $p \circ p_D = p_Z \circ f|_D$.*

In other words, we have a full commutative diagram around D and Z as follows:

$$\begin{array}{ccccc}
 ND & \xrightarrow{\text{pr}_D} & D & \xrightarrow{p_D} & T^2 \\
 f \downarrow & & f|_D \downarrow & & p \downarrow \\
 NZ & \xrightarrow{\text{pr}_Z} & Z & \xrightarrow{p_Z} & S^1
 \end{array}$$

8. *Examples and applications*

In this section we discuss some applications of the results obtained in this paper. For more examples we refer to [4], which in particular contains the classification theorem for boundary Lefschetz fibrations over \mathbb{D}^2 degenerating over its boundary (see Theorem 1.4). We start with the following immediate consequence of the results in Section 6 and Section 7.

THEOREM 8.1. *Let $f: X \rightarrow \mathbb{D}^2$ be a genus one Lefschetz fibration over the disc whose monodromy around the boundary is a power of a Dehn twist. Then all possible completions $\tilde{f}: (\tilde{X}, D) \rightarrow (\mathbb{D}^2, \partial\mathbb{D}^2)$ admit a stable generalized complex structure.*

Proof. Any Lefschetz fibration over the disc \mathbb{D}^2 is homologically essential, as it is obtained from the trivial fibration by two-handle attachments, so that it homotopy equivalent to Σ_g^2 (here $g = 1$) with some two-cells attached. By the monodromy assumption around the boundary, f admits completions \tilde{f} using Corollary 6.6. By Proposition 6.7, the map \tilde{f} is a homologically essential boundary Lefschetz fibration, so that (\tilde{X}, D) admits a stable generalized complex structure by Theorem 7.1, as the completion has D coorientable. \square

We now turn to exhibiting stable generalized complex structures on concrete four-manifolds using boundary (Lefschetz) fibrations.

EXAMPLE 8.2 ($S^1 \times S^3$). Let $I = [-1, 1]$ with coordinate t and view $S^2 \subseteq \mathbb{R}^3$ with north and south pole p_N and p_S . Consider the standard height function $p: (S^2, p_N \cup p_S) \rightarrow (I, \partial I)$ given by $(x, y, z) \mapsto z$. The map $h: (I, \partial I) \rightarrow (\mathbb{R}, 0)$ given by $h(t) := 1 - t^2$ is a defining function for ∂I . Then $p^*(h)(x, y, z) = 1 - (1 - x^2 - y^2) = x^2 + y^2$, so that p is a boundary map by the description of Proposition 5.8. In fact, p is a boundary fibration. Let $\varphi: S^3 \rightarrow S^2$ be the Hopf fibration, making $p \circ \varphi: (S^3, S_N^1 \cup S_S^1) \rightarrow (I, \partial I)$ a boundary fibration, where $S_N^1 = \varphi^{-1}(p_N)$ and similarly for S_S^1 . Finally, define the boundary fibration $f: (S^3 \times S^1, D) \rightarrow (I \times S^1, Z)$, where $Z = \partial I \times S^1$ and $D = (S_N^1 \cup S_S^1) \times S^1 = f^{-1}(Z)$, by $f(x, \theta) := ((p \circ \varphi)(x), \theta)$. Using Lemma 3.34, $(I \times S^1, Z)$ admits a log-symplectic structure, while the fibers of f are clearly homologically essential. By Theorem 7.2 applied to f , we conclude that $(S^1 \times S^3, D)$ admits a stable generalized complex structure whose type-change locus has two connected components. This structure is integrable with respect to the zero three-form due to Remark 7.4.

REMARK 8.3. We can make the previous example more explicit by introducing coordinates. On $\mathbb{C}^2 \setminus \{0\}$ with complex coordinates (z_1, z_2) , we can deform the complex structure by the holomorphic Poisson bivector $P = z_1 z_2 \partial_{z_1} \partial_{z_2}$ to obtain a stable generalized complex structure (see Proposition 2.8). Since this structure is invariant under scalar multiplication, it descends to a stable structure on the Hopf surface. This structure is compatible with the boundary fibration of the previous example. Indeed, the boundary fibration is given by the composition

$$f: S^3 \times S^1 \rightarrow S^2 \times S^1 \rightarrow I \times S^1, \quad (z_1, z_2) \mapsto \left(\frac{|z_1/z_2|^2 - 1}{|z_1/z_2|^2 + 1}, |z_1|^2 + |z_2|^2 \right).$$

The map then becomes a symplectic boundary fibration as one computes that

$$\omega \wedge f^*(dx) \wedge f^*(dy) = -2 \frac{z_1 \bar{z}_1 + z_2 \bar{z}_2}{|z_2|^4} dz_1 \wedge dz_2 \wedge d\bar{z}_1 \wedge d\bar{z}_2,$$

where $\omega = (z_1 z_2)^{-1} dz_1 dz_2$ is the associated singular two-form dual to P .

Using a slightly different map, we can ensure that the type change locus is connected.

EXAMPLE 8.4 ($S^1 \times S^3$). View $S^3 \subseteq \mathbb{C}^2$ with coordinates (z_1, z_2) in the standard way and consider the map $\varphi: S^3 \rightarrow \mathbb{D}^2$ given by $\varphi(z_1, z_2) = z_2$, viewing $\mathbb{D}^2 \subseteq \mathbb{C}$. In the interior $\mathbb{D}^2 \setminus \partial\mathbb{D}^2$, the map φ admits a section $s(z) = (\sqrt{1 - |z|^2}, z)$. As $\varphi \circ s = \text{id}$, we see that φ is submersive on $S^3 \setminus D$, where D is the circle $D = \varphi^{-1}(\partial\mathbb{D}^2) = \{(0, z_2) \in \mathbb{C}^2 : |z_2| = 1\}$. Moreover, $\varphi|_D$ is the identity from D to $Z = \partial\mathbb{D}^2$, which in particular implies that $\varphi|_D$ is submersive. Let $h(z) := 1 - |z|^2$ be a defining function for $\partial\mathbb{D}^2$ on \mathbb{D}^2 . Then $\varphi^*(h)(z_1, z_2) = 1 - |z_2|^2$, which shows that φ is a boundary map using Proposition 5.8. But then φ is in fact a boundary fibration. Noting that $\varphi|_{S^3 \setminus D}$ admits a section, $\varphi: S^3 \setminus D \rightarrow \mathbb{D}^2 \setminus \partial\mathbb{D}^2$ is a trivial S^1 -bundle. Now define a map of pairs $f: (S^3 \times S^1, D \times S^1) \rightarrow (\mathbb{D}^2, \partial\mathbb{D}^2)$ by $f(z_1, z_2, x) := \varphi(z_1, z_2)$. This is also a boundary fibration, which is homologically essential as $f|_{(S^3 \setminus D) \times S^1}$ is given by the projection $S^1 \times (\mathbb{D}^2 \setminus \partial\mathbb{D}^2) \times S^1 \rightarrow (\mathbb{D}^2, \partial\mathbb{D}^2)$. Finally, $(\mathbb{D}^2, \partial\mathbb{D}^2)$ admits a log-symplectic structure by Lemma 3.34. Using Theorem 7.2 we conclude that $(S^3 \times S^1, D \times S^1)$ admits a stable generalized complex structure with connected type-change locus which is integrable with respect to a nonzero degree three cohomology class. This agrees with [7, Example 4.1].

Before we consider actual boundary Lefschetz fibrations, it is convenient to reinterpret Proposition 6.5 with a Kirby calculus point of view. There, the completion of each end of X was done by gluing in a copy of the total space of L_n , the disc bundle over the torus with Euler class n . Observe that $\text{Tot}(L_n)$ is a four-manifold built out of one 0-handle, two 1-handles and one 2-handle. The process of capping off an end of X with $\text{Tot}(L_n)$ amounts to inverting the handle structure of $\text{Tot}(L_n)$ and adding it to X . That is, to cap each end of X we add one 2-handle, two 3-handles and one 4-handle. Further, due to inversion, the 2-handle of $\text{Tot}(L_n)$ has its original core and co-core interchanged and once this 2-handle is added there is a unique way to complete with the 3- and 4-handles, up to diffeomorphism. Therefore we only have to describe how to attach the 2-handle, whose core is the circle fiber of the projection $M \rightarrow T^2$, where M is the ε -sphere bundle of L_n . From the discussion in Proposition 6.3, this fiber is a circle left fixed by the monodromy map, which, if the monodromy is nontrivial, is determined by the monodromy map and is essentially unique. Further, since the circle is the fiber of the projection $M \rightarrow T^2$, the framing, in double strand notation, is given by the nearby fibers.

EXAMPLE 8.5 ($S^1 \times S^3 \# \overline{CP}^2$). Consider the genus one Lefschetz fibration over \mathbb{D}^2 with one singular fiber with vanishing cycle b , where $b \in H^1(T^2)$ is a generator. Then the monodromy around $\partial\mathbb{D}^2$ is b , which is clearly the power of a Dehn twist. The resulting completion \tilde{X} thus admits a stable generalized complex structure by Theorem 8.1, and we are left with determining its diffeomorphism type. Draw a Kirby diagram for the trivial T^2 -bundle over \mathbb{D}^2 in the standard way (see [13]) and attach a -1 -framed two-handle to represent the vanishing cycle b . The completion process adds a single 0-framed two-handle along the same cycle (see Figure 1). Slide the -1 -framed two-handle over this to split off a -1 -framed unknot, representing a copy of \overline{CP}^2 . The remaining diagram collapses to a single one-handle after two further handle slides, which shows that $\tilde{X} = S^1 \times S^3 \# \overline{CP}^2$. For more details, see [4, 8].

We can also recover the examples of [8], showing that $mCP^2 \# n\overline{CP}^2$ admits a stable generalized complex structure if and only if m is odd (that is, if it admits an almost-complex structure).

EXAMPLE 8.6 ($mCP^2 \# n\overline{CP}^2$). Let $m, n \in \mathbb{N}$ and assume that $m = 2k + 1$ is odd. Choose generators $a, b \in H^1(T^2)$ and consider $(\mathbb{D}^2, \partial\mathbb{D}^2)$ with counterclockwise assignment of singular fibers given by $a + (4k - 1)b$, followed by $a + 4jb$ for j from $k - 1$ up to $1 - k$, then $a - (4k - 1)b$, and finally n copies of b . The monodromy around the boundary can be computed to be $(10k - 1 - n)b$ [8]. Therefore the associated genus one Lefschetz fibration admits a completion




FIGURE 1. Figure (a) shows the base of a boundary Lefschetz fibration over the disc with a single nodal fiber, a choice of regular fiber for reference and loop around the boundary used to compute the monodromy and (b) shows the corresponding Kirby diagram.




FIGURE 2 (colour online). Figure (a) shows the base of a boundary Lefschetz fibration over the disc with $m + n$ singular fibers and a choice of regular fiber for reference at the center of disc. Figure (b) shows the corresponding Kirby diagram.

to $f: (\tilde{X}, D) \rightarrow (\mathbb{D}^2, \partial\mathbb{D}^2)$ (see Figure 2), resulting in a stable generalized complex structure on \tilde{X} by Theorem 8.1. As shown in [8, Proposition 5.4] we have $\tilde{X} = m\mathbb{C}P^2 \# n\overline{\mathbb{C}P}^2$.

Note that not every stable generalized complex structure comes from a boundary Lefschetz fibration, similarly to the case for symplectic structures and Lefschetz fibrations, or log-symplectic structures and achiral Lefschetz fibrations (see [10]).

EXAMPLE 8.7. The symplectic manifold $(\mathbb{C}P^2, \omega_{FS})$ carries a stable generalized complex structure with $D = \emptyset$. As $(\mathbb{C}P^2, \omega_{FS})$ is not a symplectic fibration over any surface, this stable generalized complex structure cannot be obtained through our construction.

EXAMPLE 8.8. Let Σ_g be the genus g surface and $g > 1$. Then $X = S^2 \times \Sigma_g$ has negative Euler characteristic so cannot admit a boundary Lefschetz fibration by Proposition 6.8. Hence X does not admit a stable generalized complex structure obtained through our methods. However, X is symplectic hence carries a stable generalized complex structure with $D = \emptyset$.

Acknowledgements. The authors would like to thank Selman Akbulut, Stefan Behrens, Robert Gompf and András Stipsicz for useful discussions.

References

1. D. AUROUX, S. K. DONALDSON and L. KATZARKOV, ‘Singular Lefschetz pencils’, *Geom. Topol.* 9 (2005) 1043–1114.
2. M. BAILEY, G. R. CAVALCANTI and M. GUALTIERI, ‘Type one generalized Calabi-Yaus’, *J. Geom. Phys.* 120 (2017) 89–95.
3. R. İ. BAYKUR, ‘Topology of broken Lefschetz fibrations and near-symplectic four-manifolds’, *Pacific J. Math.* 240 (2009) 201–230.
4. S. BEHRENS, G. R. CAVALCANTI and R. L. KLAASSE, ‘Classification of boundary Lefschetz fibrations over the disc’, *Geometry and Physics: Volume 2, Chapter 15* (Oxford University Press, 2018).
5. H. BURSZTYN, H. LIMA and E. MEINRENKEN, ‘Splitting theorems for Poisson and related structures’, *J. reine angew. Math.* 0 (0), <https://doi.org/10.1515/crelle-2017-0014>.
6. G. R. CAVALCANTI, ‘Examples and counter-examples of log-symplectic manifolds’, *J. Topol.* 10 (2017) 1–21.
7. G. R. CAVALCANTI and M. GUALTIERI, ‘A surgery for generalized complex structures on 4-manifolds’, *J. Differential Geom.* 76 (2007) 35–43.
8. G. R. CAVALCANTI and M. GUALTIERI, ‘Blow-up of generalized complex 4-manifolds’, *J. Topol.* 2 (2009) 840–864.
9. G. R. CAVALCANTI and M. GUALTIERI, ‘Stable generalized complex structures’, *Proc. Lond. Math. Soc.* 116 (2018) 1075–1111.
10. G. R. CAVALCANTI and R. L. KLAASSE, ‘Fibrations and log-symplectic structures’, *J. Symplectic Geom.*, to appear.
11. R. E. GOMPF, ‘The topology of symplectic manifolds’, *Turkish J. Math.* 25 (2001) 43–59.
12. R. E. GOMPF, ‘Toward a topological characterization of symplectic manifolds’, *J. Symplectic Geom.* 2 (2004) 177–206.
13. R. E. GOMPF and A. I. STIPSICZ, *4-manifolds and Kirby calculus*, Graduate Studies in Mathematics 20 (American Mathematical Society, Providence, RI, 1999).
14. R. GOTO, ‘Unobstructed deformations of generalized complex structures induced by C^∞ logarithmic symplectic structures and logarithmic Poisson structures’, *Geometry and topology of manifolds*, Springer Proceedings in Mathematics and Statistics 154 (eds A. Futaki, R. Miyaoki, Z. Tang, and W. Zhang; Springer, Tokyo, 2016) 159–183.
15. R. GOTO and K. HAYANO, ‘ C^∞ -logarithmic transformations and generalized complex structures’, *J. Symplectic Geom.* 14 (2016) 341–357.
16. M. GUALTIERI, ‘Generalized complex geometry’, PhD Thesis, Oxford University, Oxford, 2004.
17. M. GUALTIERI, ‘Generalized complex geometry’, *Ann. of Math.* (2) 211 (2011) 726–765.
18. M. GUALTIERI and S. LI, ‘Symplectic groupoids of log symplectic manifolds’, *Int. Math. Res. Not. IMRN* 11 (2014) 3022–3074.
19. V. GUILLEMIN, E. MIRANDA and A. R. PIRES, ‘Symplectic and Poisson geometry on b -manifolds’, *Adv. Math.* 264 (2014) 864–896.
20. N. HITCHIN, ‘Generalized Calabi-Yau manifolds’, *Q. J. Math.* 54 (2003) 281–308.
21. R. L. KLAASSE, ‘Geometric structures and Lie algebroids’, PhD Thesis, Utrecht University, Utrecht, 2017.
22. J. VAN DER LEER DURÁN, ‘Blow-ups in generalized complex geometry’, PhD Thesis, Utrecht University, Utrecht, 2016.
23. H. LI, ‘Topology of co-symplectic/co-Kähler manifolds’, *Asian J. Math.* 12 (2008) 527–543.
24. K. C. H. MACKENZIE, *General theory of Lie groupoids and Lie algebroids*, London Mathematical Society Lecture Note Series 213 (Cambridge University Press, Cambridge, 2005).
25. I. MĂRCUȚ and B. OSORNO TORRES, ‘Deformations of log-symplectic structures’, *J. Lond. Math. Soc.* (2) 90 (2014) 197–212.
26. I. MĂRCUȚ and B. OSORNO TORRES, ‘On cohomological obstructions for the existence of log-symplectic structures’, *J. Symplectic Geom.* 12 (2014) 863–866.
27. R. B. MELROSE, *The Atiyah-Patodi-Singer index theorem*, Research Notes in Mathematics 4 (A K Peters, Wellesley, MA, 1993).
28. R. NEST and B. TSYGAN, ‘Formal deformations of symplectic manifolds with boundary’, *J. reine angew. Math.* 481 (1996) 27–54.
29. L. NICOLAESCU, *An invitation to Morse theory*, 2nd edn, Universitext (Springer, New York, 2011).
30. B. OSORNO TORRES, ‘Codimension-one symplectic foliations: constructions and examples’, PhD Thesis, Utrecht University, Utrecht, 2015.
31. W. P. THURSTON, ‘Some simple examples of symplectic manifolds’, *Proc. Amer. Math. Soc.* 55 (1976) 467–468.
32. R. TORRES, ‘Constructions of generalized complex structures in dimension four’, *Comm. Math. Phys.* 314 (2012) 351–371.
33. R. TORRES and J. YAZINSKI, ‘On the number of type change loci of a generalized complex structure’, *Lett. Math. Phys.* 104 (2014) 451–464.
34. F. WALDHAUSEN, ‘Eine Klasse von 3-dimensionalen Mannigfaltigkeiten. I’, *Invent. Math.* 3 (1967) 308–333.
35. A. WEINSTEIN, ‘The local structure of Poisson manifolds’, *J. Differential Geom.* 18 (1983) 523–557.

Gil R. Cavalcanti
Department of Mathematics
Utrecht University
3508 TA Utrecht
The Netherlands

gil.cavalcanti@gmail.com

Ralph L. Klaasse
Department of Mathematics
Utrecht University
3508 TA Utrecht
The Netherlands

and

Département de Mathématique
Université libre de Bruxelles
CP 218, Boulevard du Triomphe
Brussels 1050
Belgium

r.l.klaasse@gmail.com