

LETTER • OPEN ACCESS

Global simulations of marine plastic transport show plastic trapping in coastal zones

To cite this article: Victor Onink *et al* 2021 *Environ. Res. Lett.* **16** 064053

View the [article online](#) for updates and enhancements.

ENVIRONMENTAL RESEARCH
LETTERS

LETTER

Global simulations of marine plastic transport show plastic trapping in coastal zones

OPEN ACCESS

RECEIVED

6 November 2020

REVISED

1 March 2021

ACCEPTED FOR PUBLICATION

8 March 2021

PUBLISHED

2 June 2021

Original Content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Victor Onink^{1,2,3,*} , Cleo E Jongedijk⁴ , Matthew J Hoffman⁵ , Erik van Sebille³ 
and Charlotte Laufkötter^{1,2} 

¹ Climate and Environmental Physics, Physics Institute, University of Bern, Bern, Switzerland

² Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

³ Institute for Marine and Atmospheric Research, Utrecht University, Utrecht, The Netherlands

⁴ Department of Civil and Environmental Engineering, Imperial College London, London, United Kingdom

⁵ School of Mathematical Sciences, Rochester Institute of Technology, Rochester, NY, United States of America

* Author to whom any correspondence should be addressed.

E-mail: victor.onink@climate.unibe.ch

Keywords: ocean modeling, marine plastic pollution, Lagrangian modeling, global ocean plastic transport, beached marine plastic

Supplementary material for this article is available [online](#)

Abstract

Global coastlines potentially contain significant amounts of plastic debris, with harmful implications for marine and coastal ecosystems, fisheries and tourism. However, the global amount, distribution and origin of plastic debris on beaches and in coastal waters is currently unknown. Here we analyze beaching and resuspension scenarios using a Lagrangian particle transport model. Throughout the first 5 years after entering the ocean, the model indicates that at least 77% of positively buoyant marine plastic debris (PBMPD) released from land-based sources is either beached or floating in coastal waters, assuming no further plastic removal from beaches or the ocean surface. The highest concentrations of beached PBMPD are found in Southeast Asia, caused by high plastic inputs from land and limited offshore transport, although the absolute concentrations are generally overestimates compared to field measurements. The modeled distribution on a global scale is only weakly influenced by local variations in resuspension rates due to coastal geomorphology. Furthermore, there are striking differences regarding the origin of the beached plastic debris. In some exclusive economic zones (EEZ), such as the Indonesian Archipelago, plastic originates almost entirely from within the EEZ while in other EEZs, particularly remote islands, almost all beached plastic debris arrives from remote sources. Our results highlight coastlines and coastal waters as important reservoirs of marine plastic debris and limited transport of PBMPD between the coastal zone and the open ocean.

1. Introduction

Marine plastic debris is found in almost all marine habitats, specifically on coastlines worldwide [1]. Coastal ecosystems can be particularly sensitive to plastic pollution [2], and plastic debris on beaches can reduce the economic value of a beach by up to 97% [3]. Furthermore, while an estimated 1.15–12.7 million tons of plastic enter the ocean per year [4–6], the amount of positively buoyant marine plastic debris (PBMPD) found floating at the ocean surface is estimated to be significantly lower [7–10]. Some of the plastic entering the ocean likely immediately sinks, as 34.5% of all plastics produced between 1950

and 2015 were made of neutrally or negatively buoyant polymers [11, 12], yet a significant amount of PBMPD is still unaccounted for. A large fraction of this missing PBMPD is potentially distributed on coastlines [12–14], with local concentrations varying between 0 and 647 kg km⁻¹ [15, 16]. However, given the scarcity of measurements relative to the length of coastlines, limited insight into local temporal and spatial fluctuations in beached PBMPD concentrations and the lack of a standardized sampling methodology, it is currently not possible to estimate the total amount of beached plastic debris from field measurements alone, or to describe the global pattern of beached plastic [1].

Using a simple box model, Lebreton *et al* [12] suggest that 66.8% of PBMPD released into the ocean since 1950 is stored on coastlines, however assuming a very high beaching probability and a resuspension probability below the observed range [17]. More complex global Lagrangian simulations of PBMPD have either not included beaching at all [18, 19] or use simple best guess implementations without considering resuspension [20–22]. Most of these studies have focused on plastic debris in the open ocean [9, 18], and do not report how global estimates of the amount and distribution of beached plastic vary with different beaching and resuspension parametrizations.

Here we present a series of idealized beaching experiments, using a Lagrangian particle tracking model with beaching and resuspension parametrizations. We estimate upper and lower bounds for the fraction of positively buoyant terrestrial plastic debris in coastal waters, on beaches and in the open ocean within the first years of release into the marine environment. Additionally, we describe the global relative distribution of beached plastic debris, and we analyze the relative amount of plastic with local versus remote origin.

2. Methods

2.1. Ocean surface current data

For the 2005–2015 global surface currents, we use the HYCOM + NCODA Global 1/12° surface current reanalysis [23] and the surface Stokes drift estimates from the WaveWatch III hindcast dataset [24, 25]. The HYCOM + NCODA Global 1/12° reanalysis [23] has a temporal resolution of 3 h and an equatorial spatial resolution of 1/12° (≈ 9.3 km). The HYCOM + NCODA Global 1/12° reanalysis does not incorporate Stokes drift, which has been shown to play an important role in shoreward surface transport [18]. Therefore, we add surface Stokes drift estimates from the WaveWatch III hindcast dataset [24, 25], which has a temporal resolution of 3 h and a spatial resolution of 1/2°. Comparison of Stokes drift estimates from the WaveWatch III dataset with *in situ* measurements from drifters have shown high correlations [26, 27], where root mean square errors have been on orders of centimeters per second [26]. Unless otherwise mentioned, simulations discussed in this paper have been done with surface currents obtained by the sum of the HYCOM currents and Stokes drift, as has been done in earlier modeling studies [10, 28, 29]. Not including Stokes drift reduces the amount of PBMPD that beaches by 6%–7% (supplementary figure 3 (available online at stacks.iop.org/ERL/16/064053/mmedia)) and reduces the trapping of PBMPD near the coast (supplementary figure 4). Stokes drift is thus an important component of the ocean circulation to consider in global PBMPD transport and beaching modeling.

PBMPD floating at the surface can be exposed to winds, with the strength of this effect depending on the size of the object that is exposed to winds above the ocean surface [30]. However, for the open ocean, the best model performance for modeling PBMPD is without including a separate windage term [10]. Furthermore, windage and Stokes drift are shown to be similar on a global scale [18]. Given that we include Stokes drift, we therefore do not consider an additional term for windage.

2.2. Lagrangian transport

We use Parcels [31, 32] to model plastic as virtual particles which are advected using surface ocean flow field data. A change in the position \vec{x} of a particle is calculated according to:

$$\vec{x}(t + \Delta t) = \vec{x}(t) + \int_t^{t+\Delta t} \vec{v}(\vec{x}(\tau), \tau) d\tau + R \sqrt{\frac{2 dt K_h}{r}}, \quad (1)$$

where $\vec{v}(\vec{x}(t), t)$ is the surface flow velocity at the particle location $\vec{x}(t)$ at time t , $R \in [-1, 1]$ is a random process representing subgrid motion with a mean of zero and variance $r = 1/3$, dt is the integration timestep, and K_h is the horizontal diffusion coefficient. The seed value of the random number generator does not influence the amount of beached plastic (supplementary figure 3). Equation (1) is integrated with a 4th order Runge–Kutta scheme with an integration timestep of $dt = 10$ min, and particle positions are saved every 24 h. We take $K_h = 10 \text{ m}^2 \text{ s}^{-1}$ [29, 33] to parameterize sub-grid processes.

2.3. Plastic emissions into the ocean

We use a terrestrial plastic input estimate based on the low end estimates of [4], where 15% of mismanaged plastic from the population living within 50 km of the coast enters the ocean. To obtain high-resolution estimates we multiply the country-specific mismanaged waste estimates with population densities [34] for 2010. This results in estimates of total mismanaged plastic for all cells on the HYCOM grid (supplementary figure 2). Polypropylene, polyethylene and polystyrene constitute 54% of primary plastic production in 2010 [11], and we assume that this fraction is indicative of how much mismanaged plastic is initially buoyant. We acknowledge that this 54% is a rough estimate, as it assumes that mismanaged plastic inputs have the same composition as global plastic production, whereas it has also been reported that heavier polymers can float with sufficient trapped air bubbles [35] and light polymers have been found submerged [36]. This leads to a total buoyant plastic input of 2.16×10^6 tons in 2010 (71.70% Asia, 4.39% North America, 3.94% South America, 2.16% Europe, 17.36% Africa and 0.45% Oceania).

The release of the virtual particles is scaled according to the estimate of buoyant plastic entering the ocean, where each particle represents up to 5.4 tons of buoyant plastic. To save computational resources we neglect sources smaller than 0.06 tons per year per grid cell, which represent 0.007% of the total input. In each run particles are released every 31 d during the first year of the simulation starting in 2010 (628,236 particles in total) and advected for 5 years. Starting the simulation in 2005 barely affects the amount of beached plastic (supplementary figure 3). We refer to this input scenario as the Jambeck input.

To test the model sensitivity to the plastic input, we calculate one simulation using a low end estimate of plastic waste entering the ocean from rivers [5]. Again assuming 54% of plastic entering the ocean is initially buoyant, we have an input of 6.21×10^5 tons for 2010 (87.04% Asia, 0.78% North America, 4.58% South America, 0.13% Europe, 7.45% Africa, 0.02% Oceania). Due to the smaller total input, no sources were neglected. We refer to this input scenario as the Lebreton input.

Particles are released in the shore-adjacent ocean cell nearest to the total mismanaged plastic cell in question. Since it is unlikely that real plastic always enters the ocean at exactly the same location, at the first time step particles are distributed randomly throughout the shore-adjacent ocean cell prior to the start of advection by the ocean currents.

2.4. Beaching parametrizations

2.4.1. Stochastic beaching and resuspension

Many processes are hypothesized to influence the amount of beached plastic on coastlines, such wind direction and speed, coast angle, aspect and morphology, local runoff, the proximity to urban centers and the degree of human usage of the beach [1, 36–42]. Many of these factors have some limited predictive power in statistical models that attempt to explain patterns of beached plastic [38, 39]. However, it is unclear from these studies whether these factors influence beaching, resuspension or both. They can also partially cancel each other out as they might work in opposite directions and in general it is unclear how these factors should be parameterized. We therefore decided to implement the simplest model possible, where we assume that on a global average, the main drivers of plastic beaching are the surface currents and the location of plastic input.

To account for the uncertainty of the ocean current data in land-adjacent ocean cells, we parametrize beaching as a stochastic process in the coastal zone, within which we consider the currents unreliable. For any given time step, we calculate the beaching probability p_B as:

$$p_b = \begin{cases} \text{if } d \leq D, p_B = 1 - \exp(-dt/\lambda_B) \\ \text{if } d > D, p_B = 0, \end{cases} \quad (2)$$

where d is the distance of particle to the nearest coastal cell, D is a predefined distance to the shore within which beaching can occur, dt is the integration timestep and λ_B is the characteristic timescale of plastic beaching. Beaching is therefore only possible within a beaching zone set by D . To account for the fact that global-scale ocean current datasets are inaccurate in ocean cells adjacent to land (referred to henceforth as coastal cells), we set D such that all coastal cells are fully contained within the beaching zone, resulting in a beaching zone of 10 km.

The probability of beaching is set by the beaching timescale λ_B , where λ_B is the number of days that a particle must spend within the beaching zone such that there is a 63.2% chance that the particle has beached. There is no experimental study to base the value of λ_B on, nor how it might vary for different types of plastic debris, so we selected a range of possibilities to investigate the sensitivity. For the sensitivity analysis we take $\lambda_B \in [1, 2, 5, 10, 26, 35, 100]$ days. Given the mean current speed in the coastal cells in the HYCOM dataset, $\lambda_B = 1$ day is the time a particle would require to travel 10 km in a straight line, representing a lower bound for the beaching probability. In the Mediterranean, analysis of GPS trajectories of drifter buoys suggests $\lambda_B = 76$ days [43], and an inverse modeling study suggests $\lambda_B = 26$ d for plastic debris [43]. We consider $\lambda_B = 100$ d to represent scenarios in which particles have very low beaching probabilities. A wooden drifter experiment in the North Sea found 46.88% of drifters beached within 91 d, traveling geodesic distances between 452 and 559 km [44]. Given that the drifters crossed the North Sea in this time and therefore spent time outside of the coastal zone, this suggests that λ_B is less than 100 d. However, we acknowledge that the values for λ_B remain a major source of uncertainty. Furthermore, unless specifically mentioned we parameterize beaching (and also resuspension) rates as global constants.

Particle resuspension is also implemented stochastically, where the resuspension probability p_R of a beached particle is defined as:

$$p_R = 1 - \exp(-dt/\lambda_R), \quad (3)$$

where dt is the time step and λ_R is the characteristic timescale of plastic resuspension. Hinata *et al* [17] has experimentally studied the resuspension timescale of plastic objects with different sizes and found $\lambda_R = 69$ –273 d. For our sensitivity analysis we take $\lambda_R \in [69, 171, 273]$ d. When a particle beaches, we save its last floating position, and when a particle resuspends it continues its trajectory from this position.

2.4.2. Coast-dependent resuspension

There have been a number of studies that have tried to explain the pattern of beached plastic using

statistical models that among others factors take geomorphology into account [38, 39]. However, it is unclear whether plastic beaching, resuspension or both are affected by geomorphology, and the influence of geomorphology likely differs for different types of plastic debris [45]. We are not aware of any studies investigating how geomorphology affects beaching probabilities. However, the dependence of resuspension probabilities on beach types have been studied with regard to the resuspension of oil [46]. Oil resuspension rates for sandy and rocky shores were found to be 24 and 18 h, respectively, and while these timescales are much shorter than the resuspension timescales for plastic [17], we use the ratio of the timescales of different coast types as a starting point for a sensitivity analysis.

For coastal geomorphology, we use data from Luijendijk *et al* [47] to determine the relative amount of sandy coastline s of each model cell of the HYCOM grid, where $s=0$ indicates a completely not-sandy coastline while $s=1$ indicates a completely sandy coastline (supplementary figure 1). Note that ‘not sandy’ covers multiple shore geomorphologies, such as rocky shores, cliffs and shorelines covered by vegetation such as mangrove forests. The resuspension timescale is determined by:

$$\lambda_R = \begin{cases} 3:4 \text{ Dependence} \rightarrow \lambda_R = 69 \times (0.75 + 0.25 \times s) \\ 1:4 \text{ Dependence} \rightarrow \lambda_R = 69 \times (0.25 + 0.75 \times s), \end{cases} \quad (4)$$

where with 3:4 Dependence we use the resuspension timescale coastline dependence for oil [46], whereas with 1:4 Dependence there a stronger dependence on the coastline type to check the sensitivity. In both cases we use $\lambda_{RT,s=1} = 69$ d.

There is currently little knowledge about how resuspension timescales vary with the coastline type and the classification of coastlines as sandy and not-sandy is overly simplistic, as coastlines such as rocky beaches, cliffs and mangroves are now considered equivalent. However, to our knowledge there have not been any studies that consider how plastic resuspension might depend on coastal geomorphology, and therefore we consider these runs as a first exploration of the potential role of coastal geomorphology on the global beached plastic budget and distribution.

2.5. Model concentration units

Model concentrations are computed by binning beached particle masses onto the same grid as the HYCOM reanalysis data, and then dividing the total beached mass in each cell by the length of model coastline (sum of cell edges shared with land cells) for that cell. This is because the beached plastic is not distributed homogeneously over the entire cell, but is instead concentrated on the shoreline interface between land and water, such as beaches. Since there is no global dataset of beach area, concentrations are

instead reported as the amount of plastic per length of model coastline (kg km^{-1}), which is commonly used for reporting field measurements (see table 1). However, due to the coarse resolution of the HYCOM grid, the length of the model coastline is an approximation of the true coastline length for a given cell.


3. Results

3.1. Global beached plastic budget

A systematic test of the effect of different beaching and resuspension probabilities on the global plastic budget is shown in figure 1. In all scenarios, the model reaches an equilibrium between the beaching and resuspension fluxes after the initial release within less than two years. At the end of our 5 years of simulations, between 31% and 95% of PBMPD is beached depending on parameter values. High beaching probabilities combined with small resuspension probabilities lead to a large amount of plastic stored on beaches and vice versa. With the Jambeck input, and assuming 54% of the input is buoyant, this corresponds to $0.72\text{--}2.06 \times 10^6$ tons of beached PBMPD originating from plastic debris released in 2010 alone.

Rather than being a function of the absolute values of λ_B and λ_R , our model shows that the average beached fraction is dependent on the ratio λ_B/λ_R (figure 2(a)). At very low ratios, i.e. high beaching but low resuspension probabilities, up to 99% of PBMPD is beached in the 5th year of the simulations. As the ratio increases, the beached fraction decreases to 31%. However, at least 77% of the PBMPD remains within 10 km of the model coastline in all scenarios (figure 2(a)) and only a small fraction escapes to the open ocean. Multiple studies report decreasing concentrations of floating PBMPD with increasing distance from shore, which is commonly attributed to PBMPD being removed from the ocean surface over time [35, 48–50]. However, such trends could also be partially due to PBMPD remaining trapped near-shore by the surface ocean currents.

While PBMPD can leave and return to the coastal zone, a large portion of PBMPD never travels far from the coastline (figure 2(b)). Over 25% of PBMPD mass never travels beyond 50 km from the nearest coastline even with the lowest beaching probability (supplementary table 1). The likelihood for plastic to leave the coastal zone is not uniform worldwide, as PBMPD that enters the ocean from island sources or from sources close to energetic boundary currents is more likely to travel further from shore (figure 2(c)). However, there are long stretches of coastline where most plastic remains near shore, as the median of the maximum distances from shore reached by particles released from those coastlines is less than 20 km offshore.




3.2. Global beached plastic distribution

Across all simulations, the highest beached PBMPD concentrations are found near regions with the largest PBMPD sources. These include areas such as Southeast Asia and the Mediterranean Sea (figure 3(a)), and have concentrations up to 10^6 kg km⁻¹. The lowest concentrations are in areas with low population densities, such as polar regions, the Chilean coastline and parts of the Australian coast. No PBMPD reaches the Antarctic mainland in any of our simulations. This is largely in line with measurements of plastic in Antarctica, which have been very low both on beaches [15] and afloat [51]. The lack of PBMPD in Antarctica is due to a lack of terrestrial input sources and the Antarctic Circumpolar Current blocking transport of PBMPD to Antarctic coastlines.

The global pattern of beached plastic is fairly robust towards the choice of beaching and resuspension probabilities (supplementary figure 5) but strongly depends on the plastic input distribution. With the Lebreton input, the relative global fraction

of PBMPD that is beached over the last year of simulation is 3% higher in comparison to using the Jambeck input (supplementary figure 3). However, absolute beached PBMPD concentrations are significantly lower, and a larger fraction of beached PBMPD is concentrated in Southeast Asia (figure 3(b)), reflecting higher inputs in this region. Therefore, detailed knowledge of the distribution and size of marine plastic debris sources are essential for understanding the global distribution of beached PBMPD.

While the overall distribution of beached plastic is largely shaped by the plastic input scenario, the ocean currents can play an important local role for beached PBMPD concentrations. For example, while higher beaching probabilities lead to higher global beached fractions, certain coastlines exhibit lower beached concentrations, such as Kenya, the Indian west coast and Libya (supplementary figure 5). More beached PBMPD globally results in less PBMPD afloat, and therefore reduced transport of PBMPD by the ocean currents to these areas.




3.3. Local versus remote origin of beached plastic debris

On a global average, 48.5% of beached PBMPD within all exclusive economic zones (EEZ, [52]) is local, in that it originates from a source within the EEZ. However, the local fraction of beached PBMPD is highly variable (figure 4). Generally, the local fraction for island EEZs is relatively low, matching recent reports for individual islands in various oceans [36, 53–55]. This is likely since their location in the open ocean exposes them to floating PBMPD originating from a wide range of EEZ's, while at the same time, PBMPD originating from an island EEZ itself is less likely to beach locally than on a mainland shore due to the comparatively small coastline of islands on which beaching can occur. In the field the fraction from local sources is further reduced due

to the contribution of maritime sources [36, 54, 55], which are not included in this model.

Higher local fractions of beached PBMPD are due to a combination of factors. Large local inputs generally lead to a higher local fractions, as a large fraction of PBMPD beaches close to its initial input. Examples of such EEZ's include China, Indonesia and Brazil. In addition, the ocean currents can play a critical role (see also figure 2(c)). Eastern Africa has relatively low local beached fractions, partly due to receiving high amounts of PBMPD from Indonesia transported by the Indian Ocean South Equatorial Current (matching observations by Ryan [56]). Meanwhile, coastlines such as the Russian Arctic and Chilean mainland have high local fractions despite low local inputs, since the prevailing local currents do not carry much PBMPD from other regions.




However, our local beached fraction estimates are only based on PBMPD that enters the ocean and subsequently beaches. Coastlines can also contain debris that is littered onto the coastline directly and never enters the ocean, and plastic debris that originates from maritime sources. Furthermore, ocean surface plastic removal processes such as sinking can further reduce the non-local fraction, but it is uncertain how large an effect this would have. As such, our estimates are only approximations of the actual local fraction of beached PBMPD.

4. Discussion

A systematic evaluation of our model based on a large number of field observations is currently impossible

due to the lack of a standardized measurement methodology of beached plastic [1], which prevents comparisons of plastic debris concentrations reported by different studies. Beached plastic concentrations are reported either as counts or masses, per unit area or unit length of coastline, and considering different debris sizes. Furthermore, our parameterizations do not account for beach cleanups, which are known to occur at many study sites (table 1). Additionally our simulations represent idealized scenarios: there is only one year of input; we do not consider loss processes such as sinking, ingestion, or burial in sediment [70]; maritime sources of PBMPD and beach littering are not considered.

Nevertheless, we compare the modeled relative distribution of beached plastic with studies that



measured beached plastic concentrations with a standardized method over multiple study sites. The modeled beached PBMPD distribution for South Africa closely resembles the distribution from field measurements [39], likewise our model captures the very low concentrations found on the Australian Northwestern coast [38]. However, the model appears to over-predict the amount of beached plastic on the Northeastern Australian coastline, potentially due an overestimated input of plastic from Polynesian islands. In the United States, the concentration ratio between the Northern Pacific coast and the Southern California Bight is approximately equal to the ratio reported in Ribic *et al* [62], but the relative amount of beached plastic in Hawaii is underestimated (table 1). PBMPD from maritime sources is often an important contributor of beached plastic on remote shores and islands [1, 55], and the lack of maritime sources in the input scenarios might partially account for this discrepancy. The overall relative similarity of the measured and modeled distribution is encouraging and indicates that we may have captured the dominant

drivers of beaching on a continental scale. However, these studies only allow comparisons of the relative patterns, as the model units (kg km^{-1}) do not match the field measurement concentrations. Compared to studies that do report concentrations in terms of mass, the model generally overestimates field concentrations by a factor of 5–560 (table 1). This either indicates that substantial losses of beached and floating plastic occur on short timescales that our model does not account for (such as burial within sediments [55, 71], sinking [35, 72, 73] and beach cleanups [74, 75]), and/or that the input estimates are too high. The sites where beached concentrations are underestimated (both in absolute and relative terms) are all islands, which could be due to neglecting maritime sources in the model.

Our results depend strongly on the representation of the ocean currents, the accuracy of the plastic input estimate, the beaching/resuspension parameterizations and the relative importance of processes that are not included. HYCOM has been shown to represent circulation patterns well in various parts

Table 1. A comparison of measurements of plastic debris with model simulations. The average, minimum and maximum model outputs are calculated over all stochastic simulations with the Jambeck input. Studies that do not report the occurrence of beach cleanups are indicated by a hyphen.

| Study | Location | Concentration (items km ⁻¹) | Cleanup | Model mean [min max] (kg km ⁻¹) |
|--|--|--|---|--|
| Barnes and Milner [58] ^{a,b} | Iceland | 40 | — | 3.28 [0.72 8.98] |
| | Faeroe Islands | 210 | — | 6.41 [0.04 30.58] |
| | La Gomera, Canary Islands | 1910 | — | 69.58 [0.55 178.99] |
| | Ascension | 3400 | — | 0.45 [0.00 6.14] |
| | Falkland Islands | 430 | — | 2.49 [0.68 7.68] |
| | Dominica | 1500 | — | 124.54 [0.00 918.14] |
| Pieper <i>et al</i> [37] ^a | Faial, Azores | 4610 | Regular during summer, none during the study period | 93.59 [0.00 285.74] |
| Ryan [59] ^a | Tristan da Cunha | 240 | — | 16.31 [0.00 100.53] |
| | Gough Island | 100 | — | 9.12 [0.00 96.57] |
| Otley and Ingham [60] ^a | Falkland Islands | 370 | — | 2.49 [0.68 7.68] |
| Convey <i>et al</i> [15] | Candlemas Island | 31 | None | 0.00 [0.00 0.00] |
| | Saunders Island | 50 | None | 0.00 [0.00 0.00] |
| | Adelaide Island | 0 | None | 0.00 [0.00 0.00] |
| Ribic <i>et al</i> [61] ^b | Northeast US Atlantic Coast | 102 | — | 12.46 [1.96 43.55] |
| | Middle US Atlantic Coast | 429 | — | 106.09 [5.86 397.56] |
| | Southeast US Atlantic Coast | 83 | — | 220.50 [64.85 683.85] |
| | Northern US Pacific Coast | 56 | Not regularly | 56.65 [35.00 116.03] |
| Ribic <i>et al</i> [62] ^b | Southern California Bight | 139 | Not regularly | 160.80 [43.92 623.58] |
| | Hawaii | 134 | Not regularly | 33.24 [4.90 120.65] |
| Study | Location | Concentration (kg km ⁻¹) | Cleanup | Model mean [min max] (kg km ⁻¹) |
| Corbin and Singh [63] | Dominica | 8 | — | 124.54 [0.00 918.14] |
| | St. Lucia | 3 | — | 110.18 [0.00 550.12] |
| Debrot <i>et al</i> [16] | Bonaire | 647 | — | 53.26 [0.00 285.57] |
| Debrot <i>et al</i> [40] | Curaçao | 506 | Occasionally | 43.12 [0.00 362.68] |
| Claereboudt [64] | Northern Oman | 15 | Occur, but frequency not specified | 86.43 [10.08 449.02] |
| Ali and Shams [65] | Clifton Beach, Karachi, Pakistan | 11 | Periodically, but frequency not specified | 4718.88 [167.62 55 688.46] |
| Hong <i>et al</i> [66] | South Korea | 262 | Not regularly | 970.65 [263.95 1611.52] |
| Pervez <i>et al</i> [67] | Shilaoren Beach, Qingdao, China | 5 | Daily | 2721.28 [327.44 13 124.59] |
| Pervez <i>et al</i> [68] | No. 1 Bathing Beach, Qingdao, China | 73 | Daily | 2721.28 [327.44 13 124.59] |
| Madzena and Lasiak [69] | Transkei Coast, South Africa | 47 | Not regularly | 1383.42 [111.45 5084.47] |
| Study | Location | Total Plastic (metric tons) | Cleanup | Model mean [min max] (kg km ⁻¹) |
| Lavers and Bond [55] | Henderson Island | 17.6 | Never | 0.08 [0.00 0.49] |

^a Concentrations as reported in Monteiro *et al* [57].

^b Total of marine debris, not solely plastic.

of the world [76–78]. However, HYCOM does not account for Stokes drift, which plays an important role in shoreward surface transport [18]. In line with earlier modeling studies [10, 28, 29], we take the

sum of the HYCOM currents and Stokes drift from the WaveWatch III reanalysis [24, 25], and we consider this the best available representation of global scale circulation. Nevertheless, we acknowledge that

trajectory modeling is more accurate in the open ocean than on the coastal shelf, where we also miss the effects of tidal currents [79–81].

With an estimated global beached fraction of 31%–95%, we show that the beached amount of plastic is a lot less constrained than suggested in Lebreton *et al* [12], whose simple 6-box model predicted that 69% of plastic that has entered the ocean since 1950 is found beached. In this box model, the authors assumed a 97% annual beaching rate of coastal PBMPD (equivalent to $\lambda_B = 104$ d), which is at the upper end of our tested range of plausible λ_B values (1–100 d), and they tuned their resuspension rate to match the global amount of floating PBMPD, resulting in a 1% annual resuspension rate of beached plastic (equivalent to $\lambda_R = 36\,317$ d) that is much slower than what is indicated by field experiments (69–273 d, [17]). In addition, the box model of Lebreton *et al* [12] assumes uniform off-shore transport, while we show that transport varies strongly in different regions (figure 2(c)).

We mostly use the Jambeck input scenario for our model [4]. There have been a number of estimates for plastic inputs into the ocean [4–6, 82], but it is unclear which are most accurate. Furthermore, all alternative estimates also neglect contributions from maritime sources and primary microplastics. There are indications that the amount of plastic entering the ocean is lower than estimated in the Jambeck input [83, 84] and given how strongly the modeled global distribution of PBMPD is influenced by the input scenario, it is crucial to get better estimates of plastic debris input sources, both terrestrial and maritime.

The model assumes that there are no PBMPD loss processes, or at least that they do not play a significant role during the first 5 years. For example, we assume PBMPD remains at the ocean surface, but processes such as biofouling can cause the density of PBMPD to increase until it starts to sink [85]. PBMPD can also be removed through ingestion by wildlife [86]. Experiments with tethered PBMPD biofouling show sinking of cm-sized plastic sheets after 17–66 d underneath a floating dock [85], but it is unclear how this translates to sinking rates for free floating PBMPD for different sizes, shape and regions. PBMPD has been found at the surface up to 50 years after its estimated production date [10], and while this PBMPD might not have floated at the ocean surface over this entire time period, it does suggest biofouling requires more than 66 d to sink PBMPD in the open ocean. Given these uncertainties, sinking was not included as PBMPD removal process in this study. Plastic ingestion has been found to occur with a wide range of species [87], but it is unclear how much total plastic has been ingested and at what rate this occurs. We also assume beached PBMPD remains available for resuspension indefinitely, but PBMPD can be transported towards the backshore [88] or be buried [1]. As a

consequence, the beached PBMPD budgets are upper estimates given that PBMPD is unable to exit the cycle of beaching and resuspension in our model.

Finally, we assume globally uniform beaching and resuspension probabilities. Exploratory tests with shore type dependent resuspension affects the global budget and distribution of beached plastic only minimally (supplementary figures 6 and 7), even when applying substantial differences in resuspension probability. These results indicate that on a global scale, local variations in resuspension probability driven by factors such as wind direction or coastal geomorphology play only a minor role. Nevertheless, more research is needed to understand how both beaching and resuspension are influenced by geomorphology and climatological factors, particularly on local to regional scales.

5. Conclusion

Our results indicate that part of the discrepancy between current plastic input estimates and estimates of floating plastic debris in the open ocean is due to high amounts of beached and coastal PBMPD. We have also identified coastlines where PBMPD is much more likely to reach the open ocean, such as the Eastern United States, Eastern Japan and Indonesia. Here, cleanups would be particularly effective in intercepting PBMPD before it escapes to the open ocean. However, more work needs to be done investigating the behavior of PBMPD in coastal waters, specifically the role of wind, waves, tides, and coastal morphology in PBMPD transport, beaching and resuspension. This would be strongly aided by standardized beached PBMPD field measurements, allowing comparisons of PBMPD concentrations at different measurement sites. Furthermore, future studies ought to consider the influence of maritime sources on beached PBMPD, as this study only considers terrestrial inputs.

Data availability statement

The data that support the findings of this study are available upon reasonable request from the authors.

Acknowledgments

Calculations were performed on UBELIX (www.id.unibe.ch/hpc), the HPC cluster at the University of Bern. V O and C L acknowledge support from the Swiss National Science Foundation (Project PZ00P2_174124 Global interactions between microplastics and marine ecosystems). E v S was supported by the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation programme (Grant Agreement No. 715386). C E J is supported by a Skempton Scholarship.

Author's contribution

Development of the beaching and resuspension parametrizations was done by V O, C E J, and C L, with input from M J H. The manuscript was written by V O and C L, with extensive input from C E J, M J H and E v S. Everyone contributed to the study design and discussion of the analysis, with V O carrying out the analysis.

Conflict of interests

The authors declare no competing interests.

Code availability

The code for the Lagrangian simulations is available at: <https://github.com/VictorOnink/Modeling-Global-Plastic-Beaching>.

ORCID iDs

Victor Onink  <https://orcid.org/0000-0003-4177-9893>

Cleo E Jongedijk  <https://orcid.org/0000-0001-9847-5212>

Matthew J Hoffman  <https://orcid.org/0000-0002-9430-005X>

Erik van Sebille  <https://orcid.org/0000-0003-2041-0704>

Charlotte Laufkötter  <https://orcid.org/0000-0001-5738-1121>

References

- [1] Browne M A *et al* 2015 Spatial and temporal patterns of stranded intertidal marine debris: is there a picture of global change? *Environ. Sci. Technol.* **49** 7082–94
- [2] Li R, Yu L, Chai M, Wu H and Zhu X 2020 The distribution, characteristics and ecological risks of microplastics in the mangroves of Southern China *Sci. Total Environ.* **708** 135025
- [3] Ballance A, Ryan P and Turpie J 2000 How much is a clean beach worth? The impact of litter on beach users in the Cape Peninsula, South Africa *South Afr. J. Sci.* **96** 210–30
- [4] Jambeck J R *et al* 2015 Plastic waste inputs from land into the ocean *Science* **347** 768–71
- [5] Lebreton L C *et al* 2017 River plastic emissions to the world's oceans *Nat. Commun.* **8** 15611
- [6] Schmidt C, Krauth T and Wagner S 2017 Export of plastic debris by rivers into the sea *Environ. Sci. Technol.* **51** 12246–53
- [7] Cózar A *et al* 2014 Plastic debris in the open ocean *Proc. Natl Acad. Sci.* **111** 10239–44
- [8] Eriksen M *et al* 2014 Plastic pollution in the world's oceans: more than 5 trillion plastic pieces weighing over 250,000 tons afloat at sea *PLoS One* **9** e111913
- [9] van Sebille E *et al* 2015 A global inventory of small floating plastic debris *Environ. Res. Lett.* **10** 124006
- [10] Lebreton L *et al* 2018 Evidence that the great pacific garbage patch is rapidly accumulating plastic *Sci. Rep.* **8** 4666
- [11] Geyer R, Jambeck J R and Law K L 2017 Production, use and fate of all plastics ever made *Sci. Adv.* **3** e1700782
- [12] Lebreton L, Egger M and Slat B 2019 A global mass budget for positively buoyant macroplastic debris in the ocean *Sci. Rep.* **9** 1–10
- [13] Hardesty B D *et al* 2017 Using numerical model simulations to improve the understanding of micro-plastic distribution and pathways in the marine environment *Front. Mar. Sci.* **4** 30
- [14] Schwarz A, Lighthart T, Boukris E and Van Harmelen T 2019 Sources, transport and accumulation of different types of plastic litter in aquatic environments: a review study *Mar. Pollut. Bull.* **143** 92–100
- [15] Convey P, Barnes D and Morton A 2002 Debris accumulation on oceanic island shores of the Scotia Arc, Antarctica *Polar Biol.* **25** 612–17
- [16] Debrot A O, van Rijn J, Bron P S and de León R 2013 A baseline assessment of beach debris and tar contamination in Bonaire, Southeastern Caribbean *Mar. Pollut. Bull.* **71** 325–9
- [17] Hinata H, Mori K, Ohno K, Miyao Y and Kataoka T 2017 An estimation of the average residence times and onshore-offshore diffusivities of beached microplastics based on the population decay of tagged meso- and macrolitter *Mar. Pollut. Bull.* **122** 17–26
- [18] Onink V, Wichmann D, Delandmeter P and van Sebille E 2019 The role of Ekman currents, geostrophy and Stokes drift in the accumulation of floating microplastic *J. Geophys. Res. Oceans* **124** 1474–90
- [19] Miladinova S, Macías D, Stips A and Garcia-Gorrioz E 2020 Identifying distribution and accumulation patterns of floating marine debris in the black sea *Mar. Pollut. Bull.* **153** 110964
- [20] Lebreton L-M, Greer S and Borrero J C 2012 Numerical modelling of floating debris in the world's oceans *Mar. Pollut. Bull.* **64** 653–61
- [21] Critchell K *et al* 2015 Modelling the fate of marine debris along a complex shoreline: lessons from the great barrier reef *Estuar. Coast. Shelf Sci.* **167** 414–26
- [22] Carlson D F *et al* 2017 Combining litter observations with a regional ocean model to identify sources and sinks of floating debris in a semi-enclosed basin: the adriatic sea *Front. Mar. Sci.* **4** 78
- [23] Bleck R 2002 An oceanic general circulation model framed in hybrid isopycnic-cartesian coordinates *Ocean Model.* **4** 55–88
- [24] Tolman H L 1997 *User manual and system documentation of WAVEWATCH-III version 1.15* (US Department of Commerce, National Oceanic and Atmospheric Administration, National Weather Service, National Centers for Environmental Prediction)
- [25] Tolman H L 2009 User manual and system documentation of WAVEWATCH III™ version 3.14 *Technical Note, MMAB Contribution vol 276* p 220
- [26] Tamura H, Miyazawa Y and Oey L-Y 2012 The Stokes drift and wave induced-mass flux in the North Pacific *J. Geophys. Res. Oceans* **117** C08021
- [27] Rasche N and Ardhuin F 2013 A global wave parameter database for geophysical applications. Part 2: model validation with improved source term parameterization *Ocean Model.* **70** 174–88
- [28] Fraser C I *et al* 2018 Antarctica's ecological isolation will be broken by storm-driven dispersal and warming *Nat. Clim. Change* **8** 704–8
- [29] Lacerda A L d F *et al* 2019 Plastics in sea surface waters around the Antarctic Peninsula *Sci. Rep.* **9** 3977
- [30] Van Den Bremer T and Breivik Ø 2018 Stokes drift *Phil. Trans. R. Soc. A* **376** 20170104
- [31] Lange M and van Sebille E 2017 Parcels v0.9: prototyping a Lagrangian ocean analysis framework for the petascale age *Geosci. Model Dev. Discuss.* **10** 4175–86
- [32] Delandmeter P and Van Sebille E 2019 The parcels v2.0 Lagrangian framework: new field interpolation schemes *Geosci. Model Dev.* **12** 3571–84

- [33] Liubartseva S, Coppini G, Lecci R and Clementi E 2018 Tracking plastics in the Mediterranean: 2D Lagrangian model *Mar. Pollut. Bull.* **129** 151–62
- [34] Center for International Earth Science Information Network—CIESIN—Columbia University 2016 Gridded population of the world, version 4 (gpwv4): population density
- [35] Ryan P G 2015 Does size and buoyancy affect the long-distance transport of floating debris? *Environ. Res. Lett.* **10** 084019
- [36] Brignac K C *et al* 2019 Marine debris polymers on main Hawaiian island beaches, sea surface and seafloor *Environ. Sci. Technol.* **53** 12218–26
- [37] Pieper C, Ventura M A, Martins A and Cunha R T 2015 Beach debris in the azores (NE Atlantic): Faial island as a first case study *Mar. Pollut. Bull.* **101** 575–82
- [38] Hardesty B D, Lawson T, van der Velde T, Lansdell M and Wilcox C 2017 Estimating quantities and sources of marine debris at a continental scale *Front. Ecol. Environ.* **15** 18–25
- [39] Ryan P G, Perold V, Osborne A and Moloney C L 2018 Consistent patterns of debris on South African beaches indicate that industrial pellets and other mesoplastic items mostly derive from local sources *Environ. Pollut.* **238** 1008–16
- [40] Debrot A O, Tiel A B and Bradshaw J E 1999 Beach debris in Curacao *Mar. Pollut. Bull.* **38** 795–801
- [41] Smith S D 2012 Marine debris: a proximate threat to marine sustainability in bootless bay, Papua New Guinea *Mar. Pollut. Bull.* **64** 1880–3
- [42] Thiel M *et al* 2013 Anthropogenic marine debris in the coastal environment: a multi-year comparison between coastal waters and local shores *Mar. Pollut. Bull.* **71** 307–16
- [43] Kaandorp M L A, Dijkstra H A and van Sebille E 2020 Closing the Mediterranean marine floating plastic mass budget: inverse modeling of sources and sinks *Environ. Sci. Technol.* **54** 11980–9
- [44] Stanev E *et al* 2019 Extreme westward surface drift in the North Sea: public reports of stranded drifters and Lagrangian tracking *Cont. Shelf Res.* **177** 24–32
- [45] Weideman E A, Perold V, Ouardien A, Smyth L K and Ryan P G 2020 Quantifying temporal trends in anthropogenic litter in a rocky intertidal habitat *Mar. Pollut. Bull.* **160** 111543
- [46] Samaras A G, De Dominicis M, Archetti R, Lamberti A and Pinardi N 2014 Towards improving the representation of beaching in oil spill models: a case study *Mar. Pollut. Bull.* **88** 91–101
- [47] Luijendijk A *et al* 2018 The state of the world's beaches *Sci. Rep.* **8** 6641
- [48] Ryan P G 2013 A simple technique for counting marine debris at sea reveals steep litter gradients between the Straits of Malacca and the Bay of Bengal *Mar. Pollut. Bull.* **69** 128–36
- [49] Pedrotti M L *et al* 2016 Changes in the floating plastic pollution of the Mediterranean sea in relation to the distance to land *PLoS One* **11** e0161581
- [50] Ruiz-Orejón L F, Sardá R and Ramis-Pujol J 2018 Now, you see me: high concentrations of floating plastic debris in the coastal waters of the Balearic Islands (Spain) *Mar. Pollut. Bull.* **133** 636–46
- [51] Suaria G *et al* 2020 Floating macro- and microplastics around the Southern Ocean: results from the Antarctic circumnavigation expedition *Environ. Int.* **136** 105494
- [52] Institute F M 2019 Union of the ESRI country shapefile and the exclusive economic zones (version 3) (available at: <https://doi.org/10.14284/403>)
- [53] Pieper C, Amaral-Zettler L, Law K L, Loureiro C M and Martins A 2019 Application of matrix scoring techniques to evaluate marine debris sources in the remote islands of the Azores Archipelago *Environ. Pollut.* **249** 666–75
- [54] Ryan P G, Dille B J, Ronconi R A and Connan M 2019 Rapid increase in Asian bottles in the south Atlantic ocean indicates major debris inputs from ships *Proc. Natl Acad. Sci.* **116** 20892–7
- [55] Lavers J L and Bond A L 2017 Exceptional and rapid accumulation of anthropogenic debris on one of the world's most remote and pristine islands *Proc. Natl Acad. Sci.* **114** 6052–5
- [56] Ryan P G 2020 Land or sea? what bottles tell us about the origins of beach litter in Kenya *Waste Manage.* **116** 49–57
- [57] Monteiro R C, do Sul J A I and Costa M F 2018 Plastic pollution in islands of the Atlantic ocean *Environ. Pollut.* **238** 103–10
- [58] Barnes D K and Milner P 2005 Drifting plastic and its consequences for sessile organism dispersal in the Atlantic Ocean *Mar. Biol.* **146** 815–25
- [59] Ryan P G 1987 The origin and fate of artefacts stranded on islands in the African sector of the Southern Ocean *Environ. Conserv.* **14** 341–6
- [60] Otley H and Ingham R 2003 Marine debris surveys at volunteer beach, Falkland Islands, during the summer of 2001/02 *Mar. Pollut. Bull.* **46** 1534–9
- [61] Ribic C A, Sheavly S B, Rugg D J and Erdmann E S 2010 Trends and drivers of marine debris on the Atlantic coast of the United States 1997–2007 *Mar. Pollut. Bull.* **60** 1231–42
- [62] Ribic C A, Sheavly S B, Rugg D J and Erdmann E S 2012 Trends in marine debris along the U.S. Pacific coast and Hawai'i 1998–2007 *Mar. Pollut. Bull.* **64** 994–1004
- [63] Corbin C and Singh J 1993 Marine debris contamination of beaches in St. Lucia and Dominica *Mar. Pollut. Bull.* **26** 325–8
- [64] Claereboudt M R 2004 Shore litter along sandy beaches of the Gulf of Oman *Mar. Pollut. Bull.* **49** 770–7
- [65] Ali R and Shams Z I 2015 Quantities and composition of shore debris along Clifton Beach, Karachi, Pakistan *J. Coast. Conserv.* **19** 527–35
- [66] Hong S, Lee J, Kang D, Choi H-W and Ko S-H 2014 Quantities, composition and sources of beach debris in Korea from the results of nationwide monitoring *Mar. Pollut. Bull.* **84** 27–34
- [67] Pervez R, Wang Y, Ali I, Ali J and Ahmed S 2020 The analysis of the accumulation of solid waste debris in the summer season along the Shilaoren Beach Qingdao, China *Reg. Stud. Mar. Sci.* **34** 101041
- [68] Pervez R, Wang Y, Mahmood Q, Zahir M and Jattak Z 2020 Abundance, type and origin of litter on no. 1 bathing beach of Qingdao, China *J. Coast. Conserv.* **24** 34
- [69] Madzena A and Lasiak T 1997 Spatial and temporal variations in beach litter on the Transkei coast of South Africa *Mar. Pollut. Bull.* **34** 900–7
- [70] Van Sebille E *et al* 2020 The physical oceanography of the transport of floating marine debris *Environ. Res. Lett.* **15** 023003
- [71] Martin C *et al* 2020 Exponential increase of plastic burial in mangrove sediments as a major plastic sink *Sci. Adv.* **6** eaaz5593
- [72] Choy C A *et al* 2019 The vertical distribution and biological transport of marine microplastics across the epipelagic and mesopelagic water column *Sci. Rep.* **9** 1–9
- [73] Egger M, Sulu-Gambari F and Lebreton L 2020 First evidence of plastic fallout from the North Pacific garbage patch *Sci. Rep.* **10** 1–10
- [74] Uneputtu P, Evans S and Suyoso E 1998 The effectiveness of a community education programme in reducing litter pollution on shores of Ambon Bay (Eastern Indonesia) *J. Biol. Educ.* **32** 143–7
- [75] Willis K, Maureaud C, Wilcox C and Hardesty B D 2018 How successful are waste abatement campaigns and government policies at reducing plastic waste into the marine environment? *Mar. Policy* **96** 243–9
- [76] Savage J A, Tokmakian R T and Batteen M L 2015 Assessment of the HYCOM velocity fields during Agulhas return current cruise 2012 *J. Oper. Oceanogr.* **8** 11–24
- [77] Metzger E J *et al* 2010 Simulated and observed circulation in the Indonesian seas: 1/12 global HYCOM and the instant observations *Dyn. Atmos. Oceans* **50** 275–300

- [78] Wang M *et al* 2019 Origin and formation of the Ryukyu current revealed by HYCOM reanalysis *Acta Oceanol. Sin.* **38** 1–10
- [79] Liu Y and Weisberg R H 2011 Evaluation of trajectory modeling in different dynamic regions using normalized cumulative Lagrangian separation *J. Geophys. Res. Oceans* **116** C09013
- [80] Liu Y, Weisberg R H, Vignudelli S and Mitchum G T 2014 Evaluation of altimetry-derived surface current products using Lagrangian drifter trajectories in the Eastern Gulf of Mexico *J. Geophys. Res. Oceans* **119** 2827–42
- [81] Sterl M F, Delandmeter P and van Sebille E 2020 Influence of barotropic tidal currents on transport and accumulation of floating microplastics in the global open ocean *J. Geophys. Res. Oceans* **125** e2019JC015583
- [82] Lebreton L and Andrady A 2019 Future scenarios of global plastic waste generation and disposal *Palgrave Commun.* **5** 1–11
- [83] Tramoy R *et al* 2019 Assessment of the plastic inputs from the seine basin to the sea using statistical and field approaches *Front. Mar. Sci.* **6** 151
- [84] Van Emmerik T, Loozen M, Van Oeveren K, Buschman F and Prinsen G 2019 Riverine plastic emission from Jakarta into the ocean *Environ. Res. Lett.* **14** 084033
- [85] Fazey F M and Ryan P G 2016 Biofouling on buoyant marine plastics: an experimental study into the effect of size on surface longevity *Environ. Pollut.* **210** 354–60
- [86] Van Franeker J A and Law K L 2015 Seabirds, gyres and global trends in plastic pollution *Environ. Pollut.* **203** 89–96
- [87] Derraik J G 2002 The pollution of the marine environment by plastic debris: a review *Mar. Pollut. Bull.* **44** 842–52
- [88] Pham C K *et al* 2020 The Azores archipelago as a transitory repository for small plastic fragments floating in the North-East Atlantic *Environ. Pollut.* **263** 114494