



Control of transplant tolerance and intragraft regulatory T cell localization by CCL5

Nahzli Dilek, Nicolas Poirier, Claire Usal, Bernard Martinet, Gilles Blancho,
Bernard Vanhove

► To cite this version:

Nahzli Dilek, Nicolas Poirier, Claire Usal, Bernard Martinet, Gilles Blancho, et al.. Control of transplant tolerance and intragraft regulatory T cell localization by CCL5. 7th European Workshop on Immune-Mediated Inflammatory Diseases, Nov 2012, Netherlands. 10 (Suppl 3), pp.P60, 2012. <inserm-00758212>

HAL Id: inserm-00758212

<http://www.hal.inserm.fr/inserm-00758212>

Submitted on 28 Nov 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



POSTER PRESENTATION

Open Access

Control of transplant tolerance and intragraft regulatory T cell localization by CCL5

Nahzli Dilek, Nicolas Poirier, Claire Usal, Bernard Martinet, Gilles Blancho, Bernard Vanhove*

From 7th European Workshop on Immune-Mediated Inflammatory Diseases
Noordwijk aan Zee, the Netherlands. 28-30 November 2012

Background

CCL5 (Rantes) is a chemotactic cytokine playing an active role in recruiting leukocytes into inflammatory sites and known to attract Treg cells in solid tumors where they inhibit immune responses.

Material, methods and results

In rat recipients of kidney allografts where tolerance was induced by costimulation blockade, we observed a 2-fold decrease of plasma CCL5 levels as compared to control syngeneic grafted animals (0.60 ± 0.17 vs. 1.17 ± 0.48 ng/ml; $p < 0.01$). Myeloid-derived suppressor cells (MDSC), a major cell type producing CCL5 in blood, also presented a reduced capacity to produce CCL5 in these tolerant animals. In contrast, intragraft levels of CCL5 mRNA and protein were higher in tolerated allografts, establishing an increased graft-to-periphery CCL5 gradient possibly contributing to the recruitment of Treg cells into the graft, as in solid tumor, and leading to the establishment and maintenance of tolerance. To test the hypothesis, we broke the gradient by restoring normal plasma concentrations of CCL5 by implantation of osmotic pumps. This induced a strong reduction of intragraft Treg cells (decrease of 24.40 ± 5.47 fold; $p < 0.05$ by immunohistofluorescence and of Foxp3 mRNA assessed by qRT-PCR), and led to an increase of creatinine and urea concentrations and eventually to kidney graft rejection.

Conclusions

Our data uncover a novel role of CCL5, possibly controlled by MDSC, in this rat model of kidney transplant tolerance: a graft-to-periphery gradient of CCL5 help

recruiting Treg cells into the graft where they maintain tolerance.

Published: 28 November 2012

doi:10.1186/1479-5876-10-S3-P60

Cite this article as: Dilek et al.: Control of transplant tolerance and intragraft regulatory T cell localization by CCL5. *Journal of Translational Medicine* 2012 10(Suppl 3):P60.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Institute of Transplantation Urology Nephrology, University of Nantes,
INSERM UMR 643, Nantes, France Effimune, Nantes, France CHU de Nantes,
Nantes, France