



## Morse 2-jet space and h-principle

Alain Chenciner, François Laudenbach

► **To cite this version:**

Alain Chenciner, François Laudenbach. Morse 2-jet space and h-principle. Bulletin Brazilian Mathematical Society, 2009, 40 (4), pp.455-463. <hal-00363505>

**HAL Id: hal-00363505**

**<https://hal.archives-ouvertes.fr/hal-00363505>**

Submitted on 23 Feb 2009

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# MORSE 2-JET SPACE AND $h$ -PRINCIPLE

ALAIN CHENCINER AND FRANÇOIS LAUDENBACH

ABSTRACT. A section in the 2-jet space of Morse functions is not always homotopic to a holonomic section. We give a necessary condition for being the case and we discuss the sufficiency.

## 1. INTRODUCTION

Given a submanifold  $\Sigma$  in an  $r$ -jet space (of smooth sections of a bundle over a manifold  $M$ ), it is natural to look at the associated *differential relation*  $\mathcal{R}(\Sigma)$  formed by the  $(r+1)$ -jets *transverse* to  $\Sigma$ . For  $j^r f$  being transverse to  $\Sigma$  at  $x \in M$  is detected by  $j_x^{r+1} f$ . This is an open differential relation in the corresponding  $(r+1)$ -jet space. One can ask whether the Gromov *h-principle* holds: is any section with value in  $\mathcal{R}(\Sigma)$  homotopic to a *holonomic* section of  $\mathcal{R}(\Sigma)$ ? (We recall that a holonomic section of a  $(r+1)$ -jet space is a section of the form  $j^{r+1} f$ .)

According to M. Gromov, the answer is yes when  $M$  is an open manifold and  $\Sigma$  is *natural*, that is, invariant by a lift of  $\text{Diff}(M)$  to the considered jet space (see [3] p. 79, [1] ch. 7).

The answer is also yes when the codimension of  $\Sigma$  is higher than the dimension  $n$  of  $M$ ; this case follows easily from Thom's transversality theorem in jet spaces (see [6]). In the case of jet space of functions and when  $\Sigma$  is natural and  $\text{codim} \Sigma \geq n+1$ , it also can be seen as a baby case of a theorem of Vassiliev [9].

In this note we are interested in a codimension  $n$  case when  $M$  is a compact  $n$ -dimensional manifold. Let  $J^r(M)$  denote the space of  $r$ -jets of real functions; when the boundary of  $M$  is not empty, it is meant that we speak of jets of functions which are locally constant on the boundary. We take  $\Sigma \subset J^1(M)$  the set of critical 1-jets. Then  $\mathcal{R}(\Sigma) \subset J^2(M)$  is the open set of 2-jets of Morse functions. We shall analyze the obstructions preventing the *h-principle* to hold with this differential relation.

## 2. INDEX COCYCLES

It is more convenient to work with the *reduced* jet spaces  $\tilde{J}^r(M)$ , quotient of  $J^r(M)$  by  $\mathbb{R}$  which acts by translating the value of the jet. It is a vector bundle whose linear structure is induced by that of  $C^\infty(M)$ . For instance,  $\tilde{J}^1(M)$  is isomorphic to the cotangent space  $T^*M$ . Let  $\mathcal{M}$  denote the reduced 2-jets of Morse functions, that is the 2-jets which are transverse to the zero section  $0_M$  of  $T^*M$  (in the sequel, *jet* will mean *reduced jet*). Let  $\pi : \tilde{J}^2(M) \rightarrow \tilde{J}^1(M)$  be the projection and  $\pi_0 : \tilde{J}_0^2(M) \rightarrow 0_M$  be its restriction over the zero section of the cotangent

---

*Date:* This version: February, 2009.

2000 *Mathematics Subject Classification.* 58A20, 58E05, 57R45.

*Key words and phrases.* Morse singularity, 2-jet, *h-principle*.

space. Since it is formed of critical 2-jets, it is a vector bundle whose fiber is the space of quadratic forms,  $S^2(T_x^*M)$ ,  $x \in M$ . Let  $\mathcal{M}_0 := \mathcal{M} \cap \tilde{\mathcal{J}}_0^2(M)$ ; it is a bundle over  $0_M$  whose fiber consists of non-degenerate quadratic forms. Its complement in  $\tilde{\mathcal{J}}_0^2(M)$  is denoted  $\mathcal{D}$  (like discriminant); it is formed of 2-jets which are not transverse to  $0_M$ . When  $M$  is connected,  $\mathcal{M}_0$  has a connected component  $\mathcal{M}_0^i$  for each index  $i \in \{1, \dots, n\}$  of quadratic forms.

**2.1. Transverse orientation.** Each  $\mathcal{M}_0^i$  is a proper submanifold of codimension  $n$  in  $\mathcal{M}$ . Moreover the differential  $d\pi$  gives rise to an isomorphism of normal fiber bundles

$$\nu(\mathcal{M}_0^i, \mathcal{M}) \cong \pi^*(\nu(0_M, T^*M))|_{\mathcal{M}_0^i}.$$

Of course,  $\nu(0_M, T^*M)$  is canonically isomorphic to the cotangent bundle  $\tau^*M$ , whose total space is  $T^*M$ . When  $M$  is oriented, so is the bundle  $\tau^*M$ . When  $M$  is not orientable, one has a local system of orientations of  $\tau^*M$ . Pulling it back by  $\pi$  yields a local system of orientations of  $\nu(\mathcal{M}_0^i, \mathcal{M})$  (that is, co-orientations of  $\mathcal{M}_0^i$ ). Let us denote  $\mathcal{M}_0^{even}$  (resp.  $\mathcal{M}_0^{odd}$ ) the union of the  $\mathcal{M}_0^i$ 's for  $i$  even (resp. odd). We endow  $\mathcal{M}_0^{even}$  with the above local system of co-orientations. For reasons which clearly appear below, it is more natural to equip  $\mathcal{M}_0^{odd}$  with the opposite system of co-orientations.

**Lemma 2.2.** *Let  $s = j^2f$  be a holonomic section of  $\mathcal{M}$  meeting  $\mathcal{M}_0$  transversally. Then each intersection point of  $s(M)$  with  $\mathcal{M}_0$  is positive. The same statement holds when  $s$  is a local holonomic section only.*

**Proof.** Let  $a$  be such an intersection point in  $s(M) \cap \mathcal{M}_0^i$ ; so  $i$  is the index of the corresponding critical 2-jet. We can calculate in local coordinates  $(x, y', y'')$ , where  $x = (x_1, \dots, x_n)$  are local coordinates of  $M$ ,  $y' = (y'_1, \dots, y'_n)$  (resp.  $y'' = (y''_{jk})_{1 \leq j \leq k \leq n}$ ) are the associated coordinates of  $T_x^*M$  (resp.  $S^2T_x^*M$ ). Since  $f$  is holonomic, we have  $y''_{jk}(a) = \frac{\partial y'_j}{\partial x_k}(a)$ . Finally, the sign of  $\det y''(a)$  (positive if  $i$  is even and negative if not) gives the sign of the Jacobian determinant at  $a$  of the map  $x \mapsto y'(x)$ , that is the sign of the intersection point when  $\mathcal{M}_0^i$  is co-oriented by the canonical orientation of the  $y'$ -space. As we have reversed this co-orientation when  $i$  is odd, the intersection point is positive whatever the index is.  $\square$

**Proposition 2.3.** 1) *Each  $\mathcal{M}_0^i$  defines a degree  $n$  cocycle of  $\mathcal{M}$  with coefficients in the local system  $\mathbb{Z}^{or}$  of integers twisted by the orientation of  $M$ . Let  $\mu_i$  be its cohomology class in  $H^n(\mathcal{M}, \mathbb{Z}^{or})$ ; in particular, if  $s : M \rightarrow \mathcal{M}$  is a section,  $\langle \mu_i, [s] \rangle$  is an integer.*

2) *When  $s$  is homotopic to a holonomic section  $j^2f$ , then  $\langle \mu_i, [s] \rangle$  is positive and equals the number  $c_i(f)$  of critical points of the Morse function  $f$ . In particular the total number  $|Z|$  of zeroes of the section  $\pi \circ s$  (which, by construction, is transverse to the  $0$ -section) satisfies:*

$$|Z| \geq \sum_{i=0}^n c_i(f).$$

**Proof.** 1) Let  $\sigma$  be a singular  $n$ -cycle with twisted coefficients of  $\mathcal{M}$ . It can be  $C^0$ -approximated by  $\sigma'$ , an  $n$ -cycle which is transverse to  $\mathcal{M}_0^i$ . As  $\mathcal{M}_0^i$  is a proper submanifold, there are finitely many intersections points in  $\sigma' \cap \mathcal{M}_0^i$ , each one having a sign with respect to the local system

of coefficients. The algebraic sum of these signs defines an integer  $c(\sigma')$ . One easily checks that  $c(\sigma') = 0$  if  $\sigma'$  is a boundary. As a consequence, if  $\sigma'_0$  and  $\sigma'_1$  are two approximations of  $\sigma$ , as  $\sigma'_1 - \sigma'_0$  is a boundary, we have  $c(\sigma'_1) - c(\sigma'_0) = 0$  which allows us to uniquely define  $c(\sigma)$  as the value of an  $n$ -cocycle on  $\sigma$ . Typically, the image of a section carries an  $n$ -cycle with twisted coefficients and this algebraic counting applies.

2) Since  $c$  defined in 1) is a cocycle, it takes the same value on  $s$  and on  $j^2 f$ . According to lemma 2.2, it counts  $+1$  for each intersection point in  $j^2 f \cap \mathcal{M}_0^i$ , that is, for each index  $i$  critical point of  $f$ .  $\square$

**Corollary 2.4.** *If  $s$  is a section of  $\mathcal{M}$  which is homotopic to a holonomic section, the integers  $m_i := \langle \mu_i, [s] \rangle$  fulfill the Morse inequalities*

$$\begin{aligned} m_0 &\geq \beta_0(F) \\ m_1 - m_0 &\geq \beta_1(F) - \beta_0(F) \\ &\dots \\ m_0 - m_1 + \dots + (-1)^n m_n &= \beta_0(F) - \beta_1(F) \dots + (-1)^n \beta_n(F) =: \chi(M) \end{aligned}$$

where  $F$  is a field of coefficients,  $\beta_i(F) = \dim_F H_i(M, F^{or})$  is the  $i$ -th Betti number with coefficients in  $F^{or}$  ( $F$  twisted by the orientation) and  $\chi(M)$  is the Euler characteristic (independent of the field  $F$ ).

**Corollary 2.5.** *The  $h$ -principle does not hold true for the sections of  $\mathcal{M}$ .*

**Proof.** It is sufficient to construct a section  $s$  of  $\mathcal{M}$  which violates the Morse inequalities, for example a section which does not intersect  $\mathcal{M}_0^0$ . Leaving the case of the circle as an exercise, we may assume  $n > 1$ . One starts with a section  $s_1$  of  $T^*M$  tranverse to  $O_M$ . Each zero of  $s_1$  has a sign (if the local orientation of  $M$  is changed, so are both local orientations of  $s_1$  and  $O_M$  the sign of the zero in unchanged). For each zero  $a$ , one can construct a homotopy fixing  $a$ , with arbitrary small support, which makes  $s_1$  linear in a small neighborhood of  $a$ . As  $GL(n, \mathbb{R})$  has exactly two connected components, one can even suppose that after the homotopy,  $s_1$  is near  $a$  the derivative of a non degenerate quadratic function whose index can be chosen arbitrarily provided it is even (resp. odd) if  $a$  is a positive (resp. a negative) zero. Finally, one can achieve by homotopy that near each zero  $a$ , one has  $s_1 = df$  with  $a$  a non-degenerate critical point of  $f$  of index 2 (resp. 1) if  $a$  is a positive (resp. negative) zero.

Near the zeroes  $s_1$  has a canonical lift to  $\mathcal{M}$  by  $s_2 = j^2 f$ . Away from the zeroes, the lift  $s_2$  extends as a lift of  $s_1$  since the fibers of  $\pi$  are contractible over  $T^*M \setminus O_M$ . By construction, we have  $\langle \mu_0, [s_2] \rangle = 0$ , violating the first Morse inequality.  $\square$

**Remark 2.6.** *Denote  $\mu_{even} = \mu_0 + \mu_2 + \dots$  and  $\mu_{odd} = \mu_1 + \dots$ . The following statement holds true:  $\mu_{even} = \mu_{odd}$  if and only if the Euler characteristic vanishes.*

**Proof.** Assume first  $\mu_{even} = \mu_{odd}$ . Proposition 2.3 yields for any holonomic section in  $\mathcal{M}$ :  $m_{even} = m_{odd}$ , that is  $\chi(M) = 0$ . Conversely, if  $\chi(M) = 0$ , there exists a non-vanishing 1-form on  $M$  and hence, by lifting it to  $\tilde{\mathcal{J}}^2(M)$ , a section  $v_0$  in  $\mathcal{M}$  avoiding  $\mathcal{M}_0$ . We form

$$W = \{z \in \tilde{\mathcal{J}}^2(M) \mid z = z_0 + tv_0, z_0 \in \mathcal{M}_0, t \geq 0 \text{ or } z_0 \in \mathcal{D}, t > 0\}.$$

It is a proper submanifold in  $\mathcal{M}$  whose boundary (with orientation twisted coefficients) is  $\mathcal{M}_0^{\text{even}} - \mathcal{M}_0^{\text{odd}}$ . Therefore, every cycle  $c$  satisfies  $\langle \mu_{\text{even}}, c \rangle = \langle \mu_{\text{odd}}, c \rangle$ , which implies the wanted equality.  $\square$

### 3. ARE MORSE INEQUALITIES SUFFICIENT?

This question is closely related to the problem of minimizing the number of critical points of a Morse function. This problem was solved by S. Smale in dimension higher than 5 for simply connected manifolds, as a consequence of the methods he developed for proving his famous  $h$ -cobordism theorem (see [8] or chapter 2 in [2]). Under the same topological assumptions we can answer our question positively. But there are other cases, discussed later, where the answer is negative.

**Proposition 3.1.** *Two sections  $s, t$  of  $\mathcal{M} \subset \tilde{\mathcal{J}}^2(M)$  are homotopic as sections of  $\mathcal{M}$  if and only if their algebraic intersection numbers  $m_i$  with  $\mathcal{M}_0^i$  are the same.*

**Proof.** According to proposition 2.3 1), the condition is necessary. Let us prove that it is sufficient. Leaving the 1-dimensional case to the reader, we assume  $\dim M \geq 2$ . Denote  $s^1 = \pi \circ s$ . Each zero of  $s^1$  is given an index due to its lifting by  $s$  to a point of some  $\mathcal{M}_0^i$ . For each index  $i$  choose  $|m_i|$  zeroes of  $s^1$ ,  $a_i^1, \dots, a_i^{|m_i|}$ , among its zeroes of index  $i$ ; when  $m_i > 0$  (resp.  $m_i < 0$ ), we choose the  $a_i^j$  so that the corresponding intersection points of  $s(M)$  with  $\mathcal{M}_0^i$  are positive (resp. negative). When  $m_i = 0$ , no points are selected. In the same way,  $|m_i|$  zeros  $b_i^1, \dots, b_i^{|m_i|}$  of  $t^1$  are chosen.

The intersection signs being the same, one can find a homotopy of  $t$  in  $\mathcal{M}$ , which brings the  $b_i^j$  to coincide with the  $a_i^j$  and makes the two sections  $s$  and  $t$  coincide in the neighborhood of these points.

The other zeroes of  $s^1$  of index  $i$  can be matched into pairs of points  $\{a_i^{j+}, a_i^{j-}\}$  of opposite sign. A Whitney type lemma allows us to cancel all these pairs by a suitable homotopy of  $s$  in  $\mathcal{M}$ , reducing to the case when  $s^1$  has no other zeroes than the  $a_i^j$ 's,  $j = 1, \dots, |m_i|$ . A similar reduction may be assumed for  $t$ . Let us finish the proof in this case before stating and proving this lemma.

Both sections  $s^1$  and  $t^1$  of  $T^*M$  are homotopic (among sections) by a homotopy which is stationary on a neighborhood  $N(a_i^j)$ . Making this homotopy  $h : M \times [0, 1] \rightarrow T^*M$  transverse to the zero section, the preimage of  $0_M$  consists of arcs  $\{a_i^j\} \times [0, 1]$  and finitely many closed curves  $\gamma_k$ . Each of these closed curves can be arbitrarily decorated with an index  $i$ . This choice allows us to lift  $h$  to  $\tilde{\mathcal{J}}^2(M)$  as a homotopy  $\tilde{h}$  from  $s$  to  $t$ ; this  $\tilde{h}$  is the desired homotopy. More precisely, we proceed as follows for getting  $\tilde{h}$ . First  $h|_{\gamma_k}$  is lifted to  $\mathcal{M}_0^i$  by using that the fiber of  $\pi : \mathcal{M}_0^i \rightarrow 0_M$  is connected. The transversality of  $h$  to  $0_M$  allows us to extend this lifting to a neighborhood of  $\gamma_k$ , making  $\tilde{h}$  transverse to  $\mathcal{M}_0^i$ . Now it is easy to extend  $\tilde{h}$  to  $M \times [0, 1]$ , since the fiber of  $\pi$  over any point outside  $0_M$  is contractible.

**A WHITNEY TYPE LEMMA.** *Let  $(b^+, b^-)$  be a pair of transverse intersection points of  $s$  with  $\mathcal{M}_0^i$  having opposite sign when they are thought of as zeroes of  $s_1$  in  $M$ . Let  $\alpha$  be a simple path in  $M$  joining them avoiding the other zeroes of  $s^1$  and let  $N$  be a neighborhood of  $\alpha$ . Then there exists a homotopy  $S = (s_u)_{u \in [0, 1]}$  of  $s_0 = s$  into  $\mathcal{M}$ , supported in  $N$  and cancelling the*

pair  $(b^+, b^-)$ , that is,  $\pi \circ s_1$  has no zeroes in  $N$ .

**PROOF.** We choose an embedded 2-disk (with corners)  $\Delta$  in  $N \times [0, 1[$  meeting  $N \times \{0\}$  transversally along  $\alpha$ . We first construct the homotopy  $S^1 := \pi \circ S$  of  $s^1$  among the sections of  $T^*M$ , following the cancellation process of Whitney which we are going to recall. We require  $S^1$  to be transverse to  $0_M$  with  $(S^1)^{-1}(0_M) = \beta$ , where  $\alpha \cup \beta = \partial\Delta$ . Using a trivialization of  $T^*M|N$ ,  $S^1|N \times [0, 1]$  reads  $S^1(x, u) = (x, g(x, u))$ . The requirement is that  $g$  vanishes transversally along the arc  $\beta$ ; it is possible exactly because  $\dim M \geq 2$  and the end points have opposite signs. Let  $T$  be a small tubular neighborhood of  $\beta$ ; its boundary traces an arc  $\beta'$  on  $\Delta$ , “parallel” to  $\beta$ . Let  $\alpha'$  be the subarc of  $\alpha$  whose end points are those of  $\beta'$ . The restriction  $g|T$  is required to be a trivialization of  $T$ , but this latter may be chosen freely. We choose it so that the loop  $(g|\beta') \cup (s^1|\alpha')$  be homotopic to 0 in  $(\mathbb{R}^n)^* \setminus \{0\}$ ; of course, when  $n > 2$  this condition is automatically fulfilled. Now  $g$  can be extended to the rest of  $\Delta$  as a non-vanishing map. As  $N \times [0, 1]$  collapses onto  $N \times \{0\} \cup \Delta \cup T$ , the extension of  $g$  can be completed without adding zeroes outside  $\beta$ , yielding the desired homotopy  $S^1$ .

It remains to lift  $S^1$  to  $\mathcal{M}$ . The lifting is first performed along  $\beta$  with value in  $\mathcal{M}_0^i$ . Then it is globally extended in the same way as in the above lifting process.  $\square$

**Corollary 3.2.** *Let  $s$  be a section of  $\mathcal{M} \subset \tilde{\mathcal{J}}^2(M)$  and  $m_i$  be its algebraic intersection number with  $\mathcal{M}_0^i$ . Let  $f : M \rightarrow \mathbb{R}$  be a Morse function whose number  $c_i(f)$  of critical points of index  $i$  satisfies*

$$c_i(f) = m_i$$

for all  $i \in \{0, \dots, n\}$ . Then  $s$  and  $j^2(f)$  are homotopic as sections of  $\mathcal{M}$ .

**Corollary 3.3.** *We assume  $\dim M \geq 6$  and  $\pi_1(M) = 0$ . Let  $s$  be a section of  $\mathcal{M} \subset \tilde{\mathcal{J}}^2(M)$  whose algebraic intersection numbers  $m_i$  fulfills the Morse inequalities for every field of coefficients. In particular, they are non-negative. Then  $s$  is homotopic through sections in  $\mathcal{M}$  to a holonomic section.*

**Proof.** Under these topological assumptions the following result holds true: *For any set of non-negative integers  $\{c_0, c_1, \dots, c_n\}$  satisfying the Morse inequalities for any field of coefficients, there exists a Morse function on  $M$  with  $c_i$  critical points of index  $i$  (see theorem 2.3 in [2]).* So we have a Morse function  $f : M \rightarrow \mathbb{R}$  with  $m_i$  critical points of index  $i$ . According to corollary 3.2,  $s$  is homotopic in  $\mathcal{M}$  to  $j^2f$ .  $\square$

**3.4.** We end this section by recalling that the Morse inequalities are not sharp for estimating the number of critical points of a Morse function on a non-simply connected closed manifold. Typically when  $\pi_1(M)$  equals its subgroup of commutators (perfect group), some critical points of index 1 are required for generating the fundamental group, but the Morse inequalities allow  $c_1 = 0$  (see [7] for more details). On the other hand, the only constraint for a section of  $\mathcal{M}$  with intersection numbers  $m_i$  is the Euler-Poincaré identity:

$$m_0 - m_1 + \dots = \chi(M).$$



So it is possible to find a section  $s$  whose intersection number  $m_i$  is the minimal rank in degree  $i$  of a free complex whose homology is  $H_*(M, \mathbb{Z})$ , that is,

$$m_i = \beta_i + \tau_i + \tau_{i-1},$$

where  $\beta_i$  stands for the rank of the free quotient of  $H_i(M, \mathbb{Z})$  and  $\tau_i$  denotes the minimal number of generators of its torsion subgroup ([2] p. 15). Such a set of integers satisfies the Morse inequalities but is far from being realizable by a Morse function. Finally this section  $s$  is not homotopic in  $\mathcal{M}$  to a holonomic section.

#### 4. FAILURE OF THE 1-PARAMETRIC VERSION OF THE $h$ -PRINCIPLE

We thank Yasha Eliashberg who pointed out to us the failure of the  $h$ -principle in the 1-parametric version of the problem under consideration.

Here  $M$  is assumed to be a product  $M = N \times [0, 1]$ . Let  $f_0 : M \rightarrow [0, 1]$  be the projection. When  $M$  is not 1-connected and  $\dim M \geq 6$ , according to Allen Hatcher the so-called *pseudo-isotopy problem* has always a negative answer: there exists  $f$  without critical points which is not joinable to  $f_0$  among the Morse functions (see [4]). But  $j^2 f$  can be joined to  $j^2 f_0$  by a path  $\gamma$  in  $\mathcal{M}$ . Indeed, take a generic homotopy  $\gamma^1$  joining  $df$  to  $df_0$ ; then arguing as in the proof of proposition 3.1 it is possible to lift it to  $\mathcal{M}$ . When  $M$  is the  $n$ -torus  $\mathbb{T}^n$ , A. Douady showed very simply the stronger fact that the path  $\gamma^1$  can be taken among the non-singular 1-forms (see appendix to [5]). This  $\gamma$  is not homotopic in  $\mathcal{M}$  with end points fixed to a path of holonomic sections.

#### REFERENCES

- [1] Y. Eliashberg, N. Mishachev, *Introduction to the h-principle*, GSM 48, Amer. Math. Soc., 2002.
- [2] J. Franks, *Homology and Dynamical Systems*, CBMS Regional Conf. vol. 49, Amer. Math. Soc., 1982.
- [3] M. Gromov, *Partial Differential Relations*, Springer Verlag, 1986.
- [4] A. Hatcher, *Higher simple homotopy theory*, Annals of Math. (2) 102 (1975), 101-137.
- [5] F. Laudenbach, *Formes différentielles de degré 1 fermées non singulières : classes d'homotopie de leurs noyaux*, Commentarii Math. Helvetici 51 n°3 (1976), 447-464.
- [6] F. Laudenbach, *De la transversalité de Thom au h-principe de Gromov*, Leçons de Mathématiques d'Aujourd'hui, vol. 4, Éd. Cassini, Paris (to appear).
- [7] M. Maller, *Fitted diffeomorphisms of non-simply connected manifolds*, Topology 19 (1980), 395-410.
- [8] S. Smale, *Notes on differentiable dynamical systems*, 277-287 in: Proc. Symposia Pure Math., vol. 14, Amer. Math. Soc., 1970.
- [9] V.A. Vassiliev, *Topology of spaces of functions without compound singularities*, 23 (4) (1989), 277-286.

UNIVERSITÉ PARIS 7 ET IMCCE (UMR 8025 DU CNRS)  
*E-mail address:* `chenciner@imcce.fr`

UNIVERSITÉ DE NANTES, UMR 6629 DU CNRS, 44322 NANTES, FRANCE  
*E-mail address:* `francois.laudenbach@univ-nantes.fr`