



An analysis of fungal propagules transported to the *Henryk Arctowski* Antarctic Station

Anna AUGUSTYNIUK-KRAM^{1,2}, Katarzyna J. CHWEDORZEWSKA³, Małgorzata KORCZAK-ABSHIRE³, Maria OLECH^{3,4} and Maria LITYŃSKA-ZAJĄC⁵

¹ Centrum Badań Ekologicznych PAN w Dziekanowie Leśnym,
ul. M. Konopnickiej 1, 05-092 Łomianki, Poland <aaugustyniuk-kram@cbe-pan.pl>

² Instytut Ekologii i Bioetyki, Uniwersytet Kardynała Stefana Wyszyńskiego w Warszawie,
ul. Dewajtis 5, 01-815 Warszawa, Poland

³ Instytut Biochemii i Biofizyki PAN, Zakład Biologii Antarktyki,
ul. Pawińskiego 5a, 02-106 Warszawa, Poland

⁴ Instytut Botaniki, Uniwersytet Jagielloński, ul. Kopernika 27, 31-512 Kraków, Poland

⁵ Instytut Archeologii i Etnologii PAN w Krakowie, ul. Sławkowska 17, 31-016 Kraków, Poland

Abstract: During three austral summer seasons, dust and soil from clothes, boots and equipment of members of scientific expeditions and tourists visiting the Polish Antarctic Station *Henryk Arctowski* were collected and analysed for the presence of fungal propagules. Of a total of 60 samples, 554 colonies of fungi belonging to 19 genera were identified. Colonies of the genus *Cladosporium*, *Penicillium* and non-sporulating fungus (*Mycelia sterilia*) dominated in the examined samples. The microbiological assessment of air for the presence of fungi was also conducted at two points in the station building and two others outside the station. A total of 175 fungal colonies belonging to six genera were isolated. Colonies of the genus *Penicillium* were the commonest in the air samples. The potential epidemiological consequences for indigenous species as a result of unintentional transport of fungal propagules to the Antarctic biome are discussed in the light of rapid climate change in some parts of the Antarctic and adaptation of fungi to extreme conditions.

Key words: Antarctic, fungal propagules, unintentional transport.

Introduction

Despite the extreme conditions in the Antarctic, such as low temperature, high salinity, osmotic stress and high doses of UV radiation, microorganisms (including viruses, bacteria and fungi) are the most numerous groups of organisms that colonize diverse habitats (Abyzov 1993; Pearce and Wilson 2003; Onofri *et al.* 2007; Ruisi *et al.* 2007; Finster 2008). Among them, fungi are represented by in-

digenous species, including endemic species found only in the Antarctic, and show a variety of physiological and morphological adaptations to survive in such extreme conditions (Robinson 2001). With about 1,000 species of fungi reported from the Antarctic and sub-Antarctic region, only 2–3% are considered as endemic species (Bridge *et al.* 2008, 2010). Most of the remaining species are considered to be rather cosmopolitan ones, known from other regions of the world (Vishniac 1993; Selbmann *et al.* 2005; Ruisi *et al.* 2007).

An excellent vector for the transfer of fungal and other propagules to and within Antarctic regions is via humans and associated cargo, including food (Whinam *et al.* 2005; Chwedorzewska 2009; Hughes *et al.* 2011; Lityńska-Zajac *et al.* 2012; Chwedorzewska *et al.* 2013). Hughes *et al.* (2011) identified 19 species of fungi on rotting fruit and vegetables, also on the packaging and associated soil, most of which have previously been reported in different regions of the Antarctic. However, approximately 30% of them were not recorded previously from continental Antarctica and the Antarctic Peninsula. In addition, Antarctic regions have recently experienced a growing influx of tourists (Chwedorzewska and Korczak 2010), which also favors the introduction of non-native plant (Olech and Chwedorzewska 2011) and animal species (Chwedorzewska *et al.* 2013), but also microorganisms such as bacteria and fungi, including pathogens that may threaten native organisms (Mercantini *et al.* 1993; Rogers *et al.* 2004; Cowan *et al.* 2011).

The aim of this study was to investigate the extent of the transfer of fungal propagules into King George Island (where the Polish Antarctic Station is located) on clothes, boots and equipment arriving each year in association with scientists and tourists. This is the first project allowing for qualitative and quantitative assessment of transmissible fungal spores to the Antarctic biome.

Materials and methods

The samples were collected during three austral summer seasons: 2007/08, 2008/09 and 2009/10. Members of scientific expeditions and tourists (only one season 2007/08) arriving at the Polish Antarctic Station *Henryk Arctowski* (King George Island, South Shetland Islands, 62°09'S, 58°28'W) were checked for the presence of fungi. In season 2007/08 and 2009/10 samples were taken, respectively, from 20 and 21 people. All outdoor clothing and equipment (bags, backpacks) were thoroughly vacuumed each into a separate standard synthetic vacuum bag. Each sample was tagged, placed in a separate sterile zip lock bag and preserved (by cooling to 4°C) for transportation to Poland for further analysis. In season 2008/09 samples were collected from 19 people. During this season soil, mainly from boots, clothing and other equipment, was collected. Tourists were vacuuming on the ship before going down to land, while the members of the expedition immediately after their descent to land.




Fig. 1. Map of the study area showing sites used for aerobiological sampling: moraines, penguin rookery (*) and *Arctowski* Station.

Due to a small amount of the dust sample (dust was embedded in the structure of the material), of each vacuum bag three squares were cut out with sterile scissors, each with an area of 1 cm². Then the squares were placed separately in test tubes with 10 ml aqueous solution of Triton-X (0.05%), vortexed at maximum speed for 5 min and 0.1 ml plated onto agar medium. Squares cut out from the “clear bags” (bags not used for vacuuming) were treated as a control. Fungi from soil samples were isolated using the standard dilution plate method (Pepper *et al.* 1995). Because different groups of fungi require different conditions for optimal growth (different carbon and nitrogen source, different pH), four isolation media were used for all samples: PDA (Potato Dextrose Agar), MEA (Malt Extract 4% Agar), RBA (Rose Bengal Agar with chloramphenicol) and SDA (Sabourauda Dextrose 4% Agar with chloramphenicol). Each sample was tested using three plates of each of the four media and incubated at 22°C for 5–7 days. Fungal isolates were identified to genus level using morphological taxonomic keys (Gilman 1959; Barnett and Hunter 1972; Domsch *et al.* 1980).

During the 2008/09 season, mycological analysis of the air in the vicinity and inside the station was also conducted. Air from inside of the station was sampled at two points – room in the living quarters and storeroom, and at two points outside the station – in the moraines and penguin rookery (Fig. 1). Airborne fungal spores

were determined using the Koch sedimentation method in accordance with a Polish Standard (PN-89Z-04111/03). Air microorganisms were settled gravitationally, directly on open Petri dishes with Sabouraud dextrose agar (Envirocheck Settle Plate, Merck). At each point 20 plates were used. Then the plates were incubated at 25°C for 5 to 7 days. Concentration of fungal spores in 1 m³ of the air determined as colony forming units (CFU) was calculated using Omeliański formula modified by Gogoberidze (PN-89Z-04111/03).

$$\text{CFU/m}^3 = \bar{a} \times 10^4/p \times t \times 0.2$$

\bar{a} – average number of fungal colonies on the Petri plate

p – the surface of the Petri plate (in cm)

t – the time of the Petri plate exposure (in minutes)

0.2 – exposure time conversion factor of Petri plates

Results

During the 2007/08 season, 20 samples taken from people's clothing and equipment were analyzed and in total 189 fungal colonies (isolates) belonging to 11 genera were found, in the 2008/09 (19 samples) 252 colonies were represented by 14 genera, and in the 2009/10 season (21 samples) 113 colonies belonging to eight genera were found (Table 1).

Fungi were isolated from almost all the samples. No fungi were isolated only from one sample collected during the 2008/09 season and one other collected during the 2009/10 season, which represents approximately 5% of all analyzed samples. Colonies of the genus *Cladosporium* (100 isolates) were the commonest in the examined samples in the 2007/08 season, while, during the 2008/09 and 2009/10 seasons, colonies of *Penicillium* were the most commonly isolated (103 and 46 isolates respectively). *Alternaria*, *Geotrichum*, *Aspergillus* and non-sporulating fungi (*Mycelia sterilia*) were also abundant in the samples. Fungi of other genera were found occasionally (Table 1).

The most frequently isolated fungus for each of the seasons was also observed in the largest number of samples. The only exception was the genus *Alternaria* in the 2007/08 season, where 35 colonies came from only two samples (representing 10% of all analyzed samples in this season). In general, the most commonly isolated fungi were detected in more than 50% of the samples (Table 2).

A total of 175 fungal colonies belonging to six genera were isolated from the air samples at the four sampling sites (Fig. 1, Table 3). Yeast-like fungi and non-sporulating fungi (*M. sterilia*) were also identified. Both in the room and the storeroom *Penicillium* was the most represented genus, with 16 and 78 colonies respectively. In the storeroom colonies of yeast-like fungi were also abundant (47 colonies). At the moraines sampling site (Table 3) only one colony of the genus

Table 1
The number of isolates of different fungal genera from four different media in three seasons.

genus	season		
	2007/2008	2008/2009	2009/2010
<i>Acremonium</i>		1	
<i>Alternaria</i>	35		15
<i>Aspergillus</i>	5	11	12
<i>Botritis</i>	2	3	
<i>Cladosporium</i>	100	18	1
<i>Chaetomium</i>	8		
<i>Chloridium</i>	6	3	
<i>Fusarium</i>		2	
<i>Geotrichum</i>	1	52	1
<i>Hyalodendron</i>		1	
<i>Monocillium</i>	1	1	
<i>Mucor</i>	4		13
<i>Penicillium</i>	11	103	46
<i>Torula</i>		3	
<i>Trichocladium</i>			5
<i>Trichoderma</i>	1	1	
<i>Scopulariopsis</i>		5	
<i>Verticillium</i>		2	
<i>Mycelia sterilia</i>	15	46	20
Total	189	252	113

Table 2
The frequency of isolation of most often detected fungi.

season	most frequently isolated fungi	no of samples (no of positive samples*)	no of isolates	frequency of isolation (%)**
2007/2008	<i>Cladosporium</i>	20 (15)	100	75
	<i>Alternaria</i>	20 (2)	35	10
	<i>Mycelia sterilia</i>	20 (5)	15	25
	<i>Penicillium</i>	20 (6)	11	30
2008/2009	<i>Penicillium</i>	19 (15)	103	79
	<i>Geotrichum</i>	19 (12)	52	63
	<i>Mycelia sterilia</i>	19 (14)	46	74
	<i>Cladosporium</i>	19 (11)	18	58
	<i>Aspergillus</i>	19 (8)	11	42
2009/2010	<i>Penicillium</i>	21 (14)	46	67
	<i>Mycelia sterilia</i>	21 (9)	20	43
	<i>Alternaria</i>	21 (3)	15	14
	<i>Mucor</i>	21 (2)	13	10
	<i>Aspergillus</i>	21 (10)	12	48

* – number of samples from which the fungus was isolated

** – the ratio of the number of positive samples to the total number of samples

Table 3

The number of fungal colonies isolated from the air inside and outside the station.

genus	sampling site			
	room	storeroom	moraines	penguin rookery
<i>Alternaria</i>	1			
<i>Aspergillus</i>	1			
<i>Bispora</i>		1		
<i>Cladosporium</i>	1			
<i>Penicillium</i>	16	78	1	
<i>Verticillium</i>		1		
<i>Mycelia sterilia</i>	15	2	1	
Yeast-like fungi	10	47		
Total	44	129	2	0
CFU/m ³	1.7×10^2	5.1×10^2	0.8×10^1	0

Penicillium and one belonging to *M. sterilia* was isolated. At the penguin rookery sampling site no fungal colonies were isolated.

Discussion

This study shows that fungal spores can be transported to the Antarctic biome on people's clothing and expedition equipment in a rapid manner. The most common fungal genera were *Penicillium*, *Cladosporium*, *Alternaria* and *Geotrichum*. These are cosmopolitan and ubiquitous fungi, widespread in nature, occurring in many regions of the world, including alpine and polar areas (Bridge *et al.* 2010). They grow well on many substrates including decaying organic material, plant debris, soil and food stuffs, but also occur in the air as part of the bio-aerosol (Domsch *et al.* 1980; Marshall 1997). Therefore, the question is whether fungi introduced unintentionally to Antarctic regions, on personnel clothing and equipment, should be considered as alien species? It is difficult to answer this question. Numerous mycological studies conducted in different regions of the Antarctic show that fungi of the genus *Cladosporium*, *Penicilium*, *Aspergillus* are found relatively frequently in soil and air, and are even isolated from within deep layers of polar ice that were originally deposited several thousands years ago (Del Frate and Caretta 1990; Baublis *et al.* 1991; Marshall 1997; Azmi and Seppelt 1998; Tosi *et al.* 2002; Arenz *et al.* 2006; D'Elia *et al.* 2009; Rosa *et al.* 2009; Zucconi *et al.* 2012). However, different strains of the same fungal species (strains from both within and outside the Antarctic) may have different physiological characteristics and unknown consequences on the Antarctic indigenous species and ecosystems (Litchman 2010; Cowan *et al.* 2011). Nevertheless, the fungi which are accidentally introduced to colonize a new environment, besides viable propagules, must, above all, have the ecophysiological characteristics required for survival in the polar environment and to allow the species to grow and establish (Ellis-Evans and Walton 1990; Pearce *et al.* 2009).

During the austral summer the mean air temperature in the vicinity of *Arctowski* Station may reach 2.5°C (maximum 10.4°C, minimum -1.3°C) with wide daily fluctuations including temperatures dropping below freezing (Kejna 2008; Kejna *et al.* 2013). Recently, the Antarctic Peninsula is one of the fastest warming regions on the Earth. The largest annual warming trends were found in the western and northern parts of the Antarctic Peninsula, with temperatures at *Faraday/Vernadsky* Station increasing at a rate of +0.56°C over the decade in the period 1951–2000 (Turner *et al.* 2005). During sunny days the soil surface can heat up to temperatures of 10–15°C or even higher (Davey *et al.* 1992), and thus create optimal conditions for the development of some non-native (considered as psychrophilic) species of the genus *Penicillium* or *Cladosporium*. However, most of the unintentionally introduced fungi are either psychrotolerant or psychrotrophs, and even mesophilic species. Most filamentous fungi have a short reproductive cycle that can last only a few days and their life cycle can be completed by the production of a vast number of spores that can spread, but it is rather unlikely for them to significantly contribute to any aspect of nutrient cycling and to have any impact on the microbiological community structure or function, owing to their low metabolic activities (Cowan *et al.* 2011).

In the examined samples non-sporulating colonies *M. sterilia* were found relatively often. During the 2008/09 season, 46 sterile colonies were isolated from 14 of the 19 soil samples (74% of the total). It is supposed that the formation of sterile mycelia is one of the mechanisms of fungal adaptation to low temperatures or to a lack of nutrients (Robinson 2001). Other adaptation of fungi to extreme conditions may also shorten the life cycle or, conversely, lengthen it. For example, *Penicillium hirsutum* (growing on garlic in storage) showed both delayed germination of spores and sporulation at lower temperatures (from 4 to -2°C) than in higher ones (10 and 20°C), whereas at a temperature of -4°C it showed a total loss of sporulation (Bertolini and Tian 1996). Reduction of metabolism could be the first step of adaptation to extreme condition, while physiological mechanisms conferring cold tolerance in fungi are more complex (Robinson 2001). Fungi *M. sterilia* are often isolated from various habitats and regions, within both the Arctic (Hyvärinen *et al.* 2001; Salonen *et al.* 2007) and the Antarctic. Azmi and Seppelt (1998) in the Windmill Islands region found that the non-sporulating fungi *M. sterilia* were the most frequently isolated fungi from samples of soil, mosses, algae and lichens. Similarly, in earlier studies conducted by Fletcher *et al.* (1985) in Mac. Robertson and Enderby Lands (East Antarctica), the sterile mycelial fungi comprised respectively 47 and 60% of isolates.

The austral summer, from December to March, is a period of increased influx of tourism and research groups in some parts of the Antarctic. Human activities are focused mainly on ice-free areas of the maritime Antarctic, where there are around half of the research stations, and where climatic conditions are not as severe as in the continental regions. Thus, the fungal spores unintentionally introduced into the polar stations may find appropriate conditions for development. The high density of fungal spores inside the station, where there are good conditions for their development

(i.e. on food waste), shows that these places can be a good source for them to spread to the nearby surroundings. Mycological analysis of the air was carried out in the vicinity and inside the *Arctowski* Station previously in 1985 and 1999 (Czarnecki and Białasiewicz 1987; Białasiewicz and Czarnecki 1999). These studies showed that fungi of the genus *Penicillium* were the most frequently observed inside the station, which supports other research performed at *Syowa* Station (Nakashima *et al.* 2003). In our study we received a total of 94 isolates of *Penicillium* genus, of which 78 were isolated from the fruits and vegetables storeroom. As Białasiewicz and Czarnecki (1999) suggested, the source of fungi collected inside the station could be brought by human vectors, as well as originate from the natural environment outside the station. However, other studies have also indicated that research stations and human activity in the vicinity of research stations may be a source of microorganisms released into the local Antarctic environment (Hughes *et al.* 2010; Pearce *et al.* 2010).

The fact of unintentional transport of fungal propagules along with people arriving at the polar stations, regardless of whether these fungal species are considered to be alien or not, is a dangerous phenomenon since many of the species recorded in the studied samples are potentially pathogenic to plants (Hughes *et al.* 2011) and warm-blooded animals (Wicklow 1968; Mercantini *et al.* 1989). For many indigenous populations of the Antarctic flora and fauna, geographical isolation is the reason why fungi associated with human activities in the Antarctic can be novel pathogens. Thus, the flora and fauna of the polar regions may be particularly sensitive to infections, which could have devastating consequences on the indigenous biota (Rogers *et al.* 2004; Weimerskirch 2004; Barbosa and Palacios 2009).

Acknowledgments. — This research project was supported by the Ministry of Scientific Research and Higher Education by a grant IPY/27/2007. The authors would like to thank all persons involved in collecting materials during XXX, XXXI and XXXII Polish Antarctic Expeditions. We would like to thank two anonymous reviewers for constructive advice that has improved our paper.

References

- ABYZOV S.S. 1993. Microorganisms in the Antarctic ice. In: E.I. Friedman (eds) *Antarctic Microbiology*. Wiley-Liss, Inc., New York: 265–285.
- ARENZ B.E., HELD B.W., JURGENS J.A., FARRELL R.L and BLANCHETTE R.A. 2006. Fungal diversity in soils and historic wood from the Ross Sea Region of Antarctica. *Soil Biology and Biochemistry* 38: 3057–3064.
- AZMI O.R. and SEPPELT R.D. 1998. The broad-scale distribution of microfungi in the Windmill Islands region, continental Antarctica. *Polar Biology* 19: 92–100.
- BARBOSA A. and PALACIOS M.J. 2009. Health of Antarctic birds: a review of their parasites, pathogens and diseases. *Polar Biology* 32: 1095–1115.
- BARNETT H.L. and HUNTER B.B. 1972. *Illustrated genera of Imperfect Fungi*. 3rd Ed., Burgess Publishing Co., Minneapolis: 241 pp.
- BAUBLIS J.A., WHARTON R.A. and VOLZ P.A. 1991. Diversity of micro-fungi in an Antarctic dry valley. *Journal of Basic Microbiology* 31: 3–12.
- BERTOLINI P. and TIAN S.P. 1996. Low temperature biology and pathogenicity of *Penicillium hirsutum* on garlic in storage. *Postharvest Biology and Technology* 7: 83–89.

- BIAŁASIEWICZ D. and CZARNECKI B. 1999. Microfungi in the aerosphere of the *Arctowski* Polar Station. *Polish Polar Research* 20: 319–324.
- BRIDGE P.D., SPOONER B.M. and ROBERTS P.J. 2008. Non-lichenized fungi from the Antarctic region. *Mycotaxon* 106: 485–490.
- BRIDGE P.D., SPOONER B.M. and ROBERTS P.J. 2010. *List of non-lichenized fungi from the Antarctic region* <www.antarctica.ac.uk/bas_research/data/access/fungi/> (accessed November 2012).
- CHWEDORZEWSKA K. 2009. Terrestrial Antarctic Ecosystems at the Changing World – an overview. *Polish Polar Research* 30: 263–273.
- CHWEDORZEWSKA K.J. and KORCZAK M. 2010. Human impact upon the environment in the vicinity of *Arctowski* Station, King George Island, Antarctica. *Polish Polar Research* 31: 45–60.
- CHWEDORZEWSKA K.J., KORCZAK-ABSHIRE M., OLECH M., LITYŃSKA-ZAJĄC M. and AUGUSTYNIUK-KRAM A. 2013. Alien invertebrates transported accidentally to the Polish Antarctic Station in cargo and on fresh foods. *Polish Polar Research* 34: 55–66.
- COWAN D.A., CHOWN S.L., CONVEY P., TUFFIN M., HUGHES K., POINTING S. and VINCENT W.F. 2011. Non-indigenous microorganisms in the Antarctic: assessing the risks. *Trends in Microbiology* 19: 540–548.
- CZARNECKI B. and BIAŁASIEWICZ D. 1987. Fungi as a component of the aerosphere in the *H. Arctowski* Polar Station and its vicinity (King George Island, South Shetland Islands). *Polish Polar Research* 8: 153–158.
- D'ELIA T., VEERAPANENI R., HERAISNATHAN V. and ROGERS S.O. 2009. Isolation of fungi from Lake Vostok accretion ice. *Mycologia* 101: 751–763.
- DAVEY M.C., PICKUP J. and BLOCK W. 1992. Temperature variation and its biological significance in fellfield habitats on a maritime Antarctic island. *Antarctic Science* 4: 383–388.
- DEL FRATE G. and CARETTA G. 1990. Fungi isolated from Antarctic material. *Polar Biology* 11: 1–7.
- DOMSCH K.H., GAINS W. and ANDERSON T.H. 1980. *A compendium of soil fungi*. Academic Press, London: 859 pp.
- ELLIS-EVANS J.C. and WALTON D. 1990. The process of colonization in Antarctic terrestrial and freshwater ecosystems. *Proceedings of the National Institute of Polar Research Symposium on Polar Biology* 3: 151–163.
- FINSTER K. 2008. Anaerobic bacteria and Archaea in cold ecosystems. In: R. Margesin, F. Schinner J.-C. Marx and C. Gerday (eds) *Psychrophiles: from Biodiversity to Biotechnology*. Springer-Verlag, Berlin, Heidelberg: 103–119.
- FLETCHER L.D., KERRY E.J. and WESTE G.M. 1985. Microfungi of Mac. Robertson and Enderby Lands, Antarctica. *Polar Biology* 4: 81–88.
- GILMAN J.C. 1959. *A manual of soil fungi*. Constable and Company Ltd, London: 450 pp.
- HUGHES K.A., CONVEY P., MASLEN N.R. and SMITH R.I.L. 2010. Accidental transfer of non-native soil organisms into Antarctica on construction vehicles. *Biological Invasions* 12: 875–891.
- HUGHES K.A., LEE J.E., TSUJIMOTO M., IMURA S., BERGSTROM D.M., WARE C., LÉBOUVIER M.L., HUISKES A.H.L., GREMMEN N.J.M., FRENOT Y., BRIDGE P.D. and CHOWN S.L. 2011. Food for thought: Risks of non-native species transfer to the Antarctic region with fresh produce. *Biological Conservation* 144: 1682–1689.
- HYVÄRINEN A., REPONEN T., HUSMAN T. and NEVALAINEN A. 2001. Comparison of the indoor air quality in mould damaged and reference buildings in a subarctic climate. *Central European Journal of Public Health* 9: 133–139.
- KEJNA M. 2008. Topoclimatic conditions in the vicinity of the *Arctowski* Station (King George Island, Antarctica) during the summer season of 2006/2007. *Polish Polar Research* 29: 95–116.
- KEJNA M., ARAŻNY A. and SOBOTA I. 2013. Climatic change on King George Island in the years 1948–2011. *Polish Polar Research* 34: 213–235.
- LITCHMAN E. 2010. Invisible invaders: non-pathogenic invasive microbes in aquatic and terrestrial ecosystems. *Ecology Letters* 13: 1560–1572.
- LITYŃSKA-ZAJĄC M., CHWEDORZEWSKA K.J., OLECH M., KORCZAK-ABSHIRE M. and AUGUSTYNIUK-KRAM A. 2012. Diaspores and phyto-remains accidentally transported to the Antarctic Station during three expeditions. *Biodiversity and Conservation* 21: 3411–3421.

- MARSHALL W.A. 1997. Seasonality in Antarctic Airborne Fungal Spores. *Applied and Environmental Microbiology* 63: 2240–2245.
- MERCANTINI R., MARSELLA R. and CERVELATTI M.C. 1989. Keratinophilic fungi isolated from Antarctic soil. *Mycopathologia* 106: 47–52.
- MERCANTINI R., MARSELLA R., MORETTO D. and FINOTTI E. 1993. Keratinophilic fungi in the antarctic environment. *Mycopathologia* 122: 169–175.
- NAKASHIMA H., HAMADA N. and YAMANOUCHU T. 2003. Airborne microorganisms in the indoor environment of Syowa Station in Antarctica. *Polar Meteorology and Glaciology* 17: 61–67.
- OLECH M. and CHWEDORZEWSKA K.J. 2011. The first appearance and establishment of alien vascular plant in natural habitats on the forefield of retreating glacier in Antarctica. *Antarctic Science* 23: 153–154.
- ONOFRI S., SELBMANN L., DE HOOG G.S., GRUBE M., BARRECA D., RUISI S. and ZUCCONI L. 2007. Evolution and adaptation of fungi at boundaries of life. *Advances in Space Research* 40: 1657–1664.
- PEARCE D.A., BRIDGE P.D., HUGHES K.A., SATTTLER B., PSENNER R. and RUSSELL N.J. 2009. Microorganisms in the atmosphere over Antarctica. *FEMS Microbiology Ecology* 69: 143–157.
- PEARCE D.A., HUGHES K.A., LACHLAN-COPE T., HARANGOZO S.A. and JONES A.E. 2010. Biodiversity of air-borne microorganisms at Halley station, Antarctica. *Extremophiles* 14: 145–159.
- PEARCE D.A. and WILSON W.H. 2003. Viruses in Antarctic ecosystems. *Antarctic Science* 15: 319–331.
- PEPPER I.L., GERBA C.P. and BRENDENCKE J.W. 1995. Filamentous fungi. In: P. Gerba, J. Brendencke, D.C. Johnson, K.L. Josephson, H.L. Bohn and J. Tanguay (eds) *Environmental microbiology. A laboratory manual*. Academic Press, San Diego: 11–20.
- ROBINSON C.H. 2001. Cold adaptation in Arctic and Antarctic fungi. *New Phytologist* 151: 341–353.
- ROGERS S.O., STARMER W.T. and CASTELLO J.D. 2004. Recycling of pathogenic microbes through survival in ice. *Medical Hypotheses* 63: 773–777.
- ROSA L.H., VAZ A.B.M., CALIGORNE R.B., CAMPOLINA S. and ROSA C.A. 2009. Endophytic fungi associated with the Antarctic grass *Deschampsia antarctica* Desv. (Poaceae). *Polar Biology* 32: 161–167.
- RUISI S., BARRECA D., SELBMANN L., ZUCCONI L. and ONOFRI S. 2007. Fungi in Antarctica. *Reviews in Environmental Science and Biotechnology* 6: 127–141.
- SALONEN H., LAPPALAINEN S., LINDROOS O., HARJU R. and REIJULA K. 2007. Fungi and bacteria in mould-damaged and non-damaged office environments in a subarctic climate. *Atmospheric Environment* 41: 6797–6807.
- SELBMANN L., DE HOOG G.S., MAZZAGLIA A., FRIEDMANN E.I. and ONOFRI S. 2005. Fungi at the edge of life: cryptoendolithic black fungi from Antarctic desert. *Studies in Mycology* 51: 1–32.
- TOSI S., CASADO B., GERDOL R. and CARETTA G. 2002. Fungi isolated from Antarctic mosses. *Polar Biology* 25: 262–268.
- TURNER J., COLWELL S.R., MARSHALL G.J., LACHLAN-COPE T.A., CARLETON A.M., JONES P.D., LAGUN V., REID P.A. and IAGOVKINA S. 2005. Antarctic climate change during the last 50 years. *International Journal of Climatology* 25: 279–294.
- VISHNIAC H.S. 1993. The microbiology of Antarctic soils. In: E.I. Friedmann (ed.) *Antarctic microbiology*. Wiley, New York: 297–341.
- WEIMERSKIRCH H. 2004. Diseases threaten Southern Ocean albatrosses. *Polar Biology* 27: 374–379.
- WHINAM J., CHILCOTT N. and BERGSTROM D.M. 2005. Subantarctic hitchhikers: expeditioners as vectors for the introduction of alien organisms. *Biological Conservation* 121: 207–219.
- WICKLOW D.T. 1968. *Aspergillus fumigatus* Fresenius isolated from ornithogenic soil collected at Hallett Station, Antarctica. *Canadian Journal of Microbiology* 14: 717–719.
- ZUCCONI L., SELBMANN L., BUZZINI P., TURCHETTI B., GUGLIELMIN M., FRISVAD J.C. and ONOFRI S. 2012. Searching for eukaryotic life preserved in Antarctic permafrost. *Polar Biology* 35: 749–757.

Received 15 September 2012

Accepted 10 April 2013