

Mycorrhiza (2012) 22:493–499
DOI 10.1007/s00572-011-0427-6

ORIGINAL PAPER

Effects of genetic modifications to flax (*Linum usitatissimum*) on arbuscular mycorrhiza and plant performance

Magdalena Wróbel-Kwiatkowska · Katarzyna Turnau ·
Katarzyna Góralska · Teresa Anielska · Jan Szopa

Received: 22 August 2011 / Accepted: 14 December 2011 / Published online: 5 January 2012

© The Author(s) 2012. This article is published with open access at Springerlink.com

Abstract Although arbuscular mycorrhizal fungi (AMF) are known for their positive effect on flax growth, the impact of genetic manipulation in this crop on arbuscular mycorrhiza and plant performance was assessed for the first time. Five types of transgenic flax that were generated to improve fiber quality and resistance to pathogens, through increased levels of either phenylpropanoids (W92.40), glycosyltransferase (GT4, GT5), or PR2 beta-1,3-glucanase (B14) or produce polyhydroxybutyrate (M50), were used. Introduced genetic modifications did not change the degree of mycorrhizal colonization as compared to parent cultivars Linola and Nike. Arbuscules were well developed in each tested transgenic type (except M50). In two lines (W92.40 and B14), a higher abundance of arbuscules was observed when compared to control, untransformed flax plants. However, in some cases (W92.40, GT4, GT5, and B14 Md), the mycorrhizal dependency for biomass production of transgenic plants was slightly lower when compared to the original cultivars. No significant influence of mycorrhiza on the photosynthetic activity of transformed lines was found, but in most cases P concentration in mycorrhizal plants remained

higher than in nonmycorrhizal ones. The transformed flax lines meet the demands for better quality of fiber and higher resistance to pathogens, without significantly influencing the interaction with AMF.

Keywords Mycorrhiza · *Linum usitatissimum* · Transgenic flax · Plant performance

Introduction

Although genetically modified plants (GMPs) have already been accepted by several countries for use in the open field (Liu 2010), there is still a paucity of knowledge on the potential hazards of releasing such organisms into nature, including their potential impact on non-target microbiota (Giovannetti et al. 2005; Liu 2010; Stefani and Hamelin 2010). Arbuscular mycorrhizal fungi (AMF) are among the most common microbiota in soils, and plant dependency on AMF communities is essential and extensive in both natural and disturbed habitats (Van der Heijden et al. 1998; Smith and Read 2008). AMF can be crucial in low-input agriculture, while in strongly fertilized soil they can be totally ineffective or absent (Boyetchko 1996; He and Nara 2007).

As recently reviewed by Stefani and Hamelin (2010), GMPs can alter or not affect mycorrhizal development or colonization rates. Negative effects have been observed in Bt transgenic rice (Ren 2006), corn, and defensin-expressing aubergine (Turrini et al. 2004a). This involved reduced presymbiotic hyphal growth and development of appressoria (Turrini et al. 2004a) and reduced root colonization and spore density in rhizosphere soil (Ren 2006). Over half of the studies showed no effect of Bt transgenic plants on AMF, but to avoid potential risks it nevertheless seems crucial to test every newly developed GMP for its

M. Wróbel-Kwiatkowska
Department of Pharmaceutical Biology and Botany,
Medical University of Wrocław,
Al. J. Kochanowskiego 10,
51-601 Wrocław, Poland

K. Turnau (✉) · K. Góralska · T. Anielska
Institute of Environmental Sciences, Jagiellonian University,
Gronostajowa 7,
30-387 Kraków, Poland
e-mail: katarzyna.turnau@uj.edu.pl

J. Szopa
Faculty of Biotechnology, University of Wrocław,
Przybyszewskiego 63/77,
51-148 Wrocław, Poland

effect on arbuscular mycorrhizal association before released into the field.

Genetically modified (GM) flax lines have been created as a model for studying the role of antioxidants and to improve antifungal resistance and retting efficiency for the production of fiber-rich material (Szopa et al. 2009; Gredes et al. 2010; Skórkowska-Telichowska et al. 2010, Czemplik et al. 2011). Studies on the effect of genetic modifications of flax on plant–AMF interactions are important as GM flax starts to be cultivated in the field and the commercial demand for this crop is large. GM flax lines pose little environmental risk because their seed production and adventitious presence among subsequent crops and yield are relatively easy to control (Jhala et al. 2010). Nevertheless, mechanisms that the plant uses against pathogens can have deleterious effects also on mutualistic symbionts (Whipps 2004). Thus, the aims of the present study were to investigate the effect of different genetic modifications of flax (*Linum usitatissimum*) on root colonization by *Glomus intraradices* and compare mycorrhizal dependency of individual lines.

Materials and methods

Five types of transgenic flax (Table 1) were used in the present study. The transgenic lines have been generated from Linola and Nike cultivars and characterized previously (Wróbel et al. 2004; Wróbel-Kwiatkowska et al. 2004; Łukaszewicz et al. 2004a, b; Lorenc-Kukuła et al. 2005, 2009). Three have increased levels of either phenylpropanoids (W92.40), pathogenesis-related PR-2 protein (B14), or β -hydroxybutyrate polymer (M50). W92 flax plants (*L. usitatissimum* cv. Linola) were obtained by overexpression of three *Petunia hybrida* genes from the flavonoid biosynthesis pathway encoding chalcone synthase, chalcone isomerase, and dihydroflavonol reductase. The B14 transgenic plants are obtained from fibrous cultivar of flax (Nike) and

have ectopic expression of a defense-related potato β -1,3-glucanase gene (PR-2). M50 plants are also derived from cultivar Nike and transformed with *Ralstonia eutropha* genes coding for β -ketothiolase (phbA), acetoacetyl-CoA reductase (phbB), and PHB synthase (phbC) for poly- β -hydroxybutyrate (PHB) synthesis. Two other plant lines (GT4, GT5) derived from Linola cultivar overproduce glycosyltransferase, resulting in the accumulation of proanthocyanin, lignan, phenolic acid, and unsaturated fatty acids in seeds. Plants of an MB type were also obtained, by classical crossing of lines M and B. All these transgenic flax lines show improved resistance to common flax pathogens such as *Fusarium oxysporum* and/or *Fusarium culmorum*, which cause diseases such as vascular wilt, corm rot, root rot, or damping-off (Armstrong and Armstrong 1981; Jones 1991).

Culture conditions of mycorrhizal and nonmycorrhizal plants

Seeds of the GM flax lines were germinated on wet filter paper in Petri dishes at approximately 20°C. Five-day old seedlings ($n=5$ to 9 depending on the line) were transferred into 800 ml \emptyset 13-cm pots with sterile substratum composed of a mixture of garden soil (Rolex, containing peat, sand, pine bark, compost enriched with organic fertilizers and dolomite), sand, and expanded clay in the ratio 3:3:2 (v/v/v). The substratum was amended with 25 g per kg of rock phosphate (SIARKOPOL, Tarnobrzeg). Prior to seedling transfer, ca. 15 g of AM fungal inoculum containing propagules of *G. intraradices* UNIJAG.PL24-1 isolated from a grassland close to Inowroclaw (Poland) was introduced below the seedlings. Inoculum was produced in 3-L pots of sterile substratum (sand/expanded clay 2:1, v/v) with *Zea mays* and *Plantago lanceolata* as host plants. After 4 months of cultivation, the shoots were harvested and the inoculum was dried for 30 days. Inoculum quality was assessed on *P. lanceolata*

Table 1 List of transgenic plants investigated

Transgenic line	Origin of gene	Type of gene and gene construct	Method of gene introduction and selection	References
W92.40	<i>Petunia hybrida</i>	CHS, CHI, DFR, triple gene construct, each under 35S promoter	<i>Agrobacterium</i> mediated transformation, PCR, Northern blot	Lorenc-Kukuła et al. 2005
GT4, GT5	<i>Solanum soganandinum</i>	SsGT1 gene under nap promoter	<i>Agrobacterium</i> mediated transformation, PCR, Northern blot	Lorenc-Kukuła et al. 2009
M50	<i>Ralstonia eutropha</i>	phbA, phbB, phbC, triple gene construct, phbB and C genes under 35S, phbA under 14-3-3 promoter	<i>Agrobacterium</i> mediated transformation, PCR, Northern blot	Wróbel et al. 2004
B14	<i>Solanum tuberosum</i>	beta-1,3-glucanase gene under 35S promoter	<i>Agrobacterium</i> mediated transformation, PCR, Northern blot	Wróbel-Kwiatkowska et al. 2004

seedlings grown for 6 weeks; inoculation resulted in over 90% AMF colonized root length.

The pots were placed in Sunbags (Sigma) and flax plants were grown under greenhouse conditions: 25/18°C maximum/minimum average air temperatures, 12 h light supplemented if needed ($200 \mu\text{mol} \times \text{m}^{-2} \times \text{s}^{-1}$ PAR). Plants were irrigated weekly with distilled water and once per 3 weeks with Long Ashton solution (Davies et al. 1992). After 3-month growth, when plants started to form flowers, photosynthesis was measured and plants were harvested. Length of roots and shoots and dry weight of shoots were estimated (roots were used for estimation of mycorrhizal parameters).

Mycorrhizal colonization

Roots were washed, cleared in 10% KOH for 24 h at room temperature and, after rinsing in tap water, acidified for 1 h in 5% lactic acid then stained 24 h at room temperature in 0.05% aniline blue in lactic acid, in order to visualize the fungal structures inside the roots. One-centimeter root pieces were mounted on slides in lactoglycerol (approx. 45 root pieces per plant) and mycorrhizal parameters in root systems were assessed according to Trouvelot et al. (1986; <http://www2.dijon.inra.fr/mychintec/Mycocalc-prg/download.html>): frequency of mycorrhiza presence ($F\%$), intensity of mycorrhizal colonization ($M\%$), and arbuscule abundance ($A\%$). Images were acquired using a NIKON Eclipse E800 microscope equipped with DIC, a digital image capture system and a computer image analysis system.

Plant growth assessment

Shoot length was measured, then material was dried for 2 days at 40°C and weighed (Radwag analytical balance, WPA 60/c/1). Mycorrhizal dependency (Md) of plants for growth (mass, shoot length) as well as performance index (see below) was determined according to the equation given by van der Heijden and Sanders (2002)

P and Zn concentrations in plants

Dried, grinded, and homogenized flax shoots were digested in 10-mL concentrated (65%) nitric and perchloric acids (Ultranal, POCh) according to Pilegaard (1979). Residues were dissolved in 10 mL 0.2 % nitric acid, filtered, and diluted if necessary. Zinc concentrations were analyzed by flame atomic absorption spectrometry using AAnalyst 200, Perkin-Elmer. Nitric acid was used as a blank and *Ulva lactuca* BCR-279 as a standard reference material (Promochem GmbH; Zn=51.3 mg kg⁻¹). Concentrations of phosphorus were determined by flow injection analysis system (FIA) coupled with flow colorimeter (FIA compact, MLE GmbH).

Fluorescence measurements

Chlorophyll *a* fluorescence transients of intact leaves were measured with a plant efficiency analyzer fluorimeter (Hansatech Instruments, UK) and recorded as described by Strasser et al. (1995). Each transient was analyzed according to the OJIP test (Strasser et al. 2004, 2007) and the following biophysical parameters (referring to time point zero) were calculated to characterize PSII behavior: (1) specific energy fluxes per reaction center for absorption (ABS/RC); (2) electron trapping (TRo/RC), electron dissipation (Dio/RC), and electron transport (To/RC); (3) the flux ratios or yields, i.e., the maximum quantum yield of primary photochemistry ($\varphi_{Po} = \text{TRo}/\text{ABS}$); (4) the efficiency with which a trapped exciton can move an electron into the electron transport chain further than Q_A ($\psi_o = \text{ETo}/\text{TRo}$), the quantum yield of electron transport ($\varphi_{Eo} = \text{ETo}/\text{ABS}$); (5) and the performance index (PI), which provides quantitative information about the state of plants and their vitality. The latter incorporates three independent expressions: (1) ratio of reaction center chlorophylls and the total chlorophyll of PSII; (2) an expression related to primary photochemistry $\phi_{Po}/(1-\phi_{Po})$; and (3) an expression related to electron transport $\Psi_o/(1-\Psi_o)$ (Oukarroum et al. 2007).

Data analysis

Data from mycorrhizal and nonmycorrhizal plants were compared using the *t* test or were transformed with decimal logarithms prior to analysis to meet statistical assumptions and compared using ANOVA, STATISTICA (ver. 7.0) software (significance level $P < 0.05$).

Results

AMF colonization

Arbuscular mycorrhiza of *Arum* type with well-developed arbuscules were observed in all inoculated flax plants. These structures were not found in any of the non-inoculated samples. The frequency of mycorrhiza ($F\%$), reflecting propagule density in the samples, ranged from 93% to 100% and showed the smallest differences between root samples among studied parameters. The intensity of mycorrhizal colonization ($M\%$) ranged from 40% to 65%. Significantly higher $M\%$ values were noted in the case of lines B14 and W92.40, while other transformed lines from the cultivars Linola and Nike were similar (Fig. 1). The largest differences between samples involved arbuscule abundance ($A\%$) in roots which was highest in the two transgenic lines W9240 and B14. The other lines derived from cv. Linola showed similar values for arbuscule abundance. In contrast,




Fig. 1 Relative mycorrhizal intensity (*M*) and relative arbuscule richness (*A*) in *L. usitatissimum*; the numbers represent percent values, according to the parameter definitions by Trouvelot et al. (1986); asterisk—GM lines of Linola, double asterisks—GM lines of Nike; bars labeled with the same letter are not significantly different at $P < 0.05$

the GM lines derived from cv. Nike either had higher (B14) or significantly lower (M50) *A*% (Fig. 1). Roots were free of parasites or nematodes.

Plant growth and performance

Mycorrhizal plants of cv. Nike and Linola were characterized by increased total biomass and root weight in comparison to nonmycorrhizal controls (Fig. 2). Smaller but still significant differences ($P = 0.049$) were found in root or shoot weight between mycorrhizal and nonmycorrhizal GM lines. There were no significant differences between shoot weight of mycorrhizal transgenic lines and of their parent cultivars. In the case of root weight, statistically significant differences were found only between GT4 and cv. Linola (Fig. 2). Among nonmycorrhizal plants, higher shoot weight was found in GT4 than in cv. Linola and lower weight in B14 than in cv. Nike, while the root weight of W92.40 and GT4 were 3 and 2

times higher than in cv. Linola, respectively, and in M50 and GT5 were over 1.5 and 2 times higher than in cv. Nike. Md for growth was higher in cv. Nike (30%) than in cv. Linola (18%). Transformation of these two cultivars resulted in decreased values of growth Md in comparison to the parent cultivars. Negative values were found in the lines W92.40 (−18%), GT4 (−14%), and GT5 (−12%), while the Md value of B14 was ca. 10% lower than in cv. Nike.

P concentration in flax shoots was significantly higher in mycorrhizal plants than in nonmycorrhizal ones, both in GM lines and parent cultivars, except for line M50 (Fig. 3a). On the contrary, Zn concentration depended on the flax cultivar, being significantly increased in cv. Linola, but not in cv. Nike. No significant differences were found in two modified lines, B14 and W92.40 (Fig 3b). No significant differences in Md concerning P and Zn were found between GM and non-GM lines. No differences were found in flowering time of the GM and non-GM lines or between mycorrhizal and nonmycorrhizal plants.

The measure of plant performance, PI, revealed differences between transgenic lines and the parent cultivars. PI of mycorrhizal plants of cultivars Nike and Linola were significantly higher than in nonmycorrhizal ones (Fig. 3). Md estimated on the basis of plant performances (PI) in these two cultivars were 12% and 26%, respectively.

Among transformed lines, a similar situation was found only in the case of W92.04 (Md=21%); a negative effect of mycorrhization on PI was not significant in the case of GT4 (Md=−17%), while in other cases the values were similar in mycorrhizal and nonmycorrhizal plants (Fig. 4). In addition, PI increase in non-transformed cultivars following mycorrhization was associated with statistically significant differences in several PSII parameters, such as ABS/RC, TRo/RC, DIo/RC, and To/RC, the maximum quantum yield of primary photochemistry ($\varphi_{Po} = TRo/ABS$), and the quantum yield of electron transport ($\varphi_{Eo} = ET0/ABS$) (data not shown). This was not the case for transgenic lines where PI in mycorrhizal plants was not significantly different from nonmycorrhizal ones (GT4, M50, B14, and GT5).

Fig. 2 Shoot (a) and root (b) dry weight (in grams) of mycorrhizal (*M*) and nonmycorrhizal (*NM*) plants; asterisk—GM lines of Linola, double asterisks—GM lines of Nike; different letters above columns indicate statistically significant differences ($P < 0.05$)





Fig. 3 Comparison of P (a) and Zn (b) concentration in mycorrhizal (M) and nonmycorrhizal (NM) shoots; asterisk—GM lines of Linola, double asterisks—GM lines of Nike; bars labeled with different letters are significantly different at $P < 0.05$



Discussion

AMF root colonization in flax has been previously described (Dickson et al. 2003; Vierheilig et al. 2008) and effects of the symbiosis on non-modified flax have been the subject of both field (Thingstrup et al. 1998; Olsson et al. 1999) and laboratory (Püschel et al. 2008; Rydlova et al. 2010) studies. Mycorrhizal effects have been shown on flax growth, due to increased uptake of P and Zn by fungal mycelium (Thompson 1994; Thingstrup et al. 1998; Smith et al. 2004), on fiber weight, seed production, and fatty acid composition of seed oils (Rydlova et al. 2010), on drought resistance (Druge and Schonbeck 1992; von Reichenbach and Schonbeck 1995) and on resistance to plant pathogens (Dugassa et al. 1996). The main aim of the present study was to determine whether genetic transformation of flax by different resistance-related genes affects AMF and/or their effects on host performance. Transformation did not change the degree of mycorrhizal colonization; no morphological differences were found between transformed and non-transformed flax lines and arbuscules, which are indicative of a functional symbiosis, were well developed in all cases




Fig. 4 Comparison of plant performance (PI) between mycorrhizal (M) and nonmycorrhizal (NM) plants—*t* test; asterisk—GM lines of Linola, double asterisks—GM lines of Nike; bars labeled with different letters are significantly different at $P < 0.05$

except line M50. The lines B14 and W92.40 showed even higher arbuscule abundance in comparison with the parent cultivars. These observations are in agreement with most of previously published data showing no negative effect on fungal growth in plants overexpressing defense-related genes (Liu 2010). The transformations in the present flax lines increase resistance to pathogenic fungi via increased phenylpropanoid levels (W92.40) (Lorenc-Kukula et al. 2005), antioxidant capacity (W92.40, GT) and overproduction of glycosyltransferase (GT) (Lorenc-Kukula et al. 2004) and β -1,3-glucanase ectopic expression of a potato β -1,3-glucanase gene in flax (B14). Since there is some similarity between the soil-borne pathogens and AMF in their resources such as space and photosynthate within the root, such defense “pathways” could be potentially deleterious to both groups of fungi (Whipps 2004). However, as reviewed by Dumas-Gaudot et al. (2000) and Khan et al. (2010), AMF are capable themselves of inducing hydrolytic enzymes and pathogenesis-related proteins without being affected. For example, AMF are not inhibited by Class I chitinases (Arlorio et al. 1991; Salzer et al. 1997; Vierheilig et al. 1993) that are deleterious to pathogenic fungi such as *Rhizoctonia solani*. Similarly, AMF are not affected by an antifungal protein that targets *Verticillium albo-atrum* or *Botrytis cinerea* in transformed aubergine plants (Turrini et al. 2004a, b). This differential targeting of pathogenic and mycorrhizal fungi could be due to mechanisms of recognition specificity (Vierheilig et al. 2008; Khan et al. 2010; Stefani and Hamelin 2010) but this issue requires further in-depth research.

The effect of genetically modified crops on mycorrhizal fungi has so far been mostly addressed by analysis of mycorrhizal colonization levels which, as mentioned above, are largely unaffected. However, the present data indicate that, although growth of the mycorrhizal transformed flax plants was not different from the parent lines, transformation may alter plant responses to AMF which could be relevant for biomass production and crop quality under agricultural conditions. While no significant differences were observed in Md concerning P and Zn, *G. intraradices* was less

effective in stimulating growth or photosynthetic performance and, in one case P uptake (M50), by transformed as compared to parent flax plants. The magnitude of mycorrhizal responses can be influenced not only by the plant but also environmental factors (Smith and Gianinazzi-Pearson 1988, Smith and Read 2008, Jakobsen et al. 2002). In the GM flax lines, the lower mycorrhizal dependency parameters in reference to plant growth and photosynthesis efficiency resemble the situation between modern varieties or landraces and their ancestors (Hetrick et al. 1992, 1993; Boyetchko 1996), and both might be less adapted to an agriculture turned towards low fertilizer inputs, especially in phosphorus and nitrogen, where AMF can be the key players in establishment of ecosystem function and diversity (Thompson 1994, van der Heijden et al. 1998, Klironomos et al. 2000).

In conclusion, it was shown that the genes introduced into flax lines to increase resistance against pathogens did not have a negative effect on the colonization of flax by AMF but may affect some plant responses to the symbiosis. Plant growth and photosynthetic activity were not significantly improved by *G. intraradices* in the transformed as compared to parent lines, but P nutrition was still enhanced in all but one which may be promising for better survival of seedlings due to better nutrition of seeds. The fact that the transformed flax lines, which meet the demands for better quality of fiber and higher resistance to pathogens, do not appear to be negatively affected in their interactions with AMF should make them suitable for use in agriculture.

Acknowledgments We greatly acknowledge Dr. Anna Jurkiewicz (Aarhus University, DK) for the comments on the manuscript. This study was supported by grants SPUB/COST 870 (197/N-COST/2008/0), NN 302061834, and NR 12-0009-06 from the Ministry of Science and Higher Education.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Arlorio M, Ludwig A, Boller T, Mischiati P, Bonfante P (1991) Effects of chitinase and β -1,3 glucanase from pea on the growth of saprophytic, pathogenic and mycorrhizal fungi. *Giorn Bot Ital* 125:956–959
- Armstrong GM, Armstrong JK (1981) *Formae speciales* and races of *Fusarium oxysporum* causing wilt diseases. In: Nelson PE, Toussoun TA, Cook RJ (eds) *Fusarium: Diseases, biology and taxonomy*. The Pennsylvania State University Press, University Park, pp 392–399
- Boyetchko SM (1996) Impact of soil microorganisms on weed biology and ecology. *Phytoprotection* 77(1):41–56
- Czemplik M, Boba A, Kostyn K, Kulma A, Miłucha A, Sztajnert M, Wróbel-Kwiatkowska M, Żuk M, Szopa J, Skórkowska-Telichowska K (2011) Flax engineering for biomedical application in biomedical engineering. In: Komorowska MA, Olsztyńska-Janus S (eds) *Trends, research and technologies*. Intechweb Org, Rijeka, pp 407–434
- Davies FT, Potter JR, Linderman GR (1992) Mycorrhiza and repeated drought exposure affect drought resistance and extraradical hyphae development of pepper plants independent of plant size and nutrient content. *J Plant Physiol* 139:289–294
- Dickson S, Schweiger P, Smith FA, Söderström B, Smith S (2003) Paired arbuscules in the *Arum*-type arbuscular mycorrhizal symbiosis with *Linum usitatissimum*. *Can J Bot* 81:457–463
- Drüge U, Schönbeck F (1992) Effect of vesicular-arbuscular mycorrhizal infection on transpiration, photosynthesis and growth of flax (*Linum usitatissimum* L.) in relation to cytokinin levels. *J Plant Physiol* 141:40–48
- Dugassa GD, von Alten H, Schoneck F (1996) Effects of arbuscular mycorrhiza (AM) on health of *Linum usitatissimum* L. infected by fungal pathogen. *Plant Soil* 185(2):173–182
- Dumas-Gaudot E, Gollotte A, Cordier C, Gianinazzi S, Gianinazzi-Pearson V (2000) Modulation of host defence systems. In: *Arbuscular mycorrhizas: Physiology and function*. Dordrecht, Kluwer Academic, pp 173–200
- Giovannetti M, Sbrana C, Turrini A (2005) The impact of genetically modified crops on soil microbial communities. *Riv Biol-Biol Forum* 98:393–418
- Gredes T, Dominiak M, Gedrange T, Kunert-Keil C, Wróbel-Kwiatkowska M, Szopa J (2010) The influence of biocomposites containing genetically modified flax fibers on gene expression in rat skeletal muscle. *Biomedizinische Technik. Biomed Eng* 55(6):323–329
- He X, Nara K (2007) Element biofortification: can mycorrhizas potentially offer a more effective and sustainable pathway to curb human malnutrition? *Trends Plant Sci* 12(8):331–333
- Hetrick BAD, Wilson GWT, Cox TS (1992) Mycorrhizal dependence of modern wheat cultivars, landraces and ancestors. *Can J Bot* 70:2032–2040
- Hetrick BAD, Wilson GWT, Cox TS (1993) Mycorrhizal dependence of modern wheat cultivars and ancestors: a synthesis. *Can J Bot* 71:512–518
- Jakobsen I, Smith SE, Smith FA (2002) Function and diversity of arbuscular mycorrhizas in carbon and mineral nutrition. In: van der Heijden MGA, Sanders IR (eds) *Mycorrhizal ecology*. Springer, Berlin, pp 75–92
- Jhala AJ, Raatz LL, Dexter JE, Hall LM (2010) Adventitious presence: volunteer flax (*Linum usitatissimum*) in herbicide-resistant canola (*Brassica napus*). *Weed Technol* 24:244–252
- Jones JP (1991) *Fusarium* wilt. In: Jones JB, Stall RE, Zitter TA (eds) *Compendium of tomato diseases*. APS, St Paul
- Khan HMD, Meghvansi MK, Panvar V, Gogoi HK, Singh L (2010) Arbuscular mycorrhizal fungi-induced signaling in plant defence against phytopathogens. *J Phytol* 2(7):53–69
- Klironomos JN, McCune J, Hart M, Neville J (2000) The influence of arbuscular mycorrhizae on the relationship between plant diversity and productivity. *Ecol Lett* 3(2):137–141
- Liu W (2010) Do genetically modified plants impact arbuscular mycorrhizal fungi? *Ecotoxicology* 19:229–238
- Lorenc-Kukuła K, Amarowicz R, Oszmianski J, Doermann P, Starzycki M, Skala J, Żuk M, Kulma A, Szopa J (2005) Pleiotropic effect of phenolic compounds content increases in transgenic flax plant. *J Agric Food Chem* 53:3685–3692
- Lorenc-Kukuła K, Kostyń K, Łukaszewicz M, Szopa J (2004) Glucosyltransferase, the gene arrangement and enzyme function. *Cel Mol Biol* 9:935–946
- Lorenc-Kukuła K, Żuk M, Kulma A, Czemplik M, Kostyn K, Skala J (2009) Engineering flax with the GT family 1 *Solanum* sugarandin glycosyltransferase SsGT1 confers increased resistance to *Fusarium* infection. *J Agric Food Chem* 57(15):6698–6705

- Łukaszewicz M, Matysiak-Kata I, Skala J, Fecka I, Cisowski W, Szopa J (2004a) Antioxidant capacity manipulation in transgenic potato tuber by changes in phenolic compounds content. *J Agric Food Chem* 52:1526–1533
- Łukaszewicz M, Szopa J, Krasowska A (2004b) Susceptibility of lipids from different flax cultivars to peroxidation and its lowering by added antioxidants. *Food Chem* 88:225–231
- Olsson PA, Thingstrup I, Jakobsen I, Bååth E (1999) Estimation of the biomass of arbuscular mycorrhizal fungi in a linseed field. *Soil Biol Biochem* 31(13):1879–1887
- Oukarroum A, El Madidi S, Schanser G, Strasser RJ (2007) Probing the responses of barley cultivars (*Hordeum vulgare* L.) by chlorophyll *a* fluorescence OLKJIP under drought stress and re-watering. *Environ Exp Bot* 60:438–446
- Pilegaard K (1979) Heavy metals in bulk precipitation and transplanted *Hypogymnia physodes* and *Dicranoweisia cirrata* in the vicinity of a Danish steelworks. *Water Air Soil Poll* 11:77–91
- Püschel D, Rydlová J, Sudová R, Gryndler M (2008) Cultivation of flax in spoil-bank clay: mycorrhizal inoculation vs. high organic amendments. *J Plant Nutr Soil Sci* 171(6):872–877
- Ren X (2006) Effect of Bt transgenic rice (KMD) on soil bacterial community and rhizosphere AM fungi. Dissertation, Zhejiang University. Hang Zhou, China
- Rydlová J, Püschel D, Sudová R, Gryndler M, Mikanová O, Vosátka M (2010) Interaction of arbuscular mycorrhizal fungi and rhizobia: effects on flax yield in spoil-bank clay. *J Plant Nutr Soil Sc* 174:128–134
- Salzer P, Hübner B, Sirrenberg A, Hager A (1997) Differential effect of purified spruce chitinases and B-1,3-glucanases on the activity of electors from ectomycorrhizal fungi. *Plant Physiol* 114:957–968
- Skórkowska-Telichowska K, Żuk M, Kulma A, Bugajska-Prusak A, Ratajczak K, Gąsiorowski K, Kostyn K, Szopa J (2010) New dressing materials derived from transgenic flax products to treat long-standing venous ulcers—a pilot study. *Wound Rep Reg* 18:168–179
- Smith SE, Gianinazzi-Pearson V (1988) Physiological interactions between symbionts in vesicular arbuscular mycorrhizal plants. *Annu Rev Plant Phys* 39:211–244
- Smith SE, Read DJ (2008) Mycorrhizal symbiosis. Academic, London
- Smith SE, Smith FA, Jakobsen I (2004) Functional diversity in arbuscular mycorrhizal (AM) symbioses: the contribution of the mycorrhizal P uptake pathway is not correlated with mycorrhizal responses in growth or total P uptake. *New Phytol* 162:511–524
- Stefani FOP, Hamelin RC (2010) Current state of genetically modified plant impact on target and non-target fungi. *Environ Rev* 18:441–475
- Strasser RJ, Srivastava A, Govindjee (1995) Polyphasic chlorophyll *a* fluorescence transient in plants and cyanobacteria. *Photochem Photobiol* 61:32–42
- Strasser RJ, Tsimilli-Michael M, Srivastava A (2004) Analysis of the chlorophyll *a* fluorescence transient. In: Papageorgiou GC, Govindjee (eds) Chlorophyll *a* fluorescence: A signature of photosynthesis, advances in photosynthesis and respiration series (Govindjee—Series Editor) vol 19. Kluwer Academic, Rotterdam, pp 321–362
- Strasser RJ, Tsimilli-Michael M, Dangre D, Rai M (2007) Biophysical phenomics reveals functional building blocks of plants systems biology: a case study for the evaluation of the impact of mycorrhization with *Piriformospora indica*. In: Varma A, Oelmüller R (eds) Advanced techniques in soil biology. Soil Biology Series. Springer, Germany, pp 220–221
- Szopa J, Wróbel-Kwiatkowska M, Kulma A, Żuk M, Skórkowska-Telichowska K, Dymińska L, Mączka M, Hanuza J, Zebrowski J, Preisner M (2009) Chemical composition and molecular structure of fibers from transgenic flax producing polyhydroxybutyrate, and mechanical properties and platelet aggregation of composite materials containing these fibers. *Compos Sci Technol* 69:2438–2446
- Thingstrup I, Rubæk G, Sibbesen E, Jakobsen I (1998) Flax (*Linum usitatissimum* L.) depends on arbuscular mycorrhizal fungi for growth and P uptake at intermediate but not high soil P levels in the field. *Plant Soil* 203:37–46
- Thompson JP (1994) Inoculation with vesicular-arbuscular mycorrhizal fungi from cropped soil overcomes long-fallow disorder of linseed (*Linum usitatissimum* L.) by improving P and Zn uptake. *Soil Biol Biochem* 26:1133–1144
- Trouvelot A, Kough JL, Gianinazzi-Pearson V (1986) Mesure du taux de mycorrhization VA d'un système racinaire. Recherche de méthodes d'estimation ayant une signification fonctionnelle. In: Gianinazzi-Pearson V, Gianinazzi S (eds) Physiological and genetical aspects of Mycorrhizae. INRA, Paris, pp 217–221
- Turrini A, Sbrana C, Nuti MP, Pietrangeli BM, Giovannetti (2004a) Development of a model system to assess the impact of genetically modified corn and aubergine plants on arbuscular mycorrhizal fungi. *Plant Soil* 266:69–75
- Turrini A, Sbrana C, Pitto L, Ruffini Castiglione M, Giorgetti L, Briganti R, Bracci T, Giovannetti M (2004b) The antifungal Dm-AMP1 protein from *Dahlia merckii* expressed in *Solanum melongena* is released in root exudates and differentially affects pathogenic fungi and mycorrhizal symbiosis. *New Phytol* 163(2):393–403
- Van der Heijden MGA, Klironomos JN, Ursic M, Moutoglou P, Streitwolf-Engel R, Boller T, Wiemken A, Sanders IR (1998) Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. *Nature* 396:69–72
- Van der Heijden MGA, Sanders IR (2002) Mycorrhizal Ecology. Springer, Berlin
- Vierheilig H, Alt M, Neuhaus JM, Boller T, Wiemken A (1993) Colonization of transgenic *Nicotiana sylvestris* plants, expressing different form of *Nicotiana tabacum* chitinase, by the root pathogen *Rhizoctonia solani* and by the mycorrhizal symbiont *Glomus mosseae*. *Mol Plant-Microbe Interact* 6:261–264
- Vierheilig H, Steinkellner S, Khaosaad T, Garcia-Garrido JM (2008) The biocontrol effect of mycorrhization on soilborne fungal pathogens and the autoregulation of the AM symbiosis: one symbiosis, two effects? In: Varma A (ed) Mycorrhiza. Springer-Verlag, Berlin, pp 307–320
- von Reichenbach HG, Schönbeck F (1995) Influence of VA-mycorrhiza on drought tolerance of flax (*Linum usitatissimum* L.). II. Effect of VA mycorrhiza on stomatal gas exchange, shoot water potential, phosphorus nutrition and the accumulation of stress metabolites. *Angew Bot* 69:183–188
- Whipps JM (2004) Prospects and limitations for mycorrhizas in biocontrol of root pathogens. *Can J Bot* 82(8):1198–1227
- Wróbel M, Zebrowski J, Szopa J (2004) Polyhydroxybutyrate synthesis in transgenic flax. *J Biotechnol* 107:41–54
- Wróbel-Kwiatkowska M, Lorenc-Kukula K, Starzycki M, Oszmianski J, Kepczynska E, Szopa J (2004) Expression of β -1,3-glucanase in flax causes increased resistance to fungi. *Physiol Mol Plant Pathol* 65:245–256