

RESEARCH

Open Access



Interactions of tumour-derived micro(nano)vesicles with human gastric cancer cells

Małgorzata Stec¹, Rafał Szatanek¹, Monika Baj-Krzyworzeka¹, Jarosław Baran¹, Maria Zembala², Jakub Barbasz², Agnieszka Waligórska³, Jurek W. Dobrucki³, Bożenna Mytar¹, Antoni Szczepaniak⁴, Maciej Siedlar¹, Grażyna Drabik¹, Barbara Urbanowicz⁵ and Marek Zembala^{1*}

Abstract

Background: Tumour cells release membrane micro(nano)fragments called tumour-derived microvesicles (TMV) that are believed to play an important role in cancer progression. TMV suppress/modify antitumour response of the host, but there is also some evidence for their direct interaction with cancer cells. In cancer patients TMV are present in body fluid and tumour microenvironment. The present study aimed at characterization of whole types/subpopulations, but not only exosomes, of TMV from newly established gastric cancer cell line (called GC1415) and to define their interactions with autologous cells.

Methods: TMV were isolated from cell cultures supernatants by centrifugation at 50,000×g and their phenotype was determined by flow cytometry. The size of TMV was analysed by dynamic light scattering and nanoparticle tracking analysis, while morphology by transmission electron microscopy and atomic force microscopy. Interactions of TMV with cancer cells were visualized using fluorescence-activated cell sorter, confocal and atomic force microscopy, biological effects by xenografts in NOD SCID mice.

Results: Isolated TMV showed expression of CD44H, CD44v6 (hyaluronan receptors), CCR6 (chemokine receptor) and HER-2/neu molecules, exhibited different shapes and sizes (range 60–900 nm, highest frequency of particles with size range of 80–120 nm). TMV attached to autologous cancer cells within 2 h and then were internalized by them at 24 h. CD44H, CD44v6 and CCR6 molecules may play a role in attachment of TMV to cancer cells, while HER-2 associated with CD24 be involved in promoting cancer cells growth. Pre-exposure of cancer cells to TMV resulted in enhancement of tumour growth and cancer cell-induced angiogenesis in NOD SCID mice model.

Conclusions: TMV interact directly with cancer cells serving as macro-messengers and molecular cargo transfer between gastric cancer cells resulting in enhancement of tumour growth. TMV should be considered in future as target of anticancer therapy.

Keywords: Gastric cancer, Microvesicles, NOD SCID mouse model

Background

Interactions/communications between tumour cells themselves and tumour cells with the microenvironment are responsible for cancer development and progression

[1], but the pathomechanism(s) of cancer progression are far from clear. Tumour microenvironment plays a major role in this process, as both infiltrating and tumour cells produce a range of soluble mediators with protumour activity (e.g. interleukin-6, transforming growth factor β , epidermal growth factor and its receptor, vascular endothelial growth factor, etc.). However, immunotherapy approaches employing various monoclonal antibodies (mAbs), neutralizing the mediators or blocking their

*Correspondence: mizemba@cyf-kr.edu.pl

¹ Department of Clinical Immunology and Transplantology, Jagiellonian University Medical College, Wielicka 265 Str., 30-663 Kraków, Poland
Full list of author information is available at the end of the article

receptors, are only partially effective [2]. Therefore, there must be another mechanism(s) responsible for cancer progression.

Recently, there has been an increasing interest in studying tumour cells surface micro(nano)fragments also called micro(nano)vesicles (MV). MV are defined as spherical proteolipids with sizes ranging from 30 nm to 1 μ m [3]. MV circulate in blood of healthy individuals and their elevated numbers are found in patients with various diseases, including cancer [4, 5]. They are shed by many cell types, in particular rapidly proliferating. Tumour cells, in particular more malignant, release MV called tumour-derived MV (TMV). They are present in body fluids and tumour microenvironment [1]. In some types of cancer (ovarian, breast, gastric) elevated numbers of MV, not necessarily of tumour origin, are associated with poor prognosis [4–6]. MV carry various surface determinants and mRNA/miRNA and transfer some of them to other cells (e.g. blood cells), hence are regarded as important messengers in cell to cell communication [7–9]. TMV inhibit antitumour response of the host [10] and may play a pivotal role in cancer progression [1, 11]. There is some evidence that tumour cell-derived exosomes (particles with sizes <100 nm), that TMV contain, interact with autologous tumour cells. Firstly, TMV entry to recipient cells, including prostate, lung and breast cancer [12]. Secondly, gastric cancer exosomes promote tumour cell proliferation, at least in part, by activation of PI3K/Akt and MAPK/EPK pathways [13]. Thirdly, TMV contain selectively secreted components which are required for metastasis formation, e.g. in breast and non-small lung cancer [14, 15].

We have previously described the characteristics of circulating MV in blood of gastric cancer patients, and concluded that at least some of them may be of tumour origin [5]. As little is known about TMV from gastric cancer cells, we have established the novel gastric cancer cell line (called GC1415) and characterized immunophenotype, including expression of tumour markers of cells and TMV derived from them. The aim of these studies was to characterize all particles shed/released by tumour cells, but not only isolated exosomes, i.e. particles with sizes less than 100 nm, as the latter may not be representative for whole types/subpopulations of TMV. Their size, physical properties and morphology was determined. Isolated TMV were used to define their interactions with cancer cells they derived from. The present study shows that TMV attached to the cell surface and then were internalized as visualized independently by three different methodologies. Furthermore, TMV changed some biological properties of autologous cancer cells.

Methods

Cancer cells

Tumour cell line GC1415 was recently established in our laboratory from carcinomatous ascites of a patient with advanced gastric (adenocarcinoma) cancer. These studies were approved by the Jagiellonian University Medical College Ethical Committee (KBET/491/B/2003). The cells were cultured in DMEM with high glucose content (PAA Laboratories GmbH, Pasching, Austria) supplemented with 5 % foetal bovine serum (FBS, Biowest, Nuaille, France) at 37 °C, in 5 % CO₂ atmosphere and regularly tested for *Mycoplasma* sp. contamination by PCR-ELISA kit (Roche, Mannheim, Germany) and for endotoxin contamination by the Limulus test (Charles River Laboratories, Wilmington, MA) according to the manufacturers' instruction.

TMV isolation

GC1415 cells were cultured by bi-weekly passages in DMEM with 5 % FBS previously centrifuged at 50,000 \times g for 1 h to remove bovine MV. Supernatants from the cultures at confluency were collected and spun down at 3000 \times g for 10 min to remove cellular debris. Then, supernatants were centrifuged using RC28S ultracentrifuge (Sorvall, Newton, CT) at 50,000 \times g for 1 h at 4 °C. Pellets were washed several times in serum-free DMEM to remove FBS, and finally resuspended in it. Quantification of TMV protein concentration was evaluated by the Bradford's method using Quick Start Bradford Dye Reagent (BioRad, Hercules, CA). The number of TMV was determined by flow cytometry (FACS Canto, BD Biosciences Immunocytometry Systems, San Jose, CA) and nanoparticle tracking analysis (NTA) using NanoSight system, i.e. LM10HS microscope equipped with the LM14 488 nm laser module (Malvern Instruments Ltd., Malvern, UK). TMV were also tested for *Mycoplasma* sp. and endotoxin contamination and stored at –20 °C for further usage.

Flow cytometry

The following murine IgG₁ monoclonal antibodies (mAbs) were used: phycoerythrin (PE)-conjugated anti-CD44H and fluorescein (FITC)-conjugated anti-CD44v6 (eBioscience GmbH, Vienna, Austria), PE-conjugated anti-CCR6, anti-EGFR and allophycocyanin (APC)-conjugated anti-HER-2/neu (BD Biosciences, San Diego, CA), PE-conjugated anti-Mucin1 and anti-EMMPRIN (Santa Cruz Biotechnologies, Santa Cruz, CA), FITC-conjugated anti-EpCAM (DAKO, Glostrup, Denmark). Isotype controls included appropriate FITC-, PE- or APC-labelled mouse immunoglobulins. Cancer cells (1 \times 10⁵ per sample) or TMV (1 μ g per sample) were incubated with appropriate saturating concentrations of

mAbs for 20 min at 4 °C, washed (except TMV), resuspended in phosphate buffered saline (PBS) and analysed by FACS.

Determination of HER-2/neu and MAGE-1, -2 mRNA expression

Total RNA was isolated from the GC1415 tumour cells (1×10^6) and their TMV using the RNeasy Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol. The cDNA synthesis and nested, "real-time" PCR for MAGE-1, -2 and β -actin was performed as previously described [16]. For the detection of HER-2/neu mRNA "real time" PCR was performed using the following primer pair: for HER-2/neu: sense-5'-CCTCTGACGTCCATCATCTC-3' and antisense-5'-ATCTTCTCGTGCCGTCGCTT-3'. The cycle profile for HER-2/neu PCR run was as follows: initial denaturation at 95 °C for 10 min, then denaturation at 95 °C for 0 s, annealing at 60 °C for 35 s, and elongation at 72 °C for 35 s for 35 cycles, followed by final extension at 72 °C for 2 min. All the results were normalized with β -actin data and were presented as ΔC_T . To verify the amplified product, melting curve analysis using the 2.3 LightCycler software was performed for each sample.

Determination of TMV size distribution

Size distribution using the dynamic light scattering (DLS) was determined as previously described [5]. Briefly, samples either diluted or not (depending on signal intensity) in 0.15 M NaCl were measured in Zetasizer Nano ZS apparatus (Malvern Instruments) equipped with laser of $\lambda = 633$ nm. Size distribution was obtained from measured diffusion coefficients recalculated to the size by assuming spherical shape of particles. The obtained values represent the diameter of spherical particles which move in viscous media with the same velocity as TMV. The size and distribution of TMV was also determined by NTA. For this purpose TMV samples were diluted 200 times in prefiltered (0.2 μ m), sterile PBS to the total volume of 1 ml, then aspirated into an insulin-type syringe and loaded into the sample chamber. The TMV size and distribution measurements were collected in triplicates of 1 min film intervals recorded by the sCMOS camera and calculated using the NanoSight NTA 2.3 analytical software.

Zeta potential determination

Particle electrophoretic mobility was measured with Zetasizer Nano ZS apparatus (Malvern Instruments) in samples 10 times diluted with distilled water (to final ionic strength equal to 0.015 M) and recalculated to Zeta

potential using Smoluchowsky's equation, as previously described [6].

Transmission electron microscopy (TEM)

Purified TMV were mixed with 2 % of uranyl acetate and applied to carbon coated copper grid. The microphotographs were taken with Philips EM 300 electron microscope (Philips, Eindhoven, The Netherlands) operating at 80 kV.

Atomic force microscopy (AFM)

AFM was used to determine morphology of TMV. Samples for AFM were prepared from TMV suspensions by adsorbing onto highly ordered pyrolytic graphite (ZYH—mosaic spread 3.5–5.0) supports (NT-MDT Co., Zelenograd, Russia). The measurements were performed using Ntegra Vita microscope (NT-MDT Co.). The AFM images were recorded in semicontact mode. Experiments were performed using the commercial silicon tip with polysilicon cantilever (NT-MDT Co.) and resonance frequency within the range of 140–240 kHz and spring constants $k = 3.4$ N/m or $k = 5.8$ N/m. AFM imaging of cancer cells were acquired in phase contact mode, which reflects both morphology and mechanical properties of the sample. Image Analysis 2.2.0 software from NT-MDT Co. was used for AFM image processing.

Binding and engulfment of TMV

FACS analysis

To test the ability of TMV to bind to tumour cells, TMV were stained with red PKH26 fluorescent dye (Sigma-Aldrich, St. Louis, MO) for 5 min, according to the manufacturer's instruction and then incubated with GC1415 cells (app. 10 TMV per cell) for 2 and 24 h. To discriminate between the cells which already engulfed TMV (intracellular localization) from those with TMV only attached to their surface (extracellular) quenching assay with crystal violet (Sigma-Aldrich, 1 mg/ml final concentration), was applied according to the manufacturer's instructions [17]. The samples were then analysed by FACS, where gate P1 was set according to autofluorescence of GC1415 cells incubated in the medium without TMV (negative control).

AFM analysis

For AFM analysis, tumour cells (2×10^4 per well) were cultured in DMEM with 1 % FBS for 48 h on Lab-Tek Chamber Slide (Nalgene Nunc International, Naperville, IL). Then, cells were incubated with TMV (at the ratio 1:10) for the next 24 h, washed and assessed by AFM, as described above, except that the cells were analysed directly on the slide and pictures were presented in phase

contrast mode, which reflects both morphology and mechanical properties of the sample.

Confocal microscopy analysis

GC1415 cells (1×10^6 per ml) were cultured in DMEM with 10 % FBS on Super Frost Plus microscopy slides (Menzel-Glaser, Braunschweig, Germany) for 48 h. TMV were incubated for 5 min with red PKH26 dye (Sigma-Aldrich) according to manufacturer's instruction. Then, TMV were washed with 1 % bovine serum albumin and several times in serum-free DMEM medium. PKH26-labelled TMV resuspended in the medium were added to GC1415 cells and incubated for 2 and 24 h. As control the cells were processed identically without TMV. After incubation, cells with intact plasma membrane were labelled with fluorescein, following hydrolysis of fluorescein diacetate (Sigma-Aldrich), as described by Rotman and Papermaster [18]. Life cell imaging was used to visualize extra- and intracellular localization of TMV (red fluorescence, GC1415 cells-green fluorescence). Images were acquired by Leica TCS SP5 laser scanning confocal microscope (Leica Microsystems CMS GmbH, Mannheim, Germany) equipped with 63x, NA1.4 planapochromatic oil immersion objective. During imaging cells were maintained at 37 °C in DMEM/F12 medium without phenol red (Sigma-Aldrich), supplemented with 2 % FBS. The confocal pinhole size was set at 1 Airy unit. PKH26 fluorescence was excited with 543 nm line of He/Ne diode laser and emission was recorded in the range of 570–640 nm. For excitation of fluorescein 488 nm line of argon ion laser was used and the emission was recorded in the range of 500–540 nm. Image processing and analyses were performed by LAS AF (Leica Microsystems) software.

Xenografts of GC1415 cells in NOD SCID mice

The GC1415 cells (1×10^6 per mouse) suspended in 200 μ l of PBS were injected subcutaneously into dorsal region of 8-weeks old NOD SCID mice purchased from Charles River Laboratories (Sulzfeld, Germany). Handling of mice and all procedures were performed under laminar flow conditions. In some experiments tumour cells were preincubated with autologous TMV (9×10^6) for 2 or 24 h, washed and then injected. The cells preincubated in the culture medium alone were used as control. Mice (5 per group) were checked every 7 days and the tumour diameter was measured with a caliper and its volume (v) was calculated according to the formula: $v = ab^2/2$ [19]. The tissues were examined macroscopically for metastases in the spleen, liver, lung, lymph nodes, kidney and then processed for histological examination.

Angiogenesis assay

Tumour cells, with or without TMV, were prepared as above, washed and resuspended in 100 μ l of PBS. Then, cells were mixed with 400 μ l of Matrigel Matrix (BD Biosciences) and injected subcutaneously into the abdominal middle line of NOD SCID mice. After 6 days mice were euthanized, Matrigel Matrix pellets were cut out and then 400 μ l of Cell Recovery Solution (BD Biosciences) was added and left at 4 °C for 7 days. Then, Matrigel Matrix implants were suspended in Drabkin's solution (Sigma-Aldrich) for 30 min, after that spun down and the content of haemoglobin was determined by measuring the absorbance (O.D.) at 540 nm in U-1800 spectrophotometer (Hitachi, Tokyo, Japan) [20]. Independently Matrigel plugs after cutting out were photographed and then fixed in formalin, paraffin embedded, serial 3 μ m sections were cut and stained with hematoxylin and eosin.

Statistical analysis

The nonparametric Mann–Whitney two-sided test was performed using GraphPad InStat version 2.0 software. P values <0.05 were considered significant.


Results

Immunophenotype of tumour cells and TMV

The CD44H (hyaluronian receptors), CD44v6, CCR6 (chemokine receptors) and EGFR (epithelial growth factor receptor) are involved in interactions of tumour cells with monocytes [21, 22]. The question was raised whether these receptors, by analogy, play a similar role in TMV-cancer cells interactions. Therefore, appropriate mAbs against these determinants were used to detect their expression on GC1415 cells and TMV. The presence of some tumour markers was also determined. The comparison of expression of tested determinants showed remarkable differences between tumour cells and their TMV (Fig. 1). The majority of cells were positive for CD44H and EGFR and about 50 % for CD44v6. The expression of EMMPRIN was observed on app. 20–60 % cells, CCR6 on 10–14 %, while few cells were EpCAM, EGFR and Mucin1 positive. Among analysed tumour markers, HER-2/neu was detected on all cells, but CEA and TAG-72 presence was not observed. The expression level of CD44H, EMMPRIN, HER-2/neu, EpCAM, EGFR and Mucin1 on TMV was significantly lower or absent, except CD44v6 and CCR6 which was comparable on cells and their TMV. The staining for MAGE-1, -2 was not performed due to the lack of suitable mAbs.

The MAGE-1, -2 and HER-2/neu mRNA expression

The expression of mRNA for MAGE-1, 2 and HER-2/neu was detected in both the cells and TMV, however, it was lower in TMV compared to the GC1415 tumour cells as



viewed by the threshold cycle value (ΔC_T). For the mRNA expression of MAGE-1, -2 and HER-2/neu in TMV the ΔC_T values were as follows: 6.163 ± 1.439 , 7.897 ± 1.385 , 6.890 ± 1.473 , where as in the GC1415 tumour cells they were 4.990 ± 3.167 , 2.673 ± 4.031 , 5.21 ± 2.598 , respectively. Melting curve analysis confirmed the specificity of the PCR products.

The physical characteristics of TMV

The effective size range of TMV as determined by DLS analysis was between 60–900 nm (Fig. 2a) and the most frequent events were around 80 nm (Fig. 2b). By NTA the reciprocal values were 70–520 nm and around 120 nm (Fig. 2c), respectively. Electrokinetic (Zeta) potential of TMV, which define the effective charge of objects, was




Fig. 2 The physical characteristics of TMV determined by DLS, NTA, TEM and AFM. **a** Size range and **b** size frequency of TMV by DLS. **c** Representative size frequency by NTA. **d** TEM, **e** 2D AFM and **f** 3D images ($5 \times 5 \mu\text{m}$ area) of TMV deposited onto pyrolytic graphite. Sizing of objects present along the *line* marked in part (**e**): x-axis—size, y-axis—height

-23.5 ± 1.80 mV. TEM and AFM were used to determine the morphology and size of TMV. TMV did not exhibit any characteristic features and were of different sizes and

shapes, mostly spheroid (Fig. 2d, e). Base cross-sections of selected particles (Fig. 2f) were within size limits, determined by DLS.


Binding and engulfment of TMV by autologous cancer cells



To test the ability of TMV to interact with autologous (parental) cancer cells, GC1415 cells were incubated with PKH26-labelled TMV. FACS analysis indicated that after 2 h of culture about 70 % of GC1415 cells were fluorescent, which increased to 99 % after 24 h (Fig. 3b, d). To discriminate, whether TMV only adhered to the cell surface or were engulfed by the cells, crystal violet was added. After quenching, the percentage of positive cells decreased to 16.8 and 37.7 % at 2 and 24 h, respectively (Fig. 3f, h), thus indicating intracellular localization of TMV.

To confirm localization of TMV in GC1415 cells confocal microscopy was applied. When GC1415 cells were incubated with PKH26-labelled TMV for 2 h the red fluorescence of the vesicles was associated with plasma membranes (Fig. 4a), however following a 24 h incubation the majority of TMVs were detected inside the cells (Fig. 4b). TMV localized intracellularly were of different size. AFM analysis of control and TMV-treated GC1415 cells showed at 24 h, but not 2 h, significant differences in their surface outlook. The cells not exposed to TMV (Fig. 5a) showed smooth surface and sharp contour. After incubation with TMV (Fig. 5b) the cell surface was rough, uneven with visible protuberations. The cells edges were thicker and diffused of torus shape.

Tumour growth and angiogenesis

Inoculation of NOD SCID mice with GC1415 cells led to the formation of tumours in all animals. Palpable tumours appeared around 3 weeks with a rapid growth thereafter. Observation period was 8 weeks. No metastases in the lung, liver, spleen, kidney, lymph nodes were detected either macro- or microscopically. Engraftment of GC1415 cells exposed to TMV in vitro for 2 or 24 h led in the latter case in a slight, but significant, increase of tumour volume as compared to untreated cells (Fig. 6a, b). Metastases were also not detected. Angiogenesis measured quantitatively by haemoglobin content showed significant increase when GC1415 cells were pre-exposed to TMV for 24 h, but not 2 h (Fig. 6c). Visualization of excised Matrigel plugs also showed the same (Fig. 6d). Histological analysis of the same plugs revealed that when tumour cells were used alone, examined areas show either no or few vessels (Fig. 6e, i, ii). When Matrigel Matrix containing GC1415 cells and TMV were implanted, both vessels and cells were more numerous, less when tumour cells were pre-exposed to TMV for 2 h (Fig. 6e, ii) and significantly more pronounced in case of cells preincubated with TMV for 24 h (Fig. 6e, iv). It was concluded, that TMV may enhanced tumour growth and angiogenesis.





Discussion

The role of MV and TMV in cancer is unclear. They are present in the circulation [5, 23] and probably involved in cancer progression, as shown by association of MV (not necessarily of tumour origin) level with poor survival of cancer patients [4]. They may act by inhibiting/hiding antitumour immune response [10] or like tumour-derived exosomes by interacting directly with tumour cells [7, 13, 24]. Current data clearly indicate that tumour microenvironment plays a major role in the inhibition

or enhancement of cancer progression. TMV that are present at the tumour site [1] may either induce cellular antitumour response [25] and release antitumour soluble factors [26] or, more likely, exhibit a tumour supporting role [27]. The latter may be due to the induction of growth factors production by tumour cells themselves or infiltrating cells, including macrophages, and/or by promoting the differentiation of myeloid cells into immunosuppressive cell subsets [28].

We have previously reported increased levels of circulating MV and their characteristics in blood of gastric cancer patients [5]. Since it is difficult to determine precisely which of these MV are of tumour origin, this gastric cancer cell line was established and its TMV were isolated and characterized in order to compare them with circulating MV. The immunophenotype analysis confirmed the other data showing that the expression of some determinants on TMV is much lower than on cells [26, 29, 30]. Only CCR6 and CD44v6 expression was comparable on both, cells and their TMV. CCR6 receptor was detected on some cells of pancreatic, colorectal and lung cancer cell lines and their TMV [26], circulating tumour cells [31] and plasma MV of gastric cancer patients [5]. This may suggest its involvement in cancer progression [32]. CCR6 is being identified as a potential new prognostic marker in some types of tumours and the axis of CCL20/CCR6/IL-17 may be a new therapeutic target in cancer [33]. The high expression of CCL20 may contribute to the selective recruitment of CCR6 positive cancer cells and metastases formation [32]. We have previously demonstrated that CCR6 transferred via TMV to monocytes is biologically active, as monocytes acquired chemotactic activity to CCL20 [26]. Based on that, it is reasonable to assume that CCR6 transfer between cancer cells, as exemplified by the attachment of PKH26-labelled TMV, may be a likely mechanism responsible for the enhancement of tumour growth. The presence of CD44H, CD44v6 (hyaluronian receptors) on TMV may also facilitate their interaction with cancer cells, known to abundantly express hyaluronian [34]. This may be analogous to its role in interaction of cancer cells with monocytes, resulting in the induction of reactive oxygen intermediates production [22]. Also, high expression of CD44v6 determinants on TMV may suggest their role in tumour progression, as soluble matrix delivered by CD44v6 is required for premetastatic niche formation [35]. Taken together, it is suggested that CD44H, CD44v6 and CCR6 may be involved in attachment rather than internalization of TMV. By analogy to exosomes, internalization of TMV may occur by various pathways of uptake [7], because the cleavage of exosome surface proteins reduced their association with target cells [12].




Fig. 6 TMV effect on the GC1415 tumour growth and angiogenesis in NOD SCID mice. Before engrafting into NOD SCID mice GC1415 cells were or not exposed to TMV in vitro. Kinetic curve of tumour growth of cells exposed (or not) to TMV for 2 h (**a**) and 24 h (**b**). Angiogenesis as assessed quantitatively by haemoglobin content (**c**). Bars represent the mean \pm SD from three experiments. Angiogenesis visualised as Matrigel plugs photos (**d**) and histological analysis (**e**). *i*-cells alone, *ii*-cells + TMV after 2 h incubation, *iii*- cells alone and *iv*- cells + TMV after 24 h pre-exposure. *Significantly different ($p < 0.01$) in comparison to control

The expression of tumour markers (Mucin1, EMM-PRIN, EpCAM, EGFR) on TMV was low or absent, despite their presence on cells. Although almost all cells were HER-2/neu positive, small proportion of TMV expressed this marker. The presence of mRNA for HER-2/neu, MAGE-1, -2 in TMV was also detected, although it was lower than in the tumour cells. These mRNAs were also found in circulating MV in plasma of gastric cancer patients [5]. The mRNA for interleukin-8 [8], EGFRvIII [36] and for housekeeping genes such as ACTB, GAPDH [24] were detected in the cargo of TMV and tumour-derived exosomes and shown to be transferred to monocytes [8], brain microvascular endothelial cells [36] and tumour cells [24], respectively. Moreover, TMV containing EGFRvIII released by more aggressive glioma cells were found to be responsible for the transfer of this receptor to indolent cells, which in turn caused the activation of transforming signaling pathways (MAPK and AKT) and changes in other gene expression (VEGF, Bcl-xL, p27) resulting in morphological transformation [37]. Altogether, it is reasonable to assume that the transfer of HER-2/neu molecule, its mRNA as cargo carried by engulfed TMV, and in result HER-2/neu expression by cancer cells may be associated with tumour progression, as observed in breast cancer [38].

The size range of TMV between app. 60–900 nm and the most frequent particle sizes at 80 nm determined by DLS were comparable to 70–520 and 120 nm by NTA, respectively. Similar size ranges were reported for circulating MV in patients with ovarian and gastric cancers [5, 39]. TMV size determined by AFM was also similar. The smaller values of the height in AFM can be explained as a result of probing soft material by relatively hard cantilevers. Thus, the measured altitudes provide only a comparison of the relative size of objects. Morphological characteristics of TMV by TEM and AFM indicated no major differences in comparison to circulating patients' MV [5]. Such analysis revealed that both MV and TMV particles are mostly spheroid in shape and of different sizes. TMV possessed high (negative) electrokinetic potential. This is in keeping with higher absolute zeta potential values of MV isolated from plasma of gastric cancer patients, as compared to those from donors' plasma [5] and highly metastatic sublines of rat prostate cancer cells [40]. Although we have no formal proof, similar characteristics (size, morphology, electrokinetic potential) of TMV isolated from GC1415 cell line may support our previous findings that at least some of MV found in blood of gastric cancer patients are of tumour origin [5]. Yet, their role in this type of cancer has to be established.

The present data from FACS, AFM and confocal microscopy provided unequivocal evidence that TMV

attached to and were internalized by autologous tumour cells. Both FACS and confocal microscopy indicated that at 2 h of exposure the majority of TMV were seen extracellularly, while at 24 h a large proportion of TMV was present inside the cells. Thus, between 2 and 24 h the entry of TMV occurred. TMV localized intracellularly were of different sizes, implicating their indiscriminate entry to the cells. This suggests that not only exosomes [9] may be taken up by cancer cells. There is an open question whether once engulfed TMV may be transferred to other cells. Furthermore, AFM analysis revealed that the exposure of cancer cells to TMV for 24 h, but not 2 h, induced noticeable structural changes in the cells. TMV-untreated cells had smooth surface and sharp edge, as demonstrable by rapidly changing height at this point during examination. In contrast, the cells exposed to TMV showed blurred edge. This can be explained as caused by possible stronger adherence of TMV-treated cells to the slide and may be compatible with the possible alterations in cell movement. The question arises whether similar situation occurs at the tumour site where TMV are also present [1]. The cell surface showed crumb-like roughness and some protuberations, indirectly indicating intracellular localization of TMV. The present findings are in keeping with the data on uptake of autologous exosomes by ovarian cancer cells [7]. Although, as judged by size, exosomes may be present among total TMV studied, they constitute only a part of all TMV. As PKH26 labels all TMV, it may suggest the entry to the cells both larger and smaller particles.

Altogether, the above data suggest the important role of TMV in communication between cancer cells, which may lead to changes of their biological behaviour. In fact, in vitro exposure of GC1415 cells to TMV, for 24 h, but not 2 h, enhanced tumour growth and angiogenic activity of tumour cells in NOD SCID mice. This is consistent with time of attachment to tumour cell surface (2 h) and engulfment of TMV (24 h), indicating that for these effects intracellular, rather than extracellular, localization is important. It is of interest that enhancement of tumour cells induced angiogenesis was associated with an increased number of cancer cells in Matrigel plugs, This may imply the important role of TMV in supporting tumour growth. It is debatable whether enhancement of tumour growth is connected with transfer of HER-2/neu mRNA and some TMV determinants, e.g. CD44H, CD44v6, CCR6, which are known to facilitate interaction of TMV with monocytes [8, 41]. There may be a special role for HER2 (EGFR) as molecule associated with CD24 which together with phosphorylated Akt promotes breast cancer cell survival [42]. The exact mechanism of tumour growth enhancement by TMV may be multifactorial and requires further studies (on the way).

Conclusions

TMV obtained from GC1415 cells showed significant heterogeneity in immunophenotype, size and morphology. TMV expressed some tumour markers and several molecules which may facilitate their attachment to parental cancer cells. Uptake of TMV, carrying these molecules and their mRNA, by GC1415 cells resulted in enhancement of tumour growth and angiogenesis in NOD SCID mice. The presented data support hypothesis of auto-crine tumour loops [43] in which TMV serve as macro-messengers in delivering signals and molecular cargo to the neighbour tumour cells. Therefore, the engulfment of TMV by cancer cells may qualify them as potential natural carriers of anticancer agents.

Abbreviations

TMV: tumour-derived microvesicles; mAbs: monoclonal antibodies; MV: micro(nano)vesicles; NTA: nanoparticle tracking analysis; PE: phycoerythrin; FITC: fluorescein; APC: allophycocyanin; PBS: phosphate buffered saline; FACS: fluorescence activated cell sorting; DLS: dynamic light scattering; TEM: transmission electron microscopy; AFM: atomic force microscopy.

Authors' contributions

All authors read and approved the final version of the manuscript. MS was responsible for the design and interpretation of flow cytometry analysis, execution of laboratory experiments, data analysis and drafted the manuscript, RS was responsible for NTA study and mRNA analysis, MB-K assisted in interpretation of the study and critical revision of the manuscript, JB took part in scientific discussions and revising the manuscript, MZ and JB took part in the design and interpretation of AFM experiments, AW and JWD took part in the design and interpretation of confocal microscopy experiments, BM is responsible for the design of the study, execution of laboratory experiments, data analysis and drafted the manuscript, AS was responsible for the provision of patients' materials, MS took part in scientific discussions and reviewed the manuscript, GD was responsible for histological examinations, BU took part in the design and interpretation of TEM, MZ is responsible for study design, analysis of results and preparation of the manuscript final version. All authors read and approved the final manuscript.

Author details

¹ Department of Clinical Immunology and Transplantology, Jagiellonian University Medical College, Wielicka 265 Str., 30-663 Kraków, Poland. ² Institute of Catalysis and Surface Chemistry, Polish Academy of Sciences, Kraków, Poland. ³ Division of Cell Biophysics, Faculty of Biochemistry, Biophysics and Biotechnology, Jagiellonian University, Kraków, Poland. ⁴ First Department of General and Gastrointestinal Surgery, Jagiellonian University Medical College, Kraków, Poland. ⁵ Electron Microscopy Laboratory, University Children's Hospital of Cracow, Kraków, Poland.

Acknowledgements

We wish to thank Ms Irena Ruggiero for skillful technical assistance. This work was supported by National Science Centre (Grant Numbers: N N401 290239 and 2012/07/B/NZ6/03499).

Competing interests

The authors declare that they have no competing interests.

Received: 29 May 2015 Accepted: 20 November 2015

Published online: 01 December 2015

References

1. Rak J. Extracellular vesicles—biomarkers and effectors of the cellular interactome in cancer. *Front Pharmacol*. 2013;4:1–14.

- Tol J, Punt C. Monoclonal antibodies in the treatment of metastatic colorectal cancer: a review. *Clin Ther*. 2010;32:437–53.
- Thery C, Zitvogel L, Amigorena S. Exosomes: composition, biogenesis and function. *Nat Rev Immunol*. 2002;2:569–79.
- Kim HK, Song KS, Park YS, Kang YH, Lee YJ, Lee KR, Kim HK, Ryu KW, Bae JM, Kim S. Elevated levels of circulating platelet microparticles, VEGF, IL-6 and RANTES in patients with gastric cancer; possible role of a metastasis predictor. *Eur J Cancer*. 2003;39:184–91.
- Baran J, Baj-Krzyworzeka M, Węglarczyk K, Szatanek R, Zembala M, Barbasz J, Czupryna A, Szczepanik A, Zembala M. Circulating tumour-derived microvesicles in plasma of gastric cancer patients. *Cancer Immunol Immunother*. 2010;59:841–50.
- Galindo-Hernandez O, Villegas-Comonfort S, Candanedo F, González-Vázquez MC, Chavez-Ocaña S, Jimenez-Villanueva X, Sierra-Martinez M, Salazar EP. Elevated concentration of microvesicles isolated from peripheral blood in breast cancer patients. *Arch Med Res*. 2013;44:208–14.
- Escreveute C, Keller S, Altevogt P, Costa J. Interaction and uptake of exosomes by ovarian cancer cells. *BMC Cancer*. 2011;11:108–18.
- Baj-Krzyworzeka M, Szatanek R, Węglarczyk K, Baran J, Urbanowicz B, Brański P, Ratajczak MZ, Zembala M. Tumour-derived microvesicles carry several surface determinants and mRNA of tumour cells and transfer some of these determinants to monocytes. *Cancer Immunol Immunother*. 2006;55:808–18.
- Valadi H, Ekstrom K, Bossios A, Sjostrand M, Lee JJ, Lotvall JO. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol*. 2007;9:654–9.
- Whiteside T, Mandapathi M, Szczepanski M, Szajnik M. Mechanism of tumor escape from the immune system: adenosine-producing Treg, exosomes and tumor-associated TLRs. *Bull Cancer*. 2011;98:E25–31.
- van Doormaal FF, Kleinjan A, Di Nisio M, Buller HR, Nieuwland R. Cell-derived microvesicles and cancer. *Neth J Med*. 2009;67:266–73.
- Smyth TJ, Redzic JS, Graner MW, Anchordoquy TJ. Examination of the specificity of tumor cell derived exosomes with tumor cells in vitro. *Biochim Biophys Acta*. 2014;1838:2954–65.
- Qu JL, Qu XJ, Zhao MF, Teng JE, Zhang Y, Hou KZ, Jiang YH, Yang XH, Liu YP. Gastric cancer exosomes promote tumour cell proliferation through PI3K/Akt and MAPK/ERK activation. *Dig Liver Dis*. 2009;41:875–80.
- Wang T, Glikes DM, Takano N, Xiang L, Luo W, Bishop CJ, Chaturvedi P, Green JJ, Semenza GL. Hypoxia-inducible factors and RAB22A mediate formation of microvesicles that stimulate breast cancer invasion and metastasis. *PNAS*. 2014;111:E3234–42.
- Park JE, Tan HS, Datta A, Lai RC, Zhang H, Meng W, Lim SK, Sze SK. Hypoxic tumor cell modulates its environment to enhance angiogenic and metastatic potential by secretion of proteins and exosomes. *Mol Cell Proteomics*. 2010;9:1085–99.
- Szatanek R, Drabik G, Baran J, Kolodziejczyk P, Kulig J, Stachura J, Zembala M. Detection of isolated tumour cells in the blood and bone marrow of patients with gastric cancer by combined sorting, isolation and determination of MAGE-1, -2 mRNA expression. *Oncol Rep*. 2008;19:1055–60.
- van Amersfoort ES, Van Strijp JA. Evaluation of a flow cytometric fluorescence quenching assay of phagocytosis of sensitized sheep erythrocytes by polymorphonuclear leukocytes. *Cytometry*. 1994;17:294–301.
- Rotman B, Papermaster BW. Membrane properties of living mammalian cells as studied by enzymatic hydrolysis of fluorogenic esters. *Proc Natl Acad Sci USA*. 1966;55:134–41.
- Iwanuma Y, Chen FA, Egilmez NK, Takita H, Bankert RB. Antitumor immune response of human peripheral blood lymphocytes coengrafted with tumor into severe combined immunodeficient mice. *Cancer Res*. 1997;57:2937–42.
- Passaniti A, Taylor RM, Pili R, Guo Y, Long PV, Haney JA, Pauly RR, Grant DS, Martin GR. A simple, quantitative method for assessing angiogenesis and antiangiogenic agents using reconstituted basement membrane, heparin, and fibroblast growth factor. *Lab Invest*. 1992;67:519–28.
- Zembala M, Siedlar M, Ruggiero I, Wieckiewicz J, Mytar B, Mattei M, Colizzi V. The MHC class-II and CD44 molecules are involved in the induction of tumour necrosis factor (TNF) gene expression by human monocytes stimulated with tumour cells. *Int J Cancer*. 1994;56:269–74.
- Mytar B, Siedlar M, Woloszyn M, Colizzi V, Zembala M. Cross-talk between human monocytes and cancer cells during reactive oxygen intermediates generation: the essential role of hyaluronan. *Int J Cancer*. 2001;94:727–32.

23. Castellana D, Toti F, Freyssonet JM. Membrane microvesicles: macromes-sengers in cancer disease and progression. *Thromb Res*. 2010;125(Suppl 2):S84–8.
24. Chiba M, Kimura M, Asari S. Exosomes secreted from colorectal cancer contains mRNAs, microRNAs and natural antisense RNAs, that can transfer into the human hepatoma HepG2 and lung cancer A549 cell lines. *Oncol Rep*. 2012;28:1551–8.
25. Stec M, Baran J, Szatanek R, Mytar B, Lenart M, Czupryna A, Szczepanik A, Siedlar M, Zembala M. Properties of monocytes generated from haematopoietic CD34(+) stem cells from bone marrow of colon cancer patients. *Cancer Immunol Immunother*. 2013;62:705–13.
26. Baj-Krzyworzeka M, Szatanek R, Węglarczyk K, Baran J, Zembala M. Tumour-derived microvesicles modulates biological activity of human monocytes. *Immunol Lett*. 2007;113:76–82.
27. McDaniel K, Correa R, Zhou T, Johnson C, Francis H, Glaser S, Venter J, Alpini G, Meng F. Functional role of microvesicles in gastrointestinal malignancies. *Ann Transl Med*. 2013;1:4.
28. Valenti R, Huber V, Filipazzi P, Pilla L, Sovena G, Villa A, Corbelli A, Fais S, Parmiani G, Rivoltini L. Human tumor-released microvesicles promote the differentiation of myeloid cells with transforming growth factor-beta-mediated suppressive activity on T lymphocytes. *Cancer Res*. 2006;66:9290–8.
29. Dolo V, Adobati E, Canevari S, Picone MA, Vittorelli ML. Membrane vesicles shed into the extracellular medium by human breast carcinoma cells carry tumor-associated surface antigens. *Clin Exp Metastasis*. 1995;13:277–86.
30. Dolo V, Ginestra A, Cassara D, Violini S, Torrisi MR, Nagase H, Canevari S, Pavan A, Vittorelli ML. Selective localization of matrix metalloproteinase 9, beta1 integrin, and human lymphocyte antigen class I molecules on membrane vesicles shed by 8701-BC breast carcinoma cells. *Cancer Res*. 1998;58:4468–74.
31. Pituch-Noworolska A, Drabik G, Szatanek R, Biała M, Kołodziejczyk P, Szczepanik A, Stachura J, Zembala M. Immunophenotype of isolated tumour cells in the blood, bone marrow and lymph nodes of patients with gastric cancer. *Pol J Pathol*. 2007;58:93–7.
32. Rubie C, Oliveira V, Kempf K, Wagner M, Tilton B, Rau B, Kruse B, König J, Schilling M. Involvement of chemokine receptor CCR6 in colorectal cancer metastasis. *Tumour Biol*. 2006;27:166–74.
33. Kirshberg S, Izhar U, Amir G, Demma J, Vernea F, Beider K, Shlomai Z, Wald H, Zamir G, Shapira OM, Peled A, Wald O. Involvement of CCR6/CCL20/IL-17 axis in NSCLC disease progression. *PLoS One*. 2011;6:e24856. doi:10.1371/journal.pone.0024856.
34. McBride WH, Bard JBL. Hyaluronidase-sensitive halos around adherent cells. Their role in blocking lymphocyte-mediated cytotoxicity. *J Exp Med*. 1979;149:507–15.
35. Jung T, Castellana D, Klingbeil P, Cuesta Hernandez I, Vitacolonna M, Orlicky DJ, Roffler SR, Brodt P, Zoller M. CD44v6 dependence of premetastatic niche preparation by exosomes. *Neoplasia*. 2009;11:1093–105.
36. Skog J, Würdinger T, van Rijn S, Meijer DH, Gainche L, Sena-Esteves M, Curry WT Jr, Carter BS, Krichevsky AM, Breakefield XO. Glioblastoma microvesicles transport RNA and proteins that promote tumour growth and provide diagnostic biomarkers. *Nat Cell Biol*. 2008;10:1470–6.
37. Al-Nedawi K, Meehan B, Micallef J, Lhotak V, May L, Guha A, Rak J. Inter-cellular transfer of the oncogenic receptor EGFRvIII by microvesicles derived from tumour cells. *Nat Cell Biol*. 2008;10:619–24.
38. Menard S, Tagliabue E, Campiglio M, Pupa SM. Role of HER2 gene overexpression in breast carcinoma. *J Cell Physiol*. 2000;182:150–62.
39. Gercel-Taylor C, Atay S, Tullis RH, Kesimer M, Taylor DD. Nanoparticle analysis of circulating cell-derived vesicles in ovarian cancer patients. *Anal Biochem*. 2012;428:44–53.
40. Carter HB, Coffey DS. Cell surface charge in predicting metastatic potential of aspirated cells from the Dunning rat prostatic adenocarcinoma model. *J Urol*. 1988;140:173–5.
41. Baj-Krzyworzeka M, Baran J, Węglarczyk K, Szatanek R, Szaflarska A, Siedlar M, Zembala M. Tumour-derived microvesicles (TMV) mimic the effect of tumour cells on monocyte subpopulations. *Anticancer Res*. 2010;30:3515–20.
42. Hosonaga M, Arima Y, Sugihara E, Kohno N, Saya H. Expression of CD24 is associated with HER2 expression and supports HER2-Akt signaling in HER2-positive breast cancer cells. *Cancer Sci*. 2014;105:779–87.
43. Filipazzi P, Burdek M, Villa A, Rivoltini L, Huber V. Recent advances on the role of tumor exosomes in immunosuppression and disease progression. *Semin Cancer Biol*. 2012;22:342–9.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

