



Double pion production in np and pp collisions at 1.25 GeV with HADES

G. Agakishiev⁶, C. Behnke⁷, D. Belver¹⁶, A. Belyaev⁶, J.C. Berger-Chen⁸, A. Blanco¹, C. Blume⁷, M. Böhmer⁹, P. Cabanelas¹⁶, S. Chernenko⁶, C. Drita¹⁰, A. Dybczak², E. Epple⁸, L. Fabbietti⁸, O. Fateev⁶, P. Fonte^{1,a}, J. Friese⁹, I. Fröhlich⁷, T. Galatyuk^{4,b}, J. A. Garzón¹⁶, K. Gill⁷, M. Golubeva¹¹, D. González-Díaz⁴, F. Guber¹¹, M. Gumberidze¹⁴, S. Harabasz⁴, T. Hennino¹⁴, R. Holzmann³, P. Huck⁹, C. Höhne¹⁰, A. Ierusalimov⁶, A. Ivashkin¹¹, M. Jurkovic⁹, B. Kämpfer^{5,c}, T. Karavicheva¹¹, I. Koenig³, W. Koenig³, B. W. Kolb³, G. Korcyl¹², G. Kornakov¹⁶, R. Kotte⁵, A. Krása¹⁵, E. Krebs⁷, F. Krizek¹⁵, H. Kuc^{2,14}, A. Kugler¹⁵, A. Kurepin¹¹, A. Kurilkin⁶, P. Kurilkin⁶, V. Ladygin⁶, R. Lalik⁸, S. Lang³, K. Lapidus⁸, A. Lebedev¹², L. Lopes¹, M. Lorenz⁷, L. Maier⁹, A. Mangiarotti¹, J. Markert⁷, V. Metag¹⁰, J. Michel⁷, C. Müntz⁷, R. Münzer⁸, L. Naumann⁵, M. Palka², Y. Parpottas^{13,d}, V. Pechenov³, O. Pechenova⁷, J. Pietraszko⁷, W. Przygoda², B. Ramstein¹⁴, L. Rehnisch⁷, A. Reshetin¹¹, A. Rustamov⁷, A. Sadovsky¹¹, P. Salabura², T. Scheib⁷, H. Schuldes⁷, J. Siebenson⁸, Yu.G. Sobolev¹⁵, S. Spataro^e, H. Ströbele⁷, J. Stroth^{7,3}, P. Strzempek², C. Sturm³, O. Svoboda¹⁵, A. Tarantola⁷, K. Teilab⁷, P. Tlusty¹⁵, M. Traxler³, H. Tsertos¹³, T. Vasiliev⁶, V. Wagner¹⁵, M. Weber⁹, C. Wendisch^{5,c}, J. Wüstenfeld⁵, S. Yurevich³ and Y. Zanevsky⁶

(HADES Collaboration)

¹LIP-Laboratório de Instrumentação e Física Experimental de Partículas, 3004-516, Coimbra, Portugal

²Smoluchowski Institute of Physics, Jagiellonian University of Cracow, 30-059 Kraków, Poland

³GSI Helmholtzzentrum für Schwerionenforschung GmbH, 64291 Darmstadt, Germany

⁴Technische Universität Darmstadt, 64289 Darmstadt, Germany

⁵Institut für Strahlenphysik, Helmholtz-Zentrum Dresden-Rossendorf, 01314 Dresden, Germany

⁶Joint Institute of Nuclear Research, 141980 Dubna, Russia

⁷Institut für Kernphysik, Goethe-Universität, 60438 Frankfurt, Germany

⁸Excellence Cluster 'Origin and Structure of the Universe', 85748 Garching, Germany

⁹Physik Department E12, Technische Universität München, 85748 Garching, Germany

¹⁰II. Physikalisches Institut, Justus Liebig Universität Giessen, 35392 Giessen, Germany

¹¹Institute for Nuclear Research, Russian Academy of Science, 117312 Moscow, Russia

- ¹²*Institute of Theoretical and Experimental Physics, 117218 Moscow, Russia*
- ¹³*Department of Physics, University of Cyprus, 1678 Nicosia, Cyprus*
- ¹⁴*Institut de Physique Nucléaire (UMR 8608), CNRS/IN2P3 - Université Paris Sud, F-91406 Orsay Cedex, France*
- ¹⁵*Nuclear Physics Institute, Academy of Sciences of Czech Republic, 25068 Rez, Czech Republic*
- ¹⁶*LabCAF. Dpto. Física de Partículas, Univ. de Santiago de Compostela, 15706 Santiago de Compostela, Spain*
- ^a*Also at ISEC Coimbra, Coimbra, Portugal*
- ^b*Also at ExtreMe Matter Institute EMMI, 64291 Darmstadt, Germany*
- ^c*Also at Technische Universität Dresden, 01062 Dresden, Germany*
- ^d*Also at Frederick University, 1036 Nicosia, Cyprus*
- ^e*Also at Dipartimento di Fisica Generale and INFN, Università di Torino, 10125 Torino, Italy*
E-mail: akurilkin@jinr.ru

The results of double pion production in np and pp collisions at an incident beam energy of 1.25 GeV with the HADES spectrometer at GSI are presented. The $np-$ reactions were studied in dp collisions at 1.25 GeV/ u using Forward Wall hodoscope aimed at registering spectator protons. High statistic invariant mass and angular distributions are obtained within the HADES acceptance which are compared with phase-space distributions.

*XXI International Baldin Seminar on High Energy Physics Problems,
September 10-15, 2012
JINR, Dubna, Russia*

*Speaker.

1. Introduction

Double-pion production in nucleon-nucleon (NN) collisions is of particular interest in view of studying the simultaneous excitation of the baryons and their subsequent decays. A number of low statistic experiments on pion productions in NN interactions have been performed in the past spanning the energy region from threshold to many GeV's by using bubble chamber techniques [1]-[2]. The bulk of the experimental data has come from pp collision. In contrast the data on np interaction in the low and medium energy regions are scarce. Recently, double-pion production in NN collisions has been accurately measured at CELSIUS [3], COSY [4], KEK [5], and PNPI-Gatchina [6] facilities. The differential cross section for $pp \rightarrow pp\pi^+\pi^-$ and $pp \rightarrow pp\pi^0\pi^0$ reactions have been obtained at CELSIUS and COSY for the beam energies from the threshold up to 1.4 GeV [3, 4]. The total cross section of $pn \rightarrow pn\pi^+\pi^-$ and $pn \rightarrow pp\pi^-\pi^0$ channels have been measured at KEK in the beam energy range from 698 MeV up to 1172 MeV.

On the theoretical side, the double pion production in NN collisions has been the subject of many investigations. The effective Lagrangian models (Valencia[8], XuCao[9] and modified Valencia[10]) predict that at energies near threshold the $\pi\pi$ production is dominated by the excitation of one of the nucleons into the Roper resonance $N^*(1440)P_{11}$ via σ -exchange. At higher energies the double $\Delta(1232)$ excitation is expected to be the dominant reaction mechanism for $\pi\pi$ production. The OPER model [11] based on the exchange of reggeized π have been successfully used to describe bubble chamber data[1] on $np \rightarrow np\pi^+\pi^-$ reaction at the momenta above 3 GeV/c. This model can be applied for description of $np \rightarrow np\pi^+\pi^-$ reaction at the momenta below 3 GeV/c by taking into account the mechanism of one baryon exchange(OBE).

New experimental data on double-pion production are needed to provide quantitative information on hadronic interactions, resonance excitations and resonance properties. In this work we present high statistic invariant mass and angular distributions on double pion production in np and pp collisions at an incident beam energy of 1.25 GeV obtained with the HADES spectrometer.

2. Experiment

The experimental data were collected using the High Acceptance Di-Electron Spectrometer(HADES) installed at the heavy-ion synchrotron SIS-18 at GSI in Darmstadt, Germany. The HADES is a modern multi-purpose detector currently operating in the region of kinetic beam energies of up to 2A GeV. The schematic view of the HADES spectrometer is presented in Fig. 1. Geometrically the spectrometer is divided into 6 identical sectors covering the full azimuthal angle and polar angles from 18° to 85° measured relative to the beam direction. Each sector of the spectrometer contains a Ring Imaging Cerenkov detector (RICH) operating in a magnetic field-free region, inner multi-wire drift chambers (MDCs) in front of the magnetic field, outer MDCs behind the magnetic field, TOF and TOFino time-of-flight detectors and a electromagnetic cascade detector (Pre-Shower). In order to investigate the np interaction using deuteron beam the HADES setup was upgraded with a Forward Wall (FWall) scintillator hodoscope covering the polar angles between 1° and 7° . While a detailed description of the setup can be found in [12], we summarize here only the features relevant for the present analysis.




Figure 1: Cut through two sectors of the HADES spectrometer. The magnet coils are projected onto the cut plane to visualize the toroidal magnetic field. A schematic view of the quasi-free $n + p$ reaction is shown.

In the presented experiments the proton and deuteron beams with intensities up to 10^7 particles/s and 1.25 GeV/c kinetic energy were directed to a 5 cm long liquid-hydrogen target of 1% interaction probability. The momenta of the produced particles were deduced from the hits in the four drift chamber planes (two before and two after the magnetic field zone) using a Runge-Kutta algorithm [12]. The momentum resolution was 2-3% for protons and pions and 1-2% for electrons, depending on momentum and angle [12]. The start signal for the time measurements was taken from the fastest signal from the scintillator wall. To reconstruct the time-of-flight for each particle, a dedicated method was developed [12], using the identification of one reference particle, the time-of-flight of which can be calculated. The time-of-flight reconstruction algorithm was checked in a dedicated experiment with a low beam intensity using a START detector [12]. The charge hadrons were selected from leptons by using RICH detector, together with TOF/TOFINO and Pre-Shower detectors. The selection of np collisions from dp and pp one was performed by using criteria on the momentum of the proton-spectator in FW detector. The selection of the $pp \rightarrow pp\pi^+\pi^-$ and $np \rightarrow np\pi^+\pi^-$ reactions were performed by using event hypothesis. For each hypothesis, any one of the hadrons was considered as reference particle and time-of-flight was calculated. The velocities of all the other products were then deduced, using only the time-of-flight differences to the reference particle. The correlations between velocity and momentum of all particles were then used to reject the wrong hypotheses and to assign the final PID of all particles.

3. Results

The experimental results on double pion production in np and pp collisions at incident beam energy 1.25 GeV are presented in Fig. 2 and Fig. 3 together with the phase-space distribution(PS) in

the HADES acceptance. Panels a), b), c) and d) in the Fig. 2 and Fig. 3 correspond to the invariant mass of $\pi^+\pi^-$, $p\pi^-$, $p\pi^+$ and angular distributions of the opening angle for $\pi^+\pi^-$ in the c.m. The experimental data are presented by the black points and include the statistical errors only. The systematic errors are about of 10% due to the correction on the efficiency and normalization on pp elastic scattering. Phase-space distributions(PS) in Fig. 2 and Fig. 3 are normalized on the area of experimental data.




Figure 2: The invariant mass spectra and angular distributions of the opening angle for $\pi^+\pi^-$ in the c.m.s. for the $np \rightarrow np\pi^+\pi^-$ reaction at incident beam energy 1.25 GeV. Black points and filled histogram are experimental data and phase-space(PS) distribution in the HADES acceptance, respectively.

The large discrepancies between the experimental data and phase-space distributions are observed at small angles in forward semisphere for angular distributions and in the $p\pi^+$ and $p\pi^-$ invariant mass distributions. M_{inv} of $\pi^+\pi^-$ demonstrate phase-space like behavior except the low mass region, where experimental results for np and pp reactions show clearly an enhancement in the invariant mass distributions. The similar effect was observed in the CELSIUS data on $\pi^0\pi^0$ production in pp collisions at the beam energies above 1.0 GeV [3]. Different models give the different possible interpretations of the low mass enhancement in $\pi\pi$ mass spectra for the $NN \rightarrow NN\pi\pi$ reaction[10]-[11]. CELSIUS data have been described by using the modified Valencia model[10]. This model gives reasonable agreement both for invariant mass spectra and angular distributions of




Figure 3: The invariant mass spectra and angular distributions of the opening angle for $\pi^+\pi^-$ in the c.m.s. for the $pp \rightarrow pp\pi^+\pi^-$ reaction at incident beam energy 1.25 GeV. Black points and filled histogram are experimental data and phase-space(PS) distribution in the HADES acceptance, respectively.

the opening angle in the c.m. for the $pp \rightarrow pp\pi^0\pi^0$ reactions. The ratio between decay of $N^* \rightarrow \Delta\pi$ and $N^* \rightarrow N\sigma$ was changed from 4 to 1 and ρ -exchange in the $\Delta\Delta$ excitation was suppressed by factor of 12 in this model[10]. On the other hand the OPER model, which successfully used to describe the bubble chamber data in $np \rightarrow np\pi^+\pi^-$ reactions shows the importance of the so called "hanged" diagrams[13]. The inclusion of these diagrams significantly improves the description of $\pi\pi$ invariant mass spectra in the low mass region.

4. Conclusion

The experimental results on double pion production in np and pp collisions at incident beam energy 1.25 GeV are obtained with the HADES spectrometer at GSI. The new high statistic HADES data provide precise results for mesons production measured in hadronic channels, paving the way for further theoretical or experimental studies of hadronic channels in elementary reactions.

References

- [1] C. Besliu *et al.*, *YaF*, **43** (1986) 888-892;
- [2] A. Baldini, V. Flamino, W.G. Moorhead, and D. R. O. Morrison (eds.), Landolt-Bornstein, *Numerical Data and Functional Relationships in Science and Technology*, Vol. 12, (Springer-Verlag, Berlin, 1988).
- [3] J. Johanson *et al.*, *Two-pion production in proton-proton collisions near threshold*, *Nucl. Phys.* **A712** (2002) 75; W. Brodowski *et al.*, *Exclusive Measurement of the $pp \rightarrow pp\pi^+\pi^-$ Reaction Near Threshold*, *Phys. Rev. Lett.* **88** (2002) 192301; J. Patzold *et al.*, *Study of the $pp \rightarrow pp\pi^+\pi^-$ Reaction in the Low-Energy Tail of the Roper Resonance*, *Phys. Rev. C* **67** (2003) 052202; E. Doroshkevich *et al.*, *Study of baryon and search for dibaryon resonances by the $pp \rightarrow pp\pi^+\pi^-$ reaction*, *Eur. Phys. J. A* **18** (2003) 171; T. Skorodko *et al.*, *Excitation of the Roper resonance in single- and double-pion production in nucleon-nucleon collisions*, *Eur. Phys. J. A* **35** (2008) 317;
- [4] S. Abd El-Bary *et al.*, *Two Pion Production in Proton-Proton Collisions with Polarized Beam* *Eur. Phys. J A* **18** (2008) 67; S. Abd El-Samad *et al.*, *On the production of $\pi^+\pi^-$ pairs in pp collisions at 0.8 GeV*, *Eur. Phys. J A* **42** (2009) 159;
- [5] T. Tsuboyama *et al.*, *Double-pion production induced by deuteron-proton collisions in the incident deuteron momentum range 2.1-3.8 GeV/c*, *Phys. Rev. C* **62** (2000) 034001.
- [6] V.V. Sarantsev *et al.*, *Measurement of the cross sections for $\pi(+)\pi(-)$ production in pp collisions at energies below 1 GeV*, *Phys. Atom. Nucl.* **70** (2007) 1998.
- [7] H. Clement *et al.*, *Two-pion production, gamma gamma line and aspects of sigma meson, Bose-Einstein correlations and isospin breaking*, *Int. J. Mod. Phys. A* **20** (2005) 1747.
- [8] L. Alvarez-Ruso, *et al.*, *Theoretical study of the $NN \rightarrow NN\pi\pi$ reaction*, *Nucl. Phys. A* **633** (1998) 519.
- [9] Xu. Cao, *et al.*, *Phenomenological analysis of the double pion production in nucleon-nucleon collisions up to 2.2 GeV*, *Phys. Rev. C* **81** (2010) 12.
- [10] T. Skorodko *et al.*, *Two-Pion Production in Proton-Proton Collisions: Experimental Total Cross Sections and their Isospin Decomposition*, *Phys. Lett.* **B679** (2009) 30.
- [11] A.P. Jerusalemov, *Analysis of the Reaction: $np \rightarrow np\pi^+\pi^-$ from the Point of View of Oper-Model*, (2012) arXiv:[nucl-th/1203.3330]
- [12] G. Agakishiev *et al.*, *The High-Acceptance Dielectron Spectrometer HADES*, *Eur. Phys. J.* **A41** (2009) 243.
- [13] A.P. Jerusalemov, *Contribution of the "hanged" diagrams into the reaction $np \rightarrow np\pi^+\pi^-$* , (2012) arXiv:[nucl-th/1208.3982]