

Protein Intake and Cognitive Function in Older Adults: A Systematic Review and Meta-Analysis

Hélio José Coelho-Júnior¹ , Riccardo Calvani²,
Francesco Landi¹, Anna Picca² and Emanuele Marzetti^{1,2} 

¹Università Cattolica del Sacro Cuore, Rome, Italy. ²Fondazione Policlinico Universitario "Agostino Gemelli" IRCCS, Rome, Italy.

Nutrition and Metabolic Insights
Volume 14: 1–10
© The Author(s) 2021
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/11786388211022373



ABSTRACT

INTRODUCTION: The present study investigated the association between protein intake and cognitive function in older adults.

METHODS: We performed a literature search with no restriction on publication year in MEDLINE, SCOPUS, CINAHL, AgeLine from inception up to October 2020. Observational studies that investigated as a primary or secondary outcome the association of protein intake and cognitive function in older adults aged ≥ 60 years were included.

RESULTS: Nine cross-sectional studies that investigated a total of 4929 older adults were included in the qualitative analysis. Overall cognitive function was examined in 6 studies. Four investigations reported null associations and 2 studies found that older adults with a high protein intake had higher global cognitive function than their counterparts. Results from the meta-analysis suggested that there were no significant associations between protein consumption and global cognitive function in older adults, regardless of gender. Three studies investigated other cognitive domains. Memory and protein intake were significantly and positively correlated in all studies. In addition, visuospatial, verbal fluency, processing speed, and sustained attention were positively associated with protein consumption in 1 study each.

CONCLUSION: No significant associations between protein intake and global cognitive function were observed in neither qualitative nor quantitative analyses. The association between protein consumption with multiple other cognitive domains were also tested. As a whole, 3 studies reported a positive and significant association between high protein intake and memory, while 1 study observed a significant and positive association with visuospatial, verbal fluency, processing speed, and sustained attention.

KEYWORDS: Dementia, frailty, nutrition, elderly

RECEIVED: February 8, 2021. **ACCEPTED:** May 8, 2021.

TYPE: Review

FUNDING: The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by Innovative Medicines Initiative–Joint Undertaking [IMI-JU 115621], the nonprofit research foundation "Centro Studi Achille e Linda Lorenzon", and by a scholarship to H.J.C.-J. from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior [CAPES; Finance Code 001]. The funders had no role in study design, data collection and analysis, preparation of the manuscript, or decision to publish.

DECLARATION OF CONFLICTING INTERESTS: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CORRESPONDING AUTHOR: Hélio José Coelho-Júnior, Center for Geriatric Medicine (Ce.M.I.), Fondazione Policlinico Universitario "Agostino Gemelli" IRCCS, Università Cattolica del Sacro Cuore, L.go F. Vito 1, Rome 00168, Italy.
EMAIL: coelhojunior@hotmail.com.br

Introduction

Cognition might be understood as the expression of brain activity by which the mind interacts with the world.¹ This concept involves many brain functions that allow human beings to create, execute, monitor, adjust, and perform many other tasks with a vital role in the proper accomplishment of daily activities.² Cognitive function is not homogeneous throughout life so that it expands from the gestational period until adulthood, remains constant during adult life, and declines past the age of the sixth decade of life.²

Such variations on cognition have a direct impact on older adults' quality of life, given that cognitive decline is significantly associated with depression, falls, vehicle collisions, hospitalization, disability, mild cognitive impairment (MCI), and death.^{3–8} This scenario has attracted considerable attention of the World Health Organization (WHO),⁹ which stated that the maintenance of cognitive function should be prioritized in older adults in an attempt to preserve their autonomy and avoid the genesis of chronic degenerative diseases.

Notably, lifestyle modifications have been recognized by the scientific community as possible tools to counteract age-related

cognitive decline. Physical activity and exercise, for example, have been extensively examined and investigations have indicated that older adults engaged in exercise programs might benefit from improvements in cognition,^{10–12} although these findings were not unanimous.¹³ Diet patterns have also been subject to investigation. According to recent systematic reviews,^{14,15} high adherence to the Mediterranean, Dietary Approach to Stop Hypertension (DASH), and the Mediterranean-DASH diet Intervention for Neurodegenerative Delay (MIND) diets might positively affect cognitive function in older people and likely slow or prevent cognitive decline.

However, a deeper look at the components of the diet, rather than the whole combination, might be necessary, given that the consumption of macronutrients can influence cognitive function. Particularly, the current recommendations for protein intake¹⁶ have been under intense debate.^{17–19} Most of the criticism is based on the numerous investigations that have identified that a dietary protein intake greater than the Recommended Dietary Allowances (RDA) can be required to maintain nitrogen balance and postpone age-related neuromuscular decline.^{20–23}



In the last years, the association between protein intake and cognitive function has also gained considerable attention,²⁴⁻³² and authors have claimed that a high consumption of proteins might contribute with a better cognitive functioning in older adults.³²⁻³⁴ These premises are based in the fact that increases in peripheral branched-chain amino acids (BCAA) content can contribute to the homeostasis of brain glutamate [GLU]. In addition, substantial evidence has accumulated that protein intake is significantly associated with numerous health-related parameters that might influence cognition,³⁴ including physical function,²⁰⁻²³ sleep quality,^{35,36} and microbiota.³⁷ Nonetheless, there is no consensus about the effects of consuming such amounts of protein on cognition.

Hence, the present systematic review investigated and combined the available evidence on the literature on the association between protein intake and cognitive function in older adults.

Methods

We conducted a systematic review of observational studies to assess the association between protein intake and cognitive function in older adults. The study was fully performed by investigators and no librarians were part of the team. This study complies with the criteria proposed by the Primary Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) Statement,³⁸ and the Meta-analysis of Observational Studies in Epidemiology (MOOSE) guidelines.³⁹

Eligibility criteria

The inclusion criteria consisted of: (a) Observational studies, including cross-sectional and case-control studies, which investigated as a primary or secondary outcome the association of protein intake and cognitive function in older adults free of dementia aged ≥ 60 years. Longitudinal cohort studies were also included whether crude baseline data were available; (b) assessed at least one cognitive domain via validated questionnaires and tests; and (c) published studies (English, Italian, Portuguese, and Spanish languages). To be included in the meta-analysis, investigations must provide (d) mean and standard deviations (SD) of protein intake in both case (i.e. high cognitive function) and control groups (i.e. low cognitive function). We excluded observational studies that investigated diet patterns, as well as randomized-controlled trials (RCTs), quasi-experimental, cross-over studies, and any kind of investigation that examined the effects of nutritional interventions on cognition. Studies that enrolled people with mild cognitive impairment (MCI), dementia and/or gastrointestinal and/or renal diseases, anorexia, cancer, or any kind of condition that might directly impair protein metabolism (e.g. maple syrup urine disease, tyrosinemia) were also excluded.

Search strategy and selection criteria

Studies published on or before October 2020 were retrieved from the following 4 electronic databases by 2 investigators: (1) MEDLINE (PubMed interface); (2) SCOPUS (Elsevier

Table 1. PICO strategy used for literature search.

PATIENT	INTERVENTION	OUTCOME
Aged [MESH]	Protein consumption	Cognition [MESH]
Older adults	Protein intake	Cognition
Elderly	Animal-protein	Cognitive function
Geriatrics#	Animal-based protein	Executive function
Seniors#	Plant-based protein	Problem solving
Aged 65+#	Plant-protein	Memory
	Vegetal-protein	Attention
		Language
		Visual perception
		Brain function#

[MESH] terms were only used for MEDLINE search. #These terms were only used for CINAHL and AgeLine searches.

interface); (3) CINAHL (EBSCO interface); (4) AgeLine (EBSCO interface). Reference lists for reviews and retrieved articles for additional studies were checked and citation searches on key articles were performed in Google Scholar and ResearchGate for additional reports. Initially, a search strategy was designed using keywords, MeSH terms, and free text words, such as protein consumption, cognitive function, older adults. Additionally, keywords and subject headings were exhaustively combined using Boolean operators. The PICO (i.e. Patient, Intervention, Comparison, and Outcome) method was used for literature search and the complete search strategy is shown in Table 1.

Data extraction and quality assessment

Titles and abstracts of retrieved articles were screened for eligibility by 2 researchers. If an abstract did not provide enough information for evaluation, the full-text was retrieved. Disagreements were solved by a third reviewer. Reviewers were not blinded to authors, institutions, or manuscript journals. Data extraction was independently performed by 2 reviewers using a standardized coding form. Disagreements were solved by a third reviewer. Coded variables involved the characteristics of the studies and included: year, authors, country, study design, setting, sample size, mean age, female prevalence, dietary intake assessment method, protein intake, cognitive scores, cognitive tests, and cognitive functions. The quality of reporting for each study was performed by 2 researchers using the Study Assessment Tool for Observational Cohort and Cross-Sectional Studies developed by the National Heart, Lung, and Blood Institute (NHLBI).⁴⁰ The agreement rate between reviewers for the quality assessment was $\kappa = 0.99$.

Statistical analysis

The meta-analysis was conducted using Revman 5.4.1 (Cochrane Collaboration, Copenhagen, Denmark). Effect size




Figure 1. Flowchart of the present study.

(ES) was measured using Standard Mean Differences (SMD). When data were not made available by authors as means and SDs, they were calculated according to Cochrane guidelines.⁴¹ Specifically, SDs were calculated from confidence intervals or standard errors and means were converted from medians. A single pairwise comparison was created when multiple studies referred to the same database using the formulas proposed by the Cochrane group.⁴¹ Due to the variability of sample characteristics, a random-effect model was used to calculate the pooled ES. A sensitivity analysis was performed based on the stratification technique according to gender.

Results

Literature search

Of the 3863 registers recovered from electronic databases and hand search, 3846 records were excluded based on duplicate data, title, or abstract. Seventeen studies were fully reviewed and assessed for eligibility. Of these, 4 studies included people

aged less than 60 years, 3 studies investigated people with dementia, and 1 study was a review of the literature. Finally, 9 studies were included in the qualitative synthesis. Three studies provided data to be included in the meta-analysis (Figure 1).

Characteristics of the included studies

Table 2 provides a general description of the included studies. Eight cross-sectional studies²⁴⁻³² and 1 prospective²⁴⁻³² study investigated a total of 4929 older adults from 6 different countries (i.e. France, Greece, Korea, Spain, Portugal, United States of America) between 1983 and 2020. Most studies were performed with community-dwellers,^{24-29,31,32} whereas 1 study investigated people resident in a nursing home.³⁰ The mean age of study participants was of approximately 71 years old and the mean prevalence of women in the samples ranged from 48.8% to 70.6%. Four studies reported values to calculate protein intake adjusted for body weight (g/kg of body weight [BW]/d), while the other 4 studies only provided data relative

Table 2. General description of the included studies.

AUTHORS	COUNTRY	STUDY DESIGN	SETTING	SAMPLE SIZE	MEAN AGE	FEMALE PREVALENCE (%)	DIETARY INTAKE ASSESSMENT METHOD	PROTEIN INTAKE (G/KG OF BW/DAY)	COGNITIVE SCORES	COGNITIVE TESTS	COGNITIVE FUNCTIONS
Li et al ²²	USA	Cross-sectional	Community-dwelling	2460	69.4	52.5	24-h dietary recall	0.92	Recall Score: 6.36; Delayed Recall Score: 6.01; Animal Fluency Score: 16.79; Digit Symbol Score: 46.83	(a) Word List Learning Test, (b) Word List Recall Test, (c) Animal Fluency Test, (d) Digit Symbol Substitution Test	Immediate and delayed learning ability, verbal fluency, processing speed, sustained attention and working memory
Katsiardanis et al ³¹	Greece	Cross-sectional	Community-dwelling	557	65+	57.5	Food Frequency Questionnaire	~78.4*	MMSE: 22.7 in men; 21.1 in women	Mini-Mental State Examination	Global cognitive function
Vizuite et al ³⁰	Spain	Cross-sectional	Nursing home	178	~81.6	~62.3	Precise weighing methods and food record	-1.0	SPMSQ: 1.38	Short Portable Mental State Questionnaire	Global cognitive function
Velho et al ²⁹	Portugal	Prospective	Community-dwelling	187	69.7	70.6	3-d dietary intake record	~75.1*	—	Mini-Mental State Examination	Global cognitive function
Lee et al ²⁸	Korea	Cross-sectional	Community-dwelling	449	~70.9	53.3	24-h dietary recall	-0.90	MMSE: 23.9 in men; 21.65 in women	Mini-Mental State Examination	Global cognitive function
Ortega et al ²⁷	Spain	Cross-sectional	Community-dwelling	260	~71.0	58.5	Weighed-food record	-1.1	MMSE: 28.6 in men; 26.2 in women	Mini-Mental State Examination	Global cognitive function
La Rue et al ²⁶	USA	Cross-sectional	Community-dwelling	137	76.9	51.1	3-d dietary intake record	72*	Rey-Osterrieth Complex Figure, copy: 31.0; recall: 14.7; Wechsler Memory Scale Visual Reproduction: 6.0; Logical Memory: 7.2; Shipley-Hartford Abstraction: 10.4	(a) Abstraction scale, (b) Logical memory test, (c) Visual Rey-Osterrieth Complex Figure test	Abstract reasoning, verbal and nonverbal memory, nonverbal learning and memory
Pradignac et al ²⁵	France	Cross-sectional	Community-dwelling	441	76.3	48.8	3-d dietary intake record	66.6*	—	Mini-Mental State Examination	Global cognitive function
Goodwin et al ²⁴	USA	Cross-sectional	Community-dwelling	260	~71.7	54.2	3-d dietary intake record	—	—	(a) Wechsler Memory Test and (b) Halstead-Reitan Categories Test	Short-term memory, abstract reasoning and problem solving ability

Abbreviations: BW, body weight; USA, United States of America.

*g/d.

Table 3. Study quality.

STUDY	1	2	3	4	5	6	7	8	9	10	11	12	13	14	OVERALL SCORE (0/12)	OVERALL SCORE (QUALITATIVE)
Li et al ³²	Y	Y	Y	Y	N	N	N	N	Y	NA	Y	Y	NA	Y	8	Good
Katsiardanis et al ³¹	Y	Y	Y	Y	N	N	N	N	Y	NA	Y	NR	NA	Y	7	Good
Vizuete et al ³⁰	Y	Y	NR	Y	N	N	N	N	Y	NA	Y	NR	NA	N	6	Fair
Velho et al ²⁹	Y	Y	NR	Y	N	N	N	N	Y	NA	Y	NR	Y	Y	7	Good
Lee et al ²⁸	Y	Y	NR	Y	N	N	N	Y	Y	NA	Y	NR	Y	N	7	Fair
Ortega et al ²⁷	Y	Y	NR	Y	Y	N	N	N	Y	NA	Y	NR	NR	N	9	Fair
La Rue et al ²⁶	Y	Y	NR	Y	N	N	N	N	Y	NA	Y	NR	NR	N	5	Fair
Pradignac et al ²⁵	Y	Y	NR	Y	N	N	N	Y	Y	NA	Y	NR	NR	Y	7	Good
Goodwin et al ²⁴	Y	Y	NR	Y	N	N	N	N	Y	NA	Y	NR	NR	N	5	Fair

Abbreviations: N, No; NA, not applied; NR, not reported; Y, Yes.

1. Was the research question or objective in this paper clearly stated?; 2. Was the study population clearly specified and defined?; 3. Was the participation rate of eligible persons at least 50%?; 4. Were all the subjects selected or recruited from the same or similar populations (including the same time period)? Were inclusion and exclusion criteria for being in the study prespecified and applied uniformly to all participants?; 5. Was a sample size justification, power description, or variance and effect estimates provided?; 6. For the analyses in this paper, were the exposure(s) of interest measured prior to the outcome(s) being measured?; 7. Was the timeframe sufficient so that one could reasonably expect to see an association between exposure and outcome if it existed?; 8. For exposures that can vary in amount or level, did the study examine different levels of the exposure as related to the outcome (eg, categories of exposure, or exposure measured as continuous variable)?; 9. Were the exposure measures (independent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?; 10. Was the exposure(s) assessed more than once over time?; 11. Were the outcome measures (dependent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?; 12. Were the outcome assessors blinded to the exposure status of participants?; 13. Was loss to follow-up after baseline 20% or less?; 14. Were key potential confounding variables measured and adjusted statistically for their impact on the relationship between exposure(s) and outcome(s)?.

to crude protein intake per day. Mean protein intake values were higher than the RDA¹⁶ and ranged from 0.90 g/kg of BW/d to 1.1 g/kg of BW/d. Dietary intake was assessed using 24-hour dietary recalls, 3-day dietary intake records, Food Frequency Questionnaires (FFQ), and precise weighing methods. Many cognitive tests were performed, including the Mini-Mental State Examination (MMSE), Short Portable Mental State Questionnaire, Word List Learning and Word List Recall Test from the Consortium to Establish a Registry for Alzheimer's disease (CERAD), Pfeiffer's Mental Status Questionnaire (PMSQ), Animal Fluency Test, Digit Symbol Substitution Test (DSST), The abstraction scale from the Shipley-Hartford Intelligence Test, The Logical Memory and Visual Reproduction subtests from the Wechsler Memory Scale (WMS), Rey-Osterrieth Complex Figure test, Wechsler Memory Test and Halstead-Reitan Categories Test. These tests were designed to evaluate overall cognitive function, immediate and delayed learning ability, nonverbal learning, verbal and nonverbal memory, short-term memory, working memory, verbal fluency, processing speed, problem-solving ability, sustained attention, and abstract reasoning.

Quality assessment

The qualitative analysis of the included studies is shown in Table 3. The overall quantitative score ranged from 5 to 9 of 12 possible points, while the qualitative score ranged from fair to good. All studies clearly stated their objectives (item 1), specified the studied population (item 2), recruited study participants

in the same or similar population (item 4), and described and used valid and reliable instruments to assess protein intake (item 9) and cognitive function (item 11). Items 6 and 7 refer to parameters associated with cohort studies and therefore were not reported in any of the included studies. The participation rate of the eligible person (item 3) and loss to follow-up after baseline (item 13) were only reported in 22% of the studies each. Different levels of exposure (item 8), in this case evaluated according to the sources of protein, were also only investigated in 2 studies (22%). Sample size (item 5) was only justified in one (11.1%) study. Similarly, just 11.1% of the studies described if assessors were blinded to exposure status (item 12). The most heterogeneous result is regarding the adjustment for potential confounding variables (item 14), given that this variable was controlled in 44.4% of the studies. None of the studies assessed exposure more than once (item 10).

Association between protein intake and overall cognitive function

Six studies^{25,27-31} investigated the association between protein intake and overall cognitive function in older adults. Study results were highly heterogeneous. Four studies^{25,27,29,31} observed null associations. Pradignac et al²⁵ studied apparently healthy community-dwelling French older adults and observed no significant associations between protein intake and global cognitive function. Similar results were found by Velho et al²⁹ in physically active Portuguese older people. In older adults from a rural Greece area, Katsiardanis et al³¹ did not observe




Figure 2. Standard Mean Differences (SMD) in global cognitive function according to protein intake in: (a) all participants, (b) women, and (c) men.

significant associations between diary protein intake and global cognitive scores. Ortega et al²⁷ confirmed and expanded these findings by indicating that protein consumption was not associated with neither MMSE nor PMSQ scores. In contrast, 2 studies^{28,30} reported positive associations. Lee et al²⁸ found a significant association between MMSE scores and total and vegetal protein intakes in older women, but not men, who participated of social and physical activities in a welfare center. One study investigated institutionalized older adults without a clinical diagnosis of MCI and/or dementia. In this study, Vizuite et al³⁰ reported better scores on PMSQ as protein intake increased.

Only 3 studies^{27,28,31} provided data to be included in the meta-analysis (Figure 2). No significant associations between global cognitive function and protein intake were observed, regardless of gender.

Association between protein intake and other cognitive domains

Three studies^{24,26,32} investigated the association between protein intake and immediate and delayed learning ability, nonverbal learning, verbal and nonverbal memory, short-term memory, working memory, verbal fluency, processing speed, problem-solving ability, sustained attention, and abstract reasoning in older adults. Goodwin et al²⁴ studied older adults who living independently and observed that those subjects in the bottom 5% or 10% of protein intake tended to do poorly on Halstead-Reitan Categories Test, which reflects abstract thinking and problem-solving ability, and WMS, a test designed to measure different subdomains of memory. However, results were only significant for the memory test. La Rue et al²⁶ confirmed these

findings by indicating a positive association between protein intake and Rey-Osterrieth Recall, representing a measurement of visuospatial skill, and WMS logical memory scores in apparently healthy older adults of the New Mexico Aging Process Study. However, no significant correlations were found with Rey-Osterrieth Copy, which assess visuospatial skill and non-verbal memory, and the Shipley-Hartford Intelligence Test, assessing abstract reasoning. Li et al³² investigated data of the National Health and Nutrition Examination Survey (NHANES), which contained multiple cognitive tests. Authors noted that a protein intake higher than 0.6 g/kg of BW per day was significantly associated with a better performance on the Animal Fluency Test, designed to examines categorical verbal fluency, and Digit Symbol Score, designed to assess processing speed, sustained attention and working memory regardless of gender.

Overall, 3 studies^{24,26,32} indicated that a high protein intake was positively associated with memory, assessed according to different assessment tools; 1 study²⁶ observed a significant association with visuospatial skill; and 1 study³² reported a positive correlation with categorical verbal fluency, processing speed, and sustained attention.

Discussion

The present systematic study investigated the association between protein intake and cognitive function in older adults. After screening 2915 records, 9 cross-sectional studies fulfilled all the requirements to be included in the present investigation. Examined studies varied in many methodological aspects, including sample size, dietary intake assessment method, cognitive tests, adjusting factors, and statistical approach. Results of the included investigations were also highly heterogeneous,

with 4 studies^{25,27,29,31} reporting null associations, whereas 2 investigations^{28,30} found a significant positive association between protein intake and global cognitive function. No significant associations were observed by pooling data in the meta-analysis. Though, only 3 studies were included. The association between protein consumption with multiple other cognitive domains were also tested. As a whole, 3 studies^{24,26,32} reported a positive and significant association between high protein intake and memory; 1 study²⁶ observed a significant and positive association with visuospatial skills; and 1 study³² reported a positive correlation with categorical verbal fluency, processing speed, and sustained attention. However, no significant associations were observed with many other tests, including Halstead-Reitan Categories Test, Rey-Osterrieth Copy, and the Shipley-Hartford Intelligence Test.

The quality analysis revealed that most investigations failed to report many important methodological aspects, including participation rate of the eligible person, loss to follow-up after baseline, sample size calculation, blindness, different levels of exposure, and the trustworthiness of the exposure.

Particularly, the level of exposure, which was evaluated according to the source of protein, was only investigated in 2 studies. Lee et al²⁸ examined community-dwelling Korean older adults and reported that both total and vegetal protein, but not animal protein, were significantly associated with global cognitive function in women. In contrast, Pradignac et al²⁵ did not observe a significant association between protein-related parameters and cognitive function in a sample of community-dwelling French older adults.

This nutritional aspect deserves concern, given that animal protein is expected to provide higher amounts of BCAA than vegetal protein.⁴²⁻⁴⁴ BCAAs are directly and indirectly involved in numerous metabolic processes in the neural system.⁴⁵⁻⁴⁹ However, much attention has been paid to their effects on GLU metabolism, given that it might serve as a possible mechanistic explanation for the association between protein intake and cognitive function.⁴⁹

GLU is the major excitatory transmitter in the mammalian central nervous system.⁵⁰⁻⁵³ This molecule might impact neuronal differentiation, migration, and survival during maturation and cognitive functioning throughout life.⁵⁰⁻⁵³ The hypothesis that GLU can affect cognition is based on the fact that its receptors are widely distributed in pre- and post-synaptic neurons, influencing neuronal communication and signal processing.⁵⁰⁻⁵³ Particularly, N-methyl-D-aspartate (NMDA), one of the 3 types of ionotropic GLU receptors, is the predominant molecular device for controlling synaptic plasticity and for the proper formation of memory.^{54,55}

GLU concentrations in the brain are tightly controlled.⁴⁵ In the intra-neuronal space, GLU supply must be kept constant and at optimum concentrations to maintain neuronal depolarization. On the other hand, high GLU concentrations in the extracellular space might induce excitotoxic injury and even kill

susceptible neurons.⁵⁶ Hence, the transport of GLU across the blood brain barrier (BBB) and the synaptic space is limited to promote proper cerebral functioning and minimize the risk of harmful effects.⁴⁵

GLU concentrations are also regulated by the GLU/glutamine (GLU/GLN) cycle.⁴⁵ During neurotransmission, GLU is released in the synapses through a calcium-dependent process that involves the fusion of GLU-containing presynaptic vesicles with the neuronal membrane.⁵⁶ Not all GLU is uptake into the postsynaptic compartment and reuptake into presynaptic neurons.^{45,56} This scenario is problematic to the brain, which must keep GLU levels continuous for new neurotransmission and avoid its traffic to the extracellular fluid.⁴⁵ Astrocytes, star-shaped glial cells involved in many neurological functions, such as biochemical homeostasis, sequester GLU from the synaptic space by the use of GLU-specific transporters and convert it into a non-neuroactive component GLN via the glutamine synthetase enzyme.^{45,56}

Then, GLN is released to the extracellular space when it cannot cause depolarization or harmful effects.⁴⁵ Finally, GLN is reuptake into nerve endings via sodium-dependent and independent mechanisms where it is converted back into GLU via glutaminase enzyme.^{45,56}

However, it is important to note that the current understanding of the GLU/GLA cycle has been believed to be limited, simplified, and incomplete to explain how the brain compensates GLU catabolism.⁴⁶ Hence, Yudkoff⁴⁶ proposed that peripheral BCAA is required to cross the BBB to provide the necessary amount of nitrogen to keep GLU synthesis at adequate rates. This mechanism involves the neural transamination of BCAA by cytosolic and mitochondrial enzymes, leading to the formation of GLU and α -ketoisocaproate.⁴⁶ The role of this model on proper cognitive functioning seems to increase with age, given that age-related cognitive decline can be at least partially explained by changes in GLU concentrations and transportation in central areas dedicated to learning and memory.⁵⁷⁻⁵⁹

These premises suggest that the maintaining of GLU homeostasis in the brain due to an adequate peripheral BCAA content might explain the significant and positive associations between high protein intake and some cognitive domains. Particularly, 3 studies^{24,26,32} reported that older adults with a high protein consumption had better results in memory tests, and multiple studies have shown that the activation of NMDA receptors is required to induce long-term potentiation (LTP),^{51,54,60-62} a neuronal event that represents an increased synaptic transmission among neurons argued to be the main mechanism underlying memory encoding. In contrast, LTP in cortical and hippocampal (i.e. CA1 and dental gyrus) areas was blocked by antagonists of the NMDA receptors.⁶⁰⁻⁶²

No significant associations between protein intake and global cognitive function were observed in neither qualitative nor quantitative analyses. The main reason why high protein

intake might benefit some cognitive domains, but not overall cognitive function is unknown. However, it is important to note that our results were based in a small number of studies and inferences must be made with caution, given the high variability in measuring methods, study outcomes, and participant characteristics, besides the lack of sensitivity analysis in most of the studies.

Indeed, global cognitive function were assessed using MMSE and PMSQ. Although both tests claim to measure the same cognitive domain, only a moderate agreement has been observed among the scales for screening negative events,^{63,64} which suggests that MMSE and PMSQ capture different dimensions of cognition. Moreover, each test involves recognized limitations. The MMSE, for example, has a ceiling effect and poor content validity in assessing language, visuospatial, and executive functions,^{65,66} while PMSQ presents a low sensitivity and specificity for screening some cases of dementia, which led some authors to suggest that this tool should not be used alone in certain clinical settings.^{63,64}

Another important aspect involving cognitive assessment is that most cognitive tests do not assess only 1 cognitive domain, but involves 2 or more areas of cognition, limiting conclusions about specific cognitive functions.⁶⁷ The DSST, for example, was initially designed to assess associative learning.⁶⁸ Currently, it is accepted that the performance on the DSST test correlates with memory, motor speed, attention, and visuo-perceptual functions.⁶⁸

Five different dietary assessment methods (i.e. 3-day dietary intake, 24-hour dietary recall, FFQ, weighed-food record, and precise weighing methods and food record) were used in the studies that investigated global cognitive function. All these methods have strengths and limitations, and specific guidelines for guide their application in older adults are still missing.⁶⁹ Hence, experts in the field⁶⁹ argued that the reliability of dietary information is also dependent on the approach used to collect data, so that long interviews and questionnaires may be stressful to many older adults, leading to incomplete or unreliable results.

The lack of adjust of the results for potential confounding variables by using complex statistical methods (e.g. multiple regression) is one more potential source of heterogeneity. In fact, most of the included studies based their conclusions on the differences between the means of 2 groups, whereas 1 study³² provided adjusted odds ratio. Notably, the effects of protein intake on cognition in older adults might be influenced by its relationship with numerous health-related parameters (to review see Glenn et al³⁴). Physical dysfunction^{70,71} and physical frailty,^{72,73} for example, are predictors of cognitive decline in older adults, and numerous studies have supported a significant relationship between protein intake and physical performance.²⁰⁻²³ Acute and long-term sleep deprivation impairs memory, language, executive function, and attention domains,⁷⁴ and are highly frequent in people with dementia.⁷⁵ Besides, cross-sectional studies using nationally representative samples have observed that people with

sleep disturbance had a low protein intake.^{35,36} More recently, the role of microbiota on cognitive impairment⁷⁶ and severe mental disorders⁷⁷ has received increased attention. These data have important clinical implications since protein intake and microbiota are interconnected so that the structure and function of the microbiome might be impacted by protein quality, while it influences protein catabolism, digestion, and absorption.³⁷


Taken together, these observations indicate the future studies should investigate global cognition using standard methods with high sensitivity and specificity to predict negative outcomes, utilize validated methods to assess dietary intake and provide a detailed description of the approach used to data collection, and adjust findings by potential confounding factors according to the use of sensitive statistical methods. Preferably, studies should be conducted taken into consideration the current literature and assess the main variables using the same or similar methods to allow comparisons and better inferences. Finally, studies investigating other cognitive domains (e.g. executive function) are still needed.

This study is not free of limitations. First, we only included cross-sectional studies, limiting the establishment of a cause-effect relationship. Second, our sample was composed exclusively of older adults free of dementia and inferences to other populations must be made with caution. Third, most studies provided data of total protein intake. However, numerous studies and experts in the field have suggested that other parameters, including protein intake adjusted by BW, protein distribution across the meals, and protein sources might provide a clearer understanding of protein consumption.^{20-23,78,79}

Conclusion

The association between protein intake and cognition in older adults has attracted considerable attention. However, no specific recommendations are available. In the present study, no significant associations between protein intake and global cognitive function were observed in neither qualitative nor quantitative analyses. The association between protein consumption with multiple other cognitive domains were also tested. As a whole, 3 studies reported a positive and significant association between high protein intake and memory; 1 study observed a significant and positive association with visuospatial skills; and 1 study reported a positive correlation with categorical verbal fluency, processing speed, and sustained attention.

ORCID iDs

Hélio José Coelho-Júnior  <https://orcid.org/0000-0001-7482-9514>

Emanuele Marzetti  <https://orcid.org/0000-0001-9567-6983>

REFERENCES

1. American Psychological Association (APA). Dictionary of Psychology. Accessed April 9, 2021. <https://dictionary.apa.org/cognition>

2. Murman DL. The impact of age on cognition. *Semin Hear*. 2015;36:111-121. doi:10.1055/s-0035-1555115
3. Lee JH, Sung J, Choi MK. The factors associated with subjective cognitive decline and cognitive function among older adults. *J Adv Nurs*. 2020;76:555-565. doi:10.1111/jan.14261
4. Montero-Odasso M, Speechley M. Falls in cognitively impaired older adults: implications for risk assessment and prevention. *J Am Geriatr Soc*. 2018;66:367-375. doi:10.1111/jgs.15219
5. Fraade-Blanar LA, Ebel BE, Larson EB, et al. Cognitive decline and older driver crash Risk. *J Am Geriatr Soc*. 2018;66:1075-1081. doi:10.1111/jgs.15378
6. Brigola AG, Ottaviani AC, Alexandre T da S, Luchesi BM, Pavarini SCI. Cumulative effects of cognitive impairment and frailty on functional decline, falls and hospitalization: a four-year follow-up study with older adults. *Arch Gerontol Geriatr*. 2020;87:104005. doi:10.1016/j.archger.2019.104005
7. van Harten AC, Mielke MM, Swenson-Dravis DM, et al. Subjective cognitive decline and risk of MCI: the Mayo Clinic Study of Aging. *Neurology*. 2018;91:e300-e312. doi:10.1212/WNL.0000000000005863
8. Lv X, Li W, Ma Y, et al. Cognitive decline and mortality among community-dwelling Chinese older people. *BMC Med*. 2019;17:63. doi:10.1186/s12916-019-1295-8
9. World Health Organization (WHO). Risk reduction of cognitive decline and dementia. 2019. Accessed February 2, 2021. <https://www.who.int/publications/i/item/risk-reduction-of-cognitive-decline-and-dementia>
10. Erickson KI, Hillman C, Stillman CM, et al. Physical activity, cognition, and brain outcomes. *Med Sci Sports Exerc*. 2019;51:1242-1251. doi:10.1249/MSS.0000000000001936
11. Lee J. The relationship between physical activity and dementia: a systematic review and meta-analysis of prospective cohort studies. *J Gerontol Nurs*. 2018;44:22-29. doi:10.3928/00989134-20180814-01
12. Coelho-Júnior HJ, Oliveira Gonçalves I de, Sampaio RAC, et al. Effects of combined resistance and power training on cognitive function in older women: a randomized controlled trial. *Int J Environ Res Public Health*. 2020;17:3435. doi:10.3390/ijerph17103435
13. Coelho-Junior H, Marzetti E, Calvani R, Picca A, Arai H, Uchida M. Resistance training improves cognitive function in older adults with different cognitive status: a systematic review and meta-analysis. *Aging Ment Health*. Published online December 16, 2020. doi:10.1080/13607863.2020.1857691
14. Lourida I, Soni M, Thompson-Coon J, et al. Mediterranean diet, cognitive function, and dementia: a systematic review. *Epidemiology*. 2013;24:479-489. doi:10.1097/EDE.0b013e3182944410
15. Chen X, Maguire B, Brodaty H, O'Leary F. Dietary patterns and cognitive health in older adults: a systematic review. *J Alzheimers Dis*. 2019;67:583-619. doi:10.3233/JAD-180468
16. *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (Macronutrients)*. National Academies Press; 2005. doi:10.17226/10490
17. Volpi E, Campbell WW, Dwyer JT, et al. Is the optimal level of protein intake for older adults greater than the recommended dietary allowance? *J Gerontol A Biol Sci Med Sci*. 2013;68:677-681. doi:10.1093/gerona/gls229
18. Bauer J, Biolo G, Cederholm T, et al. Evidence-based recommendations for optimal dietary protein intake in older people: a position paper from the prot-age study group. *J Am Med Dir Assoc*. 2013;14:542-559. doi:10.1016/j.jamda.2013.05.021
19. Calvani R, Miccheli A, Landi F, et al. Current nutritional recommendations and novel dietary strategies to manage sarcopenia. *J Frailty Aging*. 2013;2:38-53.
20. Coelho-Junior HJ, Calvani R, Gonçalves IO, et al. High relative consumption of vegetable protein is associated with faster walking speed in well-functioning older adults. *Aging Clin Exp Res*. 2019;31:837-844. doi:10.1007/s40520-019-01216-4
21. Coelho-Junior HJ, Calvani R, Picca A, et al. Association between dietary habits and physical function in Brazilian and Italian older women. *Nutrients*. 2020;12:1635. doi:10.3390/nu12061635
22. Coelho-Junior H, Rodrigues B, Uchida M, et al. Low protein intake is associated with frailty in older adults: a systematic review and meta-analysis of observational studies. *Nutrients*. 2018;10:1334. doi:10.3390/nu10091334
23. Coelho-Junior H, Milano-Teixeira L, Rodrigues B, Bacurau R, Marzetti E, Uchida M. Relative protein intake and physical function in older adults: a systematic review and meta-analysis of observational studies. *Nutrients*. 2018;10:1330. doi:10.3390/nu10091330
24. Goodwin JS, Goodwin JM, Garry PJ. Association between nutritional status and cognitive functioning in a healthy elderly population. *JAMA*. 1983;249:2917-2921. doi:10.1001/jama.1983.03330450047024
25. Pradignac A, Schlienger JL, Velten M, Mejean L. Relationships between macronutrient intake, handicaps, and cognitive impairments in free living elderly people. *Aging Clin Exp Res*. 1995;7:67-74. doi:10.1007/BF03324295
26. La Rue A, Koehler KM, Wayne SJ, Chiulli SJ, Haaland KY, Garry PJ. Nutritional status and cognitive functioning in a normally aging sample: a 6-y reassessment. *Am J Clin Nutr*. 1997;65:20-29. doi:10.1093/ajcn/65.1.20
27. Ortega RM, Requejo AM, Andrés P, et al. Dietary intake and cognitive function in a group of elderly people. *Am J Clin Nutr*. 1997;66:803-809. doi:10.1093/ajcn/66.4.803
28. Lee L, Kang S, Lee H, et al. Relationships between dietary intake and cognitive function level in Korean elderly people. *Public Health*. 2001;115:133-138. doi:10.1038/sj.ph.1900729
29. Velho S, Marques-Vidal P, Baptista F, Camilo ME. Dietary intake adequacy and cognitive function in free-living active elderly: a cross-sectional and short-term prospective study. *Clin Nutr*. 2008;27:77-86. doi:10.1016/j.clnu.2007.10.011
30. Vizuete AA, Robles F, Rodríguez-Rodríguez E, López-Sobaler AM, Ortega RM. Association between food and nutrient intakes and cognitive capacity in a group of institutionalized elderly people. *Eur J Nutr*. 2010;49:293-300. doi:10.1007/s00394-009-0086-y
31. Katsiardanis K, Diamantaras AA, Dessypris N, et al. Cognitive impairment and dietary habits among elders: the vellestino study. *J Med Food*. 2013;16:343-350. doi:10.1089/jmf.2012.0225
32. Li Y, Li S, Wang W, Zhang D. Association between dietary protein intake and cognitive function in adults aged 60 years and older. *J Nutr Health Aging*. 2020;24:223-229. doi:10.1007/s12603-020-1317-4
33. Koh F, Charlton K, Walton K, McMahon A-T. Role of dietary protein and thiamine intakes on cognitive function in healthy older people: a systematic review. *Nutrients*. 2015;7:2415-2439. doi:10.3390/nu7042415
34. Glenn JM, Madero EN, Bott NT. Dietary protein and amino acid intake: links to the maintenance of cognitive health. *Nutrients*. 2019;11:1315. doi:10.3390/nu11061315
35. Kant AK, Graubard BI. Association of self-reported sleep duration with eating behaviors of American adults: NHANES 2005-2010. *Am J Clin Nutr*. 2014;100:938-947. doi:10.3945/ajcn.114.085191
36. Grandner MA, Jackson N, Gerstner JR, Knutson KL. Dietary nutrients associated with short and long sleep duration. Data from a nationally representative sample. *Appetite*. 2013;64:71-80. doi:10.1016/j.appet.2013.01.004
37. Zhao J, Zhang X, Liu H, Brown MA, Qiao S. Dietary protein and gut microbiota composition and function. *Curr Protein Pept Sci*. 2018;20:145-154. doi:10.2174/1389203719666180514145437
38. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med*. 2009;6:e1000100. doi:10.1371/journal.pmed.1000100
39. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA*. 2000;283:2008-2012. doi:10.1001/jama.283.15.2008
40. NHLBI, NIH. Study quality assessment tools. Accessed October 25, 2020. <https://www.nlm.nih.gov/health-topics/study-quality-assessment-tools>
41. Green S, Higgins J. Cochrane handbook for systematic reviews of interventions. Published online 2005. Accessed July 21, 2018. <https://training.cochrane.org/handbook>
42. van Vliet S, Burd NA, van Loon LJ. The skeletal muscle anabolic response to plant- versus animal-based protein consumption. *J Nutr*. 2015;145:1981-1991. doi:10.3945/jn.114.204305
43. Landi F, Calvani R, Tosato M, et al. Protein intake and muscle health in old age: from biological plausibility to clinical evidence. *Nutrients*. 2016;8:295. doi:10.3390/nu8050295
44. Coelho-Junior HJ, Marzetti E, Picca A, Cesari M, Uchida MC, Calvani R. Protein intake and frailty: a matter of quantity, quality, and timing. *Nutrients*. 2020;12:1-20. doi:10.3390/nu12102915
45. Sperringer JE, Addington A, Hutson SM. Branched-chain amino acids and brain metabolism. *Neurochem Res*. 2017;42:1697-1709. doi:10.1007/s11064-017-2261-5
46. Yudkoff M. Interactions in the metabolism of glutamate and the branched-chain amino acids and ketoacids in the CNS. *Neurochem Res*. 2017;42:10-18. doi:10.1007/s11064-016-2057-z
47. Griffin JWD, Bradshaw PC. Amino acid catabolism in Alzheimer's disease brain: friend or foe? *Oxid Med Cell Longev*. 2017;2017:5472792. doi:10.1155/2017/5472792
48. Fernstrom JD. Large neutral amino acids: dietary effects on brain neurochemistry and function. *Amino Acids*. 2013;45:419-430. doi:10.1007/s00726-012-1330-y
49. Fernstrom JD. Branched-chain amino acids and brain function. *J Nutr*. 2005;135:1539S-1546S. doi:10.1093/jn/135.6.1539s
50. Zhou Y, Danbolt NC. Glutamate as a neurotransmitter in the healthy brain. *J Neural Transm*. 2014;121:799-817. doi:10.1007/s00702-014-1180-8
51. Riedel G, Platt B, Micheau J. Glutamate receptor function in learning and memory. *Behav Brain Res*. 2003;140:1-47. doi:10.1016/S0166-4328(02)00272-3
52. McEntee WJ, Crook TH. Glutamate: its role in learning, memory, and the aging brain. *Psychopharmacology (Berl)*. 1993;111:391-401. doi:10.1007/BF02253527
53. Meldrum BS. Glutamate as a neurotransmitter in the brain: review of physiology and pathology. *J Nutr*. 2000;130:1007S-1015S. doi:10.1093/jn/130.4.1007s

54. Nakazawa K, McHugh TJ, Wilson MA, Tonegawa S. NMDA receptors, place cells and hippocampal spatial memory. *Nat Rev Neurosci.* 2004;5:361-372. doi:10.1038/nrn1385
55. Li F, Tsien JZ. Memory and the NMDA receptors. *N Engl J Med.* 2009;361:302-303. doi:10.1056/nejmcibr0902052
56. Daikhin Y, Yudkoff M. Compartmentation of brain glutamate metabolism in neurons and glia. *J Nutr.* 2000;130:1026S-1031S. doi:10.1093/jn/130.4.1026s
57. Potier B, Billard JM, Rivière S, et al. Reduction in glutamate uptake is associated with extrasynaptic NMDA and metabotropic glutamate receptor activation at the hippocampal CA1 synapse of aged rats. *Aging Cell.* 2010;9:722-735. doi:10.1111/j.1474-9726.2010.00593.x
58. Price MT, Olney JW, Haft R. Age-related changes in glutamate concentration and synaptosomal glutamate uptake in adult rat striatum. *Life Sci.* 1981;28:1365-1370. doi:10.1016/0024-3205(81)90410-0
59. Dawson R, Wallace DR, Meldrum MJ. Endogenous glutamate release from frontal cortex of adult and aged rats. *Neurobiol Aging.* 1989;10:665-668. doi:10.1016/0197-4580(89)90002-X
60. Errington ML, Lynch MA, Bliss TVP. Long-term potentiation in the dentate gyrus: induction and increased glutamate release are blocked by d(-)aminophosphonovalerate. *Neuroscience.* 1987;20:279-284. doi:10.1016/0306-4522(87)90019-4
61. Huerta PT, Sun LD, Wilson MA, Tonegawa S. Formation of temporal memory requires NMDA receptors within CA1 pyramidal neurons. *Neuron.* 2000;25:473-480. doi:10.1016/S0896-6273(00)80909-5
62. Artola A, Singer W. Long-term potentiation and NMDA receptors in rat visual cortex. *Nature.* 1987;330:649-652. doi:10.1038/330649a0
63. Mackenzie DM, Copp P, Shaw RJ, Goodwin GM. Brief cognitive screening of the elderly: a comparison of the Mini-Mental State Examination (MMSE), Abbreviated Mental Test (AMT) and Mental Status Questionnaire (MSQ). *Psychol Med.* 1996;26:427-430. doi:10.1017/s0033291700034826
64. Malhotra C, Chan A, Matchar D, et al. Diagnostic performance of Short Portable Mental Status Questionnaire for screening dementia among patients attending cognitive assessment clinics in Singapore. *Ann Acad Med Singapore.* 2013;42:315-319.
65. Mitchell AJ, Shiri-Feshki M. Rate of progression of mild cognitive impairment to dementia - meta-analysis of 41 robust inception cohort studies. *Acta Psychiatr Scand.* 2009;119:252-265. doi:10.1111/j.1600-0447.2008.01326.x
66. Spencer RJ, Wendell CR, Giggey PP, et al. Psychometric limitations of the mini-mental state examination among nondemented older adults: an evaluation of neurocognitive and magnetic resonance imaging correlates. *Exp Aging Res.* 2013;39:382-397. doi:10.1080/0361073X.2013.808109
67. Knight A, Bryan J, Murphy K. The Mediterranean diet and age-related cognitive functioning: a systematic review of study findings and neuropsychological assessment methodology. *Nutr Neurosci.* 2017;20:449-468. doi:10.1080/1028415X.2016.1183341
68. Jaeger J. Digit symbol substitution test. *J Clin Psychopharmacol.* 2018;38:513-519. doi:10.1097/JCP.0000000000000941
69. Volkert D, Schrader E. Dietary assessment methods for older persons. *Curr Opin Clin Nutr Metab Care.* 2013;16:534-540. doi:10.1097/MCO.0b013e328363c8d1
70. Auyeung TW, Kwok T, Lee J, Leung PC, Leung J, Woo J. Functional decline in cognitive impairment - the relationship between physical and cognitive function. *Neuroepidemiology.* 2008;31:167-173. doi:10.1159/000154929
71. Clouston SAP, Brewster P, Kuh D, et al. The dynamic relationship between physical function and cognition in longitudinal aging cohorts. *Epidemiol Rev.* 2013;35:33-50. doi:10.1093/epirev/mxs004
72. Fougère B, Dumas M, Lilamand M, et al. Association between frailty and cognitive impairment: cross-sectional data from Toulouse frailty day hospital. *J Am Med Dir Assoc.* 2017;18:990.e1-990.e5. doi:10.1016/j.jamda.2017.06.024
73. Borges MK, Canevelli M, Cesari M, Aprahamian I. Frailty as a predictor of cognitive disorders: a systematic review and meta-analysis. *Front Med.* 2019;6:26. doi:10.3389/fmed.2019.00026
74. Alhola P, Polo-Kantola P. Sleep deprivation: impact on cognitive performance. *Neuropsychiatr Dis Treat.* 2007;3:553-567.
75. McCurry SM, Logsdon RG, Teri L, et al. Characteristics of sleep disturbance in community-dwelling Alzheimer's disease patients. *J Geriatr Psychiatry Neurol.* 1999;12:53-59. doi:10.1177/089198879901200203
76. Novotný M, Klimova B, Valis M. Microbiome and cognitive impairment: can any diets influence learning processes in a positive way? *Front Aging Neurosci.* 2019;11:170. doi:10.3389/fnagi.2019.00170
77. Sani G, Manchia M, Simonetti A, et al. The role of gut microbiota in the high-risk construct of severe mental disorders: a mini review. *Front Psychiatry.* 2021;11:585769. doi:10.3389/fpsy.2020.585769
78. Loenneke JP, Loprinzi PD, Murphy CH, Phillips SM. Per meal dose and frequency of protein consumption is associated with lean mass and muscle performance. *Clin Nutr.* 2016;35:1506-1511. doi:10.1016/j.clnu.2016.04.002
79. Ten Haaf DSM, van Dongen EJI, Nuijten MAH, Eijsvogels TMH, de Groot LCPGM, Hopman MTE. Protein intake and distribution in relation to physical functioning and quality of life in community-dwelling elderly people: acknowledging the role of physical activity. *Nutrients.* 2018;10:506. doi:10.3390/nut10040506