

This article was downloaded by: [Soldatini, Celcilia]

On: 14 April 2011

Access details: Access Details: [subscription number 936393826]

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Bird Study

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t904369352>

### Diet and diving behaviour of European Storm Petrels *Hydrobates pelagicus* in the Mediterranean (ssp. *melitensis*)

Y. V. Albores-Barajas<sup>a</sup>; F. Riccato<sup>a</sup>; R. Fiorin<sup>b</sup>; B. Massa<sup>c</sup>; P. Torricelli<sup>a</sup>; C. Soldatini<sup>a</sup>

<sup>a</sup> Department of Environmental Science, University Ca' Foscari of Venezia, Campo della Celestia, I-30122 Venezia, Italy <sup>b</sup> Laguna Project SNC, Venice, Italy <sup>c</sup> Department DEMETRA (Bird Ringing Unit), University of Palermo, I-90128 Palermo, Italy

First published on: 31 March 2011

**To cite this Article** Albores-Barajas, Y. V. , Riccato, F. , Fiorin, R. , Massa, B. , Torricelli, P. and Soldatini, C.(2011) 'Diet and diving behaviour of European Storm Petrels *Hydrobates pelagicus* in the Mediterranean (ssp. *melitensis*)', *Bird Study*, 58: 2, 208 – 212, First published on: 31 March 2011 (iFirst)

**To link to this Article:** DOI: 10.1080/00063657.2011.560244

**URL:** <http://dx.doi.org/10.1080/00063657.2011.560244>

## PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

# Diet and diving behaviour of European Storm Petrels *Hydrobates pelagicus* in the Mediterranean (*ssp. melitensis*)

Y.V. ALBORES-BARAJAS<sup>1\*</sup>, F. RICCATO<sup>1</sup>, R. FIORIN<sup>2</sup>, B. MASSA<sup>3</sup>, P. TORRICELLI<sup>1</sup> and C. SOLDATINI<sup>1</sup>

<sup>1</sup>Department of Environmental Science, University Ca' Foscari of Venezia, Campo della Celestia, Castello 2737/b, I-30122 Venezia, Italy, <sup>2</sup>Laguna Project SNC, Castello 6411, Venice, Italy and <sup>3</sup>Department DEMETRA (Bird Ringing Unit), University of Palermo, Viale Scienze 13, I-90128 Palermo, Italy

**Capsule** Unlike Atlantic populations, which feed on krill, Mediterranean populations feed mainly on pelagic fish *Gymnammodites cicerellus*.

**Aims** To determine the diet and dive depth of the Mediterranean subspecies of European Storm Petrels *Hydrobates pelagicus melitensis*.

**Methods** Analysis of regurgitates of adults arriving at the colony for chick feeding and by determination of dives depth using the capillary tube method.

**Results** The main prey is *Gymnammodites cicerellus*, a pelagic fish. Storm Petrels dive for their prey and can reach up to 5 m in depth. They also make short foraging trips just outside the colony where they capture Opossum Shrimps Misydacea.

**Conclusions** European Storm Petrels in the Mediterranean exploit pelagic fish which are taken by diving. This contrasts with the Atlantic populations which feed mainly on krill. Mediterranean birds also feed on Opossum Shrimps *Mysidacea* during short foraging trips made at night just outside the colony. Differences in diet between long and short foraging trips may be because adults have to forage for both themselves and their chicks.

When devising conservation strategies for seabirds, it is important to consider not only the breeding sites of the species, but also the areas in which they forage and for which prey (Hazlitt *et al.* 2010). For seabirds, this delimitation of areas may be a difficult task, as some species make foraging trips up to several hundred kilometres away from the colony (Magalhaes *et al.* 2008). Furthermore, the colonies of many seabird species are located on inaccessible sites, which make it difficult to study their biology. Among seabirds, nocturnally active species, such as European Storm Petrels *Hydrobates pelagicus* are even less accessible to study due to their habits of breeding in burrows (Brooke 2004).

European Storm Petrels have large colonies. In the Atlantic part of their range, colonies often contain several hundred thousand breeding pairs, but in the Mediterranean, the largest colonies have between 7000

and 9000 breeding pairs (del Hoyo *et al.* 1994). Genetic analysis has suggested a division between the Atlantic and the Mediterranean populations at the subspecies level (Cagnon *et al.* 2004). Based on this genetic difference and on the decreasing number of breeding pairs per colony in the Mediterranean (Lo Valvo & Massa 2000), this Mediterranean population *Hydrobates pelagicus melitensis* can be considered an endangered subspecies, given that all individuals within the Mediterranean breed in only a few colonies (Massa & Sultana 1991).

Dietary analysis of the Atlantic populations of the European Storm Petrel found two main prey groups, krill Euphausiacea and microzooplankton (D'Elbee & Hemery 1998). Other petrel species have been observed to dive in search of prey, with recorded depths of 2 m for the Madeiran Storm Petrel *Oceanodroma castro* (Bried 2005). Given the absence of krill in the Mediterranean, it can be assumed that the diet of the Mediterranean populations is different from that of

\*Correspondence author. Email: yalbores@unive.it

the Atlantic populations. As there is no information on the diet of the Mediterranean populations or on the diving behaviour of the European Storm Petrel, our objective was to determine the diet of the Mediterranean population of this species during the breeding season and to describe their feeding behaviour.

## METHODS

The Mediterranean Storm Petrel colony in Marettimo Island, Italy (37°58'20"N 12°3'20"E) has been monitored since 1985, with about 6000 individuals ringed (Lo Valvo & Massa 2000, Sanz-Aguilar *et al.* 2009). The main colony comprises between 2000 and 2500 breeding pairs and is located in a cave accessible only from the sea. The colony is distributed in two main chambers of the cave and on several terraces. Nests are all scattered inside the cave and most of them are not inside crevices or under rocks, therefore it is easy to mark nests and study the behaviour of pairs. Due to their lack of an anti-predator response, these Storm Petrels are easily caught on their nests by hand while brooding. We monitored the colony for the breeding seasons of 2007–2009.

In order to determine the diving depth of the adults, during the three breeding seasons, we attached capillary tubes (Mougin & Mougin 2000) on rump or tail feathers (two or three feathers) of 112 birds using waterproof duct tape. Birds were caught at the nest and when possible we placed capillary tubes on both individuals of a pair. A total of 56 single adults and 28 pairs were studied. During the 2007 breeding season we followed Mougin & Mougin's (2000) method for tube placement on tail feathers in their study of Bulwer's Petrel *Buweria bulwerii*. In 2008 and 2009, we modified the method by placing the tubes on rump feathers. As European Storm Petrels are smaller than Bulwer's Petrel (mean body weight 26 g compared to 99 g), we shortened the tube from the 12 cm length used by Mougin & Mougin (2000) to 7 cm in order to avoid it being too obstructive. The capillary tubes allow the determination of diving depth by using the general gas compression formula (Mougin & Mougin 2000). To assess the efficacy of the capillary tubes, we attached a capillary tube to a buoyancy compensator on a SCUBA diver and controlled the maximum diving depth recorded by the capillary tube with a SCUBA computer. The tubes proved to be accurate and during three trial dives at 10 m recorded an error of  $\pm 10$  cm.

The sex of adult birds was determined by measuring the length of the white rump band of both individuals

in each pair; females have a larger rump band compared to males, and this is an accurate indicator of sex (Albores-Barajas *et al.* 2010).

Although we visited the colony every 3–5 days during daylight for adult and chick measurements during the three breeding periods, we only visited the colony at night once per breeding season in order to get regurgitated material from the adults as they returned from foraging trips. We used these samples to determine the food they provided to chicks. Birds were attracted to the rocks illuminated by our head lamps. They sat on the rocks disoriented and it was then possible to catch them. Procellariiformes often regurgitate the contents of the upper digestive tract (Barrett *et al.* 2007) in the presence of danger. Regurgitates can provide important information of the diet of the individual at sea and of the food they provide to chicks as they enter the breeding colony (Gladbach *et al.* 2009, Votier *et al.* 2003, Kitson *et al.* 2000). We observed in video recordings that the food is provided to chicks a few minutes after the adult's arrival at the nest (unpubl. data). In this study we have assumed that the regurgitates collected in this way represented the diet of the chick. As Storm Petrels regurgitated the material before we could catch them, material recovered from bare rocks near the nesting site was analysed (only regurgitates found on bare rock were kept in order to reduce the possibility of contamination of the samples). The regurgitated material was collected and preserved in 4% buffered formalin for later identification. Visits were kept to one per season, as during the late parts of the breeding period adults make one feeding trip every two or three days. Therefore making multiple visits increases the possibility of the same chick losing multiple meals. Given that the birds captured immediately regurgitated considerable quantities of food, we assumed that the adults were intending to provide the regurgitate to a hatchling and were not just returning to the nest to replace their partner.

## Diet analysis

We analysed 25 regurgitates. Each prey item was identified to the lowest taxonomic level (Tortonese 1975). The relative importance of each item was expressed as a percent of occurrence of food items in stomachs (OF%, occurrence frequency) (Hyslop 1980). Data obtained with this kind of analysis are qualitative, not quantitative.

## RESULTS

We recovered 30% of the capillary tubes placed (33/112 tubes) and recorded a mean diving depth of 146 cm (se =  $\pm 0.25$  cm). However, we observed a large variation in mean diving depth from year to year: 190 cm in 2007, 120 cm in 2008 and 150 cm in 2009 (Table 1), but these means were not significantly different (one-way ANOVA  $F = 0.50$ ,  $P = 0.61$ ). We did not observe a significant difference in diving depth according to sex, on 17 individuals sexed ( $t$ -test,  $t = -0.749$ ,  $P = 0.46$ ). During the 2007 breeding season we recovered only 20.83% of tubes deployed and we observed in IR video recordings that the tube placed on the tail could be obstructive for the birds and could be easily lost. Thus we decided to place the tubes on rump feathers in the 2008 and 2009 seasons. This resulted in higher recovering rates (28.57% in 2008 and 34.78% in 2009).

We observed that Storm Petrels also feed during the night immediately outside of the colony. The prey available in this area consists mainly of Opossum Shrimps Mysidacea. Adults may alternate two feeding techniques: traditional long trips, where they spend all day out at sea, and short trips, where they feed immediately outside the colony during the night.

In general, dietary composition in terms of the main prey items did not differ among samples of different years. The main prey species, both in volume and in number, were fish. Otolith analysis identified the ingested species as *Gymnammodites cicerellus*, a pelagic fish species common in deeper waters in this part of the Mediterranean Sea during the sampling season. On the basis of OL (otolith length) and the ratio between OL and TL (total length), reported for the species in Tuset *et al.* (2008), it is possible to suggest that all the specimen of *G. cicerellus* found in the regurgitates analysed were about 4 cm in length. *G. cicerellus* (OF% = 0.92; mean content per regurgitate = 1.6 individuals). In one regurgitate, the head of a shrimp was found, while in another a bulk of organic matter was present (probably

algae or jellyfish). These latter items were a minor component of the whole diet.

## DISCUSSION

The recovery of capillary tubes deployed in this way on birds is a difficult task. As foraging trip length is variable, it is sometimes difficult to find the individuals with the tube attached. We also visited the colony every 3–5 days to reduce disturbance. Therefore, after three or four visits the probability of finding an adult with a capillary tube was very low, as tubes may also be lost during the foraging trips. Despite these problems, we recovered a good number of capillary tubes (30%), and these provided valuable information about the feeding trips and diving behaviour of this population of European Storm Petrels, a species thought to feed mainly on the surface (D'Elbee & Hemery 1998). The shortening of tube length and placement on rump feathers in comparison to the techniques used by Mougin & Mougin (2000) allowed an improvement in recovering rates and possibly implied lower impact on birds' flight capability.

As in other Storm Petrel species (Bried 2005), we found that *H. pelagicus melitensis* dives for its prey. Although there are no significant differences in diving depth between the three years probably due to small sample size, there was a large variation in these depths. The differences may be the effect of different sea-surface temperatures, forcing prey to move up and down in the water column (Hutchinson 1967). Many taxa of zooplankton perform diel vertical migrations with amplitudes from a few to 100 metres (Hutchinson 1967). The 'normal' pattern is an evening ascent and a morning descent, although several cases of 'reversed' migrations have been described (Ohman *et al.* 1983, Bayly 1986). Migrating animals spend the day in deep water, but stay near the surface at night, followed by fish predators. The amplitude of the movements and the shape of the vertical distribution of the population may be very different between species and between ontogenetic stages of the same species and may be influenced by factors like turbidity and food abundance (Bohrer 1980, George 1983). The *Gymnammodites cicerellus* is a pelagic fish species usually found in schools and in the upper 10 m of the water column (Bauchot 1987). Differences in diving depth in Mediterranean subspecies of European Storm Petrel may be a temporal adaptation to better exploit the resources present at the time of breeding.

**Table 1.** Diving depths of European Storm Petrels. Minimum, maximum, mean and se are given. No significant differences between years were observed.

Year <sub>(sample size)</sub>	Minimum (cm)	Maximum (cm)	Mean (cm)	se ( $\pm$ cm)
2007 <sub>(5)</sub>	65	470	190	72
2008 <sub>(12)</sub>	26	513	120	38
2009 <sub>(16)</sub>	27	556	150	38

The fact that we observed adults feeding immediately outside the cave during the night leads us to think they feed on different prey during diurnal and nocturnal foraging trips. When they first arrive adults carry material collected out in the open sea during the day. On subsequent trips, during the night, adults carry the material caught outside of the breeding grounds, which consist mainly of Mysidacea. In fact, as observed in video recordings, adults may feed an entire fish to the chick in the first feed of the night, and then make shorter foraging trips providing smaller quantities of food (C. Soldatini, unpubl. data) during subsequent visits. Mysidacea were not found in the regurgitates collected at first arrival of adults to the colony, leading us to think that this is a secondary prey as its abundance is lower, compared to the abundance of krill in Atlantic waters (Thiriot-Quievreux *et al.* 1998).

The diet of European Storm Petrels has now been determined for both the Atlantic (D'Elbee & Hemery 1998) and the Mediterranean populations (this paper). What we did not study for a Mediterranean colony was whether prey items differed by sex as suggested by Elliott *et al.* (2010). We did, however, determine diving depth of both, males and females and found no difference. It may either be that both sexes feed on the same prey, or each sex uses a different foraging area, therefore feeding on different prey.

The information presented in this paper is a first step toward determining the diet of this Mediterranean subspecies of European Storm Petrels. This information and further dietary analysis may help in determining some conservation management options for this population.

## ACKNOWLEDGEMENTS

This project was funded by the Assessorato Regionale Agricoltura e Foreste to the Bird Ringing Unit of the University of Palermo.

## REFERENCES

- Albores-Barajas, Y.V., Massa, B., Griffiths, K. & Soldatini, C. 2010. Sexual dichromatism in Mediterranean storm petrels *Hydrobates pelagicus melitensis*. *Ardeola* **57**: 333–337.
- Barrett, R.T., Camphuysen, K., Anker-Nilssen, T., Chardine, J.W., Furness, R.W., Garthe, S., Huppopp, O., Leopold, M.F., Montevecchi, W.A. & Veit, R.R. 2007. Diet studies of seabirds: a review and recommendations. *Ices J. Marine Sci.* **64**: 1675–1691.
- Bauchot, M.L. 1987. [Poissons osseux. In: Fiches FAO d'identification pour les besoins de la pêche (rev. 1). In Fischer, W., Bauchot, M. L. & Schneider, M. (eds) *Méditerranée et mer Noire. Zone de pêche* 37W.] Rome, Commission des Communautés Européennes and FAO (in French).
- Bayly, I.A.E. 1986. Aspects of diel vertical migration in zooplacton, and its enigma variations. In De Deckker, P. & Williams, W.D. (eds) *Limnology in Australia*: 349–368. Commonwealth Scientific and Industrial Research Organisation, Melbourne.
- Bohrer, R. 1980. Experimental studies on diel vertical migration. In Kerfoot, W.C. (ed.) *Ecology and Evolution of Zooplankton Communities*: 111–121. University Press of New England, Hanover, New Hampshire.
- Bried, J. 2005. Diving ability of the Madeiran Storm Petrel. *Waterbirds* **28**: 162–166.
- Brooke, M.D.L. 2004. *Albatrosses and Petrels across the World*. Oxford University Press, Oxford.
- Cagnon, C., Lauga, B., Hemery, G. & Mouches, C. 2004. Phylogeographic differentiation of storm petrels (*Hydrobates pelagicus*) based on cytochrome *b* mitochondrial DNA variation. *Marine Biol.* **145**: 1257–1264.
- D'Elbee, J. & Hemery, G. 1998. Diet and foraging behaviour of the British Storm Petrel *Hydrobates pelagicus* in the Bay of Biscay during summer. *Ardea* **86**: 1–10.
- Del Hoyo, J., Elliot, A. & Sargatal, J. 1994. *Handbook of the Birds of the World, Vol. 2, New World Vultures to Guineafowl*. Lynx Edicions, Barcelona.
- Elliott, K.H., Gaston, A.J. & Crump, D. 2010. Sex-specific behavior by a monomorphic seabird represents risk partitioning. *Behav. Ecol.* **21**: 1024–1032.
- George, D.G. 1983. Interrelations between the vertical migration of *Daphnia* and chlorophyll A in two large limnetic enclosures. *J. Plankton Res.* **5**: 457–475.
- Gladbach, A., Braun, C., Nordt, A., Peter, H.U. & Quillfeldt, P. 2009. Chick provisioning and nest attendance of male and female Wilson's storm petrels *Oceanites oceanicus*. *Polar Biol.* **32**: 1315–1321.
- Hazlitt, S.L., Martin, T.G., Sampson, L. & Arcese, P. 2010. The effects of including marine ecological values in terrestrial reserve planning for a forest-nesting seabird. *Biol. Conserv.* **143**: 1299–1303.
- Hutchinson, G.E. 1967. *A Treatise on Limnology, Vol. II: Introduction to Lake Biology and the Limnoplankton*. Wiley, New York.
- Hyslop, E.J. 1980. Stomach contents analysis – a review of methods and their application. *J. Fish Biol.* **17**: 411–419.
- Kitson, J.C., Cruz, J.B., Lalas, C., Jillett, J.B., Newman, J. & Lyver, P.O. 2000. Interannual variations in the diet of breeding sooty shearwaters (*Puffinus griseus*). *NZ J. Zool.* **27**: 347–355.
- Lo Valvo, F. & Massa, B. 2000. Some aspects of the population structure of Storm Petrels *Hydrobates pelagicus* breeding on a Mediterranean island. *Ring. Migr.* **20**: 125–128.
- Magalhaes, M.C., Santos, R.S. & Hamer, K.C. 2008. Dual-foraging of Cory's shearwaters in the Azores: feeding locations, behaviour at sea and implications for food provisioning of chicks. *Marine Ecol. Prog. Series* **359**: 283–293.
- Massa, B. & Sultana, J. 1991. Status and conservation of the Storm Petrel *Hydrobates pelagicus* in the Mediterranean. *Il Merill* **27**: 5.
- Mougin, J.L. & Mougin, M.C. 2000. Maximum diving depths for feeding attained by Bulwer's petrels (*Bulweria bulwerii*) during the incubation period. *J. Zool.* **250**: 75–77.
- Ohman, M.D., Frost, B.W. & Cohen, E.H. 1983. Reverse diel vertical migration and escape from invertebrate predators. *Science* **220**: 1404–1407.
- Sanz-Aguilar, A., Massa, B., Lo Valvo, F., Oro, D., Minguéz, E. & Tavecchia, G. 2009. Contrasting age specific recruitment and survival at different spatial scales: a case study with the European storm petrel. *Ecography* **32**: 1–10.

**Thiriot-Quievreux, C., Leitao, A. & Cuzin-Roudy, J.** 1998. Chromosome diversity in Mediterranean and Antarctic euphausiid species (Euphausiacea). *J. Crustacean Biol.* **18**: 290–297.

**Tortonese, E.** 1975. [*Osteichthyes. Fauna d'Italia (Fauna of Italy).*] Edizioni Calderini, Bologna (in Italian).

**Tuset, V.M., Lombarte, A. & Assis, C.A.** 2008. Otolith atlas for the western Mediterranean, north and central eastern Atlantic. *Scientia Marina* **72**: 7–198.

**Voüier, S.C., Bearhop, S., Maccormick, A., Ratcliffe, N. & Furness, R.W.** 2003. Assessing the diet of great skuas, *Catharacta skua*, using five different techniques. *Polar Biol.* **26**: 20–26.

(MS received 19 October 2010; revised MS accepted 31 January 2011)