

Bond University  
Research Repository



## Prediction of Site Directed miRNAs as Key Players of Transcriptional Regulators Against Influenza C Virus Infection Through Computational Approaches

Hassan, Mubashir; Iqbal, Muhammad Shahzad; Naqvi, Sawaira; Alashwal, Hany; Moustafa, Ahmed A.; Kloczkowski, Andrzej

*Published in:*  
Frontiers in Molecular Biosciences

*DOI:*  
[10.3389/fmolb.2022.866072](https://doi.org/10.3389/fmolb.2022.866072)

*Licence:*  
CC BY

[Link to output in Bond University research repository.](#)

*Recommended citation(APA):*  
Hassan, M., Iqbal, M. S., Naqvi, S., Alashwal, H., Moustafa, A. A., & Kloczkowski, A. (2022). Prediction of Site Directed miRNAs as Key Players of Transcriptional Regulators Against Influenza C Virus Infection Through Computational Approaches. *Frontiers in Molecular Biosciences*, 9, [866072].  
<https://doi.org/10.3389/fmolb.2022.866072>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.



# Prediction of Site Directed miRNAs as Key Players of Transcriptional Regulators Against Influenza C Virus Infection Through Computational Approaches

Mubashir Hassan<sup>1,2\*</sup>, Muhammad Shahzad Iqbal<sup>3</sup>, Sawaira Naqvi<sup>1</sup>, Hany Alashwal<sup>4\*</sup>, Ahmed A. Moustafa<sup>5,6</sup> and Andrzej Kloczkowski<sup>2,7\*</sup>

## OPEN ACCESS

### Edited by:

Clement Adebajo Meseko,  
National Veterinary Research Institute  
(NVR), Nigeria

### Reviewed by:

Muhammad Aleem Ashraf,  
Khwaja Fareed University of  
Engineering and Information  
Technology (KFUEIT), Pakistan  
Lei Li,  
Cornell University, United States

### \*Correspondence:

Mubashir Hassan  
mubashirhassan\_gcui@yahoo.com  
Hany Alashwal  
halashwal@uaeu.ac.ae  
Andrzej Kloczkowski  
Andrzej.kloczkowski@  
nationwidechildrens.org

### Specialty section:

This article was submitted to  
Molecular Diagnostics and  
Therapeutics,  
a section of the journal  
Frontiers in Molecular Biosciences

Received: 30 January 2022

Accepted: 16 March 2022

Published: 08 April 2022

### Citation:

Hassan M, Iqbal MS, Naqvi S,  
Alashwal H, Moustafa AA and  
Kloczkowski A (2022) Prediction of Site  
Directed miRNAs as Key Players of  
Transcriptional Regulators Against  
Influenza C Virus Infection Through  
Computational Approaches.  
Front. Mol. Biosci. 9:866072.  
doi: 10.3389/fmolb.2022.866072

<sup>1</sup>Institute of Molecular Biology and Biotechnology, The University of Lahore, Lahore, Pakistan, <sup>2</sup>The Steve and Cindy Rasmussen Institute for Genomic Medicine, Nationwide Children Hospital, Columbus, OH, United States, <sup>3</sup>Department of Biotechnology, Faculty of Life Sciences, University of Central Punjab, Lahore, Pakistan, <sup>4</sup>College of Information Technology, United Arab Emirates University, Al-Ain, United Arab Emirates, <sup>5</sup>Department of Human Anatomy and Physiology, The Faculty of Health Sciences, University of Johannesburg, Johannesburg, South Africa, <sup>6</sup>School of Psychology, Faculty of Society and Design, Bond University, Gold Coast, QLD, Australia, <sup>7</sup>Department of Pediatrics, The Ohio State University, Columbus, OH, United States

MicroRNAs (miRNAs) are small non-coding RNAs that play critical roles in gene expression, cell differentiation, and immunity against viral infections. In this study, we have used the computational tools, RNA22, RNAhybrid, and miRanda, to predict the microRNA-mRNA binding sites to find the putative microRNAs playing role in the host response to influenza C virus infection. This computational research screened the following four miRNAs: hsa-mir-3155a, hsa-mir-6796-5p, hsa-mir-3194-3p and hsa-mir-4673, which were further investigated for binding site prediction to the influenza C genome. Moreover, multiple sites in protein-coding region (HEF, CM2, M1-M2, NP, NS1- NS2, NSF, P3, PB1 and PB2) were predicted by RNA22, RNAhybrid and miRanda. Furthermore, 3D structures of all miRNAs and HEF were predicted and checked for their binding potential through molecular docking analysis. The comparative results showed that among all proteins, HEF is higher in prevalence throughout the analysis as a potential (human-derived) microRNAs target. The target-site conservation results showed that core nucleotide sequence in three different strains is responsible for potential miRNA binding to different viral strains. Further steps to use these microRNAs may lead to new therapeutic insights on fighting influenza virus infection.

**Keywords:** miRNAs, mirbase, RStudio, RNAComposer, target site prediction, influenza C virus, HEF

## INTRODUCTION

Viral outbreaks are increasingly threatening public health (Asif et al., 2020). One of the most recurrent human affecting virus is group of Influenza viruses. These viruses are the members of the orthomyxoviridae family which include A, B, C, and D viruses (Christensen et al., 2008). Influenza C virus often causes upper respiratory infections such as common cold, sinusitis, pharyngitis, epiglottitis and laryngotracheitis (Dasaraju and Liu 1996). Influenza C virus has an average length of >300 nm (filamentous rods) and is composed of seven segments of the genomic single-stranded RNAs (ssRNAs) (Pellett et al., 2014). The genomic segments range in length from ~0.9 to ~2.4 kb, with a genome bridged

gene length of ~14 kb. The viral particle is 80–120 nm in diameter so that small virions adopt an elliptical shape. Particle lengths vary widely and can exceed tens of micrometers; producing filamentous virions. The total genome size is 12,907 bp (Wagaman et al., 1989). Influenza C virion envelopes contain a single glycoprotein called hemagglutinin-esterase-fusion (HEF) protein. HEF is a typical 1 transmembrane protein with a short N-terminal, a smart signal peptide (14 amino acids), a long ectodomain (612 amino acids), a transmembrane region (26 amino acids) and a very short cytoplasmic tail (three amino acid long). The HEF present in the infectious membranes is made up of two subunits, the N-terminal containing 432 amino acids forming the HEF1 polypeptide, and a residual sequence including hydrophobic fusion peptide, a transmembrane domain (TMD) and a cytoplasmic tail called HEF2 (Wang and Veit 2016). It encapsulates RNA with a negative strand and protects it from nucleases. The synthetic genomic RNA is called ribonucleoprotein (RNP) and serves as a model for transcription and replication. Shortly after a virion infects a new cell, the matrix 1 (M1) protein breaks down the RNP during the acidification of the virion driven by the matrix 2 (M2) protein. The separation of M1 from the RNP emits nuclear localization signals, directing the RNP to the nucleus (Muraki and Hongo 2010).

Most importantly, viruses can stimulate their life cycle by altering the cellular environment by actively controlling the expression of many microRNA (miRNA) cells (Perez et al., 2009). miRNAs are small endogenous ssRNA molecules, 21–23 nucleotides in length, which are formed after processing of hairpin loop-like miRNA precursors (pre-miRNA) by an RNase-III-like enzyme (Dicer) and can regulate the gene expression through working in cohort with mRNA and related enzyme structures (Brodersen and Voinnet 2006; Ali et al., 2020). RNA silencing, through miRNAs present in humans, thus impairs natural immunity and resistance to the host against foreign genetic elements including influenza C virus (Li et al., 2004). In connection with the influenza C virus genome, 264 mature human miRNAs have been found (MacFarlane and R Murphy 2010); a subset of these mature miRNAs should have targets in the influenza C virus genome and these miRNAs, once identified, can be expressed through cloning to enhance the resistance against infection from viruses (Peng et al., 2018).

The goal of this research study was to implement the computational approaches for the identification of targets for human miRNAs in the influenza C virus genome, as a precedent for enhancing the resistance to influenza through RNA interference (RNAi). For this objective, a set of miRNAs was retrieved from miRNA databases and was tested for binding to the influenza C virus genome through in-silico experiments involving three different miRNA target sites prediction algorithms.

## COMPUTATIONAL METHODOLOGY

### Retrieval of Genome Sequence and Annotations

The data was retrieved from the National Center for Biological Information (<https://www.ncbi.nlm.nih.gov/>) and the CLC

Genome Workbench (v 9.5.2) was used to download the complete annotated genome sequence of influenza C virus with the accession numbers ranging from NC\_006306.2 to NC\_006312.2. The CLC sequence viewer was used to create alignment between different strains of influenza C virus proteins.

### Retrieval of Mature miRNA From miRBase

The miRBase (<http://www.mirbase.org/>) is a database that collects information on miRNAs sequences and their annotations (Saini et al., 2008; Kozomara and Griffiths-Jones 2014) and 264 human miRNAs were retrieved from miRBase.

### miRNA Target Sites Predictions


Three tools, RNA22 (Rigoutsos et al., 2007), miRanda (Enright et al., 2003) and RNAhybrid (Krüger and Rehmsmeier 2006) were employed for prediction of possible targets for human derived miRNAs in influenza C virus genome to locate miRNA targeting regions.

### RNA22

RNA22 (Miranda et al., 2006; Loher and Rigoutsos 2012) uses an approach different than the other miRNA prediction tools; it implements a pattern-based approach combined with estimation of the folding energy to locate the possible miRNA target sites without a cross-species conservation filter. Identification of putative miRNA target sites is also possible even without knowing the identity of the targeting miRNA. The algorithm first analyzes the sequences of known mature miRNAs and then based on pattern information from the miRNAs, predicts the putative target sites, with many aligned patterns and then identifies the miRNAs which are likely to bind to the predicted target sites. RNA22 was accessed from the web ([cm.jefferson.edu/rna22/Interactive/](http://cm.jefferson.edu/rna22/Interactive/)) and the miRNAs and the target genome were input to the algorithm. Sensitivity and specificity values were kept at 63 and 61%, respectively and seed size of 7 was selected with one unpaired base allowed in the seed region without limiting the maximum number of G:U wobbles in the seed region. Minimum number of paired-up bases was kept at 12 while the maximum folding energy was kept at -14 kcal/mol.

### miRanda

The miRanda algorithm (John et al., 2004) has been used for miRNA target prediction (Zhao and Xue 2019). miRanda generally focuses on basic parameters such as sequence complementarity, free energy of RNA-RNA duplex and cross-species conservation of the target site to produce the output which is a weighted sum of match and mismatch scores for base pairs and gap penalties. Furthermore, it also promotes the prediction of multiple miRNA target sites including the ones with imperfect binding in the seed region within the 3'UTR of the target site thereby enhancing the specificity (Betel et al., 2008; M Witkos et al., 2011). The algorithm was run after defining the settings (gap open penalty = -9.0, gap extend penalty = -4.0, score threshold = 140, energy threshold = -20 kcal/mol and scaling parameter = 4.0).



## RNAhybrid

RNAhybrid is a tool for finding the minimum free energy for long and short miRNA. Hybridization is done in the form of domain mode, i.e. the short sequence is mixed within the most appropriate region of long sequences. miRNAs are short-acting RNAs that regulate gene targeting by binding to targeted mRNAs (Maziere and Enright 2007). Typically, RNAhybrid detects excellent hybridization sites of small RNA to large RNA. In RNAhybrid default parameters (minimum free energy value  $-30.0$ ) were adjusted to predicts the miRNA targeting sites in influenza C virus. Moreover, GUULGe tool was also employed to predict the miRNAs target sites (Gerlach et al., 2006).

## RStudio and Ggplot2

RStudio (<https://rstudio.com/>) is an integrated development environment (IDE) for R language usually employed for mathematical calculations and graphics. The ggplot2 package (<https://cran.r-project.org/web/packages/ggplot2/index.html>) was used to depict the graphical representations of miRNA predictions.

## miRNA and mRNA Model Predictions

MC-Fold (<https://major.irc.ca/MC-Fold/>) and MC-Sym (<https://major.irc.ca/MC-Sym/>) is an online pipeline to produce 3D models of miRNAs. The designed miRNA models were downloaded in PDB format and used for further structural

analysis. For mRNA model prediction, RNAComposer, an online platform was employed to predict 3D structures of mRNA molecules (Biesiada et al., 2016). The mRNA sequence retrieved from NCBI database, was entered into RNAComposer to predict 3D structure of mRNA of HEF using default parameters. Moreover, UCSF Chimera, tool was used to visualize the miRNA and mRNA models and to depict their graphical images (Meng et al., 2006).


## Molecular Docking of Mature miRNA and mRNA

The PatchDock algorithm is a molecular docking algorithm based on shape complementarity principles (<https://bioinfo3d.cs.tau.ac.il/PatchDock/>). This method simultaneously addresses the problem of flexibility and scoring of models produced by fast rigid-body docking algorithms. We obtained 100 potential docking candidates; PatchDock refines and scores them according to an energy function, spending about 3.5 s per candidate pose. The bonding interaction patterns between miRNA and mRNA for predictions from all target prediction servers were analyzed by using Discovery Studio (4.1) and UCSF Chimera 1.10.1 (Meng et al., 2006), respectively. The overview of our proposed methodology has been illustrated in **Figure 1**.


## RESULTS

Identifying the sequence of miRNAs is the first step in understanding the biology of miRNAs (Saif et al., 2019). Research and testing methods revealed many aspects of miRNA-target interactions and led to effective predictions of targeted mRNAs. Information collected on the recognition of the miRNA targets revealed the principles of miRNA-mRNA duplex formation, such as the binding of hsa-mir-3155a, hsa-mir-6796-5p, hsa-mir-3194-3p and hsa-mir-4673 to the influenza virus genome. Target selection of proteins based on predicted miRNAs would be important evidence of better understanding of the functionality of miRNAs and the prevalence of influenza C virus. All the miRNA-targets in influenza virus c genome are listed in supplementary data (*see meta data*). The genomic structure of influence c virus genome has been represented in **Figure 2**.


miRNAs are small double-stranded RNAs that exert a fine-tuning sequence-specific regulation in the cellular transcriptome (Zeng et al., 2003). miRNA regulates multiple mRNAs through binding with complementary sequences at 3'-untranslated regions for triggering the mechanism of RNA interference (Titze-de-Almeida et al., 2020). In our computational studies, target sites for 264 human-derived miRNAs from miRBase have been predicted and analyzed (hairpin and genomic location), based on microRNA-mRNA complementarity. The genomic position of four predicted miRNAs is hsa-mir-3155a (Chr10: 6152196-6152277); hsa-mir-6796-5p (Chr19: 40369846-40369907) hsa-mir-3194-3p (Chr20: 51452905-51452977) and hsa-mir-4673 (Chr9: 136519568-136519626), respectively.



**FIGURE 2 |** Influenza C virus genome. Prediction of new target-sites for miRNAs in influenza C virus.



**FIGURE 3 |** Target sites for miRNAs binding to influenza C virus genome predicted by RNA22 algorithm. The color dots represent binding to genes coding the relevant proteins. The gray dots labeled as NA which indicate non-applicable.



**FIGURE 4 |** Target sites of miRNAs in genome of influenza C virus predicted by miRanda algorithm. The color key represents the respective genes/proteins in the graph. The gray dots labeled as NA which indicate non-applicable.

## RNA22 Analysis

RNA22, a gene predictor uses a pattern-based approach and folding energy to find locations that can be targeted by miRNA without applying various types of conservation filters (Mazière and Enright 2007). **Figure 3** shows that most frequently targeted sites where those with the folding energy value ranging from  $-20$  to  $-22$  kcal/mol and nucleotide position ranges from 0 to 2000 bp. Moreover, best predicted target sites belong to genes encoding proteins: HEF, M2, NP, NS1- NS2, P3, PB1, and PB2, respectively. The corresponding alignments have been shown in supplementary data (**Supplementary Figure S1**).


## miRanda

Another tool miRanda was employed and analyzed results based on miRNA-mRNA sequence similarity, free energy (ranging from

$-18$  to  $-33$  kcal/mol), and scoring values (from 140 to 185). **Figure 4** shows that most frequently targeted sites are those where the free energy value ranges from  $-20$  to  $-30$  kcal/mol and nucleotide position range is 0–2000 bp. Moreover, significantly predicted target sites belong to genes encoding proteins CM2, HEF, M1-M2, NP, NS1- NS2, NSF, P3, PB1- PB2 and NA, respectively. The alignment figure has been reported in supplementary data (**Supplementary Figure S1**).

## RNA Hybrid

The third approach which has been employed is RNAhybrid which identifies the regions in 3' UTRs which have the potential to form a thermodynamically favorable duplex with a specific miRNA. RNAhybrid uses minimum free energy (MFE) to



evaluate the predicted binding sites for miRNAs. **Figure 5** shows that most frequently targeted sites are those where the MFE ranges from  $-30$  to  $-38$  kcal/mol and nucleotide position range is 0–2000 bp. Moreover, best predicted target sites belong to genes encoding proteins CM2, HEF, M1–M2, NP, NS1–NS2, NSF, P3, PB1–PB2 and NA, respectively. The corresponding alignment (**Supplementary Figure S1**) has been reported in supplementary data.

RNA22, miRanda and RNAhybrid predicted 242, 231 and 41 miRNAs which may have possible binding sites against the influenza C virus genome and four most common hsa-mir-3155a, has-mir-6796-5p, has-mi3194-3p and has-mir-4673, respectively (**Figure 6**). Moreover, in union graph analysis 47 miRNAs were common in RNA22–miRanda predictions, 20 and 24 miRNAs were common in RNA22–RNAhybrid and RNAhybrid–miRanda predictions, respectively (**Figure 6**). The common features of all three algorithms have been reported in **Table 1** (Peterson et al., 2014). Furthermore, GUUGLe predicted results showed less consensus to our predicted results and therefore, generated result has been tabulated in supplementary data (Meta Data, sheet 3).

### Prediction of Structures of hsa-mir-4673 and hsa-mir-3194-3p and Its Cellular Functionality

The hsa-mir-4673 and hsa-mir-3194-3p play an important role in our studies. The miRNA acts as a guide for basic pairing with the target mRNA to negatively control its expression. The paired miRNAs are aimed at mRNA targeting gene mutation by mRNA cleavage or translation compression based on the degree of compatibility between miRNA and mRNA target. **Figure 7** shows the predicted 3D structures of premature and mature hsa-mir-4673 and hsa-mir-3194-3p.

### Prediction of Structures of hsa\_mir\_6796-5p and hsa-mir-3155a and Their Cellular Functionality


The hsa\_mir\_6796\_5p and hsa\_mir\_3155a are playing an important role in our studies. miRNAs are short non-coding RNAs that regulate gene expression after transcription. They usually bind 3'-UTR (uninterrupted region) of their targeted mRNAs and suppress protein production by damaging mRNA and silencing translation. **Figure 8** shows the predicted 3D structures of premature and mature hsa-mir-6796-5p and hsa-mir-3155a.

### Target-Site Conservation Analysis

Conserved miRNAs target sites tend to be enriched at the 5' end of protein-coding gene 3' UTRs and provide evidence that miRNA targets sites are enriched in genes involved in protein synthesis pathways (Xiong et al., 2019). Our results showed the conserved nucleotides in different influenza virus strains. This conservation pattern enhanced the accuracy and reliability of predicted miRNAs across the other strains (see **Figure 9** and **Supplementary Figures S3, S4**).


### Hemagglutinin-Esterase-Fusion

HEF is composed of 655 amino acids (<https://www.uniprot.org/uniprot/P07975>). The HEF is divided into couple of subunits HEF1 and HEF2 through proteolytic cleavage (Wang and Veit 2016). HEF is a glycoprotein that is involved in receptor binding, receptor destroying and membrane fusion activities. HEF, recognizes and attaches to a receptor through determinant N-acetyl-9-O-acetylneuraminic acid on the cell surface and initiates virus entry (Rogers et al., 1986). Moreover, HEF also



**TABLE 1** | Common features of miRanda, RNA22 and RNAhybrid algorithms.

Algorithms	Seed match	Conservation	Free energy	Target-site abundance
miRanda	Yes	Yes	Yes	—
RNA22	Yes	—	Yes	—
RNAhybrid	Yes	—	Yes	Yes



catalyzes the fusion of the viral envelope with endocytic vesicles. Finally, HEF is the receptor-destroying enzyme, which is the function of neuraminidase (NA) in the influenza A and B viruses. HEF does not cleave the terminal sialic acid residue from carbohydrates but has an esterase activity that removes the acetyl group from position C-9 of N-acetyl-9-O-acetylneuraminic acid (Herrler et al., 1985). This function is probably required to release freshly budded virus particles from infected cells, which would otherwise be trapped at their plasma membrane if the receptor would still be present (Wang and Veit 2016).

### Nucleoprotein (NP)

The NP containing 498 amino acids is a highly conserved multi-functional viral protein, best known as a hot antiviral drug target. It has been observed that upon attachment to the host cell through the interaction between HA from the virus and the sialic acid receptor on the host cell surface, the influenza virus enters host cells by endocytosis (Hu et al., 2017). The X-ray


crystal structures of the influenza nucleoprotein are known, the pockets in which these mutants (Y289H, N309K/N309T, and Y52H) reside might be the corresponding drug binding sites of nucleozin (Kao et al., 2010).

### Solute Carrier Family 10 Member 3

Solute Carrier Family 10 Member 3, also known as P3 protein contains 716 amino acids. It has been observed that active P3 (coded by RNA 1) and P1 protein (coded by RNA 2) are required for complementary RNA synthesis of influenza virus (Yamashita et al., 1989). Therefore, this protein could be used as a novel target to design chemical inhibitors to stop the virulence of influenza C virus.

### Polymerase Basic Protein 1 and 2

The PB1 protein consists of 757 amino acids, occurs in influenza viruses A, B and C and is involved in the RNA elongation and endonuclease activity (Bouvier and Palese 2008). The PB2 gene contains 2341 nucleotides and codes 759 amino acids-



long PB2 protein that is responsible for mRNA cap recognition (Bouvier and Palese 2008). All influenza viruses encode the polymerase subunit PB1 on segment 2; in some strains of influenza A virus, this segment also codes the accessory protein PB1-F2, a small, 87-amino acid protein with pro-apoptotic activity, in a +1 alternate reading frame (Chen et al., 2001). The predicted frequency of occurrence of all proteins has been shown in **Figure 10**.

### Molecular Docking Analysis

The predicted 3D structures of hsa-miRNA-3155a and mRNA of HEF were docked to analyze the binding potential and nucleotide region involved in the binding conformation. The docking results showed that a good scoring value has been observed with better interactive behavior. The best docking score was 19,704, which demonstrates very good geometric shape complementarity. Moreover, the approximate interface area and atomic contact energy of docked complex were 3196 and  $-2078.92$ , respectively (**Figure 11**). The detailed docking scoring values have been tabulated in supplementary data (**Supplementary Table S1**).

The interactive behavior of hsa-miRNA-3155a and mRNA of HEF was clearly observed and generated results show that multiple nucleotides of miRNA-3155a form close contacts with mRNA of HEF. The nucleotide adenine, guanine and guanine (AGG) at positions 3, 4 and 5 in miRNA-3155a formed close interaction with mRNA of HEF. Similarly, cytosine and uracil (CU) at positions 20 and 21 also showed close interactions with mRNA of HEF. Similarly, different regions in mRNA-HEF such

as AUUGC, UUU and AGCU have been predicted as targets for potential binding with hsa-miRNA-3155a (**Figure 12**). The docking results showed that binding of hsa-miRNA-3155a with mRNA-HEF may stop the translation process and can protect from influenza C virus infection.


### Mechanistic Pathways of miRNAs

miRNAs are small, highly conserved non-coding RNA molecules (containing about 22 nucleotides) involved in the regulation of differential gene expression. miRNAs are transcribed by RNA polymerases II and III, generating precursors that undergo a series of cleavage events to form mature miRNA (MacFarlane and R Murphy 2010). The pri-miRNA (primary transcript) is converted into pre-miRNA (precursor-miRNA, a stem-loop structure of about 70 base pairs long) using DROSHA, a nuclear RNase III enzyme that executes the initiation step of miRNA processing in the nucleus (Lee et al., 2003) (Nguyen et al., 2018). The overall mechanistic pathway of mature miRNA formation has been depicted in **Figure 13**.


### miRNAs and Immune Response

miRNAs regulate the immune responses (both acquired and innate immunity) against flu infection by activating the different protein signaling cascades in humans (Zhang and Li 2013). During infection, a variety of intracellular signaling pathways are being activated at different receptor sites such as TNF receptor-related substances (TRAFs), and interferon-regulatory features (IRFs) through different proteins HEF, P3, NP, PB1 and 2, CM1 and





**FIGURE 9 |** Target-site conservation pattern in different influenza strains. The target sites are highlighted in blue, and bars show the level of conservation at specific sites. Involvement of predicted proteins in influenza c virus.



**FIGURE 10 |** Bar chart of predicted proteins.



2, NS1 and 2, respectively. The HEF is involved in the influenza (Wang and Veit 2016), therefore miRNA-3155 binding to HEF may inhibit the downstream signaling pathway (see Figure 14). Moreover, PB1 and PB2 are also key players in the activated signaling pathways after the incorporation of pathogen within the host cells (Ludwig 2011). Our computational results show that miRNA-5088 has the potential to bind to complementary sites of mRNA of PB2 and stop the further transcriptional process. As a result, miRNA-5088 may play a supportive role in the prevention of influenza. Other miRNAs such as miRNA-7159, miRNA-4747

and miRNA-10394 bind to NP, P3 and PB1, respectively and may stop their signaling cascades and therefore could be used as good protective agents against influenza infection (Gaur et al., 2011). The overall mechanistic pathway of miRNA involvement in the immune response is shown in Figure 14.

## DISCUSSION

In the current study, we have designed an effective computational approach for the validation of miRNA target prediction results by utilizing RNA22, miRanda and RNAhybrid different algorithms with default parameters. These algorithms proposed an effective method to recognize the possible host-derived miRNA targets in virus genomes (Ashraf et al., 2020; Ashraf et al., 2022). The RNA22 is an algorithm for exploring new miRNA-mRNA interactions because of its unique capabilities-although it has a high likelihood of generating false-positive results (Riffo-Campos A.L et al., 2016). miRanda is mostly extensively used algorithm that includes the main aspects of miRNA-target prediction, such as the conservation level and miRNA 3'UTR site (Betel D et al., 2010). RNAhybrid is also algorithm to identify the minimum free energy for long and short miRNAs (Maziere and Enright 2007).

In our computational results, target sites for 264 human-derived miRNAs from miRbase have been predicted and analyzed (hairpin and genomic location), based on miRNA-mRNA complementarity. The comparative results showed that a total of 242, 231 and 41



miRNAs have been predicted to bind with influenza C virus genome by RNA22, miRanda and RNAhybrid, respectively. In union graph analysis among all three utilized algorithms, we have observed 47 common miRNAs inbetween RNA22-miRanda predictions, whereas 20 and 24 miRNAs common in RNA22-RNAhybrid and RNAhybrid-miRanda predictions, respectively. In intersection graph analysis, we have found that four miRNAs: hsa-mir-3155a, hsamir-6796-5p, hsami3194-3p and hsamir-4673 which were common in three algorithms methods which suggest their good binding potential with targeted sites in influenza C virus genome (**Figure 6; Table 2**).

The minimum free energy (MEF) structure of a sequence is the secondary structure that is calculated to have the lowest value of free energy. Free energy calculation is a dynamic feature of miRNA and

target binding. Prior research data showed a significant correlation of free energy between the translational repression and the hybridization binding of the seed region (Doench J.G. and Sharp PA, 2004). The thermodynamic stability of the miRNA-mRNA duplex was estimated by the assessment of free energy to monitor site accessibility for the determination of the secondary structure duplex (Peterson et al., 2014). From RNA22 results, the predicted results were analyzed based on the sensitivity (25%) and specificity (90%) and folding energy pattern. It has been observed that highest negative value of MEF is showed the most structured and stable miRNA-mRNA complexes. The most frequently targeted sites where those with the folding energy value ranging from  $-20$  to  $-22$  Kcal/mol and nucleotide position ranges from 0 to 2000 bp. Moreover, from miRanda results, it has also been observed a good




FIGURE 13 | Mechanistic pathway of miRNA formation from premature miRNA.




FIGURE 14 | Mechanistic pathway of miRNAs involvement in the immune response.

**TABLE 2** | Premature and mature miRNAs.

miRNAs	Precursor miRNAs	Mature miRNAs
hsa-mir-3155a	uccgggcaucaccuccacugcagagccuggggagccggacaaggucogugacaggugacgucucggaccuccuccucg	Ccaggcucugcagugggaacu
hsa-mir-6796-5p	uuaccuuguggguuggagagcugggagcgcgccuccuccucgaagacuccccgacc	Uugugguuggagagcuggc
hsa-mir-3194-3p	Agguggcagggccagccaccaggaggcugcgucuccacugagcggucacucggucucucgacggcc	Agcucugcugcucacuggca
hsa-mir-4673	guccaggcaggagccggacuggaccucgacggcgucucucccgccagucggagaag	Uccaggcaggagccggacugga

binding site in with good sequence similarity, free energy, and scoring values (from 140 to 185), respectively. The best targeted sites have been observed where the free energy value ranges from  $-20$  to  $-30$  Kcal/mol with nucleotides range from 0–2000 bp. RNAhybrid identifies the regions in 30 UTRs which have the potential to form a thermodynamically favorable duplex with a specific miRNA. From RNAhybrid result, it has been observed that MFE ranges from  $-30$  to  $-38$  kcal/mol and nucleotide position range is 0–2000 bp. Moreover, in protein comparison results, we have identified multiple proteins: CM2, HEF, MI, M2, NP, NS1, NS2, P3, PB1 and PB2 that may have a significant role in cell signaling and functionality of genomic virus (Furukawa et al., 2011). It has been observed that CM2 is directly involved in the influenza C virus replication cycle (Furukawa et al., 2011). HEF showed the highest value ( $<50$ ) as compared to rest of proteins whereas, M2 depicted the lowest value ( $<10$ ). Moreover, NP, P3, PB2 and PB1 showed 41, 38, 46 and 37, respectively (Figures 7, 8). The genomic region has been mentioned in **Supplementary Figure S2**. The conserved nucleotides of showed the common predicting binding site for miRNAs among different strains of influenza virus (see **Figure 9** and **Supplementary Figure S3**).

Molecular docking is a computational approach usually employed to check the binding potential of biomolecules (Hassan et al., 2022, 2021). The modelled 3D structures of hsa-miRNA-3155a and mRNA were docked to analyze the binding potential and nucleotide region involved in the binding conformations. The comparison results from all docking scores, it has been seen that 19704 was best docking score, which demonstrates very good geometric shape complementarity of miRNA against mRNA (see **Figure 11**). Furthermore, a detail conformation analysis was done to explore the nucleotides of miRNA having good binding interactions with mRNA. Our predicted results showed that adenine, guanine, and guanine (AGG) at positions 3, 4 and 5 in miRNA-3155a formed good interaction with mRNA of HEF. Similarly, cytosine and uracil (CU) at positions 20 and 21 formed hydrogen bonds with mRNA of HEF (**Figure 12**). The docking results showed that binding of hsa-miRNA-3155a with mRNA-HEF may stop the translation process and can protect from influenza virus infection.

## CONCLUSION

This study presents a planned *in silico* approach for finding host-derived miRNAs aimed at silencing the genome of influenza virus. Predicted results showed that miRNAs: hsa-mir-3155a, hsa-mir-6796-5p, hsa-mir-3194-3p and hsa-mir-4673 demonstrated good binding complementarity to influenza

genome. Based on interactive behavior multiple proteins were identified as are active targets for influenza mediating signaling cascade in humans. Among all, HEF was repeatedly predicted in our computational studies. The docking results showed the binding pattern of hsa-mir-3155a to mRNA of HEF protein which also depicts the interactive nucleotides which may involves in the transcriptional activity HEF gene. However, molecular biology techniques are required to further verify our computational results. The target-site conservation pattern in different strains ensures the significance of hsa-mir-3155a. Taken together, exploration of these small endogenous nucleotides (miRNAs) may lead to new therapeutics against influenza virus infection.

## DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**, further inquiries can be directed to the corresponding authors.

## AUTHOR CONTRIBUTIONS

MH and MSI were performed experiments and wrote the initial draft of the manuscript. SN collected data. HA, AM, and AK edited the manuscript and compiled it into final format.

## FUNDING

This work received financial support from the United Arab Emirates University (Grant No. CIT 31T129). AK acknowledges financial support from NSF grant DBI1661391, and NIH Grants R01 GM127701, and R01HG012117.

## ACKNOWLEDGMENTS

MH acknowledges the Ohio State University for “President’s Postdoctoral Scholars Program (PPSP)” fellowship for financial support.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmolb.2022.866072/full#supplementary-material>

## REFERENCES

- Ali, Q., Jamal, A., Ullah, S., and Waqar, A. B. (2020). Male Predominant Association with Apolipoprotein B mRNA-Editing Enzyme, Catalytic Polypeptide-like 3G Variants (Rs6001417, Rs35228531, Rs8177832) Predict protection against HIV-1 Infection. *Advancements Life Sci.* 7 (2), 91–97.
- Ashraf, F., Ashraf, M. A., Hu, X., and Zhang, S. (2020). A Novel Computational Approach to the Silencing of Sugarcane Bacilliform Guadeloupe A Virus Determines Potential Host-Derived MicroRNAs in Sugarcane (*Saccharum Officinarum* L.). *PeerJ* 8, e8359. doi:10.7717/peerj.8359
- Ashraf, M. A., Feng, X., Hu, X., Ashraf, F., Shen, L., Iqbal, M. S., et al. (2022). In Silico identification of Sugarcane (*Saccharum Officinarum* L.) Genome Encoded microRNAs Targeting Sugarcane Bacilliform Virus. *PLoS ONE* 17 (1), e0261807. doi:10.1371/journal.pone.0261807
- Asif, M., Nawaz, S., Bhutta, Z. A., Kulyar, M. F.-e.-A., Rashid, M., Shabir, S., et al. (2020). Viral Outbreaks: A Real Threat to the World. *Advancements Life Sci.* 8 (1), 08–19.
- Betel, D., Wilson, M., Gabow, A., Marks, D. S., and Sander, C. (2008). The microRNA.Org Resource: Targets and Expression. *Nucleic Acids Res.* 36 (Suppl. 1\_1), D149–D153. doi:10.1093/nar/gkm995
- Betel, D., Koppal, A., Agius, P., Sander, C., and Leslie, C. (2010). Comprehensive Modeling of microRNA Targets Predicts Functional Non-conserved and Non-canonical Sites. *Genome Biol.* 11 (8), R90. pmid:20799968. doi:10.1186/gb-2010-11-8-r90
- Biesiada, M., Purzycka, K. J., Szachniuk, M., Blazewicz, J., and Adamiak, R. W. (2016). “Automated RNA 3D Structure Prediction with RNAComposer,” in *Automated RNA 3D Structure Prediction with RNAComposer. RNA Structure Determination* (New York: Springer), 199–215. doi:10.1007/978-1-4939-6433-8\_13
- Bouvier, N. M., and Palese, P. (2008). The Biology of Influenza Viruses. *Vaccine* 26, D49–D53. doi:10.1016/j.vaccine.2008.07.039
- Brodersen, P., and Voinnet, O. (2006). The Diversity of RNA Silencing Pathways in Plants. *Trends Genet.* 22 (5), 268–280. doi:10.1016/j.tig.2006.03.003
- Chen, W., Calvo, P. A., Malide, D., Gibbs, J., Schubert, U., Bacik, I., et al. (2001). A Novel Influenza A Virus Mitochondrial Protein that Induces Cell Death. *Nat. Med.* 7 (12), 1306–1312. doi:10.1038/nml201-1306
- Christensen, J. P., Bojesen, A. M., and Bisgaard, M. (2008). Fowl Cholera. *Poultry Dis.* 6, 149–154.
- Dasaraju, P. V., and Liu, C. (1996). “Chapter 93: Infections of the Respiratory System,” in *Medical Microbiology*. 4th ed. (Galveston: University of Texas Medical Branch at Galveston).
- Doench, J. G., and Sharp, P. A. (2004). Specificity of microRNA Target Selection in Translational Repression. *Genes Dev.* 18 (5), 504–511. doi:10.1101/gad.1184404
- Enright, A. J., John, B., Gaul, U., Tuschl, T., Sander, C., and Marks, D. S. (2003). MicroRNA Targets in *Drosophila*. *Genome Biol.* 5 (11), R1–R27. doi:10.1186/gb-2003-5-1-r1
- Furukawa, T., Muraki, Y., Noda, T., Takashita, E., Sho, R., Sugawara, K., et al. (2011). Role of the CM2 Protein in the Influenza C Virus Replication Cycle. *J. Virol.* 85 (3), 1322–1329. doi:10.1128/jvi.01367-10
- Gaur, P., Munjhal, A., and Lal, S. K. (2011). Influenza Virus and Cell Signaling Pathways. *Med. Sci. Monit.* 17 (6), RA148–RA154. doi:10.12659/msm.881801
- Gerlach, W., and Giegerich, R. (2006). GUUGle: a Utility for Fast Exact Matching under RNA Complementary Rules Including G-U Base Pairing. *Bioinformatics* 22 (6), 762–764. doi:10.1093/bioinformatics/btk041
- Hassan, M., Zahid, S., Alashwal, H., Kloczkowski, A., and Moustafa, A. A. (2021). Mechanistic Insights into TNFR1/MADD Death Domains in Alzheimer’s Disease through Conformational Molecular Dynamic Analysis. *Sci. Rep.* 11, 12256. doi:10.1038/s41598-021-91606-4
- Hassan, M., Zahid, S., Shahzadi, S., Malik, E., Zaib, S., Iqbal, J., et al. (2022). Mechanistic Insight of DACH1 Receptor in the Development of Carcinoma Insurgence through MD Simulation Studies. *J. Biomol. Struct. Dyn.* 40 (2), 742–751. doi:10.1080/07391102.2020.1818624
- Herrler, G., Rott, R., Klenk, H. D., Müller, H. P., Shukla, A. K., and Schauer, R. (1985). The Receptor-Destroying Enzyme of Influenza C Virus Is Neuraminidase-O-Acetyltransferase. *EMBO J.* 4 (6), 1503–1506. doi:10.1002/j.1460-2075.1985.tb03809.x
- Hu, Y., Sneyd, H., Dekant, R., and Wang, J. (2017). Influenza A Virus Nucleoprotein: a Highly Conserved Multi-Functional Viral Protein as a Hot Antiviral Drug Target. *Curr. Top. Med. Chem.* 17 (20), 2271–2285. doi:10.2174/1568026617666170224122508
- John, B., Enright, A. J., Aravin, A., Tuschl, T., Sander, C., and Marks, D. S. (2004). Human microRNA Targets. *Plos Biol.* 2 (11), e363. doi:10.1371/journal.pbio.0020363
- Kao, R. Y., Yang, D., Lau, L.-S., Tsui, W. H. W., Hu, L., Dai, J., et al. (2010). Identification of Influenza A Nucleoprotein as an Antiviral Target. *Nat. Biotechnol.* 28 (6), 600–605. doi:10.1038/nbt.1638
- Kozomara, A., and Griffiths-Jones, S. (2014). miRBase: Annotating High Confidence microRNAs Using Deep Sequencing Data. *Nucl. Acids Res.* 42 (D1), D68–D73. doi:10.1093/nar/gkt1181
- Krüger, J., and Rehmsmeier, M. (2006). RNAhybrid: microRNA Target Prediction Easy, Fast and Flexible. *Nucleic Acids Res.* 34 (Suppl. 1\_2), W451–W454.
- Lee, Y., Ahn, C., Han, J., Choi, H., Kim, J., Yim, J., et al. (2003). The Nuclear RNase III Drosha Initiates microRNA Processing. *Nature* 425 (6956), 415–419. doi:10.1038/nature01957
- Li, W.-X., Li, H., Lu, R., Li, F., Dus, M., Atkinson, P., et al. (2004). Interferon Antagonist Proteins of Influenza and Vaccinia Viruses Are Suppressors of RNA Silencing. *Proc. Natl. Acad. Sci. U.S.A.* 101 (5), 1350–1355. doi:10.1073/pnas.0308308100
- Loher, P., and Rigoutsos, I. (2012). Interactive Exploration of RNA22 microRNA Target Predictions. *Bioinformatics* 28 (24), 3322–3323. doi:10.1093/bioinformatics/bts615
- Ludwig, S. (2011). Disruption of Virus-Host Cell Interactions and Cell Signaling Pathways as an Anti-viral Approach against Influenza Virus Infections. *Biol. Chem.* 392 (10), 837–847. doi:10.1515/bc.2011.121
- MacFarlane, L.-A., and Murphy, P. R. (2010). MicroRNA: Biogenesis, Function and Role in Cancer. *Cg* 11 (7), 537–561. doi:10.2174/138920210793175895
- Mazière, P., and Enright, A. J. (2007). Prediction of microRNA Targets. *Drug Discov. Today* 12 (11–12), 452–458. doi:10.1016/j.drudis.2007.04.002
- Mazière, P., and Enright, A. J. (2007). Prediction of microRNA Targets. *Drug Discov. Today* 12 (11), 452–458. doi:10.1016/j.drudis.2007.04.002
- Meng, E. C., Pettersen, E. F., Couch, G. S., Huang, C. C., and Ferrin, T. E. (2006). Tools for Integrated Sequence-Structure Analysis with UCSF Chimera. *BMC Bioinformatics* 7 (1), 339–410. doi:10.1186/1471-2105-7-339
- Miranda, K. C., Huynh, T., Tay, Y., Ang, Y.-S., Tam, W.-L., Thomson, A. M., et al. (2006). A Pattern-Based Method for the Identification of MicroRNA Binding Sites and Their Corresponding Heteroduplexes. *Cell* 126 (6), 1203–1217. doi:10.1016/j.cell.2006.07.031
- Muraki, Y., and Hongo, S. (2010). The Molecular Virology and Reverse Genetics of Influenza C Virus. *Jpn. J. Infect. Dis.* 63 (3), 157–165.
- M. Witkos, T., Koscianska, E., and J. Krzyzosiak, W. (2011). Practical Aspects of microRNA Target Prediction. *Cmm* 11 (2), 93–109. doi:10.2174/156652411794859250
- Nguyen, T. H., Liu, X., Su, Z. Z., Hsu, A. C.-Y., Foster, P. S., and Yang, M. (2018). Potential Role of MicroRNAs in the Regulation of Antiviral Responses to Influenza Infection. *Front. Immunol.* 9, 1541. doi:10.3389/fimmu.2018.01541
- Pellet, P. E., Mitra, S., and Holland, T. C. (2014). Basics of Virology. *Handbook Clin. Neurol.* 123, 45–66. doi:10.1016/b978-0-444-53488-0.00002-x
- Peng, S., Wang, J., Wei, S., Li, C., Zhou, K., Hu, J., et al. (2018). Endogenous Cellular microRNAs Mediate Antiviral Defense against Influenza A Virus. *Mol. Ther. - Nucleic Acids* 10, 361–375. doi:10.1016/j.omtn.2017.12.016
- Perez, J. T., Pham, A. M., Lorini, M. H., Chua, M. A., Steel, J., and tenOever, B. R. (2009). MicroRNA-mediated Species-specific Attenuation of Influenza A Virus. *Nat. Biotechnol.* 27 (6), 572–576. doi:10.1038/nbt.1542
- Peterson, S. M., Thompson, J. A., Ufkin, M. L., Sathyanarayana, P., Liaw, L., and Congdon, C. B. (2014). Common Features of microRNA Target Prediction Tools. *Front. Genet.* 5, 23. pmid:24600468. doi:10.3389/fgene.2014.00023
- Riffo-Campos, Á., Riquelme, I., and Brebi-Mieville, P. (2016). Tools for Sequence-Based miRNA Target Prediction: what to Choose? *Ijms* 17 (12), 1987. pmid:27941681. doi:10.3390/ijms17121987
- Rigoutsos, I., Miranda, K., and Huynh, T. (2007). *rna22: A Unified Computational Framework for Discovering miRNA Precursors, Localizing Mature miRNAs, Identifying 3’UTR Target-Islands, and Determining the Targets of mature-miRNAs*. Armonk New York: Ibm Corporation.

- Rogers, G. N., Herrler, G., Paulson, J. C., and Klenk, H. D. (1986). Influenza C Virus Uses 9-O-Acetyl-N-Acetylneuraminic Acid as a High Affinity Receptor Determinant for Attachment to Cells. *J. Biol. Chem.* 261, 5947–5951.
- Saif, R., Qazi, K., Zia, S., Mahmood, T., Ejaz, A., Tamseel, T., et al. (2019). First Step with R for Life Sciences: Learning Basics of This Tool for NGS Data Analysis. *Advancements Life Sci.* 6 (4), 187–194.
- Saini, H. K., Enright, A. J., and Griffiths-Jones, S. (2008). Annotation of Mammalian Primary microRNAs. *BMC genomics* 9 (1), 564–619. doi:10.1186/1471-2164-9-564
- Titze-de-Almeida, S. S., Soto-Sánchez, C., Fernandez, E., Koprach, J. B., Brotchie, J. M., and Titze-de-Almeida, R. (2020). The Promise and Challenges of Developing miRNA-Based Therapeutics for Parkinson's Disease. *Cells* 9 (4), 841. doi:10.3390/cells9040841
- Wagaman, P. C., Spence, H. A., and O'Callaghan, R. J. (1989). Detection of Influenza C Virus by Using an In Situ Esterase Assay. *J. Clin. Microbiol.* 27 (5), 832–836. doi:10.1128/jcm.27.5.832-836.1989
- Wang, M., and Veit, M. (2016). Hemagglutinin-esterase-fusion (HEF) Protein of Influenza C Virus. *Protein Cell* 7 (1), 28–45. doi:10.1007/s13238-015-0193-x
- Xiong, P., Schneider, R. F., Hulsey, C. D., Meyer, A., and Franchini, P. (2019). Conservation and novelty in the microRNA Genomic Landscape of Hyperdiverse Cichlid Fishes. *Sci. Rep.* 9 (1), 13848–13912. doi:10.1038/s41598-019-50124-0
- Yamashita, M., Krystal, M., and Palese, P. (1989). Comparison of the Three Large Polymerase Proteins of Influenza A, B, and C Viruses. *Virology* 171 (2), 458–466. doi:10.1016/0042-6822(89)90615-6
- Zeng, Y., Yi, R., and Cullen, B. R. (2003). MicroRNAs and Small Interfering RNAs Can Inhibit mRNA Expression by Similar Mechanisms. *Proc. Natl. Acad. Sci. U.S.A.* 100 (17), 9779–9784. doi:10.1073/pnas.1630797100
- Zhang, Y., and Li, Y.-k. (2013). MicroRNAs in the Regulation of Immune Response against Infections. *J. Zhejiang Univ. Sci. B* 14 (1), 1–7. doi:10.1631/jzus.b1200292
- Zhao, B., and Xue, B. (2019). Significant Improvement of miRNA Target Prediction Accuracy in Large Datasets Using Meta-Strategy Based on Comprehensive Voting and Artificial Neural Networks. *BMC genomics* 20 (1), 158–217. doi:10.1186/s12864-019-5528-1

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Hassan, Iqbal, Naqvi, Alashwal, Moustafa and Kloczkowski. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.