

EPOS™

Electronic Presentation Online System

Confirmation of Presentation

This is to officially confirm that

**M. Bellia¹, M. Messina², F. Cannizzaro¹, N. Scichilone², S. Battaglia², A. giardina², V. Bellia²,
G. Triolo², M. Midiri¹; ¹Palermo (PA)/IT, ²Palermo/IT**


presented the electronic poster entitled

**Pulmonary functional and radiological correlations of nutritional status and physical activity
in systemic sclerosis**

within the Electronic Presentation Online System (EPOS™)

of the European Society of Radiology. The presentation is available
at www.myESR.org/epos and can be cited through this unique DOI:

10.1594/ecr2011/C-1555



Prof. Adrian K Dixon
Chairman of the ESR Publications Committee

Vienna, January 2011



Pulmonary functional and radiological correlations of nutritional status and physical activity in systemic sclerosis

Poster No.: C-1555
Congress: ECR 2011
Type: Scientific Exhibit
Authors: M. Bellia¹, M. Messina², F. Cannizzaro¹, N. Scichilone², S. Battaglia², A. giardina², V. Bellia², G. Triolo², M. Midiri¹; ¹Palermo (PA)/IT, ²Palermo/IT
Keywords: Connective tissue disorders, CT-High Resolution, Respiratory system
DOI: 10.1594/ecr2011/C-1555

Any information contained in this pdf file is automatically generated from digital material submitted to EPOS by third parties in the form of scientific presentations. References to any names, marks, products, or services of third parties or hypertext links to third-party sites or information are provided solely as a convenience to you and do not in any way constitute or imply ECR's endorsement, sponsorship or recommendation of the third party, information, product or service. ECR is not responsible for the content of these pages and does not make any representations regarding the content or accuracy of material in this file.

As per copyright regulations, any unauthorised use of the material or parts thereof as well as commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method ist strictly prohibited.

You agree to defend, indemnify, and hold ECR harmless from and against any and all claims, damages, costs, and expenses, including attorneys' fees, arising from or related to your use of these pages.

Please note: Links to movies, ppt slideshows and any other multimedia files are not available in the pdf version of presentations.

www.myESR.org

Purpose

Systemic Sclerosis (Ssc) or scleroderma is a connective tissue disease of unknown origin.

It may affect skin, as well as blood vessels and internal organs including lungs.

In SSc patients physical activity limitation is often observed. It is unclear whether changes in nutritional status contribute to alter physical performance.

The present study was aimed at exploring the physical and nutritional status of patients with SSc.

Methods and Materials

SAMPLE:

- 27 subjects with SSc;
- 11 age- and sex-matched healthy volunteers.

MEASUREMENTS: The lung involvement was evaluated by:

- Pulmonary function tests (spirometry and lung diffusion capacity

for CO)

- HRCT (the degree of radiological involvement was evaluated by Warrick score, a semi-quantitative method based on the assessment of severity and extent of different types of lesions).

The nutritional status and the body composition were evaluated by:

- Mini Nutritional Assessment (MNA) questionnaire
- Bio Impedance Assessment (BIA)

To evaluate physical activity subjects wore, for one week, a multisensor device: - Sensewear Armband (see figure 1)



**Fig. 1: the
Armband**

Fig.: Fig.1 The armband device

References: M. Bellia; DIBIMEF sezione di scienze radiologiche, Università degli Studi di Palermo, Palermo (PA), ITALY

The multi-sensor device used five sensors:

- two accelerometers;
- skin temperature;
- near body temperature/heat flux;
- galvanic skin resistance.

The SenseWear Armband allowed to measure:

- Total energy expenditure (TEE);
- Active energy expenditure (AEE);
- Physical activity duration (PAD).

STATISTICAL ANALYSIS: Data are presented as mean±SD. Differences between groups were evaluated by Mann-Whitney test. Correlation between variables were evaluated by Spearman's rho. Probability values of $p < 0.05$ were considered to be statistically significant.

Results

In the SSc group, spirometry was normal (FEV1 % pred = $97.5 \pm 14\%$; FVC % pred = $100 \pm 12.8\%$). The diffusing capacity of the lung for CO (DLCO) was significantly lower in SSc group than in controls ($61 \pm 17\%$ pred. vs $88 \pm 4.7\%$ pred., respectively; $p=0.002$). The DLCO correlated with HRCT derived Warrick score of radiologic involvement ($r = -0.47$; $p = 0.029$).

The fat-free mass (FFM) was significantly lower in patients compared to controls (FFM= 46.8 ± 7.6 Kg vs 53.6 ± 6.3 Kg; $p=0.01$) (fig. 2)




Fig. 2

Fig.: Fig.2 Diagram showing the difference in FFM between cases and controls

References: M. Bellia; DIBIMEF sezione di scienze radiologiche, Università degli Studi di Palermo, Palermo (PA), ITALY

Physical activity duration (PAD) was significantly lower in patients than in controls (PAD= 243 ± 145 min. vs 397 ± 142 min.; p=0.004) (fig. 3).




Fig. 3

Fig.: Fig.3 Diagram showing the difference in PAD between cases and controls

References: M. Bellia; DIBIMEF sezione di scienze radiologiche, Università degli Studi di Palermo, Palermo (PA), ITALY

In the whole sample, DLCO was correlated with both PAD ($r = 0.39$; $p = 0.015$) and with FFM ($r = 0.41$; $p = 0.011$). On the contrary, these correlations were not observed for HRCT derived Warrick score. Moreover, PAD and FFM were correlated ($r = 0.48$; $p = 0.002$).

Conclusion

Although SSc patients were in apparently good nutritional and spirometric condition, they showed lower FFM compared to controls. This, together with the correlation between FFM and PAD, could account for physical activity impairment, exercise induced breathlessness and poor quality of life often shown by SSc patients. On this basis further studies are needed to confirm this relationship and to explore the possibility that physical rehabilitation programs become a part of treatment strategy in SSc patients.

References

- Naidich DP, Webb VR, Grenier PA, Harkin TJ, Gefer WB. Imaging of the airways. Functional and radiological correlations. Lippincott, Williams & Wilkins, Philadelphia 2005
- [Camiciottoli G](#), [Orlandi I](#), [Bartolucci M](#), [Meoni E](#), [Nacci F](#), [Diciotti S](#), [Barcaroli C](#), [Conforti ML](#), [Pistoiesi M](#), [Matucci-Cerinic M](#), [Mascalchi M](#). Lung CT densitometry in systemic sclerosis: correlation with lung function, exercise testing, and quality of life. *Chest*. 2007 Mar;131(3):672-81.
- Du Bois RM, Richeldi L. Interstitial lung diseases. European Respiratory Monograph 46, 2009
- [Bellia M](#), [Cannizzaro F](#), [Scichilone N](#), [Riili M](#), [Triolo G](#), [Midiri M](#), [Lagalla R](#) HRCT and scleroderma: semiquantitative evaluation of lung damage and functional abnormalities. *Radiol Med*. 2009 Mar;114(2):190-203.

Personal Information