

P05 - SYNTHESIS AND SAR STUDIES OF NEW 3-METHYL-5-(5-PROPYL-1H-1-R'-3-PYRAZOLYL)-1H-1-R-4-NITROSOPYRAZOLES AS ANTIMICOTIC AGENTS.S. Aiello^{1*}; L. C. López-Cara², F. Venturella³ and A. Licata⁴¹Unità Didattico Scientifica di Fisiologia e Farmacologia, Dip. DIGSPO, Università di Palermo²Dpto. Química Farmacéutica y Orgánica, Facultad de Farmacia, Universidad de Granada C/ Campus cartuja s/n CP.18071, Granada, Spain³Dipartimento Scienze e Tecnologie Biologiche, Chimiche e Farmaceutiche (STEBICEF), Unipa⁴Sezione di Gastroenterologia, DiBiMIS; Università di Palermo

*Corresponding author: Stefania Aiello, Phone:+39 091 6236410

e-mail: stefania.aiello@unipa.it

Many ubiquitous yeast are primarily pathogens for immunocompromised patients, individuals with AIDS and organ transplanted are at high risk of cryptococcosis and candidiasis.



In this setting, fungal infections are particularly difficult to treat because antifungal therapy usually does not eradicate the infection and require lifelong treatment with antifungal drugs.

Consequently, the need for novel antifungal agents for opportunistic infections is apparent in light of significant problems associated with current drugs and makes the development of new drug entities all more urgent.

We have reported that some 3-(3-alkyl-4-nitroso-1H-5-pyrazolyl)-5-R-isoxazoles [1] and the isomeric 5-(1-alkyl-4-nitroso-1H-3-pyrazolyl)-3-R-isoxazoles [2] showed *in vitro* potent antifungal activity at non cytotoxic concentrations.

This antifungal activity was correlated to: 1) the interaction of the isoxazolic nitrogen with the alkyl group bound to the pyrazolyl nitrogen; 2) the *cis* or *trans* configuration adopted by the nitroso group with respect to the alkyl chain bound to the pyrazolic nitrogen and perpendicularly folded to the molecular plane.

To verify this hypothesis, we synthesized compounds in which the isoxazole was substituted by a pyrazole moiety, leading to the new isomeric series **I** and **II**.



The title compounds tested *in vitro* for antifungal activity against *C. Neoformans* and *C. Krusei*, displayed an interesting antifungal activity, in particular compound **Ib** was 2 and 32 fold more potent than Amphotericin B and Fluconazole, respectively, against *C. krusei*, fungus with intrinsic resistance to many of antifungal azoles

These results suggest that, depending on the heterocyclic molecule bound to the 5 position of 1H-1-R-4-nitrosopyrazoles, it is possible to modulate the antifungal activity of 4-nitrosopyrazoles.

In vitro metabolism studies and *in vivo* assay are in progress for all described compounds.

References

1. Aiello, E.; Aiello, S.; Mingoia, F.; Bacchi, A.; Pelizzi, G.; Musiu, C.; Setzu, M. G.; Pani, A.; La Colla, P. and Marongiu, M. E.. Synthesis and Antimicrobial Activity of new 3-(1-R-3-methyl-4-nitroso-1H-5-pyrazolyl)-5-methylisoxazole derivatives. *Bioorganic and Medicinal Chemistry*, 2000, 8, 2719-2728.
2. Stefania Aiello; Enrico Aiello, Marica Orioli, Marina Carini, 3-(1-R-3-methyl-4-nitroso-1H-5-pyrazolyl)-5-methylisoxazoles: a new class of antifungal compounds. In vitro metabolism by rat liver:LC and LC-MS studies. Convegno Nazionale SCI, Sorrento 18-22 Settembre 2002.
3. S. Aiello, E. Aiello and M.E. Milici, Synthesis and Antifungal Activity of new 3(5)-methyl-5(3)-(2-thiophenyl) and -(2-quinolyl)-1H-1-R-4-nitrosopyrazoles. Part V. Polish-Austrian-German-Hungarian-Italian Joint Meeting on Medicinal Chemistry, Krakow, October 15-18, 2003.
4. Stefania Aiello, Antonio Macchiarulo, Maria Milici and Enrico Aiello, Sintesi e studi QSAR di nuovi derivati 3(5)-(2-X)-1R-1H-4-nitrosopirazoli: una classe di composti con potente attività antifungina in vitro. Parte VI. XVII Convegno Nazionale della Divisione di Chimica Farmaceutica della SCI, Pisa 6-10 settembre 2004.