

## TYPE AND COUNTER-TYPE FROM SPECIFIC CHROMOSOMAL REGIONS

FRANCESCO GRAZIANO<sup>1</sup>, VINCENZO ANTONA<sup>1</sup>, MANUELA MARTINES<sup>2</sup>, ANTONELLA BALLACCHINO<sup>3</sup>, VALERIA CONSIGLIO<sup>2</sup>, ETTORE PIRO<sup>1</sup>, GIOVANNI CORSELLO<sup>1</sup>

<sup>1</sup>Università degli Studi di Palermo, Dipartimento di Scienze per la Promozione della Salute e Materno Infantile - <sup>2</sup>CRR Malattie rare AOOR Villa Sofia-Cervello Palermo - <sup>3</sup>Università degli Studi di Palermo, Dipartimento di Biopatologia e Biotecnologie Mediche e Forenzi (Di.Bi.Me.F.), Sezione di Audiologia

*[Tipi e controtipi in specifiche regioni cromosomiche]*

**ABSTRACT**

*Several studies have shown the importance of segmental deletions/duplications in the field of chromosome pathologies. Non allelic homologous recombination, NAHR, between chromosomes or sister chromatids, mediated by segmental duplications, is the foundation of frequent mechanisms for structural chromosome mutations such as micro-deletions, micro-duplications, translocations, inversions, and marker chromosomes. We analyzed three distinct genomic regions (22q11.2, 17p11.2, 16p11.2) and we discussed how the same chromosome region can be affected by deletion or by reciprocal duplication, respectively responsible for a syndrome or for a reciprocal counter-syndrome, with different phenotypic manifestations.*

**Key words:** *Type/counter-type, a-CGH, genomic variants.*

*Received March 05, 2013; Accepted April 22, 2013*

**Introduction**

Several studies, for quite a number of years now, have demonstrated the importance of the role of segmental duplications in the field of chromosome pathologies. Segmental duplications are blocks of few low copy repeats (LCRs) that recognize a reciprocal homology higher than 90% in non-homologous 200 to 400 kb chromosome segments. They account for about 6% of the human genome and are interspersed through the genome, but especially located in the pericentromeric, telomeric and subtelomeric regions. They can include a large number of genes, but can also include broken non-processed pseudogenes. Segmental duplications are often duplicated in one chromosome region at a reciprocal distance of a few megabases (less than 10 Mb) and they mediate a possible non allelic homologous recombination, NAHR, between chromosomes or between sister chromatids: this may result in unequal crossing-over or in chromosome rearrangements and is, therefore, the basis of frequent and specific mechanisms of structural chromosome mutations such as micro-

deletions or micro-duplications, but also translocations, inversions and marker chromosomes. Micro-deletions are more frequent than micro-duplication, as widely stated in scientific literature. Micro-deletions can in fact originate from a wide range of recombination mechanisms and can be intra-chromatid, inter-chromatid or intra-chromosomal, whereas micro-duplications can only originate from an inter-chromatid or intra-chromosomal recombinations<sup>(1,2)</sup>.

The advent of array-CGH with its high image definition (it can detect chromosomal anomalies even lower than 50 kb) paves the way for an easier mapping of such submicroscopic genomic variants (micro-deletions or micro-duplications), increasing the possibility of a correct diagnosis. The introduction of array-CGH also represents a significant advancement in the diagnosis of individuals with a "chromosomal phenotype" (mental retardation, dysmorphic features, congenital anomalies)<sup>(3-7)</sup> and normal karyotype. All these conditions play a very important role in determining morbidity and mortality rates, especially in neonatal age, when many other

risk factors (prematurity<sup>(8,9)</sup>, twinning, nosocomial infections<sup>(10,11,12,13)</sup>, chemical mediators<sup>(14,15)</sup>) may add their cumulative effect influencing short- and long-term outcome. No less than 15-20 % of the patients with mental retardation, dysmorphic features, congenital anomalies, and normal karyotype are carriers of a cryptic chromosomal unbalance sometimes as extended as 100 kb, not detected by the classic cytogenetic techniques<sup>(16,17,18,19,20,21)</sup>.

Undoubtedly array-CGH allows ever more accurate and precise analyses with greater simplicity and faster response<sup>(22,23,24,25,26,27,28)</sup>. In the present study we discuss how the same chromosomal region can be affected by deletion or by duplication, respectively responsible for a syndrome (type) and for a counter-syndrome (counter-type), with different phenotypic manifestations.

### 22q11.2

Region q 11.2 of chromosome 22 shows eight LCRs organized in clusters (4 telomeric ones and 4 centromeric ones, called LCR 22s). Namely, a deletion of 3 Mb is the cause of a DiGeorge/velo-cardio-facial syndrome, while a reciprocal interstitial duplication characterizes both CES (cat eye syndrome) and dup22, precisely a tetrasomy in the first and a trisomy in the second<sup>(29,30)</sup>. Analyzing the phenotype features we can see how deletion and duplication occurring in the same region 22q11.2 (included TBX1 gene) can actually determine two distinct syndromes with different courses and prognoses, so that we can define the 22q11.2 micro-duplication syndrome as a “countersyndrome” or a “countertype” of the 22q11.2 micro-deletion syndrome<sup>(31,32)</sup>. In fact, the clinical picture leading to or hinting at a micro-deletion syndrome diagnosis is more frequently an occurrence of multiple congenital anomalies (generally cardiac conotruncal defects, cleft palate, velopharyngeal insufficiency, tonsillar hypertrophy, dental anomalies), convulsions and tremors due to neonatal hypocalcaemia, a typical facies, recurrent infections due to immune deficiency and thymus aplasia, and psychomotor and/or language retardation, hypothyroidism and behavioral disorder<sup>(33,34)</sup>. On the other hand, the most frequent features of a 22q11.2 micro-duplication syndrome are a varying degree of mental retardation/learning difficulties, attention defect and hyperactivity, hearing impairment, language disorder, psychomotor retardation, growth delay, low muscle tone, epileptic seizures<sup>(35,36,37,38,39)</sup>. Comparing the features in either

syndrome we can notice that facial dysmorphic features variably present in 22q11.2 micro-duplication syndrome are different and often slighter than those of a 22q11.2 micro-deletion syndrome.

### 17 p11.2

Region 11.2 of arm p of chromosome 17 is sided with three “proximal” “middle” and “distal” copies of “low copy repeats (LCR)” also known as “SMS-REPs” (Smith-Magenis repeats). In particular, an interstitial deletion of about 3.7 Mb determines a Smith-Magenis syndrome, whereas an interstitial reciprocal duplication is the cause of a Potocky-Lupsky syndrome<sup>(40)</sup>. Analyzing the phenotype described in literature we can observe that the Potocky-Lupsky syndrome shows different, slighter and less severe clinic and phenotypic features than the Smith-Magenis syndrome. Such phenotypic diversity allows us to define the Potocky-Lupsky syndrome as a “counter-syndrome” or a “countertype” of the Smith-Magenis syndrome, even if they both share the same genomic region and also several genes, among which the RAI-1 gene (retinoic acid 1 inducer) mainly responsible for the phenotypic manifestations in both syndromes. We can in fact, observe that, apart from delayed growth, cognitive and language deficiency, hyperactive or attention defects, EEG alterations, variably present though with different degrees of severity in both syndromes the most important features of the Potocky-Lupsky syndrome are: low muscle tone, scarce feeding and delayed growth in infancy, oropharyngeal dysphagia, dental issues, autistic disorders, central and obstructive sleep apnoea, cardiac anomalies, EEG alterations, hypermetropia, lower blood cholesterol, obsessive compulsive disorders. In addition, in the Potocky-Lupsky syndrome, variable dysmorphic features may be variably present, including triangular face, frontal bossing, microcephaly, hypertelorism, wide nasal bridge, epicanthal folds, flat philtrum<sup>(41)</sup>.

On the contrary, in the Smith-Magenis syndrome, we can variably and frequently observe: cerebral malformations, peripheral neuropathy, epileptic seizures, markedly dysmorphic features, hypercholesterolemia, self-destructive or aggressive behaviors, important sleep disturbances, obesity, otorhinolaryngologic auditory and ophthalmologic anomalies, renal and genitourinary anomalies, skeletal and digital anomalies.

## 16 p11.2

It has been noticed in a recent study that 9.89% (7.8 Mb) of chromosome 16 shows segmental duplications. Analyzing the phenotype described in literature<sup>(42,43)</sup> we can see that a 16p11.2 micro-duplication syndrome shows less serious clinical and phenotypic features with respect to a 16p11.2 micro-deletion syndrome. It is therefore possible to point out that both a 16p11.2 micro-duplication syndrome and a 16p11.2 micro-deletion syndrome share phenotypic features such as delayed growth, delayed language development, learning difficulties, cognitive disability, low muscle tone. While the micro-deletion syndrome provokes an increased tendency to overweight and obesity, a possible association between 16p11.2 micro-duplication and reduced weight and reduced body mass index has been argued.

In fact, deletion of the region 16p11.2 seems to be associated with overweight and obesity liability, due to haploinsufficiency of SH2B1 gene, which is involved in leptin and insulin signaling. Micro-deletion is more frequently associated with macrocephaly while microduplication with microcephaly. It is argued, besides, that micro-deletion might more often be associated with delayed linguistic and cognitive development, with minor dysmorphic facial features without a consistent pattern, with autistic disorders, and with minor cardiac malformations, hemivertebrae and syringomyelia, while micro-duplication is associated with psychiatric problems such as bipolar disorder, depression, psychotic disturbance, schizophrenia.

## Discussion

It is possible to highlight the differences between the phenotypic and neuro-behavioural manifestations of deletions and those of the reciprocal and complementary duplications in the same chromosomal region, especially as regards severity, frequency and recurrence. It must be noticed that the phenotypical appearance of both syndromes is often extremely variable with several degrees of severity.

Therefore micro-duplications, with respect to micro-deletion, show a slighter and milder clinical picture, sometimes even near normality, so that it is not always easy to recognize and identify such an anomaly. The identification of cryptic chromosome anomalies by means of modern molecular tech-

niques, in particular array-CGH, has enabled physicians to devise follow-up programs intended to prevent the associated clinical problems.

We can conclude that array-CGH might become a means to fill the present gap between the knowledge of the human genome and the function of its genes: it is evident that any “de novo” deletion/duplication associated with a specific malformation or syndrome suggests that, among the genes located in the region studied, at least one is responsible for that malformation and for that syndrome, either due to aplo-insufficiency (deletion or type or syndrome) or to an excessive dosage (duplication or counter-type or counter-syndrome).

## References

- 1) Sharp AJ, Hansen S, Seizer RR, et al. *Discovery of previously unidentified genomic disorders from the duplication architecture of the human genome*. Nat Genet 2006; 38: 1038-42.
- 2) Hanson E, Nasir RH, Fong A, et al. *Cognitive and behavioral characterization of 16p11.2 deletion syndrome*. J Dev Behav Pediatr 2010; 31 (8): 649-57.
- 3) Corsello G, Giuffrè M. *Congenital Malformations*. J Matern Fetal Neonatal Med 2012; 25 (S1): 25-29.
- 4) Puccio G, Giuffrè M, Piccione M, Piro E, Rinaudo G, Corsello G. *Intrauterine growth restriction and congenital malformations: a retrospective epidemiological study*. Ital J Pediatr 2013; 39:23. doi:10.1186/1824-7288-39-23.
- 5) Ghirri P, Scaramuzza RT, Bertelloni S, et al. *Prevalence of hypospadias in Italy according to severity, gestational age and birthweight: an epidemiological study*. Ital J Pediatr 2009; 35: 18.
- 6) Giuffrè M, Schierz IM, La Placa S. *Newborn with prenatal diagnosis of CAM*. Area Pediatrica 2011; 12 (1): 25-26.
- 7) Schierz IAM, Giuffrè M, Piro E, et al. *Predictive factors of abdominal compartment syndrome in neonatal age*. Am J Perinatol 2013; doi: 10.1055/s-0033-1334447.
- 8) Giuffrè M, Piro E, Corsello G. *Prematurity and Twinning*. J Matern Fetal Neonatal Med 2012; 25 (S3): 6-10.
- 9) Schierz IAM, La Placa S, Giuffrè M, et al. *Transient hepatic nodular lesions associated with patent ductus venosus in preterm infants*. Am J Perinatol 2011; 28 (3): 177-180.
- 10) Cipolla D, Giuffrè M, Mammina C, et al. *Prevention of nosocomial infections and surveillance of emerging resistances in NICU*. J Matern Fetal Neonatal Med 2011; 24 (S1): 23-26.
- 11) Mammina C, Di Carlo P, Cipolla D, et al. *Surveillance of multidrug-resistant gram-negative bacilli in a neonatal intensive care unit; prominent role of cross transmission*. Am J Infect Control 2007; 35 (4): 222-230.
- 12) Giuffrè M, Cipolla D, Bonura C, et al. *Epidemic spread of ST-1-MRSA-IVa in a neonatal intensive care unit, Italy*. BMC Pediatr 2012; 8, 12: 64.
- 13) Giuffrè M, Cipolla D, Bonura C, et al. *Outbreak of colonizations by extended-spectrum beta-lactamase-*

- producing *Escherichia coli* sequence type 131 in a neonatal intensive care unit, Italy. *Antimicrob Resist Infect Control* 2013; 2: 8; doi:10.1186/2047-2994-2-8.
- 14) David S, Bucchieri F, Corrao S, et al. *Hsp10: anatomic distribution, functions, and involvement in human disease*. *Front Biosci Elite* 2013; E5: 768-78.
  - 15) Lo Iacono M, Anzalone R, Corrao S, et al. *Perinatal and wharton's jelly-derived mesenchymal stem cells in cartilage regenerative medicine and tissue engineering strategies*. *Open Tissue Engineering and Regenerative Medicine Journal* 2011; 4: 72-81.
  - 16) Viaggi CD, Cavani S, Pierluigi M, et al. *Characterization of a complex rearrangement involving chromosomes 1, 4 and 8 by FISH and array-CGH*. *J Appl Genet* 2012; 53 (3): 285-288.
  - 17) Isidor B, Bourdeaut F, Lafon D, et al. *Wilms' tumor in patients with 9q22.3 microdeletion syndrome suggests a role for PTCH1 in nephroblastomas*. *Eur J Hum Genet* 2012; doi: 10.1038/ejhg.2012.252.
  - 18) Piccione M, Serra G, Sanfilippo C, et al. *A new mutation in EDA gene in X-linked hypohidrotic ectodermal dysplasia associated with keratoconus*. *Minerva Pediatr* 2012; 64 (1): 59-64.
  - 19) Piccione M, Niceta M, Antona V, et al. *Identification of two new mutations in TRPS 1 gene leading to the tricho-rhino-phalangeal syndrome type I and III*. *Am J Med Genet A* 2009;149A(8): 1837-1841.
  - 20) Piccione M, Serra G, Consiglio V, et al. *14q13.1-21.1 deletion encompassing the HPE8 locus in an adolescent with intellectual disability and bilateral microphthalmia, but without holoprosencephaly*. *Am J Med Genet A* 2012; 158A (6): 1427-1433.
  - 21) Piccione M, Antona R, Salzano E, et al. *Array-CGH and clinical characterization in a patient with subtelomeric 6p deletion without ocular dysgenesis*. *Am J Med Genet A* 2012; 158A (1): 150-154.
  - 22) Oostlander AE, Meijer GA, Ylstra B. *Microarray-based comparative genomic hybridization and its applications in human genetics*. *Clinical Genet* 2004; 66: 488-495.
  - 23) Vissers LE, Van Ravenswaaij CM, Admiraal R, et al. *Mutations in a new member of the chromodomain gene family cause CHARGE syndrome*. *Nature Genetics* 2004; 36 (9): 955-957.
  - 24) Shaw-Smith C, Redon R, Rickman L, et al. *Microarray based comparative genomic hybridization (array-CGH) detects submicroscopic chromosomal deletions and duplications in patients with learning disability/mental retardation and dysmorphic features*. *J Med Genet* 2004; 41: 241-248.
  - 25) Gribble SM, Prigmore E, Burford DC, et al. *The complex nature of constitutional de novo apparently balanced translocations in patients presenting with abnormal phenotypes*. *J Med Genet* 2005; 42: 8-16.
  - 26) Pickering DL, Eudy JD, Olney AH, et al. *Array-based comparative genomic hybridization analysis of 1176 consecutive clinical genetics investigations*. *Genet Med* 2008; 10: 262-266.
  - 27) Piccione M, Piro E, Serraino F, et al. *Interstitial deletion of chromosome 2p15-16.1: report of two patients and critical review of current genotype-phenotype correlation*. *Eur J Med Genet* 2012; 55 (4): 238-244.
  - 28) Striano P, Coppola A, Paravidino R, et al. *Clinical significance of rare copy number variations in epilepsy: a case-control survey using microarray-based comparative genomic hybridization*. *Arch Neurol* 2012; 69 (3): 322-330.
  - 29) Portnoi MF. *Microduplication 22q11.2: A new chromosomal syndrome*. *European Journal of Medical Genetics* 2009; 52: 88-93.
  - 30) Piccione M, Vecchio D, Cavani S, et al. *The first case of myoclonic epilepsy in a child with a de novo 22q11.2 microduplication*. *Am J Med Genet Part A* 2011; 155: 3054-3059.
  - 31) Wentzel C, Fernstrom M, Ohrner Y, et al. *Clinical variability of the 22q11.2 duplication syndrome*. *Europ J Med Genet* 2008; 51: 501-510.
  - 32) De La Rochebrochard C, Joly-Helas G, Goldenberg A, et al. *The intrafamilial variability of the 22q11.2 microduplication encompasses a spectrum from minor cognitive deficits to severe congenital anomalies*. *Am J Med Genet* 2006; 140A: 1608-1613.
  - 33) Bentivegna D, Salvago P, Agrifoglio M, et al. *The linkage between upper respiratory tract infections and otitis media: evidence of the 'united airways concept'*. *Acta Medica Mediterranea*, 2012, 28: 287-290.
  - 34) Ferrara S, Di Marzo M, Martines F, et al. *Otite media atelettasica, adesiva, timpanosclerotica: update medico e chirurgico*. *Otorinolaringologia* 2011; 61: 11-17.
  - 35) Martines F, Martines E, Ballacchino A, et al. *Speech perception outcomes after cochlear implantation in prelingually deaf infants: The Western Sicily experience*. *Int J Pediatr Otorhinolaryngol*. 2013 Feb 18. doi: pii:S0165-5876(13)00037-2.
  - 36) Salvago P, Martines E, Martines F. *Prevalence and risk factors for sensorineural hearing loss: Western Sicily overview*. *Eur Arch Otorhinolaryngol*. 2013 Feb 10. doi: 10.1007/s00405-013-2379-2.
  - 37) Martines F, Bentivegna D, Cipri S, et al. *On the threshold of effective well infant nursery hearing screening in Western Sicily*. *Int J Pediatr Otorhinolaryngol*. 2012 Mar; 76(3): 423-7.
  - 38) Maggio M, Martines F, Mucia M, et al. *A Multifactorial pattern for the understanding of the psychological development of the child with impaired hearing and its clinical-therapeutic implications*. *Acta Medica Mediterranea*, 2006, 22: 41-44.
  - 39) Martines F, Salvago P, Bentivegna D, et al. *Audiologic profile of infants at risk: experience of a Western Sicily tertiary care centre*. *Int J Pediatr Otorhinolaryngol*. 2012 Sep;76(9):1285-91.
  - 40) Edelman E, Girirajan S, Finucane B et al. *Gender, genotype, and phenotype differences in Smith-Magenis syndrome: a metanalysis of 105 cases*. *Clin Genet* 2007; 71: 540-550.
  - 41) Yusupov R, Roberts AE, Lacro RV, et al. *Potocki-Lupski syndrome: an inherited dup(17)(p11.2p11.2) with hypoplastic left heart*. *Am J Med Genet Part A* 2011; 155: 367-371.
  - 42) Hannes FD, Sharp AJ, Mefford HC, et al. *Recurrent reciprocal deletions and duplications of 16p13.11; the deletion is a risk factor for MR/MCA while the duplication may be a rare benign variant*. *J Med Genet* 2009; 46: 223-232.
  - 43) Jacquemont S, Reymond A, Zufferey F, et al. *Mirror extreme BMI phenotypes associated with gene dosage at phenotypes associated with gene dosage at the chromosome 16p11.2 locus*. *Nature* 2011; 478: 97-102.

Request reprints from:  
 Dr. ETTORE PIRO  
 Via A. Giordano, 3  
 90127 Palermo  
 (Italy)