

LETTERS

humans. Additionally, *Mycobacteria* spp. can occasionally cause disease in humans through contact with fish (*M. marinum*), and pedicure treatments have previously been associated with *M. fortuitum* infections (10).

Recently, the risks associated with exposure to *G. rufa* fish were reported to be low (1). To date, there are only a limited number of reports of patients who might have been infected by this exposure route (1). However, our study raises some concerns over the extent that these fish, or their transport water, might harbor potential zoonotic disease pathogens of clinical relevance. In particular, patients with underlying conditions (such as diabetes mellitus or immunosuppression) should be discouraged from undertaking such treatments, especially if they have obvious breaks in the skin or abrasions. This risk can probably be reduced by use of certified disease-free fish reared in controlled facilities under high standards of husbandry and welfare.

The UK Department for Environment Food and Rural Affairs provided funding for this study through projects FA001 and FB002.

**David W. Verner-Jeffreys,
Craig Baker-Austin,
Michelle J. Pond,
Georgina S. E. Rimmer,
Rose Kerr, David Stone,
Rachael Griffin, Peter White,
Nicholas Stinton,
Kevin Denham, James Leigh,
Nicola Jones,
Matthew Longshaw,
and Stephen W. Feist**

Author affiliations: Centre for Environment, Fisheries & Aquaculture Science Weymouth Laboratory, Weymouth, UK (D.W. Verner-Jeffreys, C. Baker-Austin, M.J. Pond, G.S.E. Rimmer, R. Kerr, D. Stone, R. Griffin, P. White, N. Stinton, K. Denham, M. Longshaw, S.W. Feist); University of

Nottingham, Sutton Bonington, UK (J. Leigh); and Oxford Radcliffe University Hospitals, Headington, UK (N. Jones)

DOI: <http://dx.doi.org/10.3201/eid1806.111782>

References

1. Health Protection Agency Fish Spa Working Group. Guidance on the management of the public health risks from fish pedicures: draft for consultation. 2011 Aug 31 [cited 2012 Mar 21]. http://www.hpa.org.uk/webc/HPAwebFile/HPAweb_C/1317131045549
2. Jones N, Bohnsack JF, Takahashi S, Oliver KA, Chan MS, Kunst F, et al. Multilocus sequence typing system for group B *Streptococcus*. J Clin Microbiol. 2003;41:2530–6. <http://dx.doi.org/10.1128/JCM.41.6.2530-2536.2003>
3. Evans JJ, Bohnsack JF, Klesius PH, Whiting AA, Garcia JC, Shoemaker CA, et al. Phylogenetic relationships among *Streptococcus agalactiae* isolated from piscine, dolphin, bovine and human sources: a dolphin and piscine lineage associated with a fish epidemic in Kuwait is also associated with human neonatal infections in Japan. J Med Microbiol. 2008;57:1369–76. <http://dx.doi.org/10.1099/jmm.0.47815-0>
4. Janda JM, Abbott SL. The genus *Aeromonas*: taxonomy, pathogenicity, and infection. Clin Microbiol Rev. 2010;23:35–73. <http://dx.doi.org/10.1128/CMR.00039-09>
5. Jones MK, Oliver, JD. *Vibrio vulnificus*: disease and pathogenesis. Infect Immun. 2009;77:1723–33. <http://dx.doi.org/10.1128/IAI.01046-08>
6. Morris JG. Non-O group-1 *Vibrio cholera*—a look at the epidemiology of an occasional pathogen. Epidemiol Rev. 1990;12:179–91.
7. Wagner D, Young LS. Nontuberculous mycobacterial infections: a clinical review. Infection. 2004;32:257–70. <http://dx.doi.org/10.1007/s15010-004-4001-4>
8. Skoff TH, Farley MM, Petit S, Craig AS, Schaffner W, Gershman K, et al. Increasing burden of invasive group B streptococcal disease in non-pregnant adults, 1990–2007. Clin Infect Dis. 2009;49:85–92. <http://dx.doi.org/10.1086/599369>
9. Verner-Jeffreys DW, Welch TJ, Schwarz T, Pond MJ, Woodward MJ, Haig SJ, et al. High prevalence of multidrug-tolerant bacteria and associated antimicrobial resistance genes isolated from ornamental fish and their carriage water. PLoS ONE. 2009;4:e8388. <http://dx.doi.org/10.1371/journal.pone.0008388>
10. De Groot MA, Huit G. Infections due to rapidly growing mycobacteria. Clin Infect Dis. 2006;42:1756–63. <http://dx.doi.org/10.1086/504381>

Address for correspondence: David W. Verner-Jeffreys, Cefas Weymouth Laboratory, The Nothe, Barrack Rd, Weymouth, Dorset DT4 8UB, UK; email: david.verner-jeffreys@cefas.co.uk

Rickettsia conorii Indian Tick Typhus Strain and *R. slovaca* in Humans, Sicily

To the Editor: Rickettsiae are vector-borne pathogens that affect humans and animals worldwide (1). Pathogens in the *Rickettsia conorii* complex are known to cause Mediterranean spotted fever (MSF) (*R. conorii* Malish strain), Astrakhan fever (*R. conorii* Astrakhan strain), Israeli spotted fever (*R. conorii* Israeli spotted fever strain), and Indian tick typhus (*R. conorii* Indian tick typhus strain) in the Mediterranean basin and Africa, southern Russia, the Middle East, and India and Pakistan, respectively (2). These rickettsioses share some clinical features, such as febrile illness and generalized cutaneous rash, and are transmitted to humans by *Rhipicephalus* spp. ticks (2).

MSF is endemic to Sicily (Italy); fatal cases occur each year, and the prevalence of *R. conorii* in dogs is high (3–6). Recently, *R. conorii* Malish strain and *R. conorii* Israeli spotted fever strain were confirmed in humans in Sicily in whom MSF was diagnosed (4), which suggests that other *R. conorii* strains might be present and diagnosed as causing MSF. The rickettsiae within the *R. conorii* complex, which are relevant for the study of bacterial evolution and epidemiology, can be properly identified only by appropriate genetic analyses.

We analyzed 15 blood and 19 inoculation eschar samples collected during 2005–2009 from 31 patients in Palermo Province and 2 in Catania Province, none of whom had recently traveled. None were severely ill, but all 33 had clinical manifestations and laboratory results compatible with MSF: 1-week incubation after tick bite, fever, headache, myalgia, papulonodular rash that started on the upper limbs and spread centripetally with or without tache noire, and detection of antibody titers ≥ 180 to *R. conorii* by indirect immunofluorescence antibody test (bioMérieux, Marcy L'Etoile, France).

Total DNA was extracted by using the GeneElute Mammalian Genomic DNA Miniprep Kit (Sigma-Aldrich, Milan, Italy) and used to analyze *Rickettsia* spp. sequences by PCR, cloning, and sequence analysis of the amplicons. At least 3 clones were sequenced for each amplicon. Genes targeted by PCR included ATP synthase α subunit (*atpA*) (7), heat-shock protein 70 (*dnaK*) (7), outer membrane protein A (*ompA*) (primers Rr190.70p and 190–701 [8]), outer membrane protein B (*ompB*) (primers rompBSFGIF and rompBSFG/TGIR [9]), citrate synthase (*gltA*) (2), and 17-kDa protein (primers TZ15–19 and TZ16–20 [6]). Nucleotide sequence identity to reference strains (2), multilocus analysis by *atpA*–*dnaK*–*ompA*–*ompB*–*gltA*–17-kDa and *ompA*–*ompB* sequences and in silico *PstI*–*RsaI* restriction analysis of *ompA* sequences (8) were used to characterize *Rickettsia* spp. and *R. conorii* strains.

Results for 15 (45%) patients were positive for *Rickettsia* spp. Thirteen isolates were confirmed as *R. conorii* Malish strain (identification [ID] nos. 44, 45, 47, 49, 54, 55, 57, 59, 61, 66, 68, 92, 112) and 1 each as *R. conorii* Indian tick typhus strain (ID no. 58) and *R. slovaca* (ID no. 50). *R. slovaca* DNA was also found in a *Dermacentor marginatus* tick removed from the

patient who had confirmed *R. slovaca* infection. *R. conorii* Malish strains showed 99.9%–100%, 100%, 100%, 98.7%–100%, 100%, and 97.8%–100% pairwise nt sequence identity to reference strain Malish 7 (AE006914) *atpA*, *dnaK*, *ompA*, *ompB*, *gltA*, and 17-kDa protein, respectively.

The *R. conorii* Indian tick typhus strain showed 100%, 100%, 99.4%, 100%, 100%, and 99.9% pairwise nt sequence identity to *R. conorii* strain Malish 7 (AE006914) *atpA*, *dnaK*, 17-kDa protein, and *R. conorii* Indian tick typhus reference strain *ompA* (U43794), *ompB* (AF123726), and

gltA (U59730), respectively. The *R. slovaca* strain showed 99.4%, 97.8%, 100%, 93.7%, 99.7%, and 99.4% pairwise nt sequence identity to *R. slovaca atpA* (AY124734), *dnaK* (DQ821824), *ompA* (HM149286), *ompB* (HQ232242), *gltA* (AY129301), and *R. conorii* strain Malish 7 (AE006914) 17-kDa protein, respectively. The sequences were deposited in GeneBank under accession nos. JN182782–JN182804.

Multilocus sequence analysis (Figure, panel A) and in silico *PstI*–*RsaI* restriction analysis of *ompA* sequences also confirmed the identity




Figure. Multilocus sequence analysis of *Rickettsia* spp. Evolutionary history was inferred by using the neighbor-joining method for ATP synthase α subunit (*atpA*)–heat shock protein 70 (*dnaK*)–outer membrane protein A (*ompA*)–*ompB*–citrate synthase (*gltA*)–17-kDa (A) and *ompA*–*ompB* sequences (B). The optimal tree with the sum of branch length = 0.06205323 (A) and 0.11097561 (B) is shown. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1,000 replicates) are shown next to the branches. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic relationship. Evolutionary distances were computed by using the Kimura 2-parameter method and are in the units of the number of base substitutions per site. All ambiguous positions were removed for each sequence pair. Evolutionary analyses were conducted in MEGA5 (www.megasoftware.net). Identification numbers of strains detected are shown with the species/strain name next to them. Rc, *R. conorii* strain Malish 7; Ra, *R. africae* strain ESF-5; Rr, *R. rickettsii* strain Iowa; Rs, *R. slovaca*; Rm, *R. massiliae* strain MTU5; Rcind, *R. conorii* Indian tick typhus strain.

of the *Rickettsia* spp. we identified. As shown (2), multilocus analysis with *ompA-ompB* sequences was highly informative about the phylogenetic relationship between *Rickettsia* spp. and *R. conorii* strains (Figure, panel B).

In Sicily, *R. conorii* Malish strain has been characterized in MSF patients (4), and *R. slovaca* DNA was identified in ixodid ticks (5). However, to our knowledge, *R. slovaca* in humans in Sicily and *R. conorii* Indian tick typhus strain infection in Sicily and Europe have not been reported. The only previous report outside India and Pakistan was documented in a traveler with severe clinical manifestations in France (10). Differences were not observed between *R. conorii* Indian tick typhus strain and *R. slovaca*-infected patients. Both patients had similar clinical symptoms compatible with MSF; in both, only IgM for rickettsiae was detected at hospital admission, but IgM and IgG were detected during convalescence. Tache noire were detected in the neck and right arm of patients with *R. conorii* Indian tick typhus strain and *R. slovaca*, respectively.

These results demonstrated that new rickettsiae, such as *R. conorii* Indian tick typhus strain, of public health relevance are emerging in Europe. The widespread distribution of tick vectors in Europe and the transtadial and transovarial transmission of the pathogen in ticks might favor transmission to humans.

This research was supported by the Italian Ministry of Health, project IZSSI 08/08.

**Alessandra Torina,
Isabel G. Fernández de Mera,
Angelina Alongi,
Atilio J. Mangold,
Valeria Blanda,
Francesco Scarlata,
Vincenzo Di Marco,
and José de la Fuente**

Author affiliations: Istituto Zooprofilattico Sperimentale della Sicilia, Palermo, Sicily, Italy (A. Torina, A. Alongi, V. Blanda, V. Di Marco); University of Palermo, Palermo (V. Blanda); Instituto de Investigación en Recursos Cinegéticos (IREC-CSIC-UCLM-JCCM) Ciudad Real, Spain (I.G. Fernández de Mera, J. de la Fuente); Universidad Complutense de Madrid, Madrid, Spain (I.G. Fernández de Mera); Estación Experimental Agropecuaria Rafaela, Santa Fe, Argentina (A.J. Mangold); Istituto di Patologia Infettiva e Virologia dell'Università di Palermo, Palermo (F. Scarlata); and Oklahoma State University, Stillwater, Oklahoma, USA (J. de la Fuente)

DOI: <http://dx.doi.org/10.3201/eid1806.110966>

References

- Nicholson WL, Allen KE, McQuiston JH, Breitschwerdt EB, Little SE. The increasing recognition of rickettsial pathogens in dogs and people. *Trends Parasitol.* 2010;26:205–12. <http://dx.doi.org/10.1016/j.pt.2010.01.007>
- Zhu Y, Fournier PE, Ereemeeva M, Raoult D. Proposal to create subspecies of *Rickettsia conorii* based on multilocus sequence typing and an emended description of *Rickettsia conorii*. *BMC Microbiol.* 2005;5:11. <http://dx.doi.org/10.1186/1471-2180-5-11>
- Ciceroni L, Pinto A, Ciarrocchi S, Ciervo A. Current knowledge of rickettsial diseases in Italy. *Ann N Y Acad Sci.* 2006;1078:143–9. <http://dx.doi.org/10.1196/annals.1374.024>
- Giammanco GM, Vitale G, Mansueto S, Capra G, Caleca MP, Ammatuna P. Presence of *Rickettsia conorii* subsp. *israelensis*, the causative agent of Israeli spotted fever, in Sicily, Italy, ascertained in a retrospective study. *J Clin Microbiol.* 2005;43:6027–31. <http://dx.doi.org/10.1128/JCM.43.12.6027-6031.2005>
- Beninati T, Genchi C, Torina A, Caracappa S, Bandi C, Lo N. Rickettsiae in ixodid ticks, Sicily. *Emerg Infect Dis.* 2005;11:509–11. <http://dx.doi.org/10.3201/eid1103.040812>
- Tzianabos T, Anderson BE, McDade JE. Detection of *Rickettsia rickettsii* DNA in clinical specimens by using polymerase chain reaction technology. *J Clin Microbiol.* 1989;27:2866–8.
- Fernández de Mera IG, Zivkovic Z, Bolaños M, Carranza C, Pérez-Arellano JL, Gutiérrez C, et al. *Rickettsia massiliae* in the Canary Islands. *Emerg Infect Dis.* 2009;15:1869–70.
- Roux V, Fournier PE, Raoult D. Differentiation of spotted fever group rickettsiae by sequencing and analysis of restriction fragment length polymorphism of PCR-amplified DNA of the gene encoding the protein rOmpA. *J Clin Microbiol.* 1996;34:2058–65.
- Choi YJ, Jang WJ, Ryu JS, Lee SH, Park KH, Paik HS, et al. Spotted fever group and typhus group rickettsioses in humans, South Korea. *Emerg Infect Dis.* 2005;11:237–44. <http://dx.doi.org/10.3201/eid1102.040603>
- Parola P, Fenollar F, Badiaga S, Brouqui P, Raoult D. First documentation of *Rickettsia conorii* infection (strain Indian tick typhus) in a traveler. *Emerg Infect Dis.* 2001;7:909–10. <http://dx.doi.org/10.3201/eid0705.010527>

Address for correspondence: José de la Fuente, Instituto de Investigación en Recursos Cinegéticos IREC-CSIC-UCLM-JCCM, Ronda de Toledo s/n, 13005 Ciudad Real, Spain; email: jose_delafuente@yahoo.com

Detection of European Strain of *Echinococcus multilocularis* in North America

To the Editor: In 2009, an alveolar hydatid cyst, the intermediate stage of the cestode *Echinococcus multilocularis*, was detected in the liver of a dog from Quesnel, British Columbia (BC), Canada (1), 600 km west of the nearest known record of this parasite in central North America (Figure). Alveolar hydatid cysts normally occur in rodent intermediate hosts. However, humans can serve as aberrant intermediate hosts; cysts generally originate in the liver and, in about one third of cases, metastasize throughout the body (2). Detection of the larval stage of this pathogen in an unusual host in a new geographic region required application of multiple molecular epidemi-