

[About](#) | [Submit](#) | [Journals](#)

Search for articles, people, events and more.

[Login](#) | [Register](#)

EVENT ABSTRACT

[Back to Event](#)

Granzyme A is an additional marker for tuberculosis (TB) to discriminate between patients with active disease and subjects with latent infection.

Nadia Caccamo¹, Valentina Orlando¹, Marco Pio La Manna¹, Diana Di Liberto¹, Stella Cutrera¹, Francesco Dieli¹, Valentina Vanini², Elisa Petruccioli² and Delia Goletti²

¹ Dipartimento di Biopatologia e Biotecnologie Mediche e Forensi, University of Palermo, Italy

² L. Spallanzani National Institute for Infectious Diseases (INMI), IRCCS, Translational Research Unit, Department of Epidemiology and Preclinical Research, Italy

Tuberculosis is a major global health problem, with an estimated 8.7 million cases and 1.4 million deaths in 2011. Cytotoxic molecules such as granulysin, perforin and granzymes produced by cytolytic T cells directly contributes to immune defense against tuberculosis (TB). It is still difficult to discriminate active disease, latent infection (LTBI), past TB patients subjects based on Quantiferon TB-Gold in tube (QFT-IT) response; therefore the aim of this study was to evaluate if the responses to granzyme A could help to discriminate between the different TB stages. We compared the release of granzyme A in whole blood cells stimulated with QFT-IT. We found significant higher level of this cytotoxic molecule in LTBI and past TB subjects compared to the level found in patients with active disease (at least $p < 0.001$).

These results suggest that the evaluation of the level of cytotoxic molecules such as granzyme A could be considered as a biomarker of TB to better discriminate among LTBI, active disease, past TB.

Acknowledgements

This work was supported by grants from the European Commission within the 6th Framework Programme, TB-VAC contract no. LSHP-CT-2003-503367 and the 7th Framework Programme, NEWTBVAC contract no. HEALTH-F3-2009-241745 (The text represents the authors' views and does not necessarily represent a position of the Commission who will not be liable for the use made of such information).

Keywords: Tuberculosis, granzyme A, ELISA, LTBI, Cytotoxic molecules, TB patients

Conference: 15th International Congress of Immunology (ICI), Milan, Italy, 22 Aug - 27 Aug, 2013. **Presentation Type:** Abstract **Topic:** Host-pathogen interactions

Citation: Caccamo N, Orlando V, La Manna M, Di Liberto D, Cutrera S, Dieli F, Vanini V, Petruccioli E and Goletti D (2013). Granzyme A is an additional marker for tuberculosis (TB) to discriminate between patients with active disease and subjects with latent infection.. *Front. Immunol. Conference Abstract: 15th International Congress of Immunology (ICI)*. doi: 10.3389/conf.fimmu.2013.02.00683

Received: 13 Jun 2013; **Published Online:** 22 Aug 2013.

* **Correspondence:** Prof. Nadia Caccamo, Dipartimento di Biopatologia e Biotecnologie Mediche e Forensi, University of Palermo, Palermo, Italy, nadia.caccamo@unipa.it

